

Identification of 370 loci for age at onset of sexual and reproductive behaviour, highlighting common aetiology with reproductive biology, externalizing behaviour and longevity

Melinda C. Mills^{1,2,†,*}, Felix C. Tropf^{1,2,3,4,†}, David M. Brazel^{1,2,†}, Natalie van Zuydam⁵, Ahmad Vaez^{6,7}, *eQTLGen Consortium*, *BIOS Consortium*, Tune H. Pers^{8,9}, Harold Snieder⁶, John R.B. Perry¹⁰, Ken K. Ong^{10,†}, Marcel den Hoed^{5,†}, Nicola Barban^{11,†}, and Felix R. Day^{10,†,*} on behalf of the *Human Reproductive Behaviour Consortium*

¹ Leverhulme Centre for Demographic Science, University of Oxford, Oxford, United Kingdom

² Nuffield College, University of Oxford, Oxford, United Kingdom

³ École Nationale de la Statistique et de L'administration Économique (ENSAE), Paris, France

⁴ Center for Research in Economics and Statistics (CREST), Paris, France

⁵ The Beijer Laboratory and Department of Immunology, Genetics and Pathology, Uppsala University and SciLifeLab, Uppsala, Sweden

⁶ Department of Epidemiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

⁷ Department of Bioinformatics, Isfahan University of Medical Sciences, Isfahan, Iran

⁸ The Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

⁹ Department of Epidemiology Research, Statens Serum Institut, Copenhagen, Denmark

¹⁰ MRC Epidemiology Unit, Institute of Metabolic Science, University of Cambridge, Cambridge, United Kingdom

¹¹ Institute of Social and Economic Research, University of Essex, Essex, United Kingdom

[†] Denotes equal contribution

* Correspondence to Melinda C. Mills, melinda.mills@nuffield.ox.ac.uk, and Felix R. Day, felix.day@mrc-epid.cam.ac.uk

Abstract. The timing of reproductive behaviour – age at first sexual intercourse (AFS) and age at first birth (AFB) – has implications for reproductive health, adolescent development and evolutionary fitness. In the largest genome-wide association study to date (AFS, N=387,338; AFB, N=542,901), we identify 370 independent signals, 11 which are sex-specific, with a 5-6% polygenic score prediction. Heritability shifted from 10% for those born in 1940 to 23% for the 1965 birth cohort. Using Genomic SEM, we show that signals are largely driven by the genetics of reproductive biology and externalizing behaviour. This is supported by extensive biological follow-up that isolates key genes related to follicle stimulating hormone (FSHB), implantation (ESR1), infertility (endometriosis, spontaneous abortion) and spermatid differentiation, morphogenesis and binding (KLF17, ZPBP). Later AFB is protective against later-life disease (type 2 diabetes, cardiovascular) and associated with longevity. Those from higher childhood socioeconomic circumstances and polygenic scores in the highest deciles (90%+) experience markedly later reproductive onset. Results are relevant for interventions in teenage sexual, reproductive and mental health, deepen our understanding of the drivers of later-life health and longevity, and fuel infertility and functional follow-up experiments.

The timing of onset of human reproductive behaviour – age at first sexual intercourse (AFS) and age at first birth (AFB) – has implications for reproductive health, adolescent development and evolutionary fitness. First sexual intercourse has occurred increasingly earlier, by the age of 16 for one-third of contemporary UK teenagers.¹ Early reproductive onset is linked to teenage pregnancy² but also adverse health outcomes such as cervical cancer, depression, sexually transmitted diseases² and substance use disorders.^{3,4} In contrast to earlier sexual debut, we have witnessed progressively later ages at first birth for women, now reaching an average of 30 years in many modern societies and even later for men (**Supp Note Fig S1**).⁵ Late reproductive behaviour is associated with lower fecundity and subfertility⁶ and infertility traits such as endometriosis and early menopause,^{7,8} with over 20% of women born after 1970 in many modern countries now remaining childless.⁹ Earlier ages of sexual debut and later ages at first birth has marked the decoupling of reproduction from sexual behaviour in many contemporary societies, with implications for sexual, reproductive and later-life health (**Supp Note Fig S2**).

Since reproductive behaviour is shaped by biology, disease and behaviour, a multidisciplinary approach is required to understand the common genetic aetiology and how it relates to health, reproductive biology and externalizing behaviour. Since the onset of reproductive behaviour generally occurs in adolescence to early adulthood, it is often linked to externalizing behaviour such as self-control and psychiatric (e.g., ADHD) and substance use disorders (e.g., smoking, alcohol use), often mediated by the environment (e.g., childhood socioeconomic conditions) (**Supp Note Fig S3**). Furthermore, it may be that individuals inherit a common genetic liability for a spectrum of interlinked complex traits related to reproduction, health and longevity. There has also been limited attention to

understanding how these genetic effects are stratified by sex or across different socioeconomic and historical contexts.


In a previous GWAS of AFS ($n=125,667$)¹⁰ and AFB ($n=343,072$),⁸ we identified 38 and 10 novel independently-associated single-nucleotide polymorphisms (SNPs), respectively. The current study comprises a markedly expanded sample size for AFS ($N=387,338$) and AFB ($N=542,901$), uncovering 370 independent autosomal or X chromosomal loci, some of which are sex-specific, with 99 candidate genes expressed at the protein level in the brain, glands and reproductive organs. With methods and main findings summarized in **Fig 1**, this study reveals underlying genetic drivers, common genetic liabilities, heterogeneity by childhood socioeconomic status and historical period and further evidence of the relationship of later reproductive onset with fewer later-life metabolic life diseases and increased longevity.

Results

Phenotypic changes in human reproductive behaviour and heritability over time.


Descriptive analyses using the UK Biobank (see **Online Methods**) illustrate shifts in mean AFS and AFB, changes in the shape of the distribution by birth cohort, and a bi-modal distribution of AFS in earlier cohorts (**Fig 2A, Supp Note Fig S1**). Whereas AFB was often in the early 20s for older birth cohorts, this distribution has spread and shifted to older ages over time, with a marked drop in Pearson's correlation between AFS and AFB from those born <1941 (0.60) to those born >1960 (0.31) (**Supp Note Fig S2**). Using GREML,^{11,12} we found a steady increase in SNP-heritability by birth cohort for AFB for women from just under 10% for those born in 1940, climbing to around 23% for the latest cohorts born in

91 **Fig 1.** Summary and description of methods and main results, onset of timing of human reproductive behaviour GWAS




92

93 Note: Dark grey indicates analysis method, light grey the purpose of the analysis and white the main results.



Note: Dark grey indicates analysis method, light grey the purpose of the analysis and white the main results.

97 **Fig 2.** Age at first sex (AFS) and age at first birth (AFB) changes over time, polygenic score (PGS) prediction and bi-directional MR



1965. For AFS, heritability ranges between 13 and 23% with a trend for women similar to AFB and a U-shaped trend for men (**Supp Note Fig 4A-B**).

Meta-analysis GWAS Human Reproductive Behaviour. We conducted a meta-analysis of GWAS results from 36 cohorts for AFS and AFB in individuals of European ancestry. We imputed to the 1000 Genomes Project reference panel in a pooled sample and then stratified the analysis by sex (**Supp Note Tables S1-8**). In total, we discovered 370 associated loci. The GWAS of AFS identified 281 (271 pooled of which 4 on the X chromosome; 2 women; 8 men) independent SNPs at genome-wide significance ($p < 5 \times 10^{-8}$, **Fig S5; Table S10**). The GWAS of AFB identified 89 (84 pooled of which 4 on the X chromosome; 1 women) independent SNPs at genome-wide significance ($p < 5 \times 10^{-8}$, **Fig S6; Table S9**). The distribution of genome-wide test statistics for AFS and AFB showed significant inflation ($\lambda_{GC} = 1.84$ and 1.47 , respectively), however LD score regression showed that this could be attributed almost entirely to polygenicity rather than to population substructure (LD intercept AFS 1.07 (SE = 0.01); AFB 1.03 (SE = 0.01 , **Supp Note**). The LD Score intercept test confirmed that only a very small percentage (5.5%) of the observed inflation in the mean χ^2 statistic was due to population stratification or other confounders, rather than to a polygenic signal.


Polygenic prediction. We then calculated polygenic scores (PGSs) using three different specifications (**Supp Note, Sect 4**). To validate the performance of the PGSs, we performed out-of-sample prediction in the AddHealth and UKHLS cohorts using ordinary least-squares (OLS) regression models and report the R^2 as a measure of goodness-of-fit of the model (**Supp Note 4; Fig S7**). PGSs including all SNPs explain up to 5.8% of the variance for AFS and 4.8% for AFB. A 1 SD change in the AFS/AFB PGS is associated with a 7.3 and 6.3 month

delay in AFS and AFB, respectively. We then ran survival models to account for right-censoring, which occurs when an individual does not experience the event of first sex or birth by the time of the interview.¹³ Using AddHealth data, we estimated nonparametric hazard functions and then compared individuals at the top and bottom 5% of the PGS (see **Fig 2B, 2C** women, **Supp Note Fig S8-9** men). Those in the top 5% PGS for AFS (i.e., genetic predisposition for later AFS) are less likely to have their sexual debut before age 19. AFS PGSs appear more relevant in explaining women's AFS in comparison to men. Those in the top 5% PGS for AFB (i.e., genetic predisposition for later AFB) postpone AFB across all ages until approximately age 27, with similar curves for both sexes.

Environmentally mediated childhood socioeconomic status. Disadvantaged socioeconomic status has been shown to be highly related to early sexual behaviour and teenage pregnancy.¹⁴ To explore the impact of environmentally mediated parental genetic effects on our PGSs, we examined PGS prediction across low (0-10%), medium (50-60%) and high (90-100%) PGS percentiles by parents' education (college versus no college) as a proxy for childhood socioeconomic status (**Supp Note Fig 10A-B**). Indeed, those in the highest decile of the PGS (90-100%) for later AFB have a higher AFB, particularly past age 27, which is accentuated for those with highly educated parents (**Fig 2D; Fig S10A**). Likewise, being in the highest PGS decile for AFS is associated with later sexual intercourse, especially for those from highest socioeconomic childhood households (**Supp Note Fig 10B**).

Genetic correlations. To test the relationships of AFS and AFB with related phenotypes, we calculated genetic correlations using LD score regression¹⁵ (**Fig 3, Supp Note Fig S11, S13, Tab S11**). Given previous evidence,⁸ we examined 28 traits by sex from six relevant categories including: reproductive (e.g., age at menarche, miscarriage or stillbirth, number

145 **Fig 3.** Genetic correlations of age at first birth (A) and age at first sex (B) with a selection of related traits.



146

147 Note: Brief definitions and full results are in Supplementary Table S13. *P*-values for the genetic correlations are reported above each symbol and the horizontal bars
148 represent 95% confidence intervals. If the trait was initially assessed separately for males and females, this is indicated on the left in brackets.

of sexual partners), behavioural (e.g., educational attainment, risk tolerance), psychiatric disorders (e.g., ADHD, schizophrenia), substance use (e.g., age of initiation of smoking, cannabis use), personality (e.g., openness to experience) and anthropometric (e.g., BMI, height). The strongest genetic correlations were observed for reproductive traits, followed by behavioural traits, particularly with AFB and educational attainment in women (0.74, ± 0.01), compared to AFS (0.53, ± 0.01). There was also a negative genetic correlation between adult risk tolerance and AFS/AFB (AFS ~ -0.40 ; AFB ~ -0.25); i.e., those less genetically prone to risk are more genetically predisposed to postpone reproductive behaviour. Amongst psychiatric traits, the strongest correlation was with ADHD (AFS females -0.58 , ± 0.03 , males -0.61 , ± 0.03 ; AFB females -0.63 , ± 0.03 ; males -0.68 , ± 0.09) and Major Depressive Disorder (MDD) (AFS females -0.37 , ± 0.03 , males -0.32 , ± 0.03 ; AFB females, -0.42 , ± 0.03 ; AFB males, -0.33 , ± 0.08). Previous studies have linked MDD to the serotonin transporter gene locus.¹⁶ We also observed strong genetic correlations with age at onset of smoking (AFS ~ 0.68 , ± 0.03 ; AFB ~ 0.74 , ± 0.03), a trait that provides a unique window into adolescent substance use behaviour around the same time of early reproductive behaviour. Genetic factors influencing early smoking, early sexual debut and teenage pregnancy are thus – to some extent – shared. As shown in **Fig 3**, there are few sex differences in these correlations, with the exception of small variations in number of children, anorexia and openness to experience.

GenomicSEM, Exploratory Factor Analysis and Bi-Directional Mendelian Randomization to explore aetiology and causality. To understand the relationships underlying these genetic correlations, we first used GenomicSEM.¹⁷ GenomicSEM uses structural equation modelling to decompose the genetic covariance matrix, calculated using multivariate LD score

regression, of a set of traits. Parameters are estimated by minimizing the difference between the observed genetic covariance matrix and the covariance matrix derived from the model (**Supp Note**). We fit a series of genetic regression models in which AFB (or AFS) was regressed on both years of education and one other possible mediating trait, such as openness, cognitive performance, ADHD and age of initiation of smoking (**Supp Note Tab S12A-L, Fig S12A-B**). In other words, we wanted to test whether the strong genetic correlation of AFS/AFB with education was the result of another mediating trait such as personality, ADHD or substance use. We found that the genetic correlation of years of education with AFB and AFS was independent of factors like risk tolerance, substance use, and psychiatric disorders. This suggests that the genetic correlation between years of education and AFB is largely a product of direct coupling between these traits, rather than being both downstream of a common identified cause. The exception was age at initiation of smoking – as noted previously, a window into risky adolescent behaviour – which partially mediated the relationship of AFB and AFS with years of education.

Exploratory factor analysis (EFA) was then used to examine whether the genetic signal of the onset of reproductive behaviour originated from two genetically distinguishable subclusters of reproductive biology versus externalizing behaviour. Using a two-factor EFA model (**Methods**) to fit the genetic covariance matrix AFS and AFB with these two additional traits, we found that the entire model accounted for 47% of the overall variance, with 22% attributed to risk tolerance and 4% to age at menarche. In a more robust analysis we fit a Genomic SEM for AFB in women and regressed several genetic measures of reproductive biology (age at menarche, age at menopause) and a latent factor representing a common genetic tendency for externalizing behaviour (age at initiation of smoking, age

first used oral contraception, ADHD) (**Fig S14**). These genetic factors predicted 88% of the variance, with the majority of variance significantly predicted by externalizing factors (0.90,±0.02), followed by age at menopause (0.20, ±0.04) and age at menarche (0.16,±0.03). We note that selection bias, induced by the fact that AFB can only be measured among individuals with at least one live birth, may have inflated this estimate.

Given the strong genetic correlations between the phenotypes discussed above, we used Mendelian Randomization (MR)-based analyses¹⁸ to explore causality and assess the direction of effect between AFB, AFS and years of education¹⁹ as well as risk taking (measured in adulthood)⁴ and age at smoking initiation²⁰ (**Supp Note Sect 8, Tab S13A**). For the majority of pairs of phenotypes we found strong evidence of bi-directionality, which was also seen after applying Steiger fitting. The relationship between AFB and years in education appeared to be the explanatory factor that linked AFB to the two risk taking phenotypes. This was not the case, however, for AFS where the analysis suggests that age at initiation of smoking (and the environment and processes that lead to this) are upstream of the start of AFS. In that case the relationship was significant when assessed as age at smoking to AFS but not the other way round. Of note, associations were much stronger for age at initiation of smoking initiation than for risk-taking behaviour assessed in adulthood, suggesting that the timing of this behaviour is key.

A second set of MR analyses examined whether AFS and AFB PGSs have effects on type 2 diabetes (T2D)²¹ and coronary artery disease (CAD),²² independently of years of education (**Fig 1E, Tab S13B, Fig S15**). T2D and CAD were chosen since they are two common major diseases, with broadly defined behavioural risk factors. Findings show that the association with years of education and later life diseases are substantially attenuated

by the effects of AFB. This concurs with a large body of research that has established a biological association with the timing of AFB and metabolic diseases including early AFB linked to high blood pressure,²³ obesity²⁴ and diabetes.²⁵ Reproductive timing thus appears to capture a latent variable that detects these metabolic effects but also years of education and other behavioural traits and can therefore serve as a more powerful predictor of later life disease than years of education alone. This also suggests that many of the associations with diseases that have previously been ascribed to years of education, may result from a more broadly defined socio-behavioural trajectory.


Cox proportional hazard models of a polygenic score for AFB on longevity. The disposable soma theory of evolution hypothesizes that longevity demands investments in somatic maintenance – such as remaining in education – that in turn reduces resources available for reproduction. To test trade-offs between reproductive behaviour and senescence as argued in the ageing and longevity literature,²⁶ we conducted additional analyses to test whether our AFB PGS was associated with (parental) longevity (**Supp Note, Tab S14**). We first estimated a baseline Cox proportional hazard model of our AFB PGS on parental longevity and then included the EA3 PGS and risk covariates followed by a final model including number of siblings as a proxy for parental fertility. We found that a genetically predicted 1 SD increase in AFB is associated with a 2-4% lower mortality, suggesting that there is likely a trade-off between reproduction and longevity.

Gene prioritization. To understand the biology represented by the variants associated with AFS and/or AFB, we performed a gene prioritization analysis that connected variants to genes and prioritized candidate genes based on likely involvement reproductive biology or psychiatric traits. To this end, we used predicted gene function,²⁷ single-cell RNA sequencing

data in mice,^{28,29} literature mining,³⁰ *in silico* sequencing,³¹ and Summary-data based Mendelian Randomization(SMR)³² using eQTL data from brain and blood.³³ Integrating results across all approaches resulted in the prioritization of 386 unique genes; 314 genes in 159 loci for AFS and 106 genes in 42 loci for AFB (**Supp Tab 15A-19C**). Of these, 99 were expressed at the protein level in cell types of brain, glands, and/or (fe)male reproductive organs³⁴ (**Fig 4**). Gene prioritization in sex-specific loci resulted in the prioritization of 11 genes for AFB in women, one gene for AFS in women and 23 genes for AFS in men. Of these, 12 genes at three loci were expressed at the protein level in relevant tissues (**Supp Note, Fig S16**).

Genes that play a role in follicle stimulating hormone (*CGA*³⁵), oocyte development (*KLF17*³⁶), and implantation and placental growth (*ESR1*, *SUMO1*³⁷, *ARNT*,³⁸ *CAV1*,³⁹ *E2F1*⁴⁰) were prioritized for AFS in data from men and women combined, while *FSHB*⁴¹ and *ESR1* were (also) prioritized for AFB. Other genes prioritized in loci identified in the pooled meta-analyses were expressed at the protein level in (developing) sperm – highlighting a role for spermatid differentiation (*KLF17*⁴²) for AFS – as well as for sperm morphogenesis and binding between acrosome-reacted sperm and the zona pellucida (*ZPBP*⁴³) for AFB. The meta-analysis in women only yielded genes related to endometriosis (*CCR1*)⁴⁴ and spontaneous abortion (*CXCR6*) for AFB (**Supp Note Fig S18**).⁴⁵ Taken together, these results suggest that intrinsic biological processes that influence fertility also influence the onset of sexual behaviour in men and women. Interestingly, *NUP210L* – prioritized for AFS and highly expressed in developing and mature sperm³⁴ – is normally testis-specific, but was recently shown to be expressed in prefrontal cortex neurons of G allele carriers in rs114697636 (MAF 3%, D' 0.90 with AFS lead SNP rs113142203), attributed to allele-specific activation through

Fig 4. Gene prioritization of age at first sex (AFS) and age at first birth (AFB)



Note: Information for 99 genes prioritized in loci identified by GWAS for age at first sex and/or age at first birth that are located within 1 million bp of lead SNPs and are highly expressed at the protein level in brain, glands and/or reproductive organs. Blue and orange indicate transitions from one locus to the next. The first panel indicates if the locus was identified as being associated at genome-wide significance with age at first sex (AFS), age at first birth (AFB). The second panel shows which bioinformatic approaches highlighted the gene as a candidate. The third panel shows – from left to right - the cell types in brain, glands, female reproductive organs, and male reproductive organs in which the genes are expressed at a low, moderate or high level (small, medium and large circles) based on data from the Human Protein Atlas. The fourth panel shows gene functions as extracted from Entrez, Uniprot and GeneCards. The fifth panel indicates which phenotypes were observed in mutant mice, as reported by the Mouse Genome Informatics (MGI) database.

improved binding affinity for testis receptor 2.⁴⁶ Methylation of, and variants near *NUP210L* have been associated with psychologic development disorders, intelligence, and

mathematical ability,⁴⁷ illustrating how a testis-specific gene can influence the brain in some individuals.

Several genes prioritized in AFS-associated loci in data from men and women combined have previously been implicated in risk seeking behaviour, sociability and anxiety (*GTF2I*,⁴⁸ *TOP2B*,⁴⁹ *E2F1*⁵⁰, *NCAM1*,⁵¹ *NFASC*,⁵² *MEF2C*⁵³). In the sex-specific meta-analysis for AFS, a role for externalizing behaviour was supported through *ERBB4* in women; and through *SLC44A1* and *NR1H3* in men. *ERBB4* has previously been linked to fear, anxiety,⁵⁴ schizophrenia,⁵⁵ and polycystic ovary syndrome (PCOS);⁵⁶ *SLC44A1* encodes a choline transporter that plays a key role in cerebral inhibition related to substance use and depressive disorders⁵⁷; and *NR1H3* has been implicated in major depressive disorder (MDD).⁵⁸ These genes provide concrete examples of how an innate predisposition for externalizing behaviour can influence initiation of reproductive behaviour.

The gene prioritization results partly mirror and compliment the rigorous post-GWAS *in silico* association analyses we performed for loci identified for AFS and AFB. However, experimental validation is required before firm conclusions can be drawn about the involvement of, and mechanisms through which prioritized candidate genes influence AFS and AFB. More information on protein-protein interaction hubs, as well as on genes highlighted by literature mining³⁰ are provided in the supplementary information.


Discussion

In this study, we presented the results of the largest GWAS to date of the onset of human reproductive behaviour in the form of age at first sex (AFS) (N=387,338) and age at first birth (AFB) (N=542,901). We identified 370 independent signals harbouring at least 386

prioritized candidate genes, using 1000G imputed genotype data and an X-Chromosome analysis, which allowed us to detect considerably more signals than ever before (**Fig 5**). In comparison, a recent GWAS for type 2 diabetes,⁵⁹ for instance, detected 243 loci. Similar to previous work, we showed that the total SNP heritability accounted for 10-22% of phenotypic variance and varied by birth cohort.^{12,60} The incremental R^2 of our PGSs based on significantly associated loci is around 5-6%, similar what is observed for common demographic and social variables (e.g., years of education, age at marriage), which have been classically used to explain the timing of human reproductive behaviour. Comparatively, 5-6% is in the range observed for other complex traits, like BMI (5.8%)⁶¹ and schizophrenia (8.4%).⁶² The number of signals also opened up opportunities for functional follow-up analyses which suggested a role for spermatid differentiation and oocyte development. The analyses of the correlation and underlying aetiology of these traits revealed a common genetic basis of both AFS and AFB with externalizing behaviour and substance use and links to internalizing traits and infertility. Finally, we showed that AFB is an important predictor for late age at onset of disease and longevity, and that it substantially attenuates the effect of years in education.

Although we opened many new avenues for research, the present GWAS still faces certain limitations. First, the sample sizes for men were still appreciably smaller than for women since reproductive and fertility data is routinely collected less often from men. In order to understand the causes of infertility in men this needs to be taken into consideration in future data collection. Initial within-family analyses showed that our discovery GWAS may actually overestimate causal effects (**Supp Note**), genotypes associated with later onset of reproductive behaviour genotypes are also associated with

323 **Fig 5.** Summary genome-wide association study of timing of onset of reproductive behaviour: age at first sex (AFS) and age at first birth (AFB)



parental reproductive genotypes, likely leading to a social environment that affects reproductive and other behaviours. Collection and analysis of family data is clearly a future area of research for reproductive and related behaviour. The lack of accessibility of publically available summary statistics from published research, meant that we were unable to examine the relationship with other traits, particularly with infertility related traits (e.g., PCOS). Future data collection could benefit from focussing on behavioural disinhibition markers that appear to be highly related to self-control, which has implications for disease prevention and behavioural interventions into lifestyle factors related to obesity, Type 2 diabetes or substance use disorders. A glaring limitation is our focus on European-ancestry individuals in Western countries. Whilst common in this area of research,⁶³ extension to other ancestries and geographical contexts is required in the future. This is particularly relevant in the context of parent gene-environment interactions, which may be specific to the social background of the sample.

Our detailed correlation, GenomicSEM and MR analyses also provided a deeper understanding of the underlying aetiology of related traits and pleiotropy and the associations between human reproductive behaviour and disease risk. We anticipate that our results will provide leads to address important interventions in infertility, teenage sexual and mental health, as well as for functional follow-up experiments that will likely yield targets that can be translated in efficient medication to improve fertility (e.g., in IVF) but also for interventions on reproductive health related to earlier sexual debut and teenage pregnancy.

References

1. Mercer, C. H. *et al.* Changes in sexual attitudes and lifestyles in Britain through the life course and over

- time: findings from the National Surveys of Sexual Attitudes and Lifestyles (Natsal). *Lancet* **382**, 1781–1794 (2013).
2. Lara, L. A. S. & Abdo, C. H. N. Age at Time of Initial Sexual Intercourse and Health of Adolescent Girls. *J. Pediatr. Adolesc. Gynecol.* **29**, 417–423 (2016).
3. Polimanti, R. *et al.* The Interplay between Risky Sexual Behaviors and Alcohol Dependence: Genome-Wide Association and Neuroimaging Support for LHPP as a Risk Gene. *Neuropsychopharmacology* **42**, 598–605 (2017).
4. Karlsson Linnér, R. *et al.* Genome-wide association analyses of risk tolerance and risky behaviors in over 1 million individuals identify hundreds of loci and shared genetic influences. *Nat. Genet.* **51**, 245–257 (2019).
5. Balbo, N., Billari, F. C. & Mills, M. Fertility in Advanced Societies: A Review of Research. *Eur. J. Popul. / Rev. Eur. Démographie* **29**, 1–38 (2013).
6. Mills, M. C. *et al.* Why do people postpone parenthood? Reasons and social policy incentives. *Hum. Reprod. Update* **17**, 848–860 (2011).
7. Rahmioglu, N. *et al.* Genome-wide enrichment analysis between endometriosis and obesity-related traits reveals novel susceptibility loci. *Hum. Mol. Genet.* **24**, 1185–1199 (2015).
8. Barban, N. *et al.* Genome-wide analysis identifies 12 loci influencing human reproductive behavior. *Nat. Genet.* **48**, (2016).
9. Balbo, N., Billari, F. C. & Mills, M. C. Fertility in advanced societies: A review of research. *Eur. J. Popul. Eur. Démographie* **29**, 1–38 (2013).
10. Day, F. R. *et al.* Physical and neurobehavioral determinants of reproductive onset and success. *Nat. Genet.* doi:10.1038/ng.3551 (2016).
11. Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: a tool for genome-wide complex trait analysis. *Am. J. Hum. Genet.* **88**, 76–82 (2011).
12. Tropf, F. C. *et al.* Hidden heritability due to heterogeneity across seven populations. *Nat. Hum. Behav.* **1**, 757–765 (2017).
13. Mills, M. C. Introducing Survival and Event History Methods. in (Sage Publications, 2011).
14. Singh, S., Darroch, J. E. & Frost, J. J. Socioeconomic Disadvantage and Adolescent Women’s Sexual and Reproductive Behavior: The Case of Five Developed Countries. *Fam. Plann. Perspect.* **33**, 251 (2001).
15. Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide association summary statistics. *Nat. Genet.* **47**, 1228–1235 (2015).
16. Seripa, D. *et al.* Role of the serotonin transporter gene locus in the response to SSRI treatment of major depressive disorder in late life. *J. Psychopharmacol.* **29**, 623–633 (2015).
17. Grotzinger, A. D. *et al.* Genomic structural equation modelling provides insights into the multivariate genetic architecture of complex traits. *Nat. Hum. Behav.* **3**, 513–525 (2019).
18. Davey Smith, G. What can mendelian randomisation tell us about modifiable behavioural and environmental exposures? *BMJ* (2005). doi:10.1136/bmj.330.7499.1076
19. Lee, J. J. *et al.* Gene discovery and polygenic prediction from a genome-wide association study of educational attainment in 1.1 million individuals. *Nat. Genet.* **50**, 1112–1121 (2018).
20. Liu, M. *et al.* Association studies of up to 1.2 million individuals yield new insights into the genetic etiology of tobacco and alcohol use. *Nat. Genet.* **51**, 237–244 (2019).
21. McCarthy, M. & Mahajan, A. Genome-wide association analyses in Type 2 Diabetes: The gift that keeps on giving. *Blog* (2018). Available at: <http://mccarthy.well.ox.ac.uk/2018/10/gwas-gift-keeps-giving/>.
22. Nelson, C. P. *et al.* Association analyses based on false discovery rate implicate new loci for coronary artery disease. *Nat. Genet.* **49**, 1385–1391 (2017).
23. Lind, J. M., Hennessy, A. & Chiu, C. L. Association Between a Woman’s Age at First Birth and High Blood Pressure. *Medicine (Baltimore)*. **94**, e697 (2015).
24. Patchen, L., Leoutsakos, J.-M. & Astone, N. M. Early Parturition: Is Young Maternal Age at First Birth Associated with Obesity? *J. Pediatr. Adolesc. Gynecol.* **30**, 553–559 (2017).
25. Kim, J. H., Jung, Y., Kim, S. Y. & Bae, H. Y. Impact of Age at First Childbirth on Glucose Tolerance Status in Postmenopausal Women: The 2008–2011 Korean National Health and Nutrition Examination Survey. *Diabetes Care* **37**, 671–677 (2014).
26. Eisenberg, D. T. A., Hayes, M. G. & Kuzawa, C. W. Delayed paternal age of reproduction in humans is associated with longer telomeres across two generations of descendants. *Proc. Natl. Acad. Sci.* **109**, 10251–10256 (2012).
27. Pers, T. H. *et al.* Biological interpretation of genome-wide association studies using predicted gene functions. *Nat. Commun.* **6**, 5890 (2015).

28. Zeisel, A. *et al.* Molecular Architecture of the Mouse Nervous System. *Cell* **174**, 999–1014.e22 (2018).
29. Tabula Muris Consortium *et al.* Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris. *Nature* **562**, 367–372 (2018).
30. Yang, H., Robinson, P. N. & Wang, K. Phenolyzer: phenotype-based prioritization of candidate genes for human diseases. *Nat. Methods* **12**, 841–3 (2015).
31. Vaez, A. *et al.* In Silico Post Genome-Wide Association Studies Analysis of C-Reactive Protein Loci Suggests an Important Role for Interferons. *Circ. Cardiovasc. Genet.* **8**, 487–497 (2015).
32. Zhu, Z. *et al.* Integration of summary data from GWAS and eQTL studies predicts complex trait gene targets. *Nat. Genet.* **48**, 481–7 (2016).
33. van der Wijst, M. G. P. *et al.* Single-cell RNA sequencing identifies celltype-specific cis-eQTLs and co-expression QTLs. *Nat. Genet.* **50**, 493–497 (2018).
34. Uhlén, M. *et al.* Proteomics. Tissue-based map of the human proteome. *Science* **347**, 1260419 (2015).
35. Ellsworth, B. S. *et al.* FOXL2 in the pituitary: molecular, genetic, and developmental analysis. *Mol. Endocrinol.* **20**, 2796–805 (2006).
36. van Vliet, J. *et al.* Human KLF17 is a new member of the Sp/KLF family of transcription factors. *Genomics* **87**, 474–82 (2006).
37. Governini, L. *et al.* FOXL2 in human endometrium: hyperexpressed in endometriosis. *Reprod. Sci.* **21**, 1249–55 (2014).
38. Rico, C. *et al.* HIF1 activity in granulosa cells is required for FSH-regulated Vegfa expression and follicle survival in mice. *Biol. Reprod.* **90**, 135 (2014).
39. Dai, Z. *et al.* Caveolin-1 promotes trophoblast cell invasion through the focal adhesion kinase (FAK) signalling pathway during early human placental development. *Reprod. Fertil. Dev.* (2019). doi:10.1071/RD18296
40. Artini, P. G. *et al.* Cumulus cells surrounding oocytes with high developmental competence exhibit down-regulation of phosphoinositol 1,3 kinase/protein kinase B (PI3K/AKT) signalling genes involved in proliferation and survival. *Hum. Reprod.* **32**, 2474–2484 (2017).
41. Zheng, J. *et al.* Novel FSH β mutation in a male patient with isolated FSH deficiency and infertility. *Eur. J. Med. Genet.* **60**, 335–339 (2017).
42. Yan, W., Burns, K. H., Ma, L. & Matzuk, M. M. Identification of Zfp393, a germ cell-specific gene encoding a novel zinc finger protein. *Mech. Dev.* **118**, 233–9 (2002).
43. Lin, Y.-N., Roy, A., Yan, W., Burns, K. H. & Matzuk, M. M. Loss of zona pellucida binding proteins in the acrosomal matrix disrupts acrosome biogenesis and sperm morphogenesis. *Mol. Cell. Biol.* **27**, 6794–805 (2007).
44. Wieser, F. *et al.* Expression and regulation of CCR1 in peritoneal macrophages from women with and without endometriosis. *Fertil. Steril.* **83**, 1878–1881 (2005).
45. Mei, J. *et al.* CXCL16/CXCR6 interaction Promotes Endometrial Decidualization via the PI3K/ AKT Pathway. *Reproduction* (2019). doi:10.1530/REP-18-0417
46. Gusev, F. E. *et al.* Epigenetic-genetic chromatin footprinting identifies novel and subject-specific genes active in prefrontal cortex neurons. *FASEB J.* **33**, 8161–8173 (2019).
47. Quinn, J. P., Savage, A. L. & Bubb, V. J. Non-coding genetic variation shaping mental health. *Curr. Opin. Psychol.* **27**, 18–24 (2019).
48. Barak, B. *et al.* Neuronal deletion of Gtf2i, associated with Williams syndrome, causes behavioral and myelin alterations rescuable by a remyelinating drug. *Nat. Neurosci.* **22**, 700–708 (2019).
49. Li, Y. *et al.* Topoisomerase II β is required for proper retinal development and survival of postmitotic cells. *Biol. Open* **3**, 172–84 (2014).
50. Athanasiou, M. C. *et al.* The transcription factor E2F-1 in SV40 T antigen-induced cerebellar Purkinje cell degeneration. *Mol. Cell. Neurosci.* **12**, 16–28 (1998).
51. Yang, X. *et al.* The association between NCAM1 levels and behavioral phenotypes in children with autism spectrum disorder. *Behav. Brain Res.* **359**, 234–238 (2019).
52. Locke, A. E. *et al.* Genetic studies of body mass index yield new insights for obesity biology. *Nature* **518**, 197–206 (2015).
53. Tu, S. *et al.* NitroSynapsin therapy for a mouse MEF2C haploinsufficiency model of human autism. *Nat. Commun.* **8**, 1488 (2017).
54. Shamir, A. *et al.* The Importance of the NRG-1/ErbB4 Pathway for Synaptic Plasticity and Behaviors Associated with Psychiatric Disorders. *J. Neurosci.* **32**, 2988–2997 (2012).
55. Yang, J.-M. *et al.* erbb4 Deficits in Chandelier Cells of the Medial Prefrontal Cortex Confer Cognitive Dysfunctions: Implications for Schizophrenia. *Cereb. Cortex* **29**, 4334–4346 (2019).

56. Day, F. R. *et al.* Causal mechanisms and balancing selection inferred from genetic associations with polycystic ovary syndrome. *Nat. Commun.* **6**, 8464 (2015).
57. Baumgartner, H. K. *et al.* Characterization of choline transporters in the human placenta over gestation. *Placenta* **36**, 1362–9 (2015).
58. Peng, Z. *et al.* Liver X receptor β in the hippocampus: A potential novel target for the treatment of major depressive disorder? *Neuropharmacology* **135**, 514–528 (2018).
59. Mahajan, A. *et al.* Fine-mapping type 2 diabetes loci to single-variant resolution using high-density imputation and islet-specific epigenome maps. *Nat. Genet.* **50**, 1505–1513 (2018).
60. Tropf, F. C. *et al.* Human Fertility, Molecular Genetics, and Natural Selection in Modern Societies. *PLoS One* **10**, e0126821 (2015).
61. Warren, E. B. & *et al.* Heterogeneity in polygenic scores for common human traits. *bioRxiv* (2017). doi:10.1101/106062
62. Ripke, S. *et al.* Biological insights from 108 schizophrenia-associated genetic loci. *Nature* **511**, 421–427 (2014).
63. Mills, M. C. & Rahal, C. The GWAS Diversity Monitor tracks diversity by disease in real time. *Nat. Genet.* **52**, 242–243 (2020).

Acknowledgements

Acknowledgments and grant details can be found in the Supplementary Note Section 12.2.

Author Contributions

A full list of all authors, including the Human Reproductive Behaviour Consortium, eQTLGen and BIOS Consortium can be founded in the Supplementary Note 12.1. MCM and FRD designed and led the study. MCM wrote the paper and supplementary note with contributions by authors for respective analyses and comments by all authors. DMB conducted phenotypic changes, phenotype preparation, LD Score and genetic correlations, Genomic SEM and exploratory factor analysis and sex-specific effects. NB conducted GWAS meta-analysis, MTAG, PGS prediction, survival models, and Cox models of longevity. FCT and FRD conducted the cohort QC. FCT conducted GREML cohort heritability analysis and phenotype preparation in UKBB. FRD ran Mendelian Randomization, conducted GWAS analyses and JRBP conducted COJO and X-Chromosome analysis. NvZ conducted DEPICT and Phenolyzer analyses. AV and HS conducted *in silico* sequencing and SMR analyses. TP conducted cell type enrichment analyses. MdH integrated gene prioritization results and

performed downstream analyses, e.g. Human Protein Atlas; Entrez, GeneCards and Uniprot mining; and STRING Protein-Protein interaction analyses. Authors in the Human Reproductive Behaviour Consortium provided data and cohort analyses. The eQTLGen and BIOS Consortia provided data for additional analyses. All authors reviewed and approved the final version of the paper.

Competing interests

The main authors declare no competing interests. The views expressed in this article are those of the author(s) and not necessarily those of the NHS, the NIHR, or the Department of Health. MMCC (Mark McCarthy) has served on advisory panels for Pfizer, NovoNordisk and Zoe Global, has received honoraria from Merck, Pfizer, Novo Nordisk and Eli Lilly, and research funding from Abbvie, Astra Zeneca, Boehringer Ingelheim, Eli Lilly, Janssen, Merck, NovoNordisk, Pfizer, Roche, Sanofi Aventis, Servier, and Takeda. As of June 2019, MMCC is an employee of Genentech, and a holder of Roche stock.

Additional Information

This work was conducted under UK Biobank application 11425, 12514, 22276 and 9797.

Data and materials availability

Upon publication, GWAS summary statistics are available at the GWAS Catalog (www.ebi.ac.uk/gwas/downloads/summary-statistics). Access to individual level data from multiple sources used in this GWAS can be obtained by bona fide scientists through application to the specific data providers. The analysis plan was pre-deposited in the Open Science Framework website: <https://osf.io/b4r4b/>

Supplementary information is available at: <https://doi.org/xxxxxx>

Reprints and permissions information is available at: xxxx

Correspondence and requests for material should be addressed to M.C.M. and F.R.D.

Authors

Melinda C. Mills^{1,2,*}, Felix C. Tropf^{1,2,3,4,†}, David M. Brazeal^{1,2,†}, Natalie van Zuydam⁵, Ahmad Vaez^{6,7}, eQTLGen Consortium, BIOS Consortium, Tune H. Pers^{8,9}, Harold Snieder⁶, John R.B. Perry¹⁰, Ken K. Ong^{10,†}, Marcel den Hoed^{5,†}, Nicola Barban^{11,†}, and Felix R. Day^{10,†,*} on behalf of the *Human Reproductive Behaviour Consortium*

¹ Leverhulme Centre for Demographic Science, University of Oxford, Oxford, United Kingdom

² Nuffield College, University of Oxford, Oxford, United Kingdom

³ École Nationale de la Statistique et de L'administration Économique (ENSAE), Paris, France

⁴ Center for Research in Economics and Statistics (CREST), Paris, France

⁵ The Beijer Laboratory and Department of Immunology, Genetics and Pathology, Uppsala University and SciLifeLab, Uppsala, Sweden

⁶ Department of Epidemiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

⁷ Department of Bioinformatics, Isfahan University of Medical Sciences, Isfahan, Iran

⁸ The Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

⁹ Department of Epidemiology Research, Statens Serum Institut, Copenhagen, Denmark

¹⁰ MRC Epidemiology Unit, Institute of Metabolic Science, University of Cambridge, Cambridge, United Kingdom

¹¹ Institute of Social and Economic Research, University of Essex, Essex, United Kingdom

[†] Denotes equal contribution

* Correspondence to Melinda C. Mills, melinda.mills@nuffield.ox.ac.uk, and Felix R. Day, felix.day@mrc-epid.cam.ac.uk

Human Reproductive Behaviour Consortium - Author information

Evelina T. Akimova¹, Sven Bergmann^{2,3,4}, Jason D. Boardman⁵, Dorret I. Boomsma⁶, Marco Brumat⁷, Julie E. Buring^{8,9}, David Cesarini^{10,11,12}, Daniel I. Chasman^{8,9}, Jorge E. Chavarro^{13,14,15}, Massimiliano Cocca¹⁶, Maria Pina Concas¹⁶, George Davey-Smith¹⁷, Gail Davies¹⁸, Ian J. Deary¹⁸, Tõnu Esko^{19,20}, Oscar Franco²¹, Audrey J. Gaskins^{14,15,22}, Eco J.C. de Geus⁶, Christian Gieger²³, Giorgia Grotto^{7,16}, Hans Jörgen Grabe²⁴, Erica P. Gunderson²⁵, Kathleen Mullan Harris²⁶, Fernando P. Hartwig^{17,27}, Chunyan He^{28,29}, Diana van Heemst³⁰, W. David Hill¹⁸, Georg Homuth³¹, Bernardo Lessa Horta²⁷, Jouke Jan Hottenga⁶, Hongyang Huang¹³, Elina Hyppönen^{32,33}, M. Arfan Ikram²¹, Rick Jansen³⁴, Magnus Johannesson³⁵, Zoha Kamali³⁶, Maryam Kavousi²¹, Peter Kraft^{13,37}, Brigitte Kühnel²³, Claudia Langenberg³⁸, Lifelines Cohort Study^{39,40}, Penelope A. Lind⁴¹, Jian'an Luan³⁸, Reedik Mägi¹⁹, Patrik K.E. Magnusson⁴², Anubha Mahajan^{43,44}, Nicholas G. Martin⁴⁵, Hamdi Mbarek^{6,46}, Mark I. McCarthy^{43,44}, George McMahon⁴⁷, Matthew B. McQueen⁴⁸, Sarah E. Medland⁴¹, Thomas Meitinger⁴⁹, Andres Metspalu^{19,50}, Evelin Mihailov¹⁹, Lili Milani¹⁹, Stacey A. Missmer^{13,51,52}, Stine Møllegaard⁵³, Dennis O. Mook-Kanamori^{54,55}, Anna Morgan¹⁶, Peter J. van der Most³⁹, Renée de Mutsert⁵⁴, Matthias Nauck⁵⁶, Ilja M. Nolte³⁹, Raymond Noordam³⁰, Brenda W.J.H. Penninx⁵⁷, Annette Peters⁵⁸, Chris Power⁵⁹, Paul Redmond¹⁸, Janet W. Rich-Edwards^{13,15,60}, Paul M. Ridker^{8,9}, Cornelius A. Rietveld^{61,62}, Susan M. Ring¹⁷, Lynda M. Rose⁸, Rico Rueedi^{2,3}, Kári Stefánsson⁶³, Doris Stöckl⁵⁸, Konstantin Strauch^{64,65,66}, Morris A. Swertz⁴⁰, Alexander Teumer⁶⁷, Gudmar Thorleifsson⁶³, Unnur Thorsteinsdottir⁶³, A. Roy Thurik^{61,62,68}, Nicholas J. Timpson¹⁷, Constance Turman¹³, André G. Uitterlinden^{61,69}, Melanie Waldenberger^{23,58}, Nicholas J. Wareham³⁸, Gonneke Willemsen⁶, and Jing Hau Zhao³⁸

Author list ordered alphabetically.

¹ Leverhulme Centre for Demographic Science, Department of Sociology, St. Antony's College, University of Oxford, Oxford, United Kingdom

² Department of Computational Biology, University of Lausanne, Lausanne, Switzerland

³ Swiss Institute of Bioinformatics, Lausanne, Switzerland

⁴ Department of Integrative Biomedical Sciences, University of Cape Town, Cape Town, South Africa

⁵ Department of Sociology and Institute of Behavioral Science, University of Colorado at Boulder, Boulder, CO, United States of America

⁶ Department of Biological Psychology, Amsterdam Public Health Research Institute, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

⁷ Department of Medical, Surgical and Health Sciences, University of Trieste, Trieste, Italy

⁸ Brigham and Women's Hospital, Boston, MA, United States of America

⁹ Harvard Medical School, Boston, MA, United States of America

¹⁰ Department of Economics, New York University, New York, NY, United States of America

¹¹ Research Institute for Industrial Economics, Stockholm, Sweden

¹² National Bureau of Economic Research, Cambridge, MA, United States of America

¹³ Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, United States of America

¹⁴ Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA, United States of America

¹⁵ Channing Division of Network Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, United States of America

¹⁶ Institute for Maternal and Child Health IRCCS "Burlo Garofolo", Trieste, Italy

¹⁷ MRC Integrative Epidemiology Unit, University of Bristol, Bristol, United Kingdom

¹⁸ Lothian Birth Cohorts, Department of Psychology, University of Edinburgh, Edinburgh, United Kingdom

¹⁹ Estonian Genome Center, University of Tartu, Tartu, Estonia

- 599 ²⁰ Broad Institute of the Massachusetts Institute of Technology and Harvard University, Cambridge,
600 MA, United States of America
- 601 ²¹ Department of Epidemiology, Erasmus Medical Center, Rotterdam, The Netherlands
- 602 ²² Department of Epidemiology, Rollins School of Public Health, Emory University, Atlanta, GA, United
603 States of America
- 604 ²³ Research Unit of Molecular Epidemiology, Helmholtz Zentrum München, German Research Center
605 for Environmental Health, Neuherberg, Germany
- 606 ²⁴ Department of Psychiatry and Psychotherapy, University Medicine Greifswald, Greifswald, Germany
- 607 ²⁵ Division of Research, Kaiser Permanente Northern California, Oakland, CA, United States of America
- 608 ²⁶ Department of Sociology, Carolina Population Center, University of North Carolina at Chapel Hill,
609 Chapel Hill, NC, United States of America
- 610 ²⁷ Postgraduate Program in Epidemiology, Federal University of Pelotas, Pelotas, Brazil
- 611 ²⁸ University of Kentucky Markey Cancer Center, Lexington, KY, United States of America
- 612 ²⁹ Department of Internal Medicine, Division of Medical Oncology, University of Kentucky College of
613 Medicine, Lexington, KY, United States of America
- 614 ³⁰ Department of Internal Medicine, Section of Gerontology and Geriatrics, Leiden University Medical
615 Center, Leiden, The Netherlands
- 616 ³¹ Interfaculty Institute for Genetics and Functional Genomics, University of Greifswald, Greifswald,
617 Germany
- 618 ³² Australian Centre for Precision Health, University of South Australia Cancer Research Institute,
619 Adelaide, Australia
- 620 ³³ South Australian Health and Medical Research Institute, Adelaide, Australia
- 621 ³⁴ Department of Psychiatry, Amsterdam Public Health and Amsterdam Neuroscience, Amsterdam
622 UMC, Vrije Universiteit, Amsterdam, The Netherlands
- 623 ³⁵ Department of Economics, Stockholm School of Economics, Stockholm, Sweden
- 624 ³⁶ Department of Bioinformatics, Isfahan University of Medical Sciences, Isfahan, Iran
- 625 ³⁷ Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA, United States
626 of America
- 627 ³⁸ MRC Epidemiology Unit, Institute of Metabolic Science, Cambridge Biomedical Campus, University
628 of Cambridge School of Clinical Medicine, Cambridge, United Kingdom
- 629 ³⁹ Department of Epidemiology, University of Groningen, University Medical Center Groningen,
630 Groningen, The Netherlands
- 631 ⁴⁰ Department of Genetics, University of Groningen, University Medical Center Groningen, Groningen,
632 The Netherlands
- 633 ⁴¹ Psychiatric Genetics, QIMR Berghofer Medical Research Institute, Herston Brisbane, Queensland,
634 Australia
- 635 ⁴² Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden
- 636 ⁴³ Wellcome Centre for Human Genetics, University of Oxford, Oxford, United Kingdom
- 637 ⁴⁴ Oxford Centre for Diabetes, Endocrinology and Metabolism, Radcliffe Department of Medicine,
638 University of Oxford, Oxford, United Kingdom
- 639 ⁴⁵ Genetic Epidemiology, QIMR Berghofer Medical Research Institute, Herston Brisbane, Queensland,
640 Australia
- 641 ⁴⁶ Qatar Genome Programme, Qatar Foundation, Doha, Qatar
- 642 ⁴⁷ School of Social and Community Medicine University of Bristol, Bristol, United Kingdom
- 643 ⁴⁸ Department of Integrative Physiology, University of Colorado at Boulder, Boulder, CO, United States
644 of America
- 645 ⁴⁹ Institute of Human Genetics, Helmholtz Zentrum München, German Research Center for
646 Environmental Health, Neuherberg, Germany
- 647 ⁵⁰ Institute of Molecular and Cell Biology, University of Tartu, Tartu, Estonia
- 648 ⁵¹ Division of Adolescent and Young Adult Medicine, Department of Medicine, Boston Children's
649 Hospital and Harvard Medical School, Boston, MA, United States of America

- ⁵² Department of Obstetrics, Gynecology, and Reproductive Biology, College of Human Medicine, Michigan State University, Grand Rapids, MI, United States of America
- ⁵³ Department of Sociology, University of Copenhagen, Copenhagen, Denmark
- ⁵⁴ Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, The Netherlands
- ⁵⁵ Department of Public Health and Primary Care, Leiden University Medical Center, Leiden, The Netherlands
- ⁵⁶ Institute of Clinical Chemistry and Laboratory Medicine, University Medicine Greifswald, Greifswald, Germany
- ⁵⁷ Department of Psychiatry, EMGO Institute for Health and Care Research and Neuroscience Campus Amsterdam, VU University Medical Center/GGZ inGeest, Amsterdam, The Netherlands
- ⁵⁸ Institute of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany
- ⁵⁹ Population, Policy and Practice Research and Teaching Department, UCL Great Ormond Street Institute of Child Health, London, United Kingdom
- ⁶⁰ Division of Women's Health, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, United States of America
- ⁶¹ Erasmus University Rotterdam Institute for Behavior and Biology, Rotterdam, The Netherlands
- ⁶² Department of Applied Economics, Erasmus School of Economics, Rotterdam, The Netherlands
- ⁶³ deCODE Genetics/Amgen Inc., Reykjavik, Iceland
- ⁶⁴ Institute of Medical Biostatistics, Epidemiology and Informatics (IMBEI), University Medical Center, Johannes Gutenberg University, Mainz, Germany
- ⁶⁵ Institute of Genetic Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany
- ⁶⁶ Chair of Genetic Epidemiology, IBE, Faculty of Medicine, LMU Munich, Germany
- ⁶⁷ Institute for Community Medicine, University Medicine Greifswald, Greifswald, Germany
- ⁶⁸ Montpellier Business School, Montpellier, France
- ⁶⁹ Department of Internal Medicine, Erasmus University Medical Center, Rotterdam, The Netherlands

eQTLGen Consortium – Author information

Mawussé Agbessi¹, Habibul Ahsan², Isabel Alves¹, Anand Kumar Andiappan³, Wibowo Arindrarto⁴, Philip Awadalla¹, Alexis Battle^{5,6}, Frank Beutner⁷, Marc Jan Bonder^{8,9}, Dorret I. Boomsma¹⁰, Mark W. Christiansen¹¹, Anniq Claringbould^{8,12}, Patrick Deelen^{8,13,12,14}, Tõnu Esko¹⁵, Marie-Julie Favé¹, Lude Franke^{8,12}, Timothy Frayling¹⁶, Sina A. Gharib^{11,17}, Greg Gibson¹⁸, Bastiaan T. Heijmans⁴, Gibran Hemani¹⁹, Rick Jansen²⁰, Mika Kähönen²¹, Anette Kalnapenkis¹⁵, Silva Kasela¹⁵, Johannes Kettunen²², Yungil Kim^{23,5}, Holger Kirsten²⁴, Peter Kovacs²⁵, Knut Krohn²⁶, Jaanika Kronberg¹⁵, Viktorija Kukushkina¹⁵, Zoltan Kutalik²⁷, Bennett Lee³, Terho Lehtimäki²⁸, Markus Loeffler²⁴, Urko M. Marigorta^{18,29,30}, Hailang Mei³¹, Lili Milani¹⁵, Grant W. Montgomery³², Martina Müller-Nurasyid^{33,34,35}, Matthias Nauck^{36,37}, Michel G. Nivard³⁸, Brenda Penninx²⁰, Markus Perola³⁹, Natalia Pervjakova¹⁵, Brandon L. Pierce², Joseph Powell⁴⁰, Holger Prokisch^{41,42}, Bruce M. Psaty^{11,43}, Olli T. Raitakari⁴⁴, Samuli Ripatti⁴⁵, Olaf Rotzschke³, Sina Rüeger²⁷, Ashis Saha⁵, Markus Scholz²⁴, Katharina Schramm^{46,34}, Ilkka Seppälä²⁸, Eline P. Slagboom⁴, Coen D.A. Stehouwer⁴⁷, Michael Stumvoll⁴⁸, Patrick Sullivan⁴⁹, Peter A.C. 't Hoen⁵⁰, Alexander Teumer⁵¹, Joachim Thiery⁵², Lin Tong², Anke Tönjes⁴⁸, Jenny van Dongen¹⁰, Maarten van Iterson⁴, Joyce van Meurs⁵³, Jan H. Veldink⁵⁴, Joost Verlouw⁵³, Peter M. Visscher³², Uwe Völker⁵⁵, Urmo Vösa^{8,15}, Harm-Jan Westra^{8,12}, Cisca Wijmenga⁸, Hanieh Yaghootkar^{16,56,57}, Jian Yang^{32,58}, Biao Zeng¹⁸, Futao Zhang³²

Author list ordered alphabetically.

1. Computational Biology, Ontario Institute for Cancer Research, Toronto, Canada
2. Department of Public Health Sciences, University of Chicago, Chicago, United States of America

- 701 3. Singapore Immunology Network, Agency for Science, Technology and Research, Singapore,
702 Singapore
- 703 4. Leiden University Medical Center, Leiden, The Netherlands
- 704 5. Department of Computer Science, Johns Hopkins University, Baltimore, United States of America
- 705 6. Departments of Biomedical Engineering, Johns Hopkins University, Baltimore, United States of
706 America
- 707 7. Heart Center Leipzig, Universität Leipzig, Leipzig, Germany
- 708 8. Department of Genetics, University Medical Centre Groningen, Groningen, The Netherlands
- 709 9. European Molecular Biology Laboratory, Genome Biology Unit, 69117 Heidelberg, Germany
- 710 10. Netherlands Twin Register, Department of Biological Psychology, Vrije Universiteit Amsterdam,
711 Amsterdam Public Health research institute and Amsterdam Neuroscience, the Netherlands
- 712 11. Cardiovascular Health Research Unit, University of Washington, Seattle, United States of America
- 713 12. Onco Institute
- 714 13. Genomics Coordination Center, University Medical Centre Groningen, Groningen, The Netherlands
- 715 14. Department of Genetics, University Medical Centre Utrecht, P.O. Box 85500, 3508 GA, Utrecht,
716 The Netherlands
- 717 15. Estonian Genome Center, Institute of Genomics, University of Tartu, Tartu 51010, Estonia
- 718 16. Genetics of Complex Traits, University of Exeter Medical School, Royal Devon & Exeter Hospital,
719 Exeter, United Kingdom
- 720 17. Department of Medicine, University of Washington, Seattle, United States of America
- 721 18. School of Biological Sciences, Georgia Tech, Atlanta, United States of America
- 722 19. MRC Integrative Epidemiology Unit, University of Bristol, Bristol, United Kingdom
- 723 20. Amsterdam UMC, Vrije Universiteit, Department of Psychiatry, Amsterdam Public Health research
724 institute and Amsterdam Neuroscience, The Netherlands
- 725 21. Department of Clinical Physiology, Tampere University Hospital and Faculty of Medicine and
726 Health Technology, Tampere University, Tampere, Finland
- 727 22. University of Helsinki, Helsinki, Finland
- 728 23. Genetics and Genomic Science Department, Icahn School of Medicine at Mount Sinai, New York,
729 United States of America
- 730 24. Institut für Medizinische Informatik, Statistik und Epidemiologie, LIFE – Leipzig Research Center
731 for Civilization Diseases, Universität Leipzig, Leipzig, Germany
- 732 25. IFB Adiposity Diseases, Universität Leipzig, Leipzig, Germany
- 733 26. Interdisciplinary Center for Clinical Research, Faculty of Medicine, Universität Leipzig, Leipzig,
734 Germany
- 735 27. Lausanne University Hospital, Lausanne, Switzerland
- 736 28. Department of Clinical Chemistry, Fimlab Laboratories and Finnish Cardiovascular Research
737 Center-Tampere, Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland
- 738 29. Integrative Genomics Lab, CIC bioGUNE, Bizkaia Science and Technology Park, Derio, Bizkaia,
739 Basque Country, Spain
- 740 30. IKERBASQUE, Basque Foundation for Science, Bilbao, Spain
- 741 31. Department of Medical Statistics and Bioinformatics, Leiden University Medical Center, Leiden,
742 The Netherlands
- 743 32. Institute for Molecular Bioscience, University of Queensland, Brisbane, Australia
- 744 33. Institute of Genetic Epidemiology, Helmholtz Zentrum München - German Research Center for
745 Environmental Health, Neuherberg, Germany
- 746 34. Department of Medicine I, University Hospital Munich, Ludwig Maximilian's University, München,
747 Germany
- 748 35. DZHK (German Centre for Cardiovascular Research), partner site Munich Heart Alliance, Munich,
749 Germany
- 750 36. Institute of Clinical Chemistry and Laboratory Medicine, Greifswald University Hospital,
751 Greifswald, Germany

37. German Center for Cardiovascular Research (partner site Greifswald), Greifswald, Germany
38. Department of Biological Psychology, Faculty of Behaviour and Movement Sciences, VU, Amsterdam, The Netherlands
39. National Institute for Health and Welfare, University of Helsinki, Helsinki, Finland
40. Garvan Institute of Medical Research, Garvan-Weizmann Centre for Cellular Genomics, Sydney, Australia
41. Institute of Human Genetics, Helmholtz Zentrum München, Neuherberg, Germany
42. Institute of Human Genetics, Technical University Munich, Munich, Germany
43. Kaiser Permanente Washington Health Research Institute, Seattle, WA, United States of America
44. Centre for Population Health Research, Department of Clinical Physiology and Nuclear Medicine, Turku University Hospital and University of Turku, Turku, Finland
45. Statistical and Translational Genetics, University of Helsinki, Helsinki, Finland
46. Institute of Genetic Epidemiology, Helmholtz Zentrum München - German Research Center for Environmental Health, Neuherberg, Germany
47. Department of Internal Medicine and School for Cardiovascular Diseases (CARIM), Maastricht University Medical Center, Maastricht, The Netherlands
48. Department of Medicine, Universität Leipzig, Leipzig, Germany
49. Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden
50. Center for Molecular and Biomolecular Informatics, Radboud Institute for Molecular Life Sciences, Radboud University Medical Center Nijmegen, Nijmegen, The Netherlands
51. Institute of Clinical Chemistry and Laboratory Medicine, University Medicine Greifswald, Greifswald, Germany
52. Institute for Laboratory Medicine, LIFE – Leipzig Research Center for Civilization Diseases, Universität Leipzig, Leipzig, Germany
53. Department of Internal Medicine, Erasmus Medical Centre, Rotterdam, The Netherlands
54. UMC Utrecht Brain Center, University Medical Center Utrecht, Department of Neurology, Utrecht University, Utrecht, The Netherlands
55. Interfaculty Institute for Genetics and Functional Genomics, University Medicine Greifswald, Greifswald, Germany
56. School of Life Sciences, College of Liberal Arts and Science, University of Westminster, 115 New Cavendish Street, London, United Kingdom
57. Division of Medical Sciences, Department of Health Sciences, Luleå University of Technology, Luleå, Sweden
58. Institute for Advanced Research, Wenzhou Medical University, Wenzhou, Zhejiang 325027, China

BIOS Consortium (Biobank-based Integrative Omics Study) – Author information

Management Team Bastiaan T. Heijmans (chair)¹, Peter A.C. 't Hoen², Joyce van Meurs³, Aaron Isaacs⁴, Rick Jansen⁵, Lude Franke⁶.

Cohort collection Dorret I. Boomsma⁷, René Pool⁷, Jenny van Dongen⁷, Jouke J. Hottenga⁷ (Netherlands Twin Register); Marleen MJ van Greevenbroek⁸, Coen D.A. Stehouwer⁸, Carla J.H. van der Kallen⁸, Casper G. Schalkwijk⁸ (Cohort study on Diabetes and Atherosclerosis Maastricht); Cisca Wijmenga⁶, Lude Franke⁶, Sasha Zhernakova⁶, Ettje F. Tigchelaar⁶ (LifeLines Deep); P. Eline Slagboom¹, Marian Beekman¹, Joris Deelen¹, Diana van Heemst⁹ (Leiden Longevity Study); Jan H. Veldink¹⁰, Leonard H. van den Berg¹⁰ (Prospective ALS Study Netherlands); Cornelia M. van Duijn⁴, Bert A. Hofman¹¹, Aaron Isaacs⁴, André G. Uitterlinden³ (Rotterdam Study).

Data Generation Joyce van Meurs (Chair)³, P. Mila Jhamai³, Michael Verbiest³, H. Eka D. Suchiman¹, Marijn Verkerk³, Ruud van der Breggen¹, Jeroen van Rooij³, Nico Lakenberg¹.

Data management and computational infrastructure Hailiang Mei (Chair)¹², Maarten van Iterson¹, Michiel van Galen², Jan Bot¹³, Dasha V. Zhernakova⁶, Rick Jansen⁵, Peter van 't Hof¹², Patrick Deelen⁶, Irene Nooren¹³, Peter A.C. 't Hoen², Bastiaan T. Heijmans¹, Matthijs Moed¹.

Data Analysis Group Lude Franke (Co-Chair)⁶, Martijn Vermaat², Dasha V. Zhernakova⁶, René Luijk¹, Marc Jan Bonder⁶, Maarten van Iterson¹, Patrick Deelen⁶, Freerk van Dijk¹⁴, Michiel van Galen², Wibowo Arindrarto¹², Szymon M. Kielbasa¹⁵, Morris A. Swertz¹⁴, Erik. W. van Zwet¹⁵, Rick Jansen⁵, Peter-Bram 't Hoen (Co-Chair)², Bastiaan T. Heijmans (Co-Chair)¹.

1. Molecular Epidemiology Section, Department of Medical Statistics and Bioinformatics, Leiden University Medical Center, Leiden, The Netherlands
2. Department of Human Genetics, Leiden University Medical Center, Leiden, The Netherlands
3. Department of Internal Medicine, ErasmusMC, Rotterdam, The Netherlands
4. Department of Genetic Epidemiology, ErasmusMC, Rotterdam, The Netherlands
5. Department of Psychiatry, VU University Medical Center, Neuroscience Campus Amsterdam, Amsterdam, The Netherlands
6. Department of Genetics, University of Groningen, University Medical Centre Groningen, Groningen, The Netherlands
7. Department of Biological Psychology, VU University Amsterdam, Neuroscience Campus Amsterdam, Amsterdam, The Netherlands
8. Department of Internal Medicine and School for Cardiovascular Diseases (CARIM), Maastricht University Medical Center, Maastricht, The Netherlands
9. Department of Gerontology and Geriatrics, Leiden University Medical Center, Leiden, The Netherlands
10. Department of Neurology, Brain Center Rudolf Magnus, University Medical Center Utrecht, Utrecht, The Netherlands
11. Department of Epidemiology, ErasmusMC, Rotterdam, The Netherlands
12. Sequence Analysis Support Core, Leiden University Medical Center, Leiden, The Netherlands
13. SURFsara, Amsterdam, The Netherlands
14. Genomics Coordination Center, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands
15. Medical Statistics Section, Department of Medical Statistics and Bioinformatics, Leiden University Medical Center, Leiden, The Netherlands

Online Methods

This article has a **Supplementary Note** with more details.

Samples. For Age at First Sexual intercourse (AFS), we included 397,338 pooled individuals (n=182,791 males; n=214,547 females) from the UK Biobank. For Age at First Birth (AFB), we included 542,901 individuals (n=124,088 males; n=418,758 females) from 36 studies. We performed a GWAS separately restricted to European ancestry individuals that passed quality control. European ancestry was chosen in this discovery study due to the availability of samples¹ and for no biological or substantive reason. We acknowledge that social science research has found large differences in the earlier initiation of AFS and AFB by lower socioeconomic status, which often coincides with societal inequality^{2,3} and the socially (not biologically) constructed categories of race and ethnicity. Socioeconomic differences are examined in this article, but results are only applicable to European Ancestry groups, with a need for further cross-ancestry discovery research.

The Human Reproductive Behaviour Consortium. This consortium is a collaboration studying the GWAS of human reproductive behaviour including age at first sex and birth, number of children ever born, childlessness and related traits. In some cases we used summary statistics from our first GWAS of AFB and NEB⁴ on discovery cohorts (see Supp Note Tables S1-S3b).

Phenotype measurements, genotyping, imputation and meta-analysis. AFS is treated as a continuous measure with individuals considered as eligible if they had given a valid answer and ages lower than 12 excluded (see Supp Note 1.2). Since AFS has a non-normal distribution, a within-sex inverse rank normal transformation is required. AFB is also treated as a continuous measure, assessed for those who have ever given birth to a child. Details about participating cohorts, sample inclusion criteria, genotyping and imputation, models used to test for association, X chromosome analysis, quality control filters and diagnostics, and meta-analysis are in the Supp Note. A sample-size weighted meta-analysis of quality-controlled cohort-level results was performed using the METAL software.⁵ We performed conditional and joint multiple SNP analyses (COJO) to identify further independent SNPs and sex-specific analyses.

Sex-specific genetic effects. We used LD score bivariate regression⁶ to estimate the genetic correlation between men and women based on the sex-specific summary statistics from the meta-analysis results. There was a large genetic overlap among the sexes for AFB (0.95) and a somewhat lower overlap for AFS (0.79), suggesting sex-specific effects would be important to examine. In order to determine if there was evidence for sex-specific effects, we compared the allelic effects for these SNPs between men and women and derived a *p*-value for heterogeneity.⁷ A multiple testing correction was applied ($0.05/242=2 \times 10^{-4}$) to identify sex-specific associations. We then selected a region of $\pm 1\text{Mb}$ around these lead SNPs to

identify the genes that may be represented by these lead SNPs, followed by gene prioritization as we did for the main AFB and AFS analyses.

X chromosome analysis. For AFS, the UK Biobank provided results for between 977,536 and 990,735 variants on the X chromosome after QC (see Table S8). For AFB, 13 cohorts provided information on the X chromosome. Overall, we received 23 files, 13 for women, 8 for men and 2 for the pooled analysis in case there were individuals who were relatives in the data. On average, 275,023 variants survived QC with a minimum of 99,794 in women from WLS to 998,304 for the women in the UK Biobank sample (see TableS7 for full descriptives). Association analyses on the X chromosome were performed using software suggested in the analysis plan (XWAS, SNPtest or BOLT-LMM) using BOLT-LMM for AFS as this was only assessed in the UK Biobank data, for AFB, METAL was used as described above (see sup. note 3.5)

Phenotypic and genotypic historical changes. Descriptive analyses and correlations were undertaken using the UK Biobank data to illustrate phenotypic shifts in the age of AFS and AFB by birth cohort, in addition to changes in the spread of the distribution. Pearson's correlation coefficients were calculated and correlation graphs illustrate the changing relationship between the two phenotypes over time. Genotypic changes and SNP-heritability by birth cohort were quantified in UK Biobank data using GREML⁸ as described earlier.⁹

MTAG. MTAG results¹⁰ were calculated using GWA meta-analysis results of the following related phenotypes: AFS, AFB, number of children ever born, childlessness. Using summary statistics from the pooled GWAS of each of the traits, MTAG uses bivariate LD score regression to account for unobserved sample overlap.

Polygenic score prediction. We performed out-of-sample prediction in two cohorts, the National Longitudinal Study of Adolescence to Adult Health (Add Health),¹¹ based in the US and the UK Household Longitudinal Study - Understanding Society (UKHLS).¹² We calculated three sets of polygenic risk scores (PGS) with weights based on meta-analysis results excluding the specific cohort from the calculation. First, pruning and thresholding of all SNPs was performed (250kb window; $r^2=.1$) using PRSice¹³. Second, LDpred PGSs¹⁴ with the LD reference were calculated from the same genotyped files, using prior distributions for the causal fraction of SNPs equal to 1 and LDpred weights calculated under the infinitesimal model. Third, MTAG + LDpred PGSs were calculated using the same methodology as in the second PGSs, but this time based on MTAG results¹⁰. For both traits, we ran ordinary least-squares (OLS) regression models and report the incremental R^2 as a measure of goodness-of-fit of the model. Confidence intervals are based on 1,000 bootstrapped samples.

Testing population stratification, survival models and environmentally mediated parental genetic effects of childhood socioeconomic status. To test whether population stratification biased our results or lead to false positives, we used the LD Score intercept

method.¹⁵ For each phenotype, we used the “eur_w_ld_chr” files of LD Scores.¹⁶ These LD Scores were computed with genotypes from the European-ancestry samples in the 1000 Genomes Project using only HapMap3 SNPs with MAF > 0.01. We then ran survival models to account for right-censoring, which occurs when an individual does not experience the event of first sex or birth by the time of the interview.¹⁷ Using Add Health data, we estimated nonparametric hazard functions based on Nelson-Aalen estimates and then compared individuals at the top and bottom 5% of the PGS and plotted the estimated hazards. To further explore the impact of environmentally mediated parental genetic effects on our PGSs, we examined PGS prediction across low (0-10%), medium (50-60%) and high (90-100%) PGS percentiles by parent’s education status (college versus no college), which serves as a proxy for childhood socioeconomic status.

Genetic correlations. Genetic correlation (r_g) values were computed to estimate the genetic correlation between the two traits using all polygenic effects captured by the SNPs and LD-score regression.¹⁸ We used summary statistics and the 1000 Genomes reference set, and restricted the analysis to European populations. We also followed the common convention of restricting our analyses to SNPs with MAF > 0.01, thus ensuring that all analyses were performed using a set of SNPs that were imputed with reasonable accuracy across all cohorts. The standard errors (SEs) were produced by the LDSC python software package that uses a block jackknife over the SNPs. We estimated the genetic correlation between 28 different traits, pooled by both sexes and then divided by sex. Traits were divided into the six categories of: reproductive, behavioural, psychiatric disorders, substance use disorders, personality and anthropometric.

Genomic SEM (structural equation modelling). In an attempt to understand the aetiology of the correlations, we used the R package GenomicSEM to fit genetic multivariable regression models. GenomicSEM¹⁹ uses structural equation modelling to decompose the genetic covariance matrix, calculated using multivariate LD score regression, of a set of traits. Formally, structural equation models subsume many statistical methods and are quite flexible. We fit a series of genetic multivariable regression models, in which AFB was regressed on EA (educational attainment) and a trait X, in which we modelled various relevant traits such as openness, cognitive performance and AI (age initiation smoking). We also fit an analogous series of models in which AFS was regressed on EA.

Exploratory factor analysis (EFA) and Genomic SEM by reproductive biology and externalizing behaviour. EFA was used to examine whether the genetic signal of the onset of reproductive behaviour originated from two genetically distinguishable sub-clusters of a biological component and an externalizing behaviour component. This would suggest distinct causal mechanisms and subtypes of individuals. We tested this by fitting a two factor EFA model to the genetic covariance matrix of AFB, AFS, NEB, and the proxies age at menarche (biological component) and risk tolerance (externalizing behaviour). To test this further, we estimated a more robust and additional measures of reproductive biology and

externalizing behaviour and a sex-specific analysis of AFB for women. We fit a genomic structural equation model (Genomic SEM) where AFB in women is regressed on age at menopause, age at menarche, and a latent factor representing the common genetic tendency to externalizing behaviour. The factor is measured by AFS in women, age at initiation of smoking, age first used oral contraception, and ADHD, with the model scaled to unit variance for the latent factor.

Bi-directional Mendelian Randomization. We then tested whether causal pathways linking these phenotypes are potentially bidirectional and whether our phenotypes might offer distinct contributions. We identified 1000 Genomes proxies for our SNPs and used these in multivariate Mendelian Randomisation (MR) models. First, we modelled the interplay between AFB, AFS and EA (educational attainment)²⁰ as well as risk taking (measured in adulthood)²¹ and age at smoking initiation (AI).²² In each case IVW²³ and MR-EGGER²⁴ methods were performed, with an additional round of IVW performed once a Steiger filter²⁵ had been applied to remove SNPs that appears to show a primary association with the outcome rather than the exposure. Multivariate MR was used to try to dissect causal pathways.²⁶ A second set of MR analyses focused on links to late life diseases, namely type 2 diabetes (T2D)²⁷ and coronary artery disease (CAD)²⁸, using the same methods. In particular, we use multivariate methods to test whether AFS or AFB had independent effects once the well-established links to length of educational attainment were controlled for.

Cox proportional hazard models of AFB polygenic score on longevity. To test trade-offs between reproductive behaviour and senescence, we conducted additional analyses to test whether our PGS for AFB was predictive of (parental) longevity. We restricted our models to mortality after age 60 to limit the possibility that early mortality affects parental fertility (i.e., collider bias).²⁹ We calculated PGSs for AFB, Educational attainment (EA)³⁰ and risky behaviour²¹ from the UK Biobank adopting the following procedure. We first split the sample in 10 random groups. We then iteratively estimated genome-wide association results for 9/10th of the sample and used these association results as weights for the calculation of polygenic scores in the remaining 1/10th of the sample. Polygenic scores were calculated using PRSice on a set of independent genotyped SNPs. We then estimated three sets of Cox Proportional hazard models to estimate the effect of the PGS of AFB on maternal and paternal age at death. All models control for the first 10 Genetic Principal Components, sex and year of birth, and are stratified by Local Authority District at birth calculated using the geo-coordinates provided in the UK Biobank due to differences in mortality related to material deprivation.³¹ We first estimated a baseline model and then included PGSs for EA and risk as covariates, followed by a final model including number of sibling (proxy for parental fertility).

Gene prioritization. We prioritized candidate genes in pooled and sex-specific GWAS-identified loci using predicted gene functions,³² single-cell RNA sequencing data in mice,^{33,34}

literature mining,³⁵ *in silico* sequencing,³⁶ and synthetic Mendelian Randomization³⁷ using eQTL data from brain and blood.^{38,39}

DEPICT, RNAseq data from mouse brain and Tabula muris RNAseq data. First, DEPICT was used to perform pathway analyses, identify enrichment for cell types and tissues, and prioritize candidate genes.³² DEPICT is agnostic to the outcomes analyzed in the GWAS and employs predicted gene functions. For both AFS and AFB, all SNPs with $p < 1 \times 10^{-5}$ in the pooled GWAS meta-analysis were used as input. Based on the results of the tissue enrichment analysis, we used DEPICT to identify nervous system cell types that are enriched for expression of genes in loci reaching $p < 1 \times 10^{-5}$ in the GWAS, using RNAseq data from mouse brain.³³ A similar approach using tabula muris RNAseq data³⁴ helped prioritize additional central nervous system and pancreatic cell types for AFS. For enriched cell types from mouse brain and tabula muris, the top-10 contributing genes were selected as candidate genes resulting in the prioritization of 296 genes for AFS and 95 for AFB based on mouse brain; and 97 genes for AFS based on tabula muris data.

Phenolyzer to integrate prior knowledge and phenotype information. We used Phenolyzer (v1.1) to prioritize candidate genes by integrating prior knowledge and phenotype information.⁴⁰ Here we used the regions defined by DEPICT v1.1, reflecting loci reaching $P < 1 \times 10^{-5}$ in first instance. Phenolyzer takes free text input and interprets these as disease names by using a word cloud to identify synonyms. It then queries precompiled databases for the disease names to find and score relevant seed genes. The seed genes are subsequently expanded to include related (predicted) genes based on several types of relationships, e.g., protein-protein interactions, transcriptional regulation and biological pathways. Phenolyzer uses machine-learning techniques on seed genes and predicted gene rankings to produce an integrated score for each gene. We used search terms capturing three broad areas, i.e., (in)fertility, congenital neurological disorders and psychological traits, based on results from pathway, tissue and cell type enrichment analyses.

***In silico* sequencing.** We used *in silico* sequencing to identify non-synonymous variants with an R^2 for LD > 0.7 with the lead SNPs in AFS and AFB-associated loci,³⁶ which yielded genes that may drive the GWAS associations through direct effects on protein function.

Summary data-based Mendelian Randomization (SMR) and Heterogeneity in Dependent Instruments (HEIDI).³⁷ We conducted this using eQTL data from brain⁴¹ and whole blood.³⁹ This approach provided a list of genes that showed Bonferroni corrected significant evidence (thresholds for blood $< 3.2 \times 10^{-6}$ brain $< 6.7 \times 10^{-6}$) of mediating the association between our phenotypes and GWAS-identified loci based on results from brain and blood.

Integration of findings across all functional approaches. We integrated findings across all approaches and retained genes in loci that reached genome-wide significance, and that were located within 1M bp of a GWAS lead SNP. We next used data from the Human Protein

Atlas⁴² to identify genes amongst 387 genes that are expressed at a low, medium or high protein level in brain, glands, and/or reproductive organs at a ‘supported’ or ‘enhanced’ degree of reliability. For the 97 genes that fulfilled these criteria, we mapped the brain, glandular and reproductive cell types in which they are highly expressed at the protein level;⁴³ used a text-mining approach to extract functions from entries in Entrez, GeneCards and Uniprot; and identified phenotypes in mutant mice from the Mouse Genome Informatics (MGI) database⁴⁴.

Ethics statement. All research was approved by the appropriate institutional review boards and participants of all studies provided informed consent to participate in those studies.

Data availability. Our policy is to make genome-wide summary statistics widely and publically available. Upon publication, summary statistics will be available on the GWAS Catalog website: <https://www.ebi.ac.uk/gwas/downloads/summary-statistics>

The phenotype and genotype data are available upon application from each of the participating cohorts and should be contacted directly regarding their different data access policies. Access to the UK Biobank is available through application with information available at: <http://www.ukbiobank.ac.uk>).

References

1. Mills, M. C. & Rahal, C. The GWAS Diversity Monitor tracks diversity by disease in real time. *Nat. Genet.* **52**, 242–243 (2020).
2. Chen, X.-K. *et al.* Teenage pregnancy and adverse birth outcomes: a large population based retrospective cohort study. *Int. J. Epidemiol.* **36**, 368–373 (2007).
3. Bongaarts, J., Mensch, B. S. & Blanc, A. K. Trends in the age at reproductive transitions in the developing world: The role of education. *Popul. Stud. (NY)*. **71**, 139–154 (2017).
4. Barban, N. *et al.* Genome-wide analysis identifies 12 loci influencing human reproductive behavior. *Nat. Genet.* **48**, (2016).
5. Willer, C. J., Li, Y. & Abecasis, G. R. METAL: Fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* **26**, 2190–2191 (2010).
6. Finucane, H. K. *et al.* Partitioning heritability by functional category using GWAS summary statistics. *bioRxiv*. (2015).
7. Altman, D. G. & Bland, J. M. Interaction revisited: the difference between two estimates. *BMJ* **326**, 219 (2003).
8. Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: a tool for genome-wide complex trait analysis. *Am. J. Hum. Genet.* **88**, 76–82 (2011).
9. Tropf, F. C. *et al.* Hidden heritability due to heterogeneity across seven populations. *Nat. Hum. Behav.* **1**, 757–765 (2017).
10. Turley, P. *et al.* Multi-trait analysis of genome-wide association summary statistics using MTAG. *Nat. Genet.* **50**, 229–237 (2018).
11. Harris, K. M. & *et al.* The National Longitudinal Study of Adolescent to Adult Health: Research Design. (2009). Available at: <http://www.cpc.unc.edu/projects/addhealth/design>.

12. Buck, N. & McFall, S. Understanding Society: design overview. *Longit. Life Course Stud.* **3**, 5–17 (2012).
13. Euesden, J., Lewis, C. M. & O'Reilly, P. F. PRSice: Polygenic Risk Score software. *Bioinformatics* **31**, btu848-1468 (2014).
14. Vilhjálmsson, B. J. *et al.* Modeling Linkage Disequilibrium Increases Accuracy of Polygenic Risk Scores. *Am. J. Hum. Genet.* **97**, 576–592 (2015).
15. Bulik-Sullivan, B. *et al.* An Atlas of Genetic Correlations across Human Diseases and Traits. *bioRxiv* **47**, 1–44 (2015).
16. Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide association summary statistics. *Nat. Genet.* **47**, 1228–1235 (2015).
17. Mills, M. C. *Introducing survival and event history analysis.* (Sage Publications, 2011).
18. Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat. Genet.* **47**, 291–295 (2015).
19. Grotzinger, A. D. *et al.* Genomic structural equation modelling provides insights into the multivariate genetic architecture of complex traits. *Nat. Hum. Behav.* **3**, 513–525 (2019).
20. Lee, J. J. *et al.* Gene discovery and polygenic prediction from a genome-wide association study of educational attainment in 1.1 million individuals. *Nat. Genet.* (2018). doi:10.1038/s41588-018-0147-3
21. Karlsson Linnér, R. *et al.* Genome-wide association analyses of risk tolerance and risky behaviors in over 1 million individuals identify hundreds of loci and shared genetic influences. *Nat. Genet.* **51**, 245–257 (2019).
22. Liu, M. *et al.* Association studies of up to 1.2 million individuals yield new insights into the genetic etiology of tobacco and alcohol use. *Nat. Genet.* **51**, 237–244 (2019).
23. Burgess, S., Butterworth, A. & Thompson, S. G. Mendelian randomization analysis with multiple genetic variants using summarized data. *Genet. Epidemiol.* **37**, 658–65 (2013).
24. Bowden, J., Smith, G. D. & Burgess, S. Mendelian randomization with invalid instruments: Effect estimation and bias detection through Egger regression. *Int. J. Epidemiol.* (2015). doi:10.1093/ije/dyv080
25. Hemani, G., Bowden, J. & Davey Smith, G. Evaluating the potential role of pleiotropy in Mendelian randomization studies. *Hum. Mol. Genet.* **27**, R195–R208 (2018).
26. Burgess, S. & Thompson, S. G. Multivariable Mendelian randomization: the use of pleiotropic genetic variants to estimate causal effects. *Am. J. Epidemiol.* **181**, 251–60 (2015).
27. Mahajan, A. *et al.* Fine-mapping type 2 diabetes loci to single-variant resolution using high-density imputation and islet-specific epigenome maps. *Nat. Genet.* **50**, 1505–1513 (2018).
28. Nelson, C. P. *et al.* Association analyses based on false discovery rate implicate new loci for coronary artery disease. *Nat. Genet.* **49**, 1385–1391 (2017).
29. Day, F. R., Loh, P.-R., Scott, R. A., Ong, K. K. & Perry, J. R. B. A Robust Example of Collider Bias in a Genetic Association Study. *Am. J. Hum. Genet.* **98**, 392–393 (2016).
30. Lee, J. J. *et al.* Gene discovery and polygenic prediction from a genome-wide association study of educational attainment in 1.1 million individuals. *Nat. Genet.* **50**, 1112–1121 (2018).
31. Woods, L. M. Geographical variation in life expectancy at birth in England and Wales is largely explained by deprivation. *J. Epidemiol. Community Heal.* **59**, 115–120 (2005).

32. Pers, T. H. *et al.* Biological interpretation of genome-wide association studies using predicted gene functions. *Nat. Commun.* **6**, 5890 (2015).
33. Zeisel, A. *et al.* Molecular Architecture of the Mouse Nervous System. *Cell* **174**, 999–1014.e22 (2018).
34. Tabula Muris Consortium *et al.* Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris. *Nature* **562**, 367–372 (2018).
35. Yang, H., Robinson, P. N. & Wang, K. Phenolyzer: phenotype-based prioritization of candidate genes for human diseases. *Nat. Methods* **12**, 841–3 (2015).
36. Vaez, A. *et al.* In Silico Post Genome-Wide Association Studies Analysis of C-Reactive Protein Loci Suggests an Important Role for Interferons. *Circ. Cardiovasc. Genet.* **8**, 487–497 (2015).
37. Zhu, Z. *et al.* Integration of summary data from GWAS and eQTL studies predicts complex trait gene targets. *Nat. Genet.* **48**, 481–7 (2016).
38. van der Wijst, M. G. P. *et al.* Single-cell RNA sequencing identifies celltype-specific cis-eQTLs and co-expression QTLs. *Nat. Genet.* **50**, 493–497 (2018).
39. Vosa, U. & Al., E. Unraveling the polygenic architecture of complex traits using blood eQTL metaanalysis. *bioRxiv* (2018). doi:<https://doi.org/10.1101/447367>
40. Yang, H., Robinson, P. N. & Wang, K. Phenolyzer: phenotype-based prioritization of candidate genes for human diseases. *Nat. Methods* **12**, 841–3 (2015).
41. Qi, T. *et al.* Identifying gene targets for brain-related traits using transcriptomic and methylomic data from blood. *Nat. Commun.* **9**, 2282 (2018).
42. Uhlén, M. *et al.* Proteomics. Tissue-based map of the human proteome. *Science* **347**, 1260419 (2015).
43. Uhlén, M. *et al.* Proteomics. Tissue-based map of the human proteome. *Science* **347**, 1260419 (2015).
44. Bult, C. J. *et al.* Mouse Genome Database (MGD) 2019. *Nucleic Acids Res.* **47**, D801–D806 (2019).