

1 **Title: ROS signaling in cotton ovule epidermal cells is necessary for fiber initiation**

2 **Authors:** Mingxiong Pang<sup>1\*</sup>, Nicholas Sanford<sup>1</sup>, Thea A Wilkins<sup>1</sup>

3 **Affiliations:**

4 <sup>1</sup>Department of Plant & Soil Science, Texas Tech University, Lubbock, TX 79409 USA.

5

6 \*Correspondences to: Mingxiong Pang. [mingxiong.pang@ttu.edu](mailto:mingxiong.pang@ttu.edu).

7

8 **Abstract:** Cotton (*Gossypium hirsutum*) fiber, an extremely elongated and thickened single cell  
9 of the seed epidermis, is the world's most important natural and economical textile fiber. Unlike  
10 *Arabidopsis* leaf trichomes, fiber initials are randomly developed and frequently form in adjacent  
11 seed epidermal cells and follow no apparent pattern. Numerous publications suggested cotton  
12 fiber development shares a similar mechanism with *Arabidopsis* leaf trichome development.  
13 Here we show that H<sub>2</sub>O<sub>2</sub> accumulation in cotton ovule epidermal cells by NBT staining ovules  
14 at different development stages between TM1 and N1n2, a lintless-fuzzless doubled mutant  
15 originated from TM1. In contrast, *Arabidopsis* and cotton leaf trichomes do not show H<sub>2</sub>O<sub>2</sub>  
16 content. By adding DPI (H<sub>2</sub>O<sub>2</sub> inhibitor) and SHAM (H<sub>2</sub>O<sub>2</sub> activator) in vitro ovule cultures,  
17 we show fiber initiation directly involves with H<sub>2</sub>O<sub>2</sub> accumulation. We propose that the  
18 directional accumulation of H<sub>2</sub>O<sub>2</sub> in cotton ovule epidermal cell is the drive for fiber initiation,  
19 elongation.

20

21

1 **Main Text:**

2 Seed dispersal is sometimes split into autochory (when dispersal is attained using the  
3 plant's own means) and allochory (when obtained through external means). Wind dispersal  
4 (anemochory) is one of the autochories and more primitive means of dispersal and plant seeds  
5 have a feathery tissue attached to their seeds and can be dispersed long distances. Primitively,  
6 plants distributed the seeds majorly through autochory; as animals emerge and human activity,  
7 some plants lost their ability to produce seed secondary tissues helpful for dispersal and some  
8 plants gained the production for specific seeds tissues such as cotton seed trichomes as human  
9 activity. But we do not understand how these tissues develop. Cotton (*Gossypium spp.*) fibers are  
10 unicellular and single-celled seed trichomes of economic importance (1) and provide a special  
11 single-cell functional genomics research platform (2, 3) for fundamental biological processes  
12 such as organogenesis and cell wall development in plants (4, 5). Fiber development initiates on  
13 the day of anthesis (0 dpa) and 30-40% of primordial cells (6) on the ovule epidermis form  
14 cuboidal outgrowths called fiber "initials" (7). Unlike *Arabidopsis* leaf trichomes, fiber initials is  
15 randomly distributed and follow no apparent pattern as fiber initials frequently form in adjacent  
16 epidermal cells (8,9). Fiber initials rapidly elongate to produce highly tapered cells within 2-3  
17 days of anthesis through a biased diffuse growth mechanism (10).

18 The initiation of single celled cotton cuboidal fiber cells is a biologically important and  
19 complex process, but how the fiber initiate and how the genes involved are regulated remain  
20 elusive. Zhang et al., (11) demonstrated by adding 30% H<sub>2</sub>O<sub>2</sub> to in vitro cultured ovules of  
21 XinFLM, a linted-fuzzless mutant, that the fiber initiation is partially rescued, but no evident  
22 effect on N1, the dominant naked seed mutant from TM-1 genetic background. This implies the  
23 H<sub>2</sub>O<sub>2</sub> biosynthesis pathway may be defect and the signal transduction is still functional in

1 XinFLM. Like cuboidal outgrowth fiber initials, Andriunas et al., (12) shows a flavin-containing  
2 enzyme (NAPDH oxidase) and superoxide dismutase cooperatively generate a regulatory H<sub>2</sub>O<sub>2</sub>  
3 to govern ingrowth wall formation, a transfer cell (13), using *Visia Faba* cotyledon culture.  
4 Forman et al., (14) verified that ROS accumulates in growing wild type root hairs but  
5 dramatically decreases in *rhd2* short hair mutants and the generation of ROS is inhibited and  
6 mimics *rhd2* mutant and further showed (15) a RhoGTPase GDP dissociation inhibitor controls  
7 spatial ROS production during root hair development.

8         To determine whether H<sub>2</sub>O<sub>2</sub> is necessary for cotton fiber initiation, we firstly used SEM  
9 to observe fiber initiation and development dynamics starting ovules at 2 days of pre-anthesis to  
10 2 days of post-anthesis. In the dominant N1 mutant (Fig. 1), no fiber initials are present on ovule  
11 at anthesis and very rare fiber initials occur on the ovules at 2 days of post anthesis. The  
12 recessive mutant n2 does not show significant change for fiber initials. The double mutants N1n2  
13 generated by crossing N1 and n2 and selecting till 7 generation does not show any fiber  
14 initiation. XZ142-fl, a lintless and fuzzless mutant originated from its wild type XZ142, also  
15 does not produce any fibers.

16         In higher plants ROS can probably be produced by several alternative pathways (16)  
17 including a cell-wall-localized peroxidase, amine oxidases and non-flavin NADPH oxidases, or  
18 NADPH oxidases that resemble the flavin-containing type that is activated by Rac in leukocytes  
19 (17). Simultaneously, a vast network of antioxidants is acting like ROS scavengers of H<sub>2</sub>O<sub>2</sub> and  
20 maintains ROS homeostasis (18). This scavenging system consists of catalase (CAT), ascorbate  
21 (APX) and secretory peroxidases (POX), glutathione reductases (GR) and peroxiredoxines  
22 (PRX), and non-enzymatic compounds like tocopherols, ascorbic acid and flavonoids. (19). Li et  
23 al., (20) identified a cotton cytosolic APX1 (*GhAPX1*) to be highly accumulated during cotton

1 fiber elongation by proteomic analysis and showed (21) ethylene is induced in in vitro cultured  
2 ovules by endogenous H<sub>2</sub>O<sub>2</sub> and found a feedback regulatory system between H<sub>2</sub>O<sub>2</sub> and  
3 ethylene in fiber development. Hovav et al., (22) found three genes modulating hydrogen  
4 peroxide levels were consistently expressed in domesticated and wild cotton species with long  
5 fibers, but not detected by q-PCR in wild species with short fibers. Potika et al., (23) Potikha et  
6 al., 1999) speculated H<sub>2</sub>O<sub>2</sub> may function as a developmental signal in the differentiation of  
7 secondary walls in cotton fibers with support of coincidence of H<sub>2</sub>O<sub>2</sub> generation with the onset  
8 of secondary wall deposition and prevention the wall differentiation because of inhibition of  
9 H<sub>2</sub>O<sub>2</sub> production or removing the available H<sub>2</sub>O<sub>2</sub>. H<sub>2</sub>O<sub>2</sub> modulates downstream signaling  
10 events including calcium mobilization protein phosphorylation and gene expression (24) which  
11 may lead to cotton epidermal cell fate determination. To determine whether H<sub>2</sub>O<sub>2</sub> involves fiber  
12 initiation, we used 0.1mg/mL NBT (Nitro Blue Tetrazolium), a chemical specifically staining  
13 H<sub>2</sub>O<sub>2</sub> in living cells (Fig.2). Fig.2A timely profiled a normal cotton fiber development starting 5  
14 days of pre-anthesis to fiber maturation at 50 days of post-anthesis. Fig.2B demonstrated that  
15 TM1 ovules stained darkly at 2 days of pre-anthesis to 2 days of post-anthesis has already  
16 accumulated H<sub>2</sub>O<sub>2</sub> in comparison to these from N1n2. Significantly, in contrast to TM1 ovules,  
17 no any staining was detected on *Arabidopsis* and Cotton leaf trichomes (Fig.2C). We also used  
18 cut cotton leaf edge as positive, dark staining was found at the cut leaf edge and spots with high  
19 concentration of polyphenols (Fig.2C). To rule out the possibility of staining accessibility, we  
20 conducted H<sub>2</sub>O<sub>2</sub> production measurement using an Amplex red hydrogen peroxide/peroxidase  
21 assay kit (Molecular Probes, <http://probes.invitrogen.com/>). We found significant decrease of  
22 H<sub>2</sub>O<sub>2</sub> in ovules of different stages of N1n2 comparing with those from TM, especially at the 2  
23 DPA (Fig.2D).

1           To verify whether H<sub>2</sub>O<sub>2</sub> directly influence epidermal cell differentiation and initiation,  
2 we conducted in vitro ovule cultures by adding DPI (H<sub>2</sub>O<sub>2</sub> inhibitor) and SHAM (H<sub>2</sub>O<sub>2</sub>  
3 activator. The result (Fig.3A) shows the fiber initiation normally developed in the control (with  
4 0.1% DMSO) in vitro ovule culture and NTB staining shows H<sub>2</sub>O<sub>2</sub> accumulation in the fiber  
5 initiation and epidermal cells. Fig.3B shows the fiber initiation was almost totally retarded in the  
6 in vitro ovule culture with addition of 0.3 μM DPI and the NTB staining shows barely H<sub>2</sub>O<sub>2</sub>  
7 accumulation. Interestingly, Fig.3C shows very similar results with Fig.3A. We cannot conclude  
8 SHAM enhances fiber initiation as a more controlled experiment is needed for this result. Based  
9 on this data, we can efficiently conclude that H<sub>2</sub>O<sub>2</sub> directly involves cotton fiber initiation and  
10 development.

11           To molecularly characterize the mechanism of fiber initiation, many researchers (25, 26)  
12 proposed the mechanism similarity of cotton fiber development to *Arabidopsis* trichome and  
13 study cotton fiber initiation by cloning cotton orthologs for *Arabidopsis* trichome development.  
14 Three criteria were used to characterize whether the genes of interest relate to cotton fiber  
15 initiation including (i) orthologs to *Arabidopsis* trichome related genes; (ii) highly expressed in  
16 cotton ovules and fiber initials and (iii) specific expression in *Arabidopsis* transgenic Col-0  
17 trichome by cotton orthologous promoter-driven GUS and complementation of trichome  
18 development for *Arabidopsis* trichomeless mutants by overexpression of cotton ortholog and the  
19 varied influences to cotton fiber development in loss-of-function and gain-of function transgenic  
20 cotton (27). Not like *Arabidopsis* trichome, cotton fibers are initiated from seed epidermal cells,  
21 their initiation and development highly synchronize and unbranch with extreme elongation (28).  
22 To verify the reasonability of the speculation, we roughly grouped the documented cotton fiber

1 development related genes into three groups (S-table 1) including ROS-related, fiber initiation-  
2 related and fiber elongation-related.

3 Transcript profiling for ROS-related genes including *Ghmir398*, with its two targets,  
4 *GhCSD1* and *GhCSD2*, *GhAPX*, *GhRDL1* and *GhRDL2*, only *GhRDL1* and *GhRDL2* show  
5 consistently lowest expression in the two lintless fuzzless mutants but significantly increase  
6 expression starting 2 days of pre-anthesis and post-anthesis (fig.1S). The expression of these  
7 genes helps maintain homeostasis of H<sub>2</sub>O<sub>2</sub>, an important signal for cotton fiber initiation and  
8 development. We grouped cotton orthologs based on homology to Arabidopsis leaf trichome  
9 development into cotton fiber-initiation-related genes. We do not find specific correlation of  
10 these group genes to the two lintless fuzzless mutants. Interestingly, all of these genes express in  
11 the cotton expanded leaves which produce trichomes (fig.2S). This result suggests these genes  
12 may be cotton leaf trichome orthologs but not cotton seed fiber genes.

13 Interestingly, Walford et al., report (29) RNAi cotton transgenic plants for *GhMYB25-like*  
14 resulted in fibreless seeds but developed with normal trichomes elsewhere, mimics the Xu142 fl  
15 mutant. Furthermore, these plants had decreased expression of the fibers related MYBs including  
16 *GhMYB25* and *GhMYB109*, suggesting these MYBs are downstream of *GhMYB25-like*.  
17 *GhMYB25-like*, not an ortholog to Arabidopsis trichome genes, is identified from its reduced  
18 expression in a fibreless mutant of cotton (Xu142 fl) (30) Du et al., 2001). These data implicate  
19 cotton fiber initiation and development may share similar regulatory process with *Arabidopsis*  
20 trichome but the involved genes are different. It is important to note that Arabidopsis seeds do  
21 not have seed trichomes like cotton and thus cannot be considered as an ideal surrogate model to  
22 evaluate the specificity of expression of cotton fiber genes. The search for cotton fiber initiation  
23 and development genes is still elusive. We also detect the expression of the third group of

1 documented fiber related gene, fiber elongation genes, lowest to undetectable expression of  
2 *GhMYB25* and *GhEXPI* (fig.3S) relate to the two lintless fuzzless mutants. This implies these  
3 two genes are acting downstream and necessary for fiber elongation.

4 Cell morphogenesis is the process by which cells acquire shape during development and  
5 is central to organogenesis in multicellular organisms. Based on the currently available results,  
6 Qin et.al. (31) speculated that fiber cells may elongate via a combination of both tip-growth and  
7 diffuse-growth modes and named linear cell-growth mode. Using NBT staining ovules at  
8 different development stages between TM1 and N1n2, a lintless-fuzzless doubled mutant  
9 originated from TM, we show that H<sub>2</sub>O<sub>2</sub> accumulation in cotton ovule epidermal cells but no  
10 H<sub>2</sub>O<sub>2</sub> staining in *Arabidopsis* and cotton leaf trichomes. We propose that the directional  
11 accumulation of H<sub>2</sub>O<sub>2</sub> in cotton ovule epidermal cell is the drive for cotton fiber initiation and  
12 elongation.

13

#### 14 **References and Notes**

- 15 1. L. M. A. Dirk, T. A. Wilkins, J. M. Stewart, *et al.*, Eds. *Physiology of Cotton*, (2010).
- 16 2. D. W. Galbraith, *Comp. Funct. Genom.* **4**, 208–15 (2003).
- 17 3. B. M. Lange, *Curr. Opin. Plant Biol.* **8**, 236–41 (2005).
- 18 4. T. A. Wilkins, J. A. Jernstedt, in *Cotton Fibers*, A. S. Basra, Ed. (1999), vol. 269, pp. 1–34.
- 19 5. A. B. Arpat *et al.*, *Plant Mol. Boil.* **54**, 911–29 (2004).
- 20 6. Y. L. Ruan, D. J. Llewellyn, R. T. Furbank, *Plant Cell* **13**, 47–60 (2001).
- 21 7. J. M. Stewart, *Amer. J. Bot.* **62**, 723–730 (1975).

- 1 8. N. Yu *et al.*, *Plant Cell* **22**, 2322–35 (2010).
- 2 9. J. Holt, J. M. Stewart, *Proc. Beltwide cotton conf.* 1320–1324 (1994).
- 3 10. S. C. Tiwari, T. A. Wilkins, *Canad. J. Bot.* **73**, 746–757 (1995).
- 4 11. D. Zhang, T. Zhang, W. Guo, *J. Plant Physiol.* **167**, 393–9 (2010).
- 5 12. F. A. Andriunas *et al.*, *J.Exp.Bot.* **63**, 3617–29 (2012).
- 6 13. C. E. Offler *et al.*, *Annu. Rev. Plant Biol.* **54**, 431–54 (2003).
- 7 14. J. Foreman *et al.*, *Nature* **422**, 442–6 (2003).
- 8 15. R. J. Carol *et al.*, *Nature* **438**, 1013–6 (2005).
- 9 16. A. C. Allan, R. Fluhr, *Plant Cell* **9**, 1559–1572 (1997).
- 10 17. T. Xing, V. J. Higgins, E. Blumwald, *Plant Cell* **9**, 249–59 (1997).
- 11 18. V. D. Petrov, F. Van Breusegem, *AoB Plants* **2012**, 1–13 (2012).
- 12 19. G. Miller *et al.*, *Plant, Cell & Environ.* **33**, 453–67 (2010).
- 13 20. H. B. Li *et al.*, *New Phytol.* **175**, 462–71 (2007).
- 14 21. Y.M. Qin *et al.*, *Plant Signal. & Bahav.* **3**, 194–196 (2008).
- 15 22. R. Hovav *et al.*, *PLoS Genet.* **4**, e25 (2008).
- 16 23. T. S. Potikha *et al.*, *Plant Physiol.* **119**, 849–58 (1999).
- 17 24. L. Xiong *et al.*, *Plant Cell* **14**, S165–S183 (2002).
- 18 25. L. Serna, C. Martin, *Trends Plant Sci.* **11**, 274–80 (2006).
- 19 26. S. Wang *et al.*, *Plant Cell* **16**, 2323–2334 (2004).



- 1 27. X.Y. Guan *et al.*, *Physiol. Plantar.* **134**, 174–82 (2008).
- 2 28. H. J. Kim, B. A. Triplett, *Plant Physiol.* **127**, 1361–1366 (2001).
- 3 29. S. A. Walford, *et al.*, *Plant J.* **65**, 785–97 (2011).
- 4 30. X. Du *et al.*, *Plant Breed.* **120**, 519–522 (2001).
- 5 31. Y.-M. Qin, Y.-X. Zhu, *Curr.Opin.Plant Biol.* **14**, 106–11 (2011).

6

7 **Acknowledgments:** We thank Dr. Nouredine Abidi for help with Scanning Electron  
8 Microscopy (SEM) at Fiber and Biopolymer Research Institute (FBRI) at Texas Tech University  
9 and funding by the Texas Governor's Emerging Technology Superior Research Award granted to  
10 T.A.W.

11

12 Supplementary Materials

13

14

15 Materials and Methods

16

17 Figs. S1 to S3

18

19 Tables S1

20

21

22

23

TNT

N1

e2

N1e2


N2143

N2143-48

SEM

EDS

EDS



A



B

C

D

E

F




SLP72<sup>+/+</sup> EGFP<sup>+/+</sup>

SLP72<sup>+/+</sup> EGFP<sup>+/+</sup> + 1.5 μM TCR

SLP72<sup>+/+</sup> EGFP<sup>+/+</sup> + 1.5 μM TCR + 100 nM Vβ2

TCR



TCR Staining

SLP



SLP Staining (2.5 μm)

A

B

C