

Characterization of the transcriptome, nucleotide sequence polymorphism, and natural selection in the desert adapted mouse *Peromyscus eremicus*

Matthew D. MacManes¹, Michael B. Eisen²

¹ Department of Molecular, Cellular and Biomedical Sciences, University of New Hampshire, Durham NH, USA

² HHMI and University of California, Berkeley, Berkeley, CA, USA

* E-mail: macmanes@gmail.com, @PeroMHC

1 Abstract

2 As a direct result of intense heat and aridity, deserts are thought to be among the most
3 harsh of environments, particularly for their mammalian inhabitants. Given that os-
4 moregulation can be challenging for these animals, with failure resulting in death, strong
5 selection should be observed on genes related to the maintenance of water and solute
6 balance. One such animal, *Peromyscus eremicus*, is native to the desert regions of the
7 southwest United States and may live its entire life without oral fluid intake. As a first
8 step toward understanding the genetics that underlie this phenotype, we present a char-
9 acterization of the *P. eremicus* transcriptome. We assay four tissues (kidney, liver, brain,
10 testes) from a single individual and supplement this with population level renal transcrip-
11 tome sequencing from 15 additional animals. We identified a set of transcripts undergoing
12 both purifying and balancing selection based on estimates of Tajima's D. In addition, we
13 used the branch-site test to identify a transcript – *Slc2a9*, likely related to desert os-
14 moregulation – undergoing enhanced selection in *P. eremicus* relative to a set of related
15 non-desert rodents.

16

17 Introduction

18 Deserts are widely considered one of the harshest environments on Earth. Animals living
19 in desert environments are forced to endure intense heat and drought, and as a result,
20 species living in these environments are likely to possess specialized mechanisms to deal
21 with them. While living in deserts likely involves a large number of adaptive traits, the
22 ability to osmoregulate – to maintain the proper water and electrolyte balance – appears

23 to be paramount (Walsberg, 2000). Indeed, the maintenance of water balance is one of
24 the most important physiologic processes for all organisms, whether they be desert inhab-
25 itants or not. Most animals are exquisitely sensitive to changes in osmolality, with slight
26 derangement eliciting physiologic compromise. When the loss of water exceeds dietary
27 intake, dehydration - and in extreme cases, death - can occur. Thus there has likely been
28 strong selection for mechanisms supporting optimal osmoregulation in species that live
29 where water is limited. Understanding these mechanisms will significantly enhance our
30 understanding of the physiologic processes underlying osmoregulation in extreme envi-
31 ronments, which will have implications for studies of human health, conservation, and
32 climate change.

33

34 The genes and structures responsible for the maintenance of water and electrolyte
35 balance are well characterized in model organisms such as mice (Tatum et al., 2009), rats
36 (Romero et al., 2007; Rojek et al., 2006; Nielsen et al., 1995), and humans (Mobasheri
37 et al., 2007; Bedford et al., 2003; Nielsen et al., 1999). These studies, many of which
38 have been enabled by newer sequencing technologies, provide a foundation for studies of
39 renal genomics in non-model organisms. Because researchers have long been interested
40 in desert adaptation, a number of studies have looked at the morphology or expression
41 of single genes in the renal tissues of desert adapted rodents *Phyllotis darwini* (Gallardo
42 et al., 2005), *Psammomys obesus* (Kaissling et al., 1975), and *Perognathus penicillatus*
43 (Altschuler et al., 1979). More recently, full renal transcriptomes have been generated for
44 *Dipodomys spectabilis* and *Chaetodipus baileyi*, (Marra et al., 2014) as well as *Abrothrix*
45 *olivacea* (Giorello et al., 2014).

46

47 These studies provide a rich context for current and future work aimed at developing
48 a synthetic understanding of the genetic and genomic underpinnings of desert adaptation
49 in rodents. As a first step, we have sequenced, assembled, and characterized the tran-
50 scriptome (using four tissue types - liver, kidney, testes and brain) of a desert adapted
51 cricetid rodent endemic to the southwest United States, *Peromyscus eremicus*. These an-
52 imals have a lifespan typical of small mammals (Veal and Caire, 2001), and therefore an
53 individual may live its entire life without ever drinking water. Additionally, they have a
54 distinct advantage over other desert animals (e.g. *Dipodomys*) in that they breed readily
55 in captivity, which enables future laboratory studies of the phenotype of interest. In addi-
56 tion, the focal species is positioned in a clade of well known animals (e.g. *P. californicus*,

57 *P. maniculatus*, and *P. polionotus*) (Feng et al., 2007) with growing genetic and genomic
58 resources (Shorter et al., 2014; Panhuis et al., 2011; Shorter et al., 2012). Together, this
59 suggests that future comparative studies are possible.

60

61 While the elucidation of the mechanisms underlying adaptation to desert survival is
62 beyond the scope of this manuscript, we aim to lay the groundwork by characterizing the
63 transcriptome from four distinct tissues (brain, liver, kidney, testes). These data will be
64 included in the current larger effort aimed at sequencing the entire genome. Further, via
65 sequencing the renal tissue of a total of 15 additional animals, we characterize nucleotide
66 polymorphism and genome-wide patterns of natural selection. Together, these investiga-
67 tions will aid in our overarching goal to understand the genetic basis of adaptation to
68 deserts in *P. eremicus*.

69 **Materials and Methods**

70 **Animal Collection and Study Design**

71 To begin to understand how genes may underlie desert adaptation, we collected 16 adult
72 individuals (9 male, 7 female) from a single population of *P. eremicus* over a two-year time
73 period (2012-2013). These individuals were captured in live traps and then euthanized
74 using isoflurane overdose and decapitation. Immediately post-mortem, the abdominal and
75 pelvic organs were removed, cut in half (in the case of the kidneys), placed in RNAlater and
76 flash frozen in liquid nitrogen. Removal of the brain, with similar preservation techniques,
77 followed. Time from euthanasia to removal of all organs never exceeded five minutes.
78 Samples were transferred to a -80C freezer at a later date. These procedures were approved
79 by the Animal Care and Use Committee located at the University of California Berkeley
80 (protocol number R224) and University of New Hampshire (protocol number 130902) as
81 well as the California Department of Fish and Game (protocol SC-008135) and followed
82 guidelines established by the American Society of Mammalogy for the use of wild animals
83 in research (Sikes et al., 2011).

84 **RNA extraction and Sequencing**

85 Total RNA was extracted from each tissue using a TRIzol extraction (Invitrogen) fol-
86 lowing the manufacturer's instructions. Because preparation of an RNA library suitable

87 for sequencing is dependent on having high quality, intact RNA, a small aliquot of each
88 total RNA extract was analyzed on a Bioanalyzer 2100 (Agilent, Palo Alto, CA, USA).
89 Following confirmation of sample quality, the reference sequencing libraries were made
90 using the TruSeq stranded RNA prep kit (Illumina), while an unstranded TruSeq kit was
91 used to construct the other sequencing libraries. A unique index was ligated to each
92 sample to allow for multiplexed sequencing. Reference libraries (n=4 tissue types from
93 Peer360, a male mouse used for generating a genome sequence - not part of the current
94 study) were then pooled to contain equimolar quantities of each individual library and
95 submitted for Illumina sequencing using two lanes of 150nt paired end sequencing em-
96 ploying the rapid-mode of the HiSeq 2500 sequencer at The Hubbard Center for Genome
97 Sciences (University of New Hampshire). The remaining 15 libraries were multiplexed
98 and sequenced in a mixture of 100nt paired and single end sequencing runs across several
99 lanes of an Illumina HiSeq 2000 at the Vincent G. Coates Genome Center (University of
100 California, Berkeley).

101 **Sequence Data Preprocessing and Assembly**

102 The raw sequence reads corresponding to the four tissue types were error corrected using
103 the software `bless` version 0.17 (Heo et al., 2014) using `kmer=25`, based on the developer's
104 default recommendations ([https://github.com/macmanes/pero_transcriptome/blob/](https://github.com/macmanes/pero_transcriptome/blob/master/analyses.md#error-correction)
105 `master/analyses.md#error-correction`). The error-corrected sequence reads were adapter
106 and quality trimmed following recommendations from MacManes (MacManes, 2014) and
107 Mbandi (Mbandi et al., 2014). Specifically, adapter sequence contamination and low
108 quality nucleotides (defined as PHRED <2) were removed using the program `Trimmo-`
109 `matic` version 0.32 (Bolger et al., 2014). Reads from each tissue were assembled using the
110 `Trinity` version released 17 July 2014 (Haas et al., 2013). We used flags to indicate the
111 stranded nature of sequencing reads and set the maximum allowable physical distance be-
112 tween read pairs to 999nt ([https://github.com/macmanes/pero_transcriptome/blob/](https://github.com/macmanes/pero_transcriptome/blob/master/analyses.md#trinity-assemblies)
113 `master/analyses.md#trinity-assemblies`). We elected to assembly reads derived from
114 a single deeply sequenced individual (Peer360, a male) to reduce polymorphism and thus
115 the complexity of the de Bruijn graph, which has important implications for runtime,
116 hardware requirements (Lowe et al., 2014; Pop, 2009), and assembly contiguity (Vijay
117 et al., 2013). Individual tissues were assembled independently, as we hypothesize that
118 tissue specific isoforms would be reconstructed with higher fidelity that if all tissues were
119 assembled together.

120 The assembly was conducted on a linux workstation with 64 cores and 512Gb RAM.
121 To filter the raw sequence assembly, we downloaded *Mus musculus* cDNA and ncRNA
122 datasets from Ensembl (ftp://ftp.ensembl.org/pub/release-75/fasta/mus_musculus/)
123 and the *Peromyscus maniculatus* reference transcriptome from NCBI (ftp://ftp.ncbi.nlm.nih.gov/genomes/Peromyscus_maniculatus_bairdii/RNA/). We used a blastN
124 (version 2.2.29+) procedure (default settings, evaluate set to 10^{-10}) to identify contigs in
125 the *P. eremicus* dataset likely to be biological in origin (https://github.com/macmanes/pero_transcriptome/blob/master/analyses.md#blasting). This procedure, when a
126 reference dataset is available, retains more putative transcripts than a strategy employing
127 expression-based filtering (remove if transcripts per million (TPM) <1 (MacManes and
128 Lacey, 2012)) of the raw assembly. We then concatenated the filtered assemblies from each
129 tissue into a single file and reduced redundancy using the software cd-hit-est version 4.6 (Li
130 and Godzik, 2006) using default settings, except that sequences were clustered based on
131 95% sequence similarity (https://github.com/macmanes/pero_transcriptome/blob/master/analyses.md#cd-hit-est). This multi-fasta file was used for all subsequent
132 analyses, including annotation and mapping.
133
134
135
136

137 Assembled Sequence Annotation

138 The filtered assemblies were annotated using the default settings of the blastN algorithm
139 (Camacho et al., 2009) against the Ensembl cDNA and ncRNA datasets described above,
140 downloaded on 1 August 2014. Among other things, the Ensemble transcript identifiers
141 were used in the analysis of gene ontology conducted in the PANTHER package (Mi,
142 2004). Next, because rapidly evolving nucleotide sequences may evade detection by blast
143 algorithms, we used HMMER3 version 3.1b1 (Wheeler and Eddy, 2013) to search for con-
144 served protein domains contained in the dataset using the Pfam database (Punta et al.,
145 2012) (https://github.com/macmanes/pero_transcriptome/blob/master/analyses.md#hmmer3pfam). Lastly, we extracted putative coding sequences using Transdecoder ver-
146 sion 4Jul2014 (<http://transdecoder.sourceforge.net/>) (https://github.com/macmanes/pero_transcriptome/blob/master/analyses.md#transdecoder)
147
148
149

150 To identify patterns of gene expression unique to each tissue type, we mapped sequence
151 reads from each tissue type to the reference assembly using bwa-mem (version cloned from
152 Github 7/1/2014) (Li, 2013). We estimated expression for the four tissues individually

153 using default settings of the software eXpress version 1.51 (Roberts and Pachter, 2013).
154 Interesting patterns of expression, including instances where expression was limited to a
155 single tissue type, were identified and visualized.

156

157 **Population Genomics**

158 In addition to the reference individual sequenced at four different tissue types, we se-
159 quenced 15 other conspecific individuals from the same population in Palm Desert, Cali-
160 fornia. Sequence data were mapped to the reference transcriptome using bwa-mem. The
161 alignments were sorted and converted to BAM format using the samtools software pack-
162 age (Li et al., 2009), then passed to the program ANGSD version 0.610, which was used
163 for calculating the folded site frequency spectrum (SFS) and Tajima's D (Korneliusson
164 et al., 2013) using instructions found at <http://popgen.dk/angsd/index.php/Tajima>.

165

166 **Natural Selection**

167 To characterize natural selection on several genes related to water and ion homeostasis, we
168 identified several of the transcripts identified as experiencing positive selection in a recent
169 work on desert-adapted Heteromyid rodents (Marra et al., 2014). The coding sequences
170 corresponding to these genes, Solute Carrier family 2 member 9 (Slc2a9), the Vitamin
171 D3 receptor (Vdr) and several of the Aquaporin genes (Aqp1,2,4,9), were extracted from
172 the dataset, aligned using the software MACSE version 1.01b (Ranwez et al., 2011) to
173 homologous sequences in *Mus musculus*, *Rattus norvegicus*, *Peromyscus maniculatus*, and
174 *Homo sapiens* as identified by the conditional reciprocal best blast procedure (CRBB,
175 (Aubry et al., 2014)). An unrooted gene tree with branch lengths was constructed using
176 the online resource ClustalW2-Phylogeny (http://www.ebi.ac.uk/Tools/phylogeny/clustalw2_phylogeny/), and the tree and alignment were analyzed using the branch-
178 site model (model=2, nsSites=2, fix_omega=0 versus model=2, nsSites=2, fix_omega=1,
179 omega=1) implemented in PAML version 4.8 (Yang and dos Reis, 2011; Yang, 2007).
180 Significance was evaluated via the use of the likelihood ratio test.

181

182 **Results and Discussion**

183 **RNA extraction, Sequencing, Assembly, Mapping**

184 RNA was extracted from the hypothalamus, renal medulla, testes, and liver from each
185 individual using sterile technique. TRIzol extraction resulted in a large amount of high
186 quality ($RIN \geq 8$) total RNA, which was then used as input. Libraries were constructed
187 as per the standard Illumina protocol and sequenced as described above. The number
188 of reads per library varied from 56 million strand-specific paired-end reads in Peer360
189 kidney, to 9 million single-end reads in Peer321 (Table 1, available as part BioProject
190 PRJNA242486). Adapter sequence contamination and low-quality nucleotides were elim-
191 inated, which resulted in a loss of $<2\%$ of the total number of reads. These trimmed
192 reads served as input for all downstream analyses.

193 **Table 1**

194

DATASET	NUM. RAW READS	SRA ACCESSION
PEER360 TESTES	32M PE/SS	SRR1575398
PEER360 LIVER	53M PE/SS	SRR1575397
PEER360 KIDNEY	56M PE/SS	SRR1575396
PEER360 BRAIN	23M PE/SS	SRR1575395
PEER305	19M PE	SRR1575434
PEER308	15M PE	SRR1575437
PEER319	14M PE	SRR1575439
PEER321	9M SE	SRR1575441
PEER340	16M PE	SRR1575443
195 PEER352	14M PE	SRR1575464
PEER354	9M SE	SRR1575466
PEER359	14M PE	SRR1575492
PEER365	16M PE	SRR1575493
PEER366	16M PE	SRR1575494
PEER368	14M PE	SRR1575624
PEER369	14M PE	SRR1575625
PEER372	17M SE	SRR1576070
PEER373	23M SE	SRR1576071
PEER380	16M SE	SRR1576072
PEER382	14M SE	SRR1576073


196 Table 1. The number of sequencing reads per sample, whose identity is indicated
197 by Peer[number]. PE=paired end, SS=strand specific, SE=single end sequencing.

198 Transcriptome assemblies for each tissue type was accomplished using the program
199 Trinity (Haas et al., 2013). The raw assemblies for brain, liver, testes, and kidney con-
200 tained 185425, 222096, 180233, and 514091 assembled sequences respectively. This as-
201 sembly was filtered using a blastN procedure against the *Mus* cDNA and ncRNA and
202 *P. maniculatus* cDNAs, which resulted in a final dataset containing 68331 brain-derived
203 transcripts, 71041 liver-derived transcripts, 67340 testes-derived transcripts, and 113050
204 kidney-derived transcripts. Mapping the error-corrected adapter/quality trimmed reads
205 to these datasets resulted in mapping 94.98% (87.01% properly paired) of the brain-
206 derived reads to the brain transcriptome, 96.07% (88.13% properly paired) of the liver-
207 derived reads to the liver transcriptome, 96.81% (85.10% properly paired) of the testes-
208 derived reads to the testes transcriptome, and 91.87% (83.77% properly paired) of the

209 kidney-derived reads to the kidney transcriptome. Together, these statistics suggest that
210 the tissue-specific transcriptomes are of extremely high quality. All tissue-specific assem-
211 blies are to be made available on Dryad, and until then are stored on Dropbox (https://www.dropbox.com/sh/2jwzd8p6n6eluco/AAA03nSdXb_u4wtQZRTwqW9ia?dl=0).
212

213

214 **Figure 1**



215

216 Figure 1. The Venn Diagram, which provides a visual representation of the overlap
217 of expression of the four tissue types. The majority of transcripts (66,324) are
218 expressed in all studied tissue types.

219 We then estimated gene expression on each of these tissue-specific datasets, which al-
220 lowed us to understand expression patterns in the multiple tissues (Pero.tissue.xprs, will
221 be made available on Dryad, until then on Dropbox (https://www.dropbox.com/sh/2jwzd8p6n6eluco/AAA03nSdXb_u4wtQZRTwqW9ia?dl=0)). Specifically, we constructed a
222 Venn diagram (Figure 1) that allowed us to visualize the proportion of genes whose ex-
223 pression was limited to a single tissue and those whose expression was ubiquitous. 66324
224 transcripts are expressed on all tissue types, while 13086 are uniquely expressed in the
225 kidney, 2222 in the testes, 3580 in the brain, and 2798 in the liver. The kidney appears
226 to an outlier in the number of unique sequences, though this could be the result of the
227 recovery of more lowly expressed transcripts or isoforms.
228

229

230 In addition to this, we estimated mean TPM (number of transcripts per million) for
231 all transcripts. Table 2 consists of the 10 genes whose mean TPM was the highest. Sev-
232 eral genes in this list are predominately present in a single tissue type. For instance
233 Transcript_126459, Albumin is very highly expressed in the liver, but less so in the other
234 tissues. It should be noted, however, that making inference based on uncorrected values
235 for TPM is not warranted. Statistical testing for differential expression was not imple-
236 mented due to the fact that no replicates are available.

237

238 After expression estimation, the filtered assemblies were concatenated together, and
239 after the removal of redundancy with cd-hit-est, 122,584 putative transcripts remained
240 (to be made available on Genbank, and until then are stored on Dropbox [https://www.
241 dropbox.com/sh/2jwzd8p6n6eluco/AAA03nSdXb_u4wtQZRTwqW9ia?dl=0](https://www.dropbox.com/sh/2jwzd8p6n6eluco/AAA03nSdXb_u4wtQZRTwqW9ia?dl=0)). From this fil-
242 tered concatenated dataset, we extracted 71626 putative coding sequences (72Mb, to be
243 made available on Dryad). Of these 71626 sequences, 38221 contained complete open read-
244 ing frames (containing both start and stop codons), while the others were either truncated
245 at the 5-prime end (20239 sequences), the 3-prime end (6445 sequences), or were inter-
246 nal (6721 sequencing with neither stop nor start codon). The results of a Pfam search
247 conducted on the predicted amino acid sequences will be found on Dryad, and until then
248 are stored on Dropbox [https://www.dropbox.com/sh/2jwzd8p6n6eluco/AAA03nSdXb_
249 u4wtQZRTwqW9ia?dl=0](https://www.dropbox.com/sh/2jwzd8p6n6eluco/AAA03nSdXb_u4wtQZRTwqW9ia?dl=0)..

250

251 **Table 2**

252

	Transcript ID	Testes	Liver	Kidney	Brain	Genbank ID	Gene ID
	Transcript_83842	2.05E+03	6.40E+03	1.03E+04	5.47E+03	DQ073446.1	COX2
	Transcript_126459	1.43E+01	2.22E+04	2.77E+01	6.73E+00	XM_006991665.1	Alb
	Transcript_128937	4.39E+00	1.91E+04	4.74E+02	2.23E+00	XM_007627625.1	Apoa2
	Transcript_81233	1.71E+03	5.23E+03	6.11E+03	3.08E+03	XM_006993867.1	Fth1
253	Transcript_94125	3.67E+01	1.08E+04	2.09E+03	2.75E+00	XM_006977178.1	CytP450
	Transcript_119945	5.03E+03	1.15E+03	1.33E+03	3.71E+03	XM_008686011.1	Ubb
	Transcript_5977	4.95E+00	1.01E+04	3.05E+02	3.58E+02	XM_006978668.1	Tf
	Transcript_4057	2.62E+01	9.32E+03	1.34E+02	8.38E+01	XM_006994871.1	Apoc1
	Transcript_112523	4.07E+02	7.36E+03	7.78E+02	9.54E+02	XM_006994872.1	Apoe
	Transcript_98376	1.98E+00	8.66E+03	1.02E+00	2.68E+00	XM_006970208.1	Ttr

254 Table 2. The 10 transcripts with the highest mean TPM (transcripts per million).

255 Population Genomics


256 As detailed above, RNAseq data from 15 individuals were mapped to the reference tran-
257 scriptome with the resulting BAM files being used as input to the software package
258 ANGSD. The Tajima's D statistic was calculated for all transcripts covered by at least
259 14 of the 15 individuals. In brief, a negative Tajima's D - a result of lower than expected
260 average heterozygosity - is often associated with purifying or directional selection, recent
261 selective sweep or recent population expansion, or a complex combination of these forces.
262 In contrast, a positive value for Tajima's D represents higher than expected average het-
263 erozygosity, often associated with balancing selection.

264
265 The distribution of the estimates of Tajima's D for all of the assembled transcripts is
266 shown in Figure 2. Although Tajima's D is known to be sensitive to demographic history,
267 which is largely unknown for this population, the estimates may also be drive by patterns
268 of selection. In general, the distribution is skewed toward negative values (mean=-0.89,
269 variance=0.58), which may be the result of purifying selection, a model of evolution com-
270 monly invoked for coding DNA sequences (Chamary et al., 2006). Table 3 presents the
271 10 transcripts whose estimate of Tajima's D is the greatest, while Table 4 presents the
272 10 transcripts whose estimate of Tajima's D is the least. The former list of genes may
273 contain transcripts experiencing balancing selection in the studied population. This list
274 includes, interestingly, genes obviously related to solute and water balance (e.g. Clcnkb

275 and a transmembrane protein gene) and immune function (a interferon-inducible GTPase
276 and a Class 1 MHC gene). The latter group, containing transcripts whose estimates of
277 Tajima's D are the smallest are likely experiencing purifying selection. Many of these
278 transcripts are involved in core regulatory functions where mutation may have strongly
279 negative fitness consequences.

280

281 **Figure 2**



282

283 Figure 2. The distribution of Tajima's D for all putative transcripts.

284 **Table 3**

285

Transcript ID	GenBank ID	Description	Tajima's D
Transcript_49049	XM_006533884.1	heterogeneous nuclear ribonucleoprotein H1 (Hnrnp1)	3.26
Transcript_38378	XM_006522973.1	Son DNA binding protein (Son)	3.19
Transcript_126187	NM_133739.2	transmembrane protein 123 (Tmem123)	3.02
Transcript_70953	XM_006539066.1	chloride channel Kb (Clcnkb)	2.96
Transcript_37736	XM_006997718.1	h-2 class I histocompatibility antigen	2.92
Transcript_21448	XM_006986148.1	zinc finger protein 624-like	2.84
Transcript_47450	NM_009560.2	zinc finger protein 60 (Zfp60)	2.82
Transcript_122250	XM_006539068.1	chloride channel Kb (Clcnkb)	2.81
Transcript_78367	XM_006496814.1	CDC42 binding protein kinase alpha (Cdc42bpa)	2.78
Transcript_96470	XM_006987129.1	interferon-inducible GTPase 1-like	2.77

Table 3. The 10 transcripts with the highest values for Tajima's D, which suggests balancing selection.

288 **Table 4**

289

Transcript ID	GenBank ID	Description	Tajima's D
Transcript_84359	XM_006991127.1	nuclear receptor coactivator 3 (Ncoa3)	-2.82
Transcript_87121	XM_006970128.1	methyl-CpG binding domain protein 2 (Mbd2)	-2.82
Transcript_125755	EU053203.1	alpha globin gene cluster	-2.78
Transcript_87128	XM_006976644.1	membrane-associated ring finger (March5)	-2.76
Transcript_55468	XM_006978377.1	Vpr binding protein (Vprbp)	-2.75
Transcript_116042	XM_006980811.1	membrane associated guanylate kinase (Magi3)	-2.75
Transcript_18966	XM_006982814.1	ubiquitin protein ligase E3 component n-recognin 5 (Ubr5)	-2.75
Transcript_122204	XM_008772511.1	zinc finger protein 612 (Zfp612)	-2.75
Transcript_100550	XM_006971297.1	bromodomain adjacent to zinc finger domain, 1B (Baz1b)	-2.74
Transcript_33267	XM_006975561.1	pumilio RNA-binding family member 1 (Pum1)	-2.75

Table 4. The 10 transcripts with the lowest values for Tajima's D, which suggests purifying or directional selection.

293 **Natural Selection**

294 To begin to test the hypothesis that selection on transcripts related to osmoregulation is
295 enhanced in the desert adapted *P. eremicus*, we calculated Tajima's D as described above,

296 and implemented the branch-site test using alignments produced in MACSE. These align-
297 ments were manually inspected, and were relatively free from indels and internal stop
298 codons. We set the sequence corresponding to *P. eremicus* for Slc2a9, Vdr, and several
299 of the Aquaporin genes (Aqp1,2,4,9) as the foreground lineages in six distinct program
300 executions. These transcripts Slc2a9 and Vdr were chosen specifically because they - the
301 former significantly - were recently linked to osmoregulation in a desert rodent (Marra
302 et al., 2014). The test for Slc2a9 was highly significant ($2\Delta Lnl=51.4$, $df=1$, $p=0$, Ta-
303 ble 5), indicating enhanced selection in *P. eremicus* relative to the other lineages. The
304 branch site test for positive selection conducted on the Vdr and Aquaporin genes were
305 non-significant. While the branch site test of positive selection is largely non-significant,
306 estimating Tajima's D for these few candidate loci demonstrates that either a selective
307 or demographic process may be influencing the genome at these functionally relevant sites.

308

309 **Table 5**

310

Transcript ID	Description	Tajima's D	Branch Site Test p.value
Transcript_106085	Slc2a9	2.15	p=0
Transcript_114624	Vdr	1.97	p=1
311 Transcript_128972	Aqp1	1.39	p=1
Transcript_33960	Aqp2	1.78	p=1
Transcript_22154	Aqp4	2.10	p=1
Transcript_107677	Aqp9	2.06	p=1

312 Table 5. Several candidate genes were evaluated using Tajima's D and the branch
313 site test implemented in PAML.

314 Conclusions

315 As a direct result of intense heat and aridity, deserts are thought to be amongst the harsh-
316 est environments, particularly for mammalian inhabitants. Given that osmoregulation can
317 be challenging for these animals - with failure resulting in death - strong selection should
318 be observed on genes related to the maintenance of water and solute balance. This study
319 aimed to characterize the transcriptome of a desert-adapted rodent species, *P. eremicus*.
320 Specifically, we characterized the transcriptome of four tissue types (liver, kidney, brain,
321 and testes) from a single individual and supplemented this with population-level renal

322 transcriptome sequencing from 15 additional animals. We identified a set of transcripts
323 undergoing both purifying and balancing selection based on Tajima's D. In addition,
324 we used a branch site test to identify a transcript, likely related to desert osmoregula-
325 tion, undergoing enhanced selection in *P. eremicus* relative to a set of non-desert rodents.
326

327 Acknowledgments

328 This manuscript was greatly improved by careful review from C. Titus Brown, Elijah
329 Lowe, and an anonymous reviewer, as well as by Matthew Hahn, who provided feedback
330 on an earlier version of the manuscript posted on bioRxiv.

331 References

- 332 Altschuler, E. M., Nagle, R. B., Braun, E. J., Lindstedt, S. L., and Krutzsch, P. H.
333 (1979). Morphological study of the desert heteromyid kidney with emphasis on the
334 genus *Perognathus*. *The Anatomical Record*, 194(3):461–468.
- 335 Aubry, S., Kelly, S., Kumpers, B. M. C., Smith-Unna, R. D., and Hibberd, J. M. (2014).
336 Deep Evolutionary Comparison of Gene Expression Identifies Parallel Recruitment of
337 Trans-Factors in Two Independent Origins of C4 Photosynthesis. *PLOS Genetics*,
338 10(6):e1004365.
- 339 Bedford, J. J., Leader, J. P., and Walker, R. J. (2003). Aquaporin expression in normal
340 human kidney and in renal disease. *Journal of the American Society of Nephrology*,
341 14(10):2581–2587.
- 342 Bolger, A. M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for
343 Illumina sequence data. *Bioinformatics*, 30(15):btu170–2120.
- 344 Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and
345 Madden, T. L. (2009). BLAST+: architecture and applications. *BMC Bioinformatics*,
346 10(1):421.
- 347 Chamary, J. V., Parmley, J. L., and Hurst, L. D. (2006). Hearing silence: Non-neutral
348 evolution at synonymous sites in mammals. *Nature Reviews Genetics*, 7(2):98–108.

- 349 Feng, B.-J., Sun, L.-D., Soltani-Arabshahi, R., Bowcock, A. M., Nair, R. P., Stuart,
350 P., Elder, J. T., Schrodi, S. J., Begovich, A. B., Abecasis, G. R., Zhang, X.-J., Callis-
351 Duffin, K. P., Krueger, G. G., and Goldgar, D. E. (2007). Toward a Molecular Phylogeny
352 for *Peromyscus*: Evidence from Mitochondrial Cytochrome- b Sequences. *Journal of*
353 *Mammalogy*, 88(5):1146–1159.
- 354 Gallardo, P. A., Cortés, A., and Bozinovic, F. (2005). Phenotypic flexibility at the molec-
355 ular and organismal level allows desert-dwelling rodents to cope with seasonal water
356 availability. *Physiological and Biochemical Zoology*, 78(2):145–152.
- 357 Giorello, F. M., Feijoo, M., D’Elía, G., Valdez, L., Opazo, J. C., Varas, V., Naya, D. E.,
358 and Lessa, E. P. (2014). Characterization of the kidney transcriptome of the South
359 American olive mouse *Abrothrix olivacea*. *BMC Genomics*, 15(1):446.
- 360 Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J.,
361 Couger, M. B., Eccles, D., Li, B., Lieber, M., MacManes, M. D., Ott, M., Orvis, J.,
362 Pochet, N., Strozzi, F., Weeks, N., Westerman, R., William, T., Dewey, C. N., Henschel,
363 R., Leduc, R. D., Friedman, N., and Regev, A. (2013). *De novo* transcript sequence
364 reconstruction from RNA-seq using the Trinity platform for reference generation and
365 analysis. *Nature Protocols*, 8(8):1494–1512.
- 366 Heo, Y., Wu, X.-L., Chen, D., Ma, J., and Hwu, W.-M. (2014). BLESS: Bloom filter-
367 based error correction solution for high-throughput sequencing reads. *Bioinformatics*,
368 30(10):1354–1362.
- 369 Kaissling, B., De Rouffignac, C., Barrett, J. M., and Kriz, W. (1975). The structural
370 organization of the kidney of the desert rodent *Psammomys obesus*. *Anatomy and*
371 *Embryology*, 148(2):121–143.
- 372 Korneliussen, Thorfinn Sand Moltke, I., Albrechtsen, a., and Nielsen, R. (2013). Calcula-
373 tion of Tajima’s D and other neutrality test statistics from low depth next-generation
374 sequencing data. *BMC Bioinformatics*, 14(1):289.
- 375 Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-
376 MEM. *arXiv*.
- 377 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis,

- 378 G., Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The
379 Sequence Alignment/Map format and SAMtools. *Bioinformatics*, 25(16):2078–2079.
- 380 Li, W. and Godzik, A. (2006). Cd-hit: A fast program for clustering and comparing large
381 sets of protein or nucleotide sequences. *Bioinformatics*, 22(13):1658–1659.
- 382 Lowe, E. K., Swalla, B. J., and Brown, C. T. (2014). Evaluating a lightweight transcrip-
383 tome assembly pipeline on two closely related Ascidian species. *PeerJ Preprints*, pages
384 1–11.
- 385 MacManes, M. D. (2014). On the optimal trimming of high-throughput mRNA sequence
386 data. *Frontiers in Genetics*, 5:1–13.
- 387 MacManes, M. D. and Lacey, E. A. (2012). The Social Brain: Transcriptome Assembly
388 and Characterization of the Hippocampus from a Social Subterranean Rodent, the
389 Colonial Tuco-Tuco (*Ctenomys sociabilis*). *PLOS ONE*, 7(9):e45524.
- 390 Marra, N. J., Romero, A., and DeWoody, J. A. (2014). Natural selection and the genetic
391 basis of osmoregulation in heteromyid rodents as revealed by RNA-seq. *Molecular*
392 *Ecology*, 23(11):2699–2711.
- 393 Mbandi, S. K., Hesse, U., Rees, D. J. G., and Christoffels, A. (2014). A glance at quality
394 score: Implication for *de novo* transcriptome reconstruction of Illumina reads. *Frontiers*
395 *in Genetics*, 5:1–17.
- 396 Mi, H. (2004). The PANTHER database of protein families, subfamilies, functions and
397 pathways. *Nucleic Acids Research*, 33(1):D284–D288.
- 398 Mobasher, A., Marples, D., Young, I. S., Floyd, R. V., Moskaluk, C. A., and Frigeri,
399 A. (2007). Distribution of the AQP4 Water Channel in Normal Human Tissues: Pro-
400 tein and Tissue Microarrays Reveal Expression in Several New Anatomical Locations,
401 including the Prostate Gland Seminal Vesicles. *Channels*, 1(1):30–39.
- 402 Nielsen, S., Chou, C., Marples, D., Christensen, E., Kishore, B., and Knepper, M. (1995).
403 Vasopressin increases water permeability of kidney collecting duct by inducing translo-
404 cation of Aquaporin-CD water channels to plasma-membrane. *Proceedings of The Na-
405 tional Academy of Sciences of The United States of America*, 92(4):1013–1017.

- 406 Nielsen, S., Kwon, T. H., Christensen, B. M., Promeneur, D., Frøkiaer, J., and Marples, D.
407 (1999). Physiology and pathophysiology of renal aquaporins. *Journal of the American*
408 *Society of Nephrology*, 10(3):647–663.
- 409 Panhuis, T. M., Broitman-Maduro, G., Uhrig, J., Maduro, M., and Reznick, D. N. (2011).
410 Analysis of Expressed Sequence Tags from the Placenta of the Live-Bearing Fish *Poe-*
411 *ciliopsis* (Poeciliidae). *Journal of Heredity*, 102(3):352–361.
- 412 Pop, M. (2009). Genome assembly reborn: Recent computational challenges. *Briefings*
413 *In Bioinformatics*, 10(4):354–366.
- 414 Punta, M., Cogill, P. C., Eberhardt, R. Y., Mistry, J., Tate, J., Boursnell, C., Pang,
415 N., Forslund, K., Ceric, G., Clements, J., Heger, A., Holm, L., Sonnhammer, E. L. L.,
416 Eddy, S. R., Bateman, A., and Finn, R. D. (2012). The Pfam protein families database.
417 *Nucleic Acids Research*, 40:D290–301.
- 418 Ranwez, V., Harispe, S., Delsuc, F., and Douzery, E. J. P. (2011). MACSE: Multiple
419 Alignment of Coding SEquences Accounting for Frameshifts and Stop Codons. *PLOS*
420 *ONE*, 6(9):e22594.
- 421 Roberts, A. and Pachter, L. (2013). Streaming fragment assignment for real-time analysis
422 of sequencing experiments. *Nature Methods*, 10(1):71–73.
- 423 Rojek, A., Fuchtbauer, E., Kwon, T., Frokiaer, J., and Nielsen, S. (2006). Severe urinary
424 concentrating defect in renal collecting duct-selective AQP2 conditional-knockout mice.
425 *Proceedings of The National Academy of Sciences of The United States of America*,
426 103(15):6037–6042.
- 427 Romero, D. G., Plonczynski, M. W., Welsh, B. L., Gomez-Sanchez, C. E., Zhou, M. Y.,
428 and Gomez-Sanchez, E. P. (2007). Gene expression profile in rat adrenal zona glomeru-
429 losa cells stimulated with aldosterone secretagogues. *Physiological Genomics*, 32(1):117–
430 127.
- 431 Shorter, K. R., Crossland, J. P., Webb, D., Szalai, G., Felder, M. R., and Vrana, P. B.
432 (2012). *Peromyscus* as a Mammalian Epigenetic Model. *Genetics Research Interna-*
433 *tional*, 2012:1–11.
- 434 Shorter, K. R., Owen, A., Anderson, V., Hall-South, A. C., Hayford, S., Cakora, P.,
435 Crossland, J. P., Georgi, V. R. M., Perkins, A., Kelly, S. J., Felder, M. R., and Vrana,

- 436 P. B. (2014). Natural genetic variation underlying differences in *Peromyscus* repetitive
437 and social/aggressive behaviors. *Behavior genetics*, 44(2):126–135.
- 438 Sikes, R. S., Gannon, W. L., and Animal Care and Use Committee of the American
439 Society of Mammalogists (2011). Guidelines of the American Society of Mammalogists
440 for the use of wild mammals in research. *Journal of Mammalogy*, 92(1):235–253.
- 441 Tatum, R., Zhang, Y., Salleng, K., Lu, Z., Lin, J. J., Lu, Q., Jeansonne, B. G., Ding,
442 L., and Chen, Y. H. (2009). Renal salt wasting and chronic dehydration in claudin-7-
443 deficient mice. *AJP: Renal Physiology*, 298(1):F24–F34.
- 444 Veal, R. and Caire, W. (2001). *Peromyscus eremicus*. *Mammalian Species*, 118:1–6.
- 445 Vijay, N., Poelstra, J. W., Künstner, A., Wolf, J. B. W., and Wolf, J. B. W. (2013).
446 Challenges and strategies in transcriptome assembly and differential gene expression
447 quantification. A comprehensive *in silico* assessment of RNA-seq experiments. *Molec-
448 ular Ecology*, 22(3):620–634.
- 449 Walsberg, G. (2000). Small mammals in hot deserts: Some generalizations revisited.
450 *Bioscience*, 50(2):109–120.
- 451 Wheeler, T. J. and Eddy, S. R. (2013). nhmmer: DNA homology search with profile
452 HMMs. *Bioinformatics*, 29(19):2487–2489.
- 453 Yang, Z. (2007). PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Molecular
454 Biology and Evolution*, 24(8):1586–1591.
- 455 Yang, Z. and dos Reis, M. (2011). Statistical Properties of the Branch-Site Test of Positive
456 Selection. *Molecular Biology and Evolution*, 28(3):1217–1228.