

COGNITIVE DOMAINS FUNCTION COMPLEMENTATION

BY *NTNG* GENE PARALOGS

Pavel Prosselkov^{1,2*}, Denis Polygalov³, Qi Zhang¹, Thomas J. McHugh³, and Shigeyoshi Itohara^{1*}

¹Laboratory of Behavioral Genetics, RIKEN Brain Science Institute, Wakoshi, 351-0198 Saitama, Japan

²Graduate School, Department of Veterinary Medicine, Faculty of Agriculture, Tokyo University, 113-8657 Tokyo, Japan

³Laboratory of Circuit and Behavior Physiology, RIKEN Brain Science Institute, 351-0198 Saitama, Japan

*corresponding authors:

prosselkov@brain.riken.jp

sitohara@brain.riken.jp

Keywords: gene duplication, evolution, subfunctionalisation, cognitive complement

ABSTRACT

Gene duplication was proposed by S. Ohno (Ohno, 1970) as a key mechanism of a gene function evolution. A pair of gene paralogs, *NTNG1* and *NTNG2*, sharing identical gene and protein structures and encoding similar proteins, forms a functional complement subfunctionalising (SF) within cognitive domains and forming cognitive endophenotypes, as detected by Intellectual Quotient (IQ) tests (Prosser et al., 2015). Both *NTNG* paralogs are associated with autism spectrum disorder (ASD), bipolar disorder (BD) and schizophrenia (SCZ), with unique non-overlapping segregation among the other 15 cognitive disorders (CD), emphasizing an evolutionary gain-dependent link between advanced cognitive functions and concomitant neurocognitive pathologies. Complementary expression and human brain transcriptome composition of the paralogs explains the observed phenomena of their functional complementarity. The lowest identity among *NTNGs* is found in a middle of encoded by them proteins designated as unknown (Ukd) domain. *NTNG1* contains anthropoid-specific constrained regions, and both genes contain non-coding conserved sequences underwent accelerated evolution in human. *NTNG* paralogs SF perturbs “structure drives function” concept at protein and gene levels. The paralogs function diversification forms a so-called “Cognitive Complement (CC)”, a product of gene duplication and subsequent cognitive subfunction bifurcation among the *NTNG* gene duplicates.

INTRODUCTION

Complex behaviors arise from a combination of simpler genetic modules that either have evolved separately or co-evolved. Many genes and proteins they encode have been found to be involved in cognitive information processing with a single variant or a single gene generally accounting for only a partial phenotypic variation of a complex trait. Cognitive processing as a quintessence of brain functioning can be viewed as a product of intricately interlinked networks generated by deeply embedded into it players with specific or partially overlapping functions. The robustness of the cognitive processing towards its single elements genetic eliminations (to study their function) and its simultaneous fragility expressed in the multiple forms of neurological disorders manifest the existence of cognitive domains interlocked but SF within a unit of cognition formed upon these domains interaction. Previously, we have described a function of a pair of gene paralogs, *NTNG1* and *NTNG2*, involved in human IQ tests performance, and underwent hominin-specific evolutionary changes (**Prosselkov et al., 2015**). Hereby, we report on these gene paralogs features focusing on underlying mechanisms of their function segregation and complementation within the cognitive domains.

RESULTS

The previously observed phenomena of functional complementation among the *NTNG* paralogs within cognitive domains (**Prosselkov et al., 2015**) is also manifested in *NTNG*-associated human pathologies diagnosed in most cases (if only not in all) by a cognitive decline (**Figure 1A-1** and **A-2**). Both genes are associated with BD and SCZ – devastating disorders sharing similar etiology (**Lee et al., 2013**) with genetic correlation by multivariate analysis of 0.590 (**Maier et al., 2015**), linked to human creativity (**Power et al., 2015**), and characterized by impulsiveness as a common diagnostic feature (**Reddy et al., 2014**). Recently found associations of both paralogs with ASD (**Sanders et al., 2015**) supports the reported genetic correlation of 0.194 ASD/SCZ pair (**Maier et al., 2015**) and shared module eigengenes detected by PC1 among these two disorders (**Parikshak et al., 2015**). 12 *NTNG1*-linked CDs, ranging from AD to TS, span a broad spectrum of clinical features frequently involving reduced processing speed (PS) and verbal comprehension (VC, **Figure 1A-1**). As for *NTNG2*, working memory (WM) deficit and inability “to bind” events (perceptual organization, PO) are the most prominent diagnostic traits for the SLE and TLE patients (**Figure 1A-2**), with PN characterised by indolent behavior in 90% of the cases (**Cavard et al., 2009**). Thus, both *NTNG* paralogs are associated with a variety of CDs and mostly in a non-overlapping manner, except for ASD, BD and SCZ characterized by shared and wide spectrum of cognitive abnormalities. The clinical etiology of the aforementioned diseases supports the IQ-deduced functional complementation among the *NTNG* paralogs (**Prosselkov et al., 2015**) with (VC/PS) and (WM/PO) deficits being also uniquely segregated among the associated cognitive pathologies.

Since both genes are expected to have identical gene exon/intron compositions but different in their intron lengths (**Yin et al., 2002**) we have reconstructed the paralogs transcriptomes by re-processing the publicly available RNA-seq dataset (**Wu et al., 2012**)

from healthy and SCZ human subjects superior temporal gyrus (STG) post-mortem brain tissue (Supplementary Table 1a=**ST1a**). A difference is noted instantly at the total expression levels (genes, exons, individual RNA transcripts) when two gene paralogs are compared (**Figure 1B-1** and **B-2**). *NTNG2* amount (as a whole gene) is 5 times larger comparing to *NTNG1*; exons (2-5) are 3 times, exons (8-9) are 18 times and exon 10 is 4 times higher expressed for *NTNG2* than for *NTNG1*. The only two exons outlaying the prevailing amount rule for the *NTNG2* mRNAs are exons 6 and 7, expressed nearly at the same absolute level as for the *NTNG1* exon paralogs, making them highly underrepresented within the whole *NTNG2* transcriptome. Next, distinct non-alternating splicing modules are formed by exons (2-5) for *NTNG1* (**Figure 1B-1**), while exons (4-5) and exons (8-9) for *NTNG2* (**Figure 1B-2**). Two structurally identical RNA transcript paralogs (*NTNG1a* = G1a and *NTNG2a* = G2a) have been found to exist in both *NTNG* transcriptomes with G2a being expressed at 8-9 times higher level than G1a. *NTNG1* is uniformly presented across the all analysed 16 human samples by 2 more protein coding RNAs (G1c and G1d, detected previously in mice brain, Nakashiba et al., 2000) and by 2 non-coding intron (9-10) derived transcripts (**Figure 1B-1**). At the same time, *NTNG2* transcriptome is comprised of one extra potentially coding RNA (G2a-like with exon 2 spliced out but in-frame coding preserved) and 2 assumed to be non-coding RNAs with exons 6 and 7 retained along with preceding and following them introns. Quite interesting that these two latter transcripts are the only RNA species with *NTNG2* exon 6 and 7 retained (**Figure 1B-2**). Two more coding (G1f and G1n) and 4 more non-coding for *NTNG1* and 9 extra non-coding for *NTNG2* RNA species have been also assembled from the available reads but due to inconsistency in their appearance across all 16 STG samples they are not presented on the figure but summarized in the table (**Figure 1C**, for details refer to **ST1d**). Summarising this, quantitative and qualitative complementary differences is a prominent feature characterising the brain RNA transcriptome of human *NTNG* paralogs.

However, no significant changes at the transcription level of neither whole genes, nor individual exons, nor reconstructed RNA transcripts have been found for SCZ and healthy subjects.

Upon calling the presence of IQ-affecting SNPs (**Prosser et al., 2015**) across all STG samples (**ST1c**) it has been revealed that 15 out of 16 subjects were positive for the T-allele of rs2149171 (exon 4-nested), shown above to attenuate the WM score in SCZ patients, making a comparison among the allele carrier vs non-carrier impossible. Four healthy and three SCZ samples carry a T-allele of rs3824574 (exon 3-nested, non-affecting IQ), and 1 healthy and 1 SCZ sample each contains a C-allele of rs4915045 (exon10, non-coding part-nested, and non-affecting IQ). Thus, among the eleven cognitive endophenotype-associated SNPs only 3 were possible to call out of the available *NTNG* transcriptome.

Distinctly complementary nature of the *NTNG* paralogs segregation within neurological disorders and RNA transcriptome usage in STG (**Figure 1**) has prompted us to analyse both genes expression across the entire human brain. We have reconstructed both genes expression profiles in the human brain areas over the life span from conception (pcw = post-conception week) to mature age (30-40 yrs old) using the RNA-seq data from BrainSpan (www.brainspan.org). Similarities and differences are easily noted when the age-dependent phases of *NTNG1* and *NTNG2* expression profiles are matched (**Figure 2**). Based on the visual inputs three distinct classifiers have been elaborated: 1. predominantly synchronous (**Figure 2A(1-4)**), characteristic mostly for the cortical areas; 2. predominantly mixed and asynchronous (**Figure 2B**), characteristic for the cerebellar cortex and subcortical formations; and 3. anti-phasic (complementary, **Figure 2C**), characteristic for the MD of thalamus and hippocampus. All analysed brain areas demonstrated an elevated level of *NTNG2* expression in comparison to *NTNG1* except for thalamus (**Figure 2C**) with the largest difference observed is at the time of birth (35-37 pcw) or soon after (4 mo) for the synchronous

classifiers (**Figure 2A**), oscillating increment values across the life span for the mixed (**Figure 2B**) and anti-phasic (**Figure 2C**) classifiers. It is quite intriguing to note that essentially all brain areas show a trend towards the expression difference being negated between the paralogs by reaching the mature age of 30-40 yrs old (nearly or above the mean age used for the IQ testing), except MD where the expression discrepancy is increased. Thus, the observed functional complementation among the *NTNG* paralogs is supported by the anatomical distribution of the genes in human brain and their expression pattern modality over the human subjects lifetime.

A direct comparison of the *NTNG* paralogs shows not only identical intron-exon gene structure (**Figure 1B-1, 2B-2**) but also closely matched exon sizes (**Figure 3A**). There are three exons of identical sizes (exons 4, 8 and 9), another three exons differed by one encoded aa (exons 3, 5 and 6) and there are exons of different sizes (exons 2, 7 and 10). In terms of size the largest difference among the genes is visually presented by the introns: intron (9-10) of *NTNG1* is 52.7 times larger its *NTNG2* paralogous intron with intron (6-7) of *NTNG1* being only 1.43-times larger pointing towards non-equilibria process of non-coding elements elaborations as the process of gene paralogs SF proceeded. Nevertheless, it can be generalised that in average all *NTNG1* introns are several times larger their *NTNG2* analogs (**Figure 3A**). We have shown previously that exons 6 and 7 are differentially used within the brain *NTNG* transcriptome (**Figure 1B-1 and B-2**) and to explore their potential contribution into the paralogs SF we have built identity matrices with these exons excluded and included (but still producing in-frame existing transcripts, **Figure 3B-1** left and right panels, respectively). Exclusion of both exons from the full-lengths transcripts (thus converting *NTNG1m* to *NTNG1a* and *NTNG2b* to *NTNG2a*, respectively) increases the identity of DNA on 2% (a relatively large effect since both exons together represent only 7.22 and 9.69% of the total coding part of the full-length RNA transcripts, *NTNG1m* and *NTNG2b*, respectively).

This effect becomes even stronger when the encoded by these transcripts proteins are also compared (**Figure 3B-2**). The spliced out Ukd protein domains (encoded by the exons 6 and 7) increases the proteins identity on 3.8% thus making the middle of both genes (and encoded proteins) substantially more different among the both gene paralogs. To corroborate this observation and to explore the importance of other protein parts we have directly compared the sequences encoded by the full-length transcripts and producing Netrin-G1m and Netrin-G2b (**Figure 3C**). Similarly to what has been shown on **Figure 3B-1** and **3B-2**, the lowest identity (17.5%) is represented by the Ukd domain (encoded by the exons 6 and 7) and by the preceding it exon 5 (a 3'-part of the LE1 domain). Two other areas also show a substantially low identity, namely the N-terminus (it includes the protein secretory signal indicated by an arrow) and the outmost C-terminus responsible for the unique feature of Netrin-Gs – the GPI attachment. Thus, based on the percent identity comparisons among the Netrin-G paralogs it can be predicted that there are several potential protein parts contributing to the paralogs SF. As it has been reported by **Seiradake et al. (2011)**, identical gene and protein domain compositions result in the identical structural motif with differences only in the spatial arrangement of the loops facing the post-synaptic Netrin-G's interacting partners, NGL-1 and NGL-2, respectively (**Figure 3D**). Loop I binding surfaces alignment (**Figure 3C**, blue color) shows a high level of conservation (with at least 5 amino acids 100% conserved) among the Netrin-G paralogs, indicating that it is unlikely to be responsible for the cognate ligand binding specificity. Neither Loop II (**Figure 5C**, yellow color) nor Loop III (**Figure 5C**, orange color) display a single conserved amino acid shared among the paralogous binding interfaces (as it originally has been described in **Seiradake et al., 2011**). Thus the complementary pattern of the pre-postsynaptic interactions mediated via specific Netrin-G/NGL pairs is reflected in the reciprocally different sizes of the loops binding interfaces representing another element of the *NTNG*-encoded protein paralogs SF.

DISCUSSION

Complementary contribution of *NTNG* paralogs into human cognitive pathologies.

Involvement of the pre-synaptically expressed axon-localised *NTNGs* in SCZ diagnosis supports the established view of SCZ as a result of distorted trans-synaptic signaling (**Lips et al., 2012**), with a recent study proving that axonal connectivity-associated genes form a functional network visualisable by fMRI (**Richiardi et al., 2015**), and that brain connectivity predicts the level of fluid intelligence (**Finn et al., 2015; Pamplona et al., 2015**). Both *NTNGs* have been found to participate in the brain functional connectivity by the parcellated connectome reconstruction (**Hawrylycs et al., 2015**). Most of the reported disease associations link *NTNG1* to SCZ with a variety of other neurologic pathologies (15 in total, **Figure 1A-1**), while *NTNG2* pathologic associations (6 in total, **Figure 1A-2**) are quite limited to those affecting WM or PO. Among them is SLE frequently characterized by WM deficit (**Shucard et al., 2011**) and also known to represent schizoid-type abnormalities characteristic for autoimmune pathologies (**Guilloux et al., 2010; Eaton et al., 2006**). Immune activation is known to lead to altered pre-pulse inhibition (a key diagnostic trait for SCZ) reversed by antipsychotics (**Romero et al., 2007**). The three diseases associated with both paralogs (ASD, BD and SCZ) are also a primary focus of the recently initiated PsychENCODE project (**PsychENCODE et al., 2015**). It is also worth to mention the resemblance of the reported disease associations with the behavioral phenotypes of *Ntng1* and *Ntng2* gene knockout mice (**Zhang et al., 2016**).

A gene content associated with attenuated IQ score often relates to numerous diseases, such as SCZ, ASD, depression, and others (see **Zhao et al., 2014** for ref.; **Johnson et al., 2016**). Several genes associated with SCZ have undergone positive selection following the human brain evolution (**Xu et al., 2015**). Despite the global network properties of the brain transcriptome are highly conserved among the species there are robust human-specific

disease-associated modules (Miller et al., 2010) and human accelerated regions (HARs) - highly conserved parts of genome that underwent accelerated evolution in humans (Pollard et al., 2006). HARs can serve as genomic markers for human-specific traits underlying a recent acquisition of modern human cognitive abilities by brain (Boyd et al., 2015) but that also “might have led to an increase in structural instability... resulted in a higher risk for neurodegeneration in the aging brain” (Zhou et al., 2015), rendering our intellectual abilities genetically fragile (Crabtree, 2013) and resulting in a variety of CDs. The role genomic context, epistasis (Hemani et al., 2014), plays in the evolution and pathology is manifested by frequently found disease-causing alleles present in animals without obvious pathological symptoms for the host (Jordan et al., 2015). Any CD is characterized by general intellectual disability (GID) plus psychiatric symptoms. A genetic perturbation-exerted behavioral cognitive deficit (BCD) in an animal model organism is a poor match to a human CD *per se* due to very poor contextual resemblance between the human GID and animal BCD together with the absence of interpretable psychiatric symptoms. Usefulness of animals as psychiatric models is also compromised by the fact that transcriptome differences within species tissues is smaller than among the homologous tissues of different species (Barbosa-Morais et al., 2012; Lin et al., 2014). No wonder that the compounds that “cure” mice models consistently fail in human trials (discussed in Hyman, 2014).

NTNG paralogs brain transcriptome intrinsic complementarity and possible mechanism for the IQ-affecting mutation alleles effect. There is no global change at the mRNA level between healthy subjects and SCZ patients (Figure 1B). This conclusion is supported by previously published works stating that globally altered mRNA expression of *NTNG1* or *NTNG2* is unlikely to confer disease susceptibility, at least in the temporal lobe (Eastwood and Harrison, 2008), and Brodmann’s area (Aoki-Suzuki et al., 2005). However, the original paper-source of the STG samples RNA-seq along with many other genes (>1,000)

found that *NTNG1* (but not *NTNG2*) falls under the group of genes with significant alternative promoter usage (**Wu et al., 2012: ST6**, $p < 9.05 \times 10^{-10}$ at FDR < 0.5) and *NTNG2* (but not *NTNG1*) clusters with genes (> 700) with significant alternative splicing change (**Wu et al., 2012: ST7**, $p < 6.15 \times 10^{-12}$ at FDR < 0.5) when SCZ and controls are compared. Such GWAS observation adds an extra layer of complementary regulation to both *NTNG* paralogs on a top of the described in the results section complementary usage rule for the exons, formed unspliced splicing modules, resulting transcripts and their comprising exons (**Figure 1B**). Based on the available RNA-seq dataset it was almost impossible to detect RNA with the matched position of *NTNG* SNPs used for the IQ testing (**ST2c**) except for two coding exons located (rs2149171 and rs3824574) and exon 10 non-coding area located but transcribed rs4915045 (in 2 out of 16 samples). This fact points towards indirect effect of the IQ-affecting mutation alleles potentially associated with shorter (secretable) isoforms generation (**Prosselkov et al., unpublished**) lacking two of the most prominent *NTNG* features: GPI-link and the Ukd domain through an aberrant splicing factor binding. The GPI-link is a hallmark of Netrin-G family members (**Nakashiba et al., 2000, 2002**) and without it the aberrant Netrin-G isoforms are likely to mimic the action of their releasable ancestry molecules - netrins, still being able to bind to their cognate postsynaptic ligand – NGL but without forming an axonal-postsynaptic contact. The Ukd domain of Netrin-G1, despite its so-far unknown function, is involved in lateral binding to the pre-synaptically localised LAR modulating the binding strength between NGL-1 and Netrin-G1 (**Song et al., 2013**). Work is currently underway in search for a similar lateral interaction partner for the Netrin-G2 Ukd domain (**Kim E, personal communications**). The inclusion of Ukd encoding exons 6 and 7 is regulated by the Nova splicing factor (**Ule et al., 2005**) affecting the cortex Netrin-G1 exon 7 but not exon 6, and, simultaneously, Netrin-G2 paralog exons exhibiting an opposite pattern. In general, it is tempting to speculate that deregulation of *NTNG* transcripts

processing may have a role in the brain-controlled cognitive abilities and associated CDs. Supporting such notion, a decreased level of Netrin-G1c mRNA (exons 6-9 excluded, **Figure 1B-1**) has been reported for BD and SCZ (**Eastwood and Harrison, 2008**) with Netrin-G1d (exons 6 and 7 included but 8-9 excluded, **Figure 1B-1**) and Netrin-G1f (a secretable short isoform consisted of domain VI only and lacking the Ukd and GPI-link) being increased in BD, but not in SCZ, in anterior cingulate cortex (**Eastwood and Harrison, 2010**). Higher Netrin-G1d mRNA expression in fetal brain but low for the Netrin-G1c isoform in the human adult (**Eastwood and Harrison, 2008**) indicates different functionality of these two splice variants juggling with the Ukd domain inclusion. And, according to our other data, if Netrin-G1 Ukd-containing isoforms are the dominant isoforms in adult mouse brain, Netrin-G2 Ukd-containing isoforms are present only at the trace level (**Prosselkov et al., forthcoming**), resembling a similar transcriptome pattern for the human STG samples (**Figure 1B-1 and B-2**). A similar “dynamic microexon regulation” associated with the protein interactome misregulation has been reported to be linked to ASD (**Irimia et al., 2014**).

Synchronous and complementary expression of *NTNG* paralogs in the human brain supports the IQ-associated cognitive endophenotypes. Influential parieto-frontal integration theory (P-FIT, **Jung and Haier, 2007**) states that general intelligence (“g”) is dependent on multiple brain cortical areas such as dlPFC, Broca's and Wernicke's areas, somatosensory and visual cortices (**Colom et al., 2009**). Despite “g” is widely accepted as the only correlate of the intelligence, its unitary nature was challenged by (**Hampshire et al., 2012**) claiming had identified two independent brain networks (for memory and for reasoning) responsible for the task performance, the idea later criticised for the employed data processing approach (**Haier et al., 2014**). Higher IQ scores (a composite surrogate of “g”) have been reportedly associated with the fronto-parietal network (FPN) connectivity (**Song et al., 2008; Glascher et al., 2009**). High level of *NTNG* paralogs expression within

the cognition intensively loaded areas of the brain and the distinct patterns of expression profiles (synchronous, asynchronous/mixed, and complementary, **Figure 2A**) support associations of *NTNG1* and *NTNG2* with the recorded cognitive endophenotypes (**Prossellkov et al., 2015**). Based on the expression patterning over the human life-span, among the total 16 analysed brain areas we found two falling under the same “anti-phasic (complementary)” classifier (**Figure 2C**): HIP and MD. Adding more to that, MD is the only brain area (out of the 16 presented) where *NTNG1* expression level exceeds that of *NTNG2* making it a promising candidate for the phenomena of *NTNGs* SF explanation. Two other brain areas classified by a synchronous paralogs expression deserve a special attention, dlPFC and mPFC (**Figure 2A-4**). PFC circuitry has been known as a “hub of the brain’s WM system” (**Kim et al., 2013; Markowitz et al., 2015**), which acts through direct HIP afferents (**Spellman et al., 2015**) and has many connections with other cortical and subcortical areas (**Riga et al., 2014**). mPFC may function as an intelligence-control switchboard and lPFC, part of the FPN global connectivity, predicts the WM performance and fluid intelligence (**Cole et al., 2012**). Interactions of the auditory recognition information fed by the vPFC stream with the sequence processing by the dorsal stream are crucial for the human language articulation (**Skeide and Friederici, 2015; Thothathiri and Rattinger, 2015**). The fact that both *NTNG* paralogs are extensively expressed across PFC (**Figure 2A-2 and A-4**) pinpoints this area as a key for future molecular studies of the human-unique symbolic communications. And PFC is not only implicated in many psychiatric disorders, including SCZ (**Gulsuner and McClellan, 2014**; see also **Riga et al., 2014** for ref.), but is also the only brain structure unique to primates without known homologs in the animal kingdom (**Wise, 2008**).

Evolution of the protein paralogs encoded by the *NTNGs*. Forkhead box P2 (FOXP2) – a ubiquitously expressed transcription factor that has been reported to be linked to the evolution of human language through T303N, N325S substitutions when compared to a

304 primate ortholog (**Enard et al., 2002**) and is 100% identical to Nea protein (**Krause et al.,**
305 **2007**). FOXP2 regulates expression of multiple genes in human and chimpanzee (**Konopka**
306 **et al., 2009**), and among them is an M3 gene brain module representative responsible for
307 general fluid cognitive abilities (**Johnson et al., 2016**), *LRRC4C*, a gene encoding NGL-1 – a
308 post-synaptic target of Netrin-G1. Similarly to FOXP2, Netrin-G1 is a 100% conserved
309 protein among the hominins with only 1 mutation found in chimpanzee which is absent in
310 marmoset (and other primates) and mice proteins (**Prosser et al., 2015**). On the other
311 hand, extinct hominins' Netrin-G2 relatively to modern human contains T346A point
312 mutation (as per current version of hg19), also found in primates and mouse and known as
313 rs4962173 (dbSNP missense mutation) representing an ancient substitution from Neandethal
314 genomes found in modern humans and reflecting a recent acquisition of the novel allele
315 around 5,300 yrs BC. Nothing is known regarding the functional significance of this mutation
316 but biochemically a substitution of alanine (A) on a polar threonine (T) could bring an extra
317 point of regulation, e.g. a phosphorylation or glycosylation (NetPhos2.0 (**Blom et al., 1999**)
318 assigns a low score for the T346 to be phosphorylated but NetOGlyc4.0 (**Steentoft et al.,**
319 **2013**) robustly predicts it to be glycosylated, **SM**). Another mutation S371A/V reflects a
320 selective sweep in Netrin-G2 protein from primates to hominins within a similar to T346A
321 functional context when a hydrophobic alanine (in chimpanzee, A)/valine (in marmoset, V) is
322 replaced by a polar serine (S) and a strong positive predictions for glycosylation but not
323 phosphorylation (**SM**). This poses a question whether these two human-specific protein
324 substitutions associate with advanced cognitive traits as they may represent a hidden layer of
325 poorly studied so far protein glycosylation-associated regulatome known to affect the brain
326 function and diseases (**Baenziger, 2012; Baenziger, 2013**). Adding more to this, T346 is
327 nested on exon 5 just 20 nu away from the affecting WM score rs2274855 (**Prosser et al.,**
328 **2015**), and, together with S371A/V, they are both located within the lowest percent identity

area (exons (5-7)) of Netrin-Gs (**Figure 3C**) and, proposedly, contributing to the *NTNG* duplicates SF. There are at least three more protein parts potentially contributing to the gene paralogs specialised function subdivision (based on the low identity scores, **Figure 3C**): the secretory peptide, the GPI-link, and the outmost structurally elaborated unstructured loops (I-III) responsible for the reciprocal binding of Netrin-Gs to their post-synaptic cognate partners, NGL-1 or NGL-2, both containing a C-terminal PDZ-binding domain (**Kim et al., 2006**). An interesting finding was reported in (**Arbuckle et al., 2010**) found a presence of SH3(PSD95) domain binding site (required for the phosphatidylinositol-3-kinase recruitment) in mice Netrin-G2 (100% identical to human) but not in Netrin-G1. The detected SH3 binding site overlaps with the Netrin-G2-loop III responsible for the binding specificity to NGL-2 (**Seiradake et al., 2011; Soto et al., 2013; DeNardo et al., 2012**). A plausible working hypothesis would be that while internalised (and being GPI-link naïve/immature) the pre-synaptic Netrin-G2 is bound to SH3-PSD95 via loop III but as soon as being secreted extracellularly (and being attached to the membrane) it is bound to post-synaptic NGL-2. Corroborating this, in the absence of Netrin-G2 in the KO mice NGL-2 is unstable on the post-synaptic surface and gets quickly internalised (**Zhang et al., 2016**). We can only speculate regarding the potential importance of PSD-95(SH3)-Netrin-G2-NGL-2 scaffolding loop interaction/competition but the ability for Netrin-G1 to bind to SH3 has not been reported. Following this logic, Netrin-G1 should have a similar binding partner via loop II.

The overall identical structural scaffold among the Netrin-G paralogs (**Figure 3D**) is likely to represent an anciently preserved one of the primordial protein (encoded by a single gene in the primitive urochordate *C.intestinalis*) and its contribution to the process of SF among the *NTNG* paralogs goes against the “structure drives function” concept. It looks like that it is not the “structure” but rather the “evolution” itself that drives a selection for the best structural (or unstructural in our case) fit out of the available frameworks provided by the

gene duplicates to fulfill the emerged functional demand in a new ecological niche. The intricate variability of phenotype is grounded by the conserved nature of genotype and constrained by the “structure-function” limitations of the coding DNA and is only possible due to permissive evolutionary continuing elaborations of non-coding areas able to absorb the most recently acquired elements (having a potential to become regulatory at some point, e.g. like HAR5 (Boyd et al., 2015)) and carried over by neutral drift as proposed by Kimura but for proteins (Kimura, 1983). At the same time, the multiple protein substitutions coinciding with the SF labor segregation phenomena among the Netrin-G paralogs question their neutral nature. Both of them undergo a purifying selection from mice to human through the reduction in size of non-coding DNA (introns) and encoded proteins (the mice Netrin-G2 is 2 aa longer its human ortholog) further contributing to the host-specific SF. Thus while the non-coding sequences are used to explore the evolutionary space in time, the restrictive boundaries of the paralogs SF are determined by the protein (unstructured) elements.

Molecular evolution of the Cognitive Complement (CC). Appearance of the neural crest (Abitua et al., 2012), an event that “affected the chordate evolution in the unprecedented manner” (Green et al., 2015), multipotent progenitor cells (Stolfi et al., 2015), and neurogenic placodes (suggesting a chemosensory and neurosecretory activities, Abitua et al., 2015) in first primitive urochordates/tunicates coincides with the presence of *Ntng* precursor gene (ENSCING00000024925) later undergoing two rounds of duplication events in lamprey and found to affect human cognitive abilities (Prosselkov et al., 2015). *NTNG* paralogs are expressed in the human neural crest-forming cells with *NTNG2* 10 times stronger than *NTNG1* (Rada-Iglesias et al., 2012), both are differentially expressed in human comparing to chimpanzee and rhesus monkey with *NTNG2* expression model showing stronger probability than *NTNG1* (Iskow et al., 2012), and both are stronger expressed in human telencephalon comparing to chimpanzee and macaque (Konopka et al., 2012). *NTNG1* has been classified

as a brain module hub gene “whose pattern fundamentally shifted between species” (Hawrylycs et al., 2015). Belonging to distinct modules of brain expression regulation (Liu et al., 2012, Konopka et al., 2012), *NTNGs* are classified as “genes with human-specific expression profiles” (Liu et al., 2012). The nearby gene ~260 kbp upstream of *NTNG2* is *MED27* (mediator of RNA polymerase II) has been proposed to be associated with the evolution of human-specific traits (McLean et al., 2011). *NTNG1* has also been reported among the “adaptive plasticity genes” (Ghalambor et al., 2015) potentiating rapid adaptive evolution in guppies (*NTNG2* was not found among the input RNA for analysis).

Complementarity among the *NTNG* paralogs and encoded by them proteins has been reported previously: brain expression complementary pattern (in almost self-exclusive manner) defined by the 5'-UTR-localised *cis*-regulatory elements (Yaguchi et al., 2014); complementary distribution within the hippocampal laminar structures (Nishimura-Akiyoshi et al., 2007); axon-dendrite synaptic ending resulting in differential control over the neuronal circuit plasticity (Matsukawa et al., 2014); mutually-exclusive binding pattern to post-synaptic partners, NGL-1 and NGL-2, dictated by the protein unstructural elements (Seiradake et al., 2011); alternative promoter usage vs alternative mRNA splicing (Wu et al., 2012) and increased coefficient of variation (CV, ST1d) for *NTNG1* expression but not *NTNG2* in SCZ patients (similar to Zhang et al., 2015); KO mice behavioral phenotypes and subcellular signaling partners complementarity (Zhang et al., 2016); “differential stability” brain modules expression (*NTNG1* is expressed in the dorsal thalamus (M11) as a hub gene (Pearson’s 0.92) while *NTNG2* is in neocortex and claustrum module (M6, Pearson’s 0.65)) (Hawrylycs et al., 2015); hypocretin neurons-specific expression of *NTNG1* (but not *NTNG2*) as a sleep modulator (Yelin-Bekerman et al., 2015); top-down vs bottom-up information flows gating in mice and differential responsiveness to neuronal stimuli (Prossellkov et al., forthcoming); and human IQ-compiling cognitive domains

complementation (**Prosselkov et al., 2015**). The current study reports the *NTNGs* complementarity association with the CDs (**Figure 1A**); mRNA splicing pattern complementary at the quantitative and qualitative levels via differential use of the middle-located exons (**Figure 1B**); brain complementary oscillatory expression over the human life span observed in the intensive cognitively loaded brain areas (**Figure 2**); AE of the paralogs-segregated unique non-coding elements (**Figure 3A**); complementary pattern of the protein orthologs (mice-to-human) protein sequence evolution. Such multi-level complementation is likely to reflect a shared evolutionary origin from a single gene in a primitive vertebrate organism 700 mln yrs ago and its subsequent functional segregation among the evolution-generated gene duplicates in jawless fish, such as lamprey.

Occupying independent but intercalating functional niches, *NTNG1* and *NTNG2* do not compensate but complement each other's function forming a "functional complement" of genes. Half a billion yrs ago the doubled gene dosage led to the gradual SF and manifested in a function complementation within the cognitive domains, at least in human. We would like to coin such gene pair as a Cognitive Complement (CC).

CONCLUSION

The emerged functional redundancy, as an outcome of gene duplication, leads to function subdivision and its bifurcation among the gene paralogs resulting in the paralogs SF. A functional compensation is known to exist among the evolutionary unrelated genes but has not been reported among the gene paralogs, more frequently characterized by the function complementation. Gene paralogs structural identity (at both, gene and protein levels) does not provide a substrate for functional compensation but rather for complementation, perturbing "structure drives function" rule. A gene duplication event of a tunicate *NTNG* primordial gene and the subsequent process of its function specialisation (driven by the new ecological

429 niches appearance and evolution) among the gene duplicates made them to SF into distinct
430 cognitive domains in a complementary manner forming a CC. In our forthcoming work we
431 are to describe how *Ntn* mice genes function resembles that of human orthologs (**Prosselkov**
432 **et al., forthcoming**).
433

MATERIALS AND METHODS

Human brain *NTNG* transcriptome reconstruction. Relates to **Figure 1B** and **1-C**. The original source of the dataset was produced by (Wu et al., 2012: E-MATB-1030) and the downloaded .bam files used for the re-processing are listed in **ST1a**. All reconstructed transcripts are presented in **ST1d** standalone Excel file. Two samples were excluded from the analysis due to failed “per base sequence quality” measure, and zero expression level for *NTNG1a* and *NTNG1int(9-10)* otherwise consistently expressed throughout other samples (**ST1b**). SAMtools software was used for the SNPs calling from the available RNA-seq datasets (**ST1c**). For details refer to **SM**.

Human brain expression profiling for *NTNGs* across the life span. The original source of data was www.brainspan.org. All available samples were initially included into the analysis but two of them excluded at a later stage (MD for 12-13 pcw and mPFC for 16-19 pcw) due to high deviation (6-7 times) from the mean for other replicas. The mean expression values per each brain area as RPKM were plotted against the sampling age. Profiles classification was done visually considering the trend over the all plotted points as an average.

***NTNG1 (NTNG1m)* and *NTNG2 (NTNG2b)* full-length mRNA transcripts assembly.** Relates to **Figure 3B**. Human *NTNG1m* brain transcript has been reported previously (Meerabux et al., 2005) and we have also confirmed its ortholog presence in the mice brain via full-length cloning (Prosselkov et al., unpublished). Since NCBI contains only its partial CDS (AY764265), we used the RNA-seq-generated exons (**Figure 1B**) to reconstruct its full-length and to generate an ORF of the encoded Netrin-G1m. Similarly, human *NTNG2b* was reconstructed from the RNA-seq dataset and from Ensemble as follows. Exon 5 sequence was deduced from ENST00000372179, other exons were from ENST00000467453 (no longer available on the current version of Ensemble) except for exon 6 deduced by running three independent alignments against the human genomic DNA with the mice 3'-intron (5-6),

exon 6, and 5'-intron (6-7) concomitantly confirmed by the generated full-length ORF for Netrin-G2b. The reconstructed protein was predicted to encode 587 amino acids, which is in a close proximity to the mice netrin-G2b ortholog of 589 residues (**Prosselkov et al., forth.**).

Full-lengths gene structures of *NTNG* paralogs reconstruction. Relates to **Figure 3A.**

Both, the obtained above from the STG brain samples RNA-seq and the reconstructed full-lengths transcripts carrying all stably expressed exons were used to confirm the intron-exon junctions positioning for *NTNG1* and *NTNG2*. Due to observed variability in the intron (1-2) and exon 10 sizes their boundaries were left unmarked.

SUPPLEMENTARY MATERIALS (SM)

Contain Supplementary Methods (RNA-seq of STG re-processing and SNPs detection) and Supplementary Tables (**ST1a-d, ST2**) as a single compiled pdf file. Reconstructed RNA-seq (.gtf) of the STG is presented as a standalone Excel file (**ST1d**). Also included: Netrin-G2b predicted phosphorylation and *O*-glycosylation, Netrin-G1 vs Netrin-G2 Ukd alignment (**McWilliam et al., 2013**), predicted secretory peptide cleavage and GPI attachments.

ACKNOWLEDGEMENTS

This work was in part supported by the “Funding Program for World-Leading Innovative R&D on Science and Technology (FIRST Program)” initiated by the Council for Science and Technology Policy (CSTP), and KAKENHI 15H04290 from the Japan Society for the Promotion of Science (JSPS).

COMPETING INTERESTS

Authors would like to express a lack of any competing interests associated with the work.

REFERENCES

- Abitua PB**, Wagner E, Navarrete IA, Levine M. 2012. Identification of a rudimentary neural crest in a non-vertebrate chordate. *Nature* **492**:104-107. doi: [10.1038/nature11589](https://doi.org/10.1038/nature11589).
- Abitua PB**, Gainous TB, Kaczmarczyk AN, Winchell CJ, Hudson C, Kamata K, Nakagawa M, Tsuda M, Kusakabe, TG, Levine M. 2015. The pre-vertebrate origins of neurogenic placodes. *Nature* **524**:462-465. doi: [10.1038/nature14657](https://doi.org/10.1038/nature14657).
- Akula N**, Barb, J, Jiang, X, Wendland, JR, Choi, KH, Sen, SK, Hou, L, Chen, DTW, Laje, G, Johnson, K, Lipska, BK, Kleinman, JE, Corrada-Bravo, H, Detera-Wadleigh, S, Munson, PJ, McMahon, FJ. 2014. RNA-sequencing of the brain transcriptome implicates dysregulation of neuroplasticity, circadian rhythms and GTPase binding in bipolar disorder. *Molecular Psychiatry* **19**:1179-1185. doi: [10.1038/mp.2013.170](https://doi.org/10.1038/mp.2013.170).
- Aoki-Suzuki M**, Yamada K, Meerabux J, Iwayama-Shigeno Y, Ohba H, Iwamoto K, Takao H, Toyota T, Suto Y, Nakatani N, Dean B, Nishimura S, Seki K, Kato T, Itohara S, Nishikawa T, Yoshikawa, T. 2005. A Family-Based Association Study and Gene Expression Analyses of Netrin-G1 and -G2 Genes in Schizophrenia. *Biological Psychiatry* **57**:382–393. doi: [10.1016/j.biopsych.2004.11.022](https://doi.org/10.1016/j.biopsych.2004.11.022).
- Arbuckle MI**, Komiyama NH, Delaney A, Coba M, Garry EM, Rosie R, Allchorne AJ, Forsyth LH, Bence M, Carlisle HJ, O'Dell TJ, Mitchell R, Fleetwood-Walker SM, Grant SGN. 2010. The SH3 domain of postsynaptic density 95 mediates inflammatory pain through phosphatidylinositol-3-kinase recruitment. *Embo Reports* **11**:473-478. doi: [10.1038/embor.2010.63](https://doi.org/10.1038/embor.2010.63).
- Archer HL**, Evans JC, Millar DS, Thompson PW, Kerr AM, Leonard H, Christodoulou J, Ravine D, Lazarou L, Grove L, Verity C, Whatley SD, Pitz DT, Sampson JR, Clarke AJ. 2006. *NTNG1* mutations are a rare cause of Rett syndrome. *American Journal of Medical Genetics Part A* **140A**:691-694. doi: [10.1002/ajmg.a.31133](https://doi.org/10.1002/ajmg.a.31133).

- Ayalew M**, Le-Niculescu H, Levey DF, Jain N, Changala B, Patel SD, Winiger E, Breier A, Shekhar A, Amdur R, Koller D, Nurnberger JI, Corvin A, Geyer M, Tsuang MT, Salomon D, Schork NJ, Fanous AH, O'Donovan MC, Niculescu AB. 2012. Convergent functional genomics of schizophrenia: from comprehensive understanding to genetic risk prediction. *Molecular Psychiatry* **17**:887-905. doi: [10.1038/mp.2012.37](https://doi.org/10.1038/mp.2012.37).
- Baenziger JU**. 2012. Moving the *O*-glycoproteome from form to function. *Proceedings of the National Academy of Sciences of the United States of America* **109**:9672–9673. doi: [10.1073/pnas.1206735109](https://doi.org/10.1073/pnas.1206735109).
- Baenziger JU**. 2013. *O*-mannosylation of cadherins. *Proceedings of the National Academy of Sciences of the United States of America* **110**:20858-20859. doi: [10.1073/pnas.1321827111](https://doi.org/10.1073/pnas.1321827111).
- Barbosa-Morais NL**, Irimia M, Pan Q, Xiong HY, Gueroussov S, Lee LJ, Slobodeniuc V, Kutter C, Watt S, Çolak R, Kim T, Misquitta-Ali CM, Wilson MD, Kim PM, Odom DT, Frey BJ, Blencowe BJ. 2012. The Evolutionary Landscape of Alternative Splicing in Vertebrate Species. *Science* **338**:1587-1593. doi: [10.1126/science.1230612](https://doi.org/10.1126/science.1230612).
- Bisgaard A-M**, Rasmussen LN, Moller HU, Kirchhoff M, Bryndorf T. 2007. Interstitial deletion of the short arm of chromosome 1 (1p13.1p21.1) in a girl with mental retardation, short stature and colobomata. *Clinical Dysmorphology* **16**:109-112. doi: [10.1097/01.mcd.0000228425.89660.bf](https://doi.org/10.1097/01.mcd.0000228425.89660.bf).
- Blom N**, Gammeltoft S, Brunak S. 1999. Sequence and structure-based prediction of eukaryotic protein phosphorylation sites. *Journal of Molecular Biology* **294**:1351-1362. doi: [10.1006/jmbi.1999.3310](https://doi.org/10.1006/jmbi.1999.3310).
- Boraska V**, Franklin CS, Floyd JAB, Thornton LM, Huckins LM, Southam L, Rayner NW, Tachmazidou I, Klump KL, Treasure J, Lewis CM, Schmidt U, Tozzi F, Kiezebrink K, Hebebrand J, Gorwood P, Adan RAH, Kas MJH, Favaro A, Santonastaso P, Fernández-

532 Aranda F, Gratacos M, Rybakowsk F, Dmitrzak-Weglarz M, Kaprio J, Keski-Rahkonen A,
533 Raevuori A, Van Furth EF, Slof-Op 't Landt MCT, Hudson JJ, Reichborn-Kjennerud T,
534 Knudsen GPS, Monteleone P, Kaplan AS, Karwautz A, Hakonarson H, Berrettini WH,
535 Guo Y, Li D, Schork NJ, Komaki G, Ando T, Inoko H, Esko T, Fischer K, Männik K,
536 Metspalu A, Baker JH, Cone RD, Dackor J, DeSocio JE, Hilliard CE, O'Toole JK, Pantel
537 J, Szatkiewicz JP, Taico C, Zerwas S, Trace SE, Davis OSP, Helder S, Bühren K,
538 Burghardt R, de Zwaan M, Egberts K, Ehrlich S, Herpertz-Dahlmann B, Herzog W,
539 Imgart H, Scherag A, Scherag S, Zipfel S, Boni C, Ramoz N, Versini A, Brandys MK,
540 Danner UN, de Kovel C, Hendriks J, Koeleman BPC, Ophoff RA, Strengman E, van
541 Elburg AA, Bruson A, Clementi M, Degortes D, Forzan M, Tenconi E, Docampo E,
542 Escaramís G, Jiménez-Murcia S, Lissowska J, Rajewski A, Szeszenia-Dabrowska N,
543 Slopian A, Hauser J, Karhunen L, Meulenbelt I, Slagboom PE, Tortorella A, Maj M,
544 Dedoussis G, Dikeos D, Gonidakis F, Tziouvas K, Tsitsika A, Papezova H, Slachtova L,
545 Martaskova D, Kennedy JL, Levitan RD, Yilmaz Z, Huemer J, Koubek D, Merl E,
546 Wagner G, Lichtenstein P, Breen G, Cohen-Woods S, Farmer A, McGuffin P, Cichon S,
547 Giegling I, Herms S, Rujescu D, Schreiber S, Wichmann H-E, Dina C, Sladek R,
548 Gambaro G, Soranzo N, Julia A, Marsal S, Rabionet R, Gaborieau V, Dick DM, Palotie A,
549 Ripatti S, Widén E, Andreassen OA, Espeseth T, Lundervold A, Reinvang I, Steen VM,
550 Le Hellard S, Mattingdal M, Ntalla I, Bencko V, Foretova L, Janout V, Navratilova M,
551 Gallinger S, Pinto D, Scherer SW, Aschauer H, Carlberg L, Schosser A, Alfredsson L,
552 Ding B, Klareskog L, Padyukov L, Courtet P, Guillaume S, Jaussent I, Finan C, Kalsi G,
553 Roberts M, Logan DW, Peltonen L, Ritchie GRS, Barrett JC, The Wellcome Trust Case
554 Control Consortium 3, Estivill X, Hinney A, Sullivan PF, Collier DA, Zeggini E, Bulik
555 CM. 2014. A genome-wide association study of anorexia nervosa. *Molecular Psychiatry*
556 **19**:1084-1094. doi: 1085–1094. 10.1038/mp.2013.187.

Borg I, Freude K, Kubart SK, Hoffmann K, Menzel C, Laccone F, Firth H, Ferguson-Smith MA, Tommerup N, Ropers HH, Sargan D, Kalscheuer VM. 2005. Disruption of Netrin G1 by a balanced chromosome translocation in a girl with Rett syndrome. *European Journal of Human Genetics* **13**:921-927. doi: [10.1038/sj.ejhg.5201429](https://doi.org/10.1038/sj.ejhg.5201429).

Boyd JL, Skove SL, Rouanet JP, Pilaz L-J, Bepler T, Gordân R, Wray GA, Silver DL. 2015. Human-Chimpanzee Differences in a FZD8 Enhancer Alter Cell-Cycle Dynamics in the Developing Neocortex. *Current Biology* **25**:772-779. doi: [10.1016/j.cub.2015.01.041](https://doi.org/10.1016/j.cub.2015.01.041).

Cavard C, Audebourg A, Letourneur F, Audard V, Beuvon F, Cagnard N, Radenen, B, Varlet P, Vacher-Lavenu M-C, Perret C, Terris B. 2009. Gene expression profiling provides insights into the pathways involved in solid pseudopapillary neoplasm of the pancreas. *Journal of Pathology* **218**:201-209. doi: [10.1002/path.2524](https://doi.org/10.1002/path.2524).

Cole MW, Yarkoni T, Repovs G, Anticevic A, Braver TS. 2012. Global Connectivity of Prefrontal Cortex Predicts Cognitive Control and Intelligence. *Journal of Neuroscience* **32**:8988-8999. doi: [10.1523/jneurosci.0536-12.2012](https://doi.org/10.1523/jneurosci.0536-12.2012).

Colom R, Haier RJ, Head K, Álvarez-Linerac J, Ángeles Quiroga M, Shih PC, Jung RE. 2009. Gray matter correlates of fluid, crystallized, and spatial intelligence: Testing the P-FIT model. *Intelligence* **37**:124-135. doi: [10.1016/j.intell.2008.07.007](https://doi.org/10.1016/j.intell.2008.07.007).

Crabtree GR. 2013. Our fragile intellect. Part I. *Trends in Genetics* **29**:1-3. doi: [10.1016/j.tig.2012.10.002](https://doi.org/10.1016/j.tig.2012.10.002).

DeNardo LA, de Wit J, Otto-Hitt S, Ghosh A. 2012. NGL-2 Regulates Input-Specific Synapse Development in CA1 Pyramidal Neurons. *Neuron* **76**:762-775. doi: [10.1016/j.neuron.2012.10.013](https://doi.org/10.1016/j.neuron.2012.10.013).

D’Gama AM, Pochareddy S, Li M, Jamuar SS, Reiff RE, Lam A-TN, Sestan N, Walsh CA. 2015. Targeted DNA Sequencing from Autism Spectrum Disorder Brains Implicates Multiple Genetic Mechanisms. *Neuron* **88**:910-917. doi: [10.1016/j.neuron.2015.11.009](https://doi.org/10.1016/j.neuron.2015.11.009).

- 582 **Eastwood SL**, Harrison PJ. 2008. Decreased mRNA expression of netrin-G1 and netrin-G2
583 in the temporal lobe in schizophrenia and bipolar disorder. *Neuropsychopharmacology*
584 **33**:933-945. doi: [10.1038/sj.npp.1301457](https://doi.org/10.1038/sj.npp.1301457).
- 585 **Eastwood SL**, Harrison PJ. 2010. Markers of Glutamate Synaptic Transmission and
586 Plasticity Are Increased in the Anterior Cingulate Cortex in Bipolar Disorder. *Biological*
587 *Psychiatry* **67**:1010-1016. doi: [10.1016/j.biopsych.2009.12.004](https://doi.org/10.1016/j.biopsych.2009.12.004).
- 588 **Eaton WW**, Byrne M, Ewald H, Mors O, Chen CY, Agerbo E, Mortensen PB. 2006.
589 Association of schizophrenia and autoimmune diseases: Linkage of Danish National
590 Registers. *American Journal of Psychiatry* **163**:521-528. doi: [10.1176/appi.ajp.163.3.521](https://doi.org/10.1176/appi.ajp.163.3.521).
- 591 **Eisenhaber B**, Bork P, Yuan YP, Löffler G, Eisenhaber F. 2000. Automated annotation of
592 GPI anchor sites: case study *C. elegans*. *Trends in Biochemical Sciences* **25**:340-341. doi:
593 [10.1016/s0968-0004\(00\)01601-7](https://doi.org/10.1016/s0968-0004(00)01601-7).
- 594 **Enard W**, Przeworski M, Fisher SE, Lai CSL, Wiebe V, Kitano T, Monaco AP, Pääbo S.
595 2002. Molecular evolution of *FOXP2*, a gene involved in speech and language. *Nature*
596 **418**:869-872. doi: [10.1038/nature01025](https://doi.org/10.1038/nature01025).
- 597 **Finn ES**, Shen X, Scheinost D, Rosenberg MD, Huang J, Chun MM, Papademetris X,
598 Constable RT. 2015. Functional connectome fingerprinting: identifying individuals using
599 patterns of brain connectivity. *Nature Neuroscience* **online**. doi:[10.1038/nn.4135](https://doi.org/10.1038/nn.4135).
- 600 **Forbes NF**, Carrick LA, McIntosh AM, Lawrie SM. 2009. Working memory in
601 schizophrenia: a meta-analysis. *Psychological Medicine* **39**:889-905. doi:
602 [10.1017/s0033291708004558](https://doi.org/10.1017/s0033291708004558).
- 603 **Fukasawa M**, Aoki M, Yamada K, Iwayama-Shigeno Y, Takao H, Meerabux J, Toyota T,
604 Nishikawa T, Yoshikawa T. 2004. Case-control association study of human netrin G1
605 gene in Japanese schizophrenia. *Journal of Medical and Dental Sciences* **51**: 121-128.
- 606 **Ghalambor CK**, Hoke KL, Ruell EW, Fischer EK, Reznick DN, Hughes KA. 2015. Non-

adaptive plasticity potentiates rapid adaptive evolution of gene expression in nature.

Nature **525**:372-375. doi: [10.1038/nature15256](https://doi.org/10.1038/nature15256).

Gilissen C, Hehir-Kwa JY, Thung DT, van de Vorst M, van Bon BWM, Willemsen MH, Kwint M, Janssen IM, Hoischen A, Schenck A, Leach R, Klein R, Tearle R, Bo T, Pfundt R, Yntema HG, de Vries BBA, Kleefstra T, Brunner HG, Vissers LELM, Veltman JA. 2014. Genome sequencing identifies major causes of severe intellectual disability. *Nature* **511**:344-347. doi: [10.1038/nature13394](https://doi.org/10.1038/nature13394).

Glascher J, Tranel D, Paul LK, Rudrauf D, Rorden C, Hornaday A, Grabowski T, Damasio H, Adolphs R. 2009. Lesion Mapping of Cognitive Abilities Linked to Intelligence. *Neuron* **61**:681-691. doi: [10.1016/j.neuron.2009.01.026](https://doi.org/10.1016/j.neuron.2009.01.026).

Green SA, Simoes-Costa M, Bronner ME. 2015. Evolution of vertebrates as viewed from the crest. *Nature* **520**:474-482. doi: [10.1038/nature14436](https://doi.org/10.1038/nature14436).

Guilloux JP, Gaiteri C, Sibille E. 2010. Network analysis of positional candidate genes of schizophrenia highlights ... more than ... myelin-related pathways. *Molecular Psychiatry* **15**:786-788. doi: [10.1038/mp.2009.68](https://doi.org/10.1038/mp.2009.68).

Gulsuner S and McClellan JM. 2014. *De Novo* Mutations in Schizophrenia Disrupt Genes Co-Expressed in Fetal Prefrontal Cortex. *Neuropsychopharmacology* **39**:238-239. doi: [10.1038/npp.2013.219](https://doi.org/10.1038/npp.2013.219).

Haier RJ, Karama S, Colom R, Jung R, Johnson W. 2014. A comment on "Fractionating Intelligence" and the peer review process. *Intelligence* **46**:323-332. doi: [10.1016/j.intell.2014.02.007](https://doi.org/10.1016/j.intell.2014.02.007).

Hampshire A, Highfield RR, Parkin BL, Owen AM. 2012. Fractionating Human Intelligence. *Neuron* **76**:1225-1237. doi: [10.1016/j.neuron.2012.06.022](https://doi.org/10.1016/j.neuron.2012.06.022).

Hawrylycz M, Miller JA, Menon V, Feng D, Dolbeare T, Guillozet-Bongaarts AL, Jegga AG, Aronow BJ, Lee C-K, Bernard A, Glasser MF, Dierker DL, Menche J, Szafer A,

632 Collman F, Grange P, Berman KA, Mihalas S, Yao Z, Stewart L, Barabási A-L, Schulkin
633 J, Phillips J, Ng L, Dang C, Haynor DR, Jones A, Van Essen DC, Koch C, Lein E. 2015.
634 Canonical genetic signatures of the adult human brain. *Nature Neuroscience* **18**:1832-
635 1844. doi: [10.1038/nn.4171](https://doi.org/10.1038/nn.4171).

636 **Hemani G**, Shakhbazov K, Westra H-J, Esko T, Henders AK, McRae AF, Yang J, Gibson G,
637 Martin NG, Metspalu A, Franke L, Montgomery GW, Visscher PM, Powell JE. 2014.
638 Detection and replication of epistasis influencing transcription in humans. *Nature*
639 **508**:249–253. doi: [10.1038/nature13005](https://doi.org/10.1038/nature13005).

640 **Hyman SE**. 2014. How Far Can Mice Carry Autism Research? *Cell* **158**:13-14. doi:
641 [10.1016/j.cell.2014.06.032](https://doi.org/10.1016/j.cell.2014.06.032).

642 **Iossifov I**, O’Roak BJ, Sanders SJ, Ronemus M, Krumm N, Levy D, Stessman HA,
643 Witherspoon KT, Vives L, Patterson KE, Smith JD, Paepers B, Nickerson DA, Dea J,
644 Dong S, Gonzalez LE, Mandell JD, Mane SM, Murtha MT, Sullivan CA, Walker MF,
645 Waqar Z, Wei L, Willsey AJ, Yamrom B, Lee Y, Grabowska E, Dalkic E, Wang Z, Marks
646 S, Andrews P, Leotta A, Kendall J, Hakker I, Rosenbaum J, Ma B, Rodgers L, Troge J,
647 Narzisi G, Seungtae Y, Schatz MC, Kenny Y, McCombie WR, Shendure J, Eichler EE,
648 State MW, Wigler M. 2014. The contribution of *de novo* coding mutations to autism
649 spectrum disorder. *Nature* **515**:216–221. doi: [10.1038/nature13908](https://doi.org/10.1038/nature13908).

650 **Irimia M**, Weatheritt RJ, Ellis JD, Parikshak NN, Gonatopoulos-Pournatzis T, Babor M,
651 Quesnel-Vallières M, Tapial J, Raj B, O’Hanlon D, Barrios-Rodiles M, Sternberg MJE,
652 Cordes SP, Roth FP, Wrana JL, Geschwind DH, Blencowe BJ. 2014. A Highly Conserved
653 Program of Neuronal Microexons Is Misregulated in Autistic Brains. *Cell* **159**:1511-1523.
654 doi: [10.1016/j.cell.2014.11.035](https://doi.org/10.1016/j.cell.2014.11.035).

655 **Ishigaki S**, Masuda A, Fujioka Y, Iguchi Y, Katsuno M, Shibata A, Urano F, Sobue G, Ohno
656 K. 2012. Position-dependent FUS-RNA interactions regulate alternative splicing events

and transcriptions. *Scientific Reports* **2**: A529. doi: [10.1038/srep00529](https://doi.org/10.1038/srep00529).

Iskow RC, Gokcumen O, Abyzov A, Malukiewicz J, Zhu Q, Sukumara AT, Pai AA, Mills RE, Habegger L, Cusanovich DA, Rubel MA, Perry GH, Gerstein M, Stone AC, Gilade Y, Lee C. 2012. Regulatory element copy number differences shape primate expression profiles. *Proceedings of the National Academy of Sciences of the United States of America*. **109**:12656–12661. doi: [10.1073/pnas.1205199109](https://doi.org/10.1073/pnas.1205199109).

Johnson MR, Shkura K, Langley SR, Delahaye-Duriez A, Srivastava P, Hill WD, Rackham OJL, Davies G, Harris SE, Moreno-Moral A, Rotival M, Speed D, Petrovski S, Katz A, Hayward C, Porteous DJ, Smith BH, Padmanabhan S, Hocking LJ, Starr JM, Liewald DC, Visconti A, Falchi M, Bottolo L, Rossetti T, Danis B, Mazzaferri M, Foerch P, Grote A, Helmstaedter C, Becker AJ, Kaminski RM, Deary IJ, Petretto E. 2016. Systems genetics identifies a convergent gene network for cognition and neurodevelopmental disease. *Nature Neuroscience* **19**:223-232. doi: [10.1038/nn.4205](https://doi.org/10.1038/nn.4205).

Jordan DM, Frangakis SG, Golzio C, Cassa CA, Kurtzberg J, Task Force for Neonatal Genomics, Davis EE, Sunyaev SR, Katsanis N. 2015. Identification of *cis*-suppression of human disease mutations by comparative genomics. *Nature* **524**:225-229. doi: [10.1038/nature14497](https://doi.org/10.1038/nature14497).

Jung RE and Haier RJ. 2007. The Parieto-Frontal Integration Theory (P-FIT) of intelligence: Converging neuroimaging evidence. *Behavioral and Brain Sciences* **30**:135-187. doi: [10.1017/s0140525x07001185](https://doi.org/10.1017/s0140525x07001185).

Kim J, Ghim J-W, Lee JH, Jung MW. 2013. Neural Correlates of Interval Timing in Rodent Prefrontal Cortex. *Journal of Neuroscience* **33**:13834-13847. doi: [10.1523/jneurosci.1443-13.2013](https://doi.org/10.1523/jneurosci.1443-13.2013).

Kim S, Burette A, Chung HS, Kwon SK, Woo J, Lee HW, Kim K, Kim H, Weinberg RJ, Kim E. 2006. NGL family PSD-95-interacting adhesion molecules regulate excitatory

synapse formation. *Nature Neuroscience* **9**:1294-1301. doi: [10.1038/nn1763](https://doi.org/10.1038/nn1763).

Kimura M. 1983. The neutral theory of molecular evolution. Cambridge University Press.

King IF, Yandava CN, Mabb AM, Hsiao JS, Huang HS, Pearson BL, Calabrese JM, Starmer J, Parker JS, Magnuson T, Chamberlain SJ, Philpot BD, Zylka MJ. 2013. Topoisomerases facilitate transcription of long genes linked to autism. *Nature* **501**:58-62. doi: [10.1038/nature12504](https://doi.org/10.1038/nature12504).

Konopka G, Bomar JM, Winden K, Coppola G, Jonsson ZO, Gao FY, Peng S, Preuss TM, Wohlschlegel JA, Geschwind DH. 2009. Human-specific transcriptional regulation of CNS development genes by FOXP2. *Nature* **462**:213-217. doi: [10.1038/nature08549](https://doi.org/10.1038/nature08549).

Konopka G, Friedrich T, Davis-Turak J, Winden K, Oldham MC, Gao F, Chen L, Wang G-Z, Luo R, Preuss TM, Geschwind DH. 2012. Human-Specific Transcriptional Networks in the Brain. *Neuron* **23**:601-617. doi: [10.1016/j.neuron.2012.05.034](https://doi.org/10.1016/j.neuron.2012.05.034).

Krause J, Lalueza-Fox C, Orlando L, Enard W, Green RE, Burbano HA, Hublin J-J, Hänni C, Fortea J, de la Rasilla M, Bertranpetit J, Rosas A, Pääbo S. 2007. The derived FOXP2 variant of modern humans was shared with neandertals. *Current Biology* **17**:1908-1912. doi: [10.1016/j.cub.2007.10.008](https://doi.org/10.1016/j.cub.2007.10.008).

Lagier-Tourenne C, Polymenidou M, Hutt KR, Vu AQ, Baughn M, Huelga SC, Clutario KM, Ling S-C, Liang TY, Mazur C, Wancewicz E, Kim AS, Watt A, Freier S, Hicks GG, Donohue JP, Shiue L, Bennett CF, Ravits J, Cleveland DW, Yeo GW. 2012. Divergent roles of ALS-linked proteins FUS/TLS and TDP-43 intersect in processing long pre-mRNAs. *Nature Neuroscience* **15**:1488-1497. doi: [doi:10.1038/nn.3230](https://doi.org/10.1038/nn.3230).

Lee SH, Ripke S, Neale BM, Faraone SV, Purcell SM, Perlis RH, Mowry BJ, Thapar A, Goddard ME, Witte JS, Absher D, Agartz I, Akil H, Amin F, Andreassen OA, Anjorin A, Anney R, Anttila V, Arking DE, Asherson P, Azevedo MH, Backlund L, Badner JA, Bailey AJ, Banaschewski T, Barchas JD, Barnes MR, Barrett TB, Bass N, Battaglia A,

707 Bauer M, Bayés M, Bellivier F, Bergen SE, Berrettini W, Betancur C, Bettecken T,
708 Biederman J, Binder EB, Black DW, Blackwood DHR, Bloss CS, Boehnke M, Boomsma
709 DI, Breen G, Breuer R, Bruggeman R, Cormican P, Buccola NG, Buitelaar JK, Bunney
710 WE, Buxbaum JD, Byerley WF, Byrne EM, Caesar S, Cahn W, Cantor RM, Casas M,
711 Chakravarti A, Chambert K, Choudhury K, Cichon S, Cloninger CR, Collier DA, Cook E
712 H, Coon H, Cormand B, Corvin A, Coryell WH, Craig DW, Craig IW, Crosbie J, Cuccaro
713 ML, Curtis D, Czamara D, Datta S, Dawson G, Day R, De Geus EJ, Degenhardt F,
714 Djurovic S, Donohoe GJ, Doyle AE, Duan JB, Dudbridge F, Duketis E, Ebstein RP,
715 Edenberg HJ, Elia J, Ennis S, Etain B, Fanous A, Farmer AE, Ferrier IN, Flickinger M,
716 Fombonne E, Foroud T, Frank J, Franke B, Fraser C, Freedman R, Freimer NB, Freitag C
717 M, Friedl M, Frisén L, Gallagher L, Gejman PV, Georgieva L, Gershon ES, Geschwind
718 DH, Giegling I, Gill M, Gordon SD, Gordon-Smith K, Green EK, Greenwood TA, Grice
719 DE, Gross M, Grozeva D, Guan WH, Gurling H, De Haan L, Haines JL, Hakonarson H,
720 Hallmayer J, Hamilton SP, Hamshere ML, Hansen TF, Hartmann AM, Hautzinger M,
721 Heath AC, Henders AK, Herms S, Hickie IB, Hipolito M, Hoefels S, Holmans PA,
722 Holsboer F, Hoogendijk WJ, Hottenga JJ, Hultman CM, Hus V, Ingason A, Ising M,
723 Jamain S, Jones EG, Jones I, Jones L, Tzeng JY, Kähler AK, Kahn RS, Kandaswamy R,
724 Keller MC, Kennedy JL, Kenny E, Kent L, Kim Y, Kirov GK, Klauck SM, Klei L,
725 Knowles JA, Kohli MA, Koller DL, Konte B, Korszun A, Krabbendam L, Krasucki R,
726 Kuntsi J, Kwan P, Landen M, Långström N, Lathrop M, Lawrence J, Lawson WB,
727 Leboyer M, Ledbetter DH, Lee PH, Lencz T, Lesch KP, Levinson DF, Lewis CM, Li J,
728 Lichtenstein P, Lieberman JA, Lin DY, Linszen DH, Liu CY, Lohoff FW, Loo SK, Lord
729 C, Lowe JK, Lucae S, MacIntyre DJ, Madden PAF, Maestrini E, Magnusson PKE, Mahon
730 PB, Maier W, Malhotra AK, Mane SM, Martin CL, Martin NG, Mattheisen M, Matthews
731 K, Mattingsdal M, McCarroll SA, McGhee KA, McGough JJ, McGrath PJ, McGuffin P,

732 McInnis MG, McIntosh A, McKinney R, McLean AW, McMahon FJ, McMahon WM,
733 McQuillin A, Medeiros H, Medland SE, Meier S, Melle I, Meng F, Meyer J, Middeldorp
734 CM, Middleton L, Milanova V, Miranda A, Monaco AP, Montgomery GW, Moran JL,
735 Moreno-De-Luca D, Morken G, Morris DW, Morrow EM, Moskvina V, Muglia P,
736 Mühleisen TW, Muir WJ, Muller-Myhsok B, Murtha M, Myers RM, Myin-Germeys I,
737 Neale MC, Nelson SF, Nievergelt CM, Nikolov I, Nimgaonkar V, Nolen WA, Nöthen
738 MM, Nurnberger JI, Nwulia EA, Nyholt DR, O'Dushlaine C, Oades RD, Olincy A,
739 Oliveira G, Olsen L, Ophoff RA, Osby U, Owen MJ, Palotie A, Parr JR, Paterson AD,
740 Pato CN, Pato MT, Penninx BW, Pergadia ML, Pericak-Vance MA, Pickard BS, Pimm J,
741 Piven J, Posthuma D, Potash JB, Poustka F, Propping P, Puri V, Quedstedt DJ, Quinn EM,
742 Ramos-Quiroga JA, Rasmussen HB, Raychaudhuri S, Rehnström K, Reif A, Ribasés M,
743 Rice JP, Rietschel M, Roeder K, Roeyers H, Rossin L, Rothenberger A, Rouleau G,
744 Ruderfer D, Rujescu D, Sanders AR, Sanders SJ, Santangelo SL, Sergeant JA, Schachar R,
745 Schalling M, Schatzberg AF, Scheftner WA, Schellenberg GD, Scherer SW, Schork NJ,
746 Schulze TG, Schumacher J, Schwarz M, Scolnick E, Scott LJ, Shi JX, Shilling PD, Shyn
747 SI, Silverman JM, Slager SL, Smalley SL, Smit JH, Smith EN, Sonuga-Barke EJS, St
748 Clair D, State M, Steffens M, Steinhausen HC, Strauss JS, Strohmaier J, Stroup TS,
749 Sutcliffe JS, Szatmari P, Szelinger S, Thirumalai S, Thompson RC, Todorov AA, Tozzi F,
750 Treutlein J, Uhr M, van den Oord EJCG, Van Grootheest G, Van Os J, Vicente AM,
751 Vieland VJ, Vincent JB, Visscher PM, Walsh CA, Wassink TH, Watson SJ, Weissman
752 MM, Werge T, Wienker TF, Wijsman EM, Willemsen G, Williams N, Willsey AJ, Witt
753 SH, Xu W, Young AH, Yu TW, Zammit S, Zandi PP, Zhang P, Zitman FG, Zöllner S,
754 International Inflammatory Bowel Disease Genetics Consortium (IIBDGC), Devlin B,
755 Kelsoe JR, Sklar P, Daly MJ, O'Donovan MC, Craddock N, Sullivan PF, Smoller JW,
756 Kendler KS, Wray NR. 2013. Genetic relationship between five psychiatric disorders

estimated from genome-wide SNPs. *Nature Genetics* **45**:984-994. doi: [10.1038/ng.2711](https://doi.org/10.1038/ng.2711).

Lesnick TG, Papapetropoulos S, Mash DC, Ffrench-Mullen J, Shehadeh L, de Andrade M, Henley JR, Rocca WA, Ahlskog JE, Maraganore DM. 2007. A genomic pathway approach to a complex disease: Axon guidance and Parkinson disease. *PLoS Genetics* **3**:984-995. doi: [10.1371/journal.pgen.0030098](https://doi.org/10.1371/journal.pgen.0030098).

Lin S, Lin Y, Nery JR, Urich MA, Breschi A, Davis CA, Dobin A, Zaleski C, Beer MA, Chapman WC, Gingeras TR, Ecker JR, Snyder MP. 2014. Comparison of the transcriptional landscapes between human and mouse tissues. *Proceedings of the National Academy of Sciences of the United States of America* **111**:17224-17229. doi: [10.1073/pnas.1413624111](https://doi.org/10.1073/pnas.1413624111).

Lips ES, Cornelisse LN, Toonen RF, Min JL, Hultman CM, the International Schizophrenia Consortium, Holmans PA, O'Donovan MC, Purcell SM, Smit AB, Verhage M, Sullivan PF, Visscher PM, Posthuma D. 2012. Functional gene group analysis identifies synaptic gene groups as risk factor for schizophrenia. *Molecular Psychiatry* **17**:996-1006. doi: [10.1038/mp.2012.37](https://doi.org/10.1038/mp.2012.37).

Liu XL, Somel M, Tang L, Yan Z, Jiang X, Guo S, Yuan Y, He L, Oleksiak A, Zhang Y, Li N, Hu YH, Chen W, Qiu ZL, Pääbo S, Khaitovich P. 2012. Extension of cortical synaptic development distinguishes humans from chimpanzees and macaques. *Genome Research* **22**:611-622. doi: [10.1101/gr.127324.111](https://doi.org/10.1101/gr.127324.111).

Maier R, Moser G, Chen G-B, Ripke S, Cross-disorder Working Group of the Psychiatric Genomics Consortium, Coryell W, Potash JB, Scheftner WA, Shi J, Weissman MM, Hultman CM, Landén M, Levinson DF, Kendler KS, Smoller JW, Wray NR, Lee SH. 2015. Joint Analysis of Psychiatric Disorders Increases Accuracy of Risk Prediction for Schizophrenia, Bipolar Disorder, and Major Depressive Disorder. *The American Journal of Human Genetics* **96**:283-294. doi: [10.1016/j.ajhg.2014.12.006](https://doi.org/10.1016/j.ajhg.2014.12.006).

782 **Markowitz DA**, Curtis CE, Pesaran B. 2015. Multiple component networks support working
783 memory in prefrontal cortex. *Proceedings of the National Academy of Sciences of the*
784 *United States of America* **112**:11084-11089. doi: [10.1073/pnas.1504172112](https://doi.org/10.1073/pnas.1504172112).

785 **Maurano MT**, Humbert R, Rynes E, Thurman RE, Haugen E, Wang H, Reynolds AP,
786 Sandstrom R, Qu HZ, Brody J, Shafer A, Neri F, Lee K, Kuttyavin T, Stehling-Sun S,
787 Johnson AK, Canfield TK, Giste E, Diegel M, Bates D, Hansen RS, Neph S, Sabo PJ,
788 Heimfeld S, Raubitschek A, Ziegler S, Cotsapas C, Sotoodehnia N, Glass I, Sunyaev SR,
789 Kaul R, Stamatoyannopoulos JA. 2012. Systematic Localization of Common Disease-
790 Associated Variation in Regulatory DNA. *Science* **337**:1190-1195. doi:
791 [10.1126/science.1222794](https://doi.org/10.1126/science.1222794).

792 **McLean CY**, Reno PL, Pollen AA, Bassan AI, Capellini TD, Guenther C, Indjeian VB, Lim
793 XH, Menke DB, Schaar BT, Wenger AM, Bejerano G, Kingsley DM. 2011. Human-
794 specific loss of regulatory DNA and the evolution of human-specific traits. *Nature*
795 **471**:216-219. doi: [10.1038/nature09774](https://doi.org/10.1038/nature09774).

796 **McWilliam H**, Li W, Uludag M, Squizzato S, Park YM, Buso N, Cowley AP, Lopez R. 2013.
797 Analysis Tool Web Services from the EMBL-EBI. *Nucleic Acids Research* **41**:597-600.
798 doi: [10.1093/nar/gkt376](https://doi.org/10.1093/nar/gkt376).

799 **Meerabux JMA**, Ohba H, Fukasawa M, Suto Y, Aoki-Suzuki M, Nakashiba T, Nishimura S,
800 Itohara S, Yoshikawa T. 2005. Human netrin-G1 isoforms show evidence of differential
801 expression. *Genomics* **86**:112–116. doi: [10.1016/j.ygeno.2005.04.004](https://doi.org/10.1016/j.ygeno.2005.04.004).

802 **Miller JA**, Horvath S, Geschwind DH. 2010. Divergence of human and mouse brain
803 transcriptome highlights Alzheimer disease pathways. *Proceedings of the National*
804 *Academy of Sciences of the United States of America* **107**:12698-12703. doi:
805 [10.1073/pnas.0914257107](https://doi.org/10.1073/pnas.0914257107).

806 **Nakashiba T**, Ikeda T, Nishimura S, Tashiro K, Honjo T, Culotti JG, Itohara, S. 2000.

Netrin-G1: a Novel Glycosyl Phosphatidylinositol-Linked Mammalian Netrin That Is Functionally Divergent from Classical Netrins. *The Journal of Neuroscience* **20**:6540-6550.

Nakashiba T, Nishimura S, Ikeda T, Itohara S. 2002. Complementary expression and neurite outgrowth activity of netrin-G subfamily members. *Mechanisms of Development* **111**:47-60. doi: [10.1016/S0925-4773\(01\)00600-1](https://doi.org/10.1016/S0925-4773(01)00600-1).

Nakaya T, Alexiou P, Maragkakis M, Chang A, Mourelatos Z. 2013. FUS regulates genes coding for RNA-binding proteins in neurons by binding to their highly conserved introns. *RNA* **19**:498-509. doi: [10.1261/rna.037804.112](https://doi.org/10.1261/rna.037804.112).

Nectoux J, Girard B, Bahi-Buisson N, Prieur F, Afenjar A, Rosas-Vargas H, Chelly J, Bienvenu T. 2007. *Netrin G1* mutations are an uncommon cause of atypical Rett syndrome with or without epilepsy. *Pediatric Neurology* **37**:270-274. doi: [10.1016/j.pediatrneurol.2007.06.002](https://doi.org/10.1016/j.pediatrneurol.2007.06.002).

Nishimura-Akiyoshi S, Niimi K, Nakashiba T, and Itohara S. 2007. Axonal netrin-Gs transneuronally determine lamina-specific subdendritic segments. *Proceedings of the National Academy of Sciences of the United States of America* **104**:14801-14806. doi: [10.1073/pnas.0706919104](https://doi.org/10.1073/pnas.0706919104).

O'Roak BJ, Vives L, Girirajan S, Karakoc E, Krumm N, Coe BP, Levy R, Ko A, Lee C, Smith JD, Turner EH, Stanaway IB, Vernot B, Malig M, Baker C, Reilly B, Akey JM, Borenstein E, Rieder MJ, Nickerson DA, Bernier R, Shendure J, Eichler EE. 2012a. Sporadic autism exomes reveal a highly interconnected protein network of *de novo* mutations. *Nature* **485**:246-250. doi: [10.1038/nature10989](https://doi.org/10.1038/nature10989).

O'Roak BJ, Vives L, Fu WQ, Egertson JD, Stanaway IB, Phelps IG, Carvill G, Kumar A, Lee C, Ankenman K, Munson J, Hiatt JB, Turner EH, Levy R, O'Day DR, Krumm N, Coe BP, Martin BK, Borenstein E, Nickerson DA, Mefford HC, Doherty D, Akey JM, Bernier

R, Eichler EE, Shendure J. 2012b. Multiplex Targeted Sequencing Identifies Recurrently Mutated Genes in Autism Spectrum Disorders. *Science* **338**:1619-1622. doi: [10.1126/science.1227764](https://doi.org/10.1126/science.1227764).

Ohno S. 1970. Evolution by Gene Duplication. Springer, New York.

Ohtsuki T, Horiuchi Y, Koga M, Ishiguro H, Inada T, Iwata N, Ozaki N, Ujike H, Watanabe Y, Someya T, Arinami T. 2008. Association of polymorphisms in the haplotype block spanning the alternatively spliced exons of the *NTNG1* gene at 1p13.3 with schizophrenia in Japanese populations. *Neuroscience Letters* **435**:194-197. doi: [10.1016/j.neulet.2008.02.053](https://doi.org/10.1016/j.neulet.2008.02.053).

Pamplona GSP, Neto GSS, Rosset SRE, Rogers BP, Salmon CEG. 2015. Analyzing the association between functional connectivity of the brain and intellectual performance. *Frontiers in Human Neuroscience* **9**:A61. doi: [10.3389/fnhum.2015.00061](https://doi.org/10.3389/fnhum.2015.00061).

Pan Y, Fang M, Shen L, Wang L, Lv Y, Xi Z, Chen D, Li C, Wang X. 2010. Abnormal expression of netrin-G2 in temporal lobe epilepsy neurons in humans and a rat model. *Experimental Neurology* **224**:340-346. doi: [10.1016/j.expneurol.2010.04.001](https://doi.org/10.1016/j.expneurol.2010.04.001).

Parikshak NN, Gandal MJ, Geschwind DH. 2015. Systems biology and gene networks in neurodevelopmental and neurodegenerative disorders. *Nature Reviews Genetics* **16**:441-458. doi: [10.1038/nrg3934](https://doi.org/10.1038/nrg3934).

Petersen TN, Brunak S, von Heijne G, Nielsen H. 2011. SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nature Methods* **8**:785-786. doi: [10.1038/nmeth.1701](https://doi.org/10.1038/nmeth.1701).

Pollard KS, Salama SR, King B, Kern AD, Dreszer T, Katzman S, Siepel A, Pedersen JS, Bejerano G, Baertsch R, Rosenbloom KR, Kent J, Haussler D. 2006. Forces shaping the fastest evolving regions in the human genome. *PLoS Genetics* **2**:1599-1611. doi: [10.1371/journal.pgen.0020168](https://doi.org/10.1371/journal.pgen.0020168).

857 **Power RA**, Steinberg S, Bjornsdottir G, Rietveld CA, Abdellaoui A, Nivard MM,
858 Johannesson M, Galesloot TE, Hottenga JJ, Willemsen G, Cesarini D, Benjamin DJ,
859 Magnusson PKE, Ullén F, Tiemeier H, Hofman A, van Rooij FJA, Walters GB,
860 Sigurdsson E, Thorgeirsson TE, Ingason A, Helgason A, Kong A, Kiemenev LA,
861 Koellinger P, Boomsma DI, Gudbjartsson D, Stefansson H, Stefansson K. 2015. Polygenic
862 risk scores for schizophrenia and bipolar disorder predict creativity. *Nature Neuroscience*
863 **18**:953-955. doi: [10.1038/nn.4040](https://doi.org/10.1038/nn.4040).

864 **Preuss TM**. 2012. Human brain evolution: From gene discovery to phenotype discovery.
865 *Proceedings of the National Academy of Sciences of the United States of America*
866 **109**:10709-10716. doi: [10.1073/pnas.1201894109](https://doi.org/10.1073/pnas.1201894109).

867 **Prosser P**, Hashimoto R, Polygalov D, Kazutaka O, Zhang Q, McHugh TJ, Takeda M,
868 Itohara S. 2015. Cognitive Endophenotypes of modern and extinct hominins associated
869 with NTNG gene paralogs. *BioRxiv*, doi: [10.1101/034413](https://doi.org/10.1101/034413).

870 **Rada-Iglesias A**, Bajpai R, Prescott S, Brugmann SA, Swigut T, Wysocka J. 2012.
871 Epigenomic Annotation of Enhancers Predicts Transcriptional Regulators of Human
872 Neural Crest. *Cell Stem Cell* **11**:1-16. doi: [10.1016/j.stem.2012.07.006](https://doi.org/10.1016/j.stem.2012.07.006).

873 **Reddy LF**, Lee J, Davis MC, Altshuler L, Glahn DC, Miklowitz DJ, Green MF. 2014.
874 Impulsivity and Risk Taking in Bipolar Disorder and Schizophrenia.
875 *Neuropsychopharmacology* **39**:456-463. doi: [10.1038/npp.2013.218](https://doi.org/10.1038/npp.2013.218).

876 **Richiardi J**, Altmann A, Milazzo A-C, Chang C, Chakravarty MM, Banaschewski T, Barker
877 GJ, Bokde ALW, Bromberg U, Büchel C, Conrod P, Fauth-Bühler M, Flor H, Frouin V,
878 Gallinat J, Garavan H, Gowland P, Heinz A, Lemaître H, Mann KF, Martinot J-L, Nees F,
879 Paus T, Pausova Z, Rietschel M, Robbins TW, Smolka MN, Spanagel R, Ströhle A,
880 Schumann G, Hawrylycz M, Poline J-B, Greicius MD, IMAGEN consortium. 2015.
881 Correlated gene expression supports synchronous activity in brain networks. *Science*

882 **348**:1241-1244. doi: [10.1126/science.1255905](https://doi.org/10.1126/science.1255905).

883 **Riga D**, Matos MR, Glas A, Smit AB, Spijker S, Van den Oever MC. 2014. Optogenetic
884 dissection of medial prefrontal cortex circuitry. *Frontiers in Systems Neuroscience* **8**:A230.
885 doi: [10.3389/fnsys.2014.00230](https://doi.org/10.3389/fnsys.2014.00230).

886 **Rogelj B**, Easton LE, Bogu GK, Stanton LW, Rot G, Curk T, Zupan B, Sugimoto Y, Modic
887 M, Haberman N, Tollervey J, Fujii R, Takumi T, Shaw CE, Ule J. 2012. Widespread
888 binding of FUS along nascent RNA regulates alternative splicing in the brain. *Scientific*
889 *Reports* **2**:A603. doi: [10.1038/srep00603](https://doi.org/10.1038/srep00603).

890 **Romero E**, Ali C, Molina-Holgado E, Castellano B, Guaza C, Borrell J. 2007.
891 Neurobehavioral and immunological consequences of prenatal immune activation in rats.
892 Influence of antipsychotics. *Neuropsychopharmacology* **32**:1791-1804. doi:
893 [10.1038/sj.npp.1301292](https://doi.org/10.1038/sj.npp.1301292).

894 **Scharf JM**, Yu D, Mathews CA, Neale BM, Stewart SE, Fagerness JA, Evans P, Gamazon E,
895 Edlund CK, Service SK, Tikhomirov A, Osiecki L, Illmann C, Pluzhnikov A,
896 Konkashbaev A, Davis LK, Han B, Crane J, Moorjani P, Crenshaw AT, Parkin MA, Reus
897 VI, Lowe TL, Rangel-Lugo M, Chouinard S, Dion Y, Girard S, Cath DC, Smit JH, King
898 RA, Fernandez TV, Leckman JF, Kidd KK, Kidd JR, Pakstis AJ, State MW, Herrera LD,
899 Romero R, Fournier E, Sandor P, Barr CL, Phan N, Gross-Tsur V, Benarroch F, Pollak Y,
900 Budman CL, Bruun RD, Erenberg G, Naarden AL, Lee PC, Weiss N, Kremeyer B, Berrío
901 GB, Campbell DD, Silgado JCC, Ochoa WC, Restrepo SCM, Muller H, Duarte AVV,
902 Lyon GJ, Leppert M, Morgan J, Weiss R, Grados MA, Anderson K, Davarya S, Singer H,
903 Walkup J, Jankovic J, Tischfield JA, Heiman GA, Gilbert DL, Hoekstra PJ, Robertson
904 MM, Kurlan R, Liu C, Gibbs JR, Singleton A for the North American Brain Expression
905 Consortium, Hardy J for the UK Human Brain Expression Database, Strengman E, Ophoff
906 RA, Wagner M, Moessner R, Mirel DB, Posthuma D, Sabatti C, Eskin E, Conti DV,

907 Knowles JA, Ruiz-Linares A, Rouleau GA, Purcell S, Heutink P, Oostra BA, McMahon
 908 WM, Freimer NB, Cox NJ, Pauls DL. 2013. Genome-wide association study of Tourette's
 909 syndrome. *Molecular Psychiatry* **18**:721-728. doi: [10.1038/mp.2012.69](https://doi.org/10.1038/mp.2012.69).

910 **Seiradake E**, Coles CH, Perestenko PV, Harlos K, McIlhinney RAJ, Aricescu AR, Jones EY.
 911 2011. Structural basis for cell surface patterning through NetrinG–NGL interactions. The
 912 *EMBO Journal* **30**:4479-4488. doi: [10.1038/emboj.2011.346](https://doi.org/10.1038/emboj.2011.346).

913 **Sanders SJ**, He X, Willsey AJ, Ercan-Sencicek AG, Samocha KE, Cicek AE, Murtha MT,
 914 Bal VH, Bishop SL, Dong S, Goldberg AP, Jinlu C, Keaney III JF, Klei L, Mandell JD,
 915 Moreno-De-Luca D, Poultney CS, Robinson EB, Smith L, Solli-Nowlan T, Su MY, Teran
 916 NA, Walker MF, Werling DM, Beaudet AL, Cantor RM, Fombonne E, Geschwind DH,
 917 Grice DE, Lord C, Lowe JK, Mane SM, Martin DM, Morrow EM, Talkowski ME,
 918 Sutcliffe JS, Walsh CA, Yu TW, Autism Sequencing Consortium, Ledbetter DH, Martin
 919 CL, Cook EH, Buxbaum JD, Daly MJ, Devlin B, Roeder K, State MW. 2015. Insights into
 920 Autism Spectrum Disorder Genomic Architecture and Biology from 71 Risk Loci. *Cell*
 921 **87**:1215–1233. doi: [10.1016/j.neuron.2015.09.016](https://doi.org/10.1016/j.neuron.2015.09.016).

922 **Shucard JL**, Lee WH, Safford AS, Shucard DW. 2011. The Relationship Between
 923 Processing Speed and Working Memory Demand in Systemic Lupus Erythematosus:
 924 Evidence From a Visual N-Back Task. *Neuropsychology* **25**:45-52. doi:
 925 [10.1037/a0021218](https://doi.org/10.1037/a0021218).

926 **Song M**, Zhou Y, Li J, Liu Y, Tian L, Yu C, Tianzi, J. 2008. Brain spontaneous functional
 927 connectivity and intelligence. *Neuroimage* **41**:1168-1176. doi:
 928 [10.1016/j.neuroimage.2008.02.036](https://doi.org/10.1016/j.neuroimage.2008.02.036).

929 **Skeide MA and Friederici AD**. 2015. Response to Bornkessel-Schlesewsky *et al.* - towards
 930 a nonhuman primate model of language? *Trends in Cognitive Sciences*. **9**:483. doi:
 931 [10.1016/j.tics.2015.05.011](https://doi.org/10.1016/j.tics.2015.05.011).

932 **Song YS**, Lee H-J, Prosser P, Itohara S, Kim E. 2013. Trans-induced cis interaction in the
933 tripartite NGL-1, netrin-G1 and LAR adhesion complex promotes development of
934 excitatory synapses. *Journal of Cell Science* **126**:4926-4938. doi: [10.1242/jcs.129718](https://doi.org/10.1242/jcs.129718).

935 **Soto F**, Watkins KL, Johnson RE, Schottler F, Kerschensteiner D. 2013. NGL-2 Regulates
936 Pathway-Specific Neurite Growth and Lamination, Synapse Formation, and Signal
937 Transmission in the Retina. *Journal of Neuroscience* **33**:11949-11959. doi:
938 [10.1523/jneurosci.1521-13.2013](https://doi.org/10.1523/jneurosci.1521-13.2013).

939 **Spellman T**, Rigotti M, Ahmari SE, Fusi S, Gogos JA, Gordon JA. 2015. Hippocampal–
940 prefrontal input supports spatial encoding in working memory. *Nature* **522**:309-314. doi:
941 [10.1038/nature14445](https://doi.org/10.1038/nature14445).

942 **Steentoft C**, Vakhrushev SY, Joshi HJ, Kong Y, Vester-Christensen MB, Schjoldager KT, BG,
943 Lavrsen K, Dabelsteen S, Pedersen NB, Marcos-Silva L, Gupta R, Bennett EP, Mandel U,
944 Brunak S, Wandall HH, Lavery SB, Clausen H. 2013. Precision mapping of the human O-
945 GalNAc glycoproteome through SimpleCell technology. *Embo Journal* **32**:1478-1488.
946 doi: [10.1038/emboj.2013.79](https://doi.org/10.1038/emboj.2013.79).

947 **Stepanyan A**, Zakharyan R, Boyajyan A. 2013. The netrin G1 gene rs628117 polymorphism
948 is associated with ischemic stroke. *Neuroscience Letters* **549**:74-77. doi:
949 [10.1016/j.neulet.2013.05.066](https://doi.org/10.1016/j.neulet.2013.05.066).

950 **Stewart SE**, Yu D, Scharf JM, Neale BM, Fagerness JA, Mathews CA, Arnold PD, Evans
951 PD, Gamazon ER, Osiecki L, McGrath L, Haddad S, Crane J, Hezel D, Illman C,
952 Mayerfeld C, Konkashbaev A, Liu C, Pluzhnikov A, Tikhomirov A, Edlund CK, Rauch
953 SL, Moessner R, Falkai P, Maier W, Ruhrmann S, Grabe HJ, Lennertz L, Wagner M,
954 Bellodi L, Cavallini MC, Richter MA, Cook EH, Kennedy JL, Rosenberg D, Stein DJ,
955 Hemmings SMJ, Lochner C, Azzam A, Chavira DA, Fournier E, Garrido H, Sheppard B,
956 Umana P, Murphy DL, Wendland JR, Veenstra-VanderWeele J, Denys D, Blom R,

Deforce D, Van Nieuwerburgh F, Westenberg HGM, Walitza S, Egberts K, Renner T, Miguel EC, Capi C, Hounie AG, do Rosário MC, Sampaio AS, Vallada H, Nicolini H, Lanzagorta N, Camarena B, Delorme R, Leboyer M, Pato CN, Pato MT, Voyiaziakis E, Heutink P, Cath DC, Posthuma D, Smit JH, Samuels J, Bienvenu OJ, Cullen B, Fyer AJ, Grados MA, Greenberg BD, McCracken JT, Riddle MA, Wang Y, Coric V, Leckman JF, Bloch M, Pittenger C, Eapen V, Black DW, Ophoff RA, Strengman E, Cusi D, Turiel M, Frau F, Macciardi F, Gibbs JR, Cookson MR, Singleton A for the North American Brain Expression Consortium, Hardy J for the UK Human Brain Expression Database, Crenshaw AT, Parkin MA, Mirel DB, Conti DV, Purcell S, Nestadt G, Hanna GL, Jenike MA, Knowles JA, Cox N, Pauls DL. 2013. Genome-wide association study of obsessive-compulsive disorder. *Molecular Psychiatry* **18**:788-798. doi: [10.1038/mp.2012.85](https://doi.org/10.1038/mp.2012.85).

Stolfi A, Ryan K, Meinertzhagen IA, Christiaen L. 2015. Migratory neuronal progenitors arise from the neural plate borders in tunicates. *Nature* **527**:371-374. doi: [10.1038/nature15758](https://doi.org/10.1038/nature15758).

The Japanese Schizophrenia Sib-Pair Linkage Group (JSSLG), Arinami T, Ohtsuki T, Ishiguro H, Ujike H, Tanaka Y, Morita Y, Mineta M, Takeichi M, Yamada S, Imamura A, Ohara K, Shibuya H, Ohara, K, Suzuki Y, Muratake T, Kaneko N, Someya T, Inada T, Yoshikawa T, Toyota T, Yamada K, Kojima T, Takahashi S, Osamu O, Shinkai T, Nakamura M, Fukuzako H, Hashiguchi T, Niwa S, Ueno T, Tachikawa H, Hori T, Asada T, Nanko S, Kunugi H, Hashimoto R, Ozaki N, Iwata N, Harano M, Arai H, Ohnuma T, Kusumi I, Koyama T, Yoneda H, Fukumaki Y, Shibata H, Kaneko S, Higuchi H, Yasui-Furukori N, Numachi Y, Itokawa M, Okazaki Y. 2005. Genomewide high-density SNP linkage analysis of 236 Japanese families supports the existence of schizophrenia susceptibility loci on chromosomes 1p, 14q, and 20p. *American Journal of Human Genetics* **77**:937-944. doi: [10.1086/498122](https://doi.org/10.1086/498122).

982 **The PsychENCODE Consortium**, Akbarian S, Liu C, Knowles JA, Vaccarino FM, Farnham
983 PJ, Crawford GE, Jaffe AE, Pinto D, Dracheva S, Geschwind DH, Mill J, Nairn AC,
984 Abyzov A, Pochareddy S, Prabhakar S, Weissman S, Sullivan PF, State MW, Weng Z,
985 Peters MA, White KP, Gerstein MB, Senthil G, Lehner T, Sklar P, Sestan N. 2015. The
986 PsychENCODE project. *Nature Neuroscience* **18**:1707-1712. doi: [10.1038/nn.4156](https://doi.org/10.1038/nn.4156).

987 **Thothathiri M. and Rattinger M.** 2015. Ventral and dorsal streams for choosing word order
988 during sentence production. *Proceedings of the National Academy of Sciences* **112**:15456-
989 15461. doi: [10.1073/pnas.1514711112](https://doi.org/10.1073/pnas.1514711112).

990 **Ule J, Ule A, Spencer J, Williams A, Hu JS, Cline M, Wang H, Clark T, Fraser C, Ruggiu M,**
991 **Zeeberg BR, Kane D, Weinstein JN, Blume J, Darnell RB.** 2005. Nova regulates brain-
992 specific splicing to shape the synapse. *Nature Genetics* **37**:844-852. doi: [10.1038/ng1610](https://doi.org/10.1038/ng1610).

993 **van Kuilenburg ABP, Meijer J, Mul ANPM, Hennekam RCM, Hoovers JMN, de Die-**
994 **Smulders CEM, Weber P, Mori AC, Bierau J., Fowler B, Macke K, Sass JO, Meinsma R,**
995 **Hennermann JB, Miny P, Zoetekouw L, Vijzelaar R, Nicolai J, Ylstra, B, Rubio-Gozalbo**
996 **ME.** 2009. Analysis of severely affected patients with dihydropyrimidine dehydrogenase
997 deficiency reveals large intragenic rearrangements of *DPYD* and a de novo interstitial
998 deletion del(1)(p13.3p21.3). *Human Genetics* **125**:581-590. doi: [10.1007/s00439-009-](https://doi.org/10.1007/s00439-009-0653-6)
999 [0653-6](https://doi.org/10.1007/s00439-009-0653-6).

1000 **Voineagu I, Wang X, Johnston P, Lowe JK, Tian Y, Horvath S, Mill J, Cantor RM,**
1001 **Blencowe BJ, Geschwind DH.** 2011. Transcriptomic analysis of autistic brain reveals
1002 convergent molecular pathology. *Nature* **474**: 380-384. doi: [10.1038/nature10110](https://doi.org/10.1038/nature10110).

1003 **Wang K, Zhang H, Bloss CS, Duvvuri V, Kaye W, Schork NJ, Berrettini W, Hakonarson H,**
1004 **the Price Foundation Collaborative Group.** 2011. A genome-wide association study on
1005 common SNPs and rare CNVs in anorexia nervosa. *Molecular Psychiatry* **16**:949-959.
1006 doi: [10.1038/mp.2010.107](https://doi.org/10.1038/mp.2010.107).

1007 **Wilcox JA and Quadri S.** 2014. Replication of NTNG1 association in schizophrenia.
1008 *Psychiatric Genetics* **24**:266-268. doi: [10.1097/ypg.0000000000000061](https://doi.org/10.1097/ypg.0000000000000061).
1009 **Wise SP.** 2008. Forward frontal fields: phylogeny and fundamental function. *Trends in*
1010 *Neurosciences* **31**:599-608. doi: [10.1016/j.tins.2008.08.008](https://doi.org/10.1016/j.tins.2008.08.008).
1011 **Wu JQ, Wang X, Beveridge NJ, Tooney PA, Scott RJ, Carr VJ, Cairns MJ.** 2012.
1012 Transcriptome Sequencing Revealed Significant Alteration of Cortical Promoter Usage
1013 and Splicing in Schizophrenia. *PLoS One* **7**:e36351. doi: [10.1371/journal.pone.0036351](https://doi.org/10.1371/journal.pone.0036351).
1014 **Xu K, Schadt EE, Pollard KS, Roussos P, Dudley JT.** 2015. Genomic and network patterns
1015 of schizophrenia genetic variation in human evolutionary accelerated regions. *Molecular*
1016 *Biology and Evolution* **32**:1148–1160. doi: [10.1093/molbev/msv031](https://doi.org/10.1093/molbev/msv031).
1017 **Yaguchi K, Nishimura-Akiyoshi S, Kuroki S, Onodera T, Itohara S.** 2014. Identification of
1018 transcriptional regulatory elements for *Ntng1* and *Ntng2* genes in mice. *Molecular Brain*
1019 **7**:19. doi: [10.1186/1756-6606-7-19](https://doi.org/10.1186/1756-6606-7-19).
1020 **Yelin-Bekerman L, Elbaz I, Diber A, Dahary D, Gibbs-Bar L, Alon S, Lerer-Goldshtein T,**
1021 **Appelbaum L.** 2015. Hypocretin neuron-specific transcriptome profiling identifies the
1022 sleep modulator Kcnh4a. *eLife* **4**:e08638. doi: [10.7554/eLife.08638](https://doi.org/10.7554/eLife.08638).
1023 **Yin Y, Miner JH, Sanes JR.** 2002. Laminets: Laminin- and netrin-related genes expressed in
1024 distinct neuronal subsets. *Molecular and Cellular Neuroscience* **19**:344-358. doi:
1025 [10.1006/mcne.2001.1089](https://doi.org/10.1006/mcne.2001.1089).
1026 **Yoon JH, Sheremata SL, Rokem A, Silver MA.** 2014. Windows to the soul: vision science as
1027 a tool for studying biological mechanisms of information processing deficits in
1028 schizophrenia. *Frontiers in Psychology* **4**:A681. doi: [10.3389/fpsyg.2014.00681](https://doi.org/10.3389/fpsyg.2014.00681).
1029 **Zakharyan R, Boyajyan A, Arakelyan A, Gevorgyan A, Mrazek F, Petrek M.** 2011.
1030 Functional variants of the genes involved in neurodevelopment and susceptibility to
1031 schizophrenia in an Armenian population. *Human Immunology* **72**:746-748. doi:

1032 [10.1016/j.humimm.2011.05.018](https://doi.org/10.1016/j.humimm.2011.05.018).

1033 **Zhang B**, Gaiteri C, Bodea LG, Wang Z, McElwee J, Podtelezhnikov AA, Zhang CS, Xie T,
1034 Tran L, Dobrin R, Fluder E, Clurman B, Melquist S, Narayanan M, Suver C, Shah H,
1035 Mahajan M, Gillis T, Mysore J, MacDonald ME, Lamb JR, Bennett DA, Molony C, Stone
1036 DJ, Gudnason V, Myers AJ, Schadt EE, Neumann H, Zhu J, Emilsson V. 2013. Integrated
1037 Systems Approach Identifies Genetic Nodes and Networks in Late-Onset Alzheimer's
1038 Disease. *Cell* **153**:707-720. doi: [10.1016/j.cell.2013.03.030](https://doi.org/10.1016/j.cell.2013.03.030).

1039 **Zhang F**, Shugart YY, Yue W, Cheng Z, Wang G, Zhou Z, Jin C, Yuan J, Liu S, Xu Y. 2015.
1040 Increased Variability of Genomic Transcription in Schizophrenia. *Scientific Reports*
1041 **5**:e17995. doi: [10.1038/srep17995](https://doi.org/10.1038/srep17995).

1042 **Zhang Q**, Goto H, Akiyoshi-Nishimura S, Prosser P, Sano C, Matsukawa H, Yaguchi K,
1043 Nakashiba T, Itohara S. 2016. Diversification of behavior and postsynaptic properties
1044 by netrin-G presynaptic adhesion family proteins. *Molecular Brain* **9**:e6. doi:
1045 [10.1186/s13041-016-0187-5](https://doi.org/10.1186/s13041-016-0187-5).

1046 **Zhao M**, Kong L, Qu H. 2014. A systems biology approach to identify intelligence quotient
1047 score-related genomic regions, and pathways relevant to potential therapeutic treatments.
1048 *Scientific Reports* **4**:A4176. doi: [10.1038/srep04176](https://doi.org/10.1038/srep04176).

1049 **Zhou H**, Hu S, Matveev R, Yu Q, Li J, Khaitovich P, Jin L, Lachmann M, Stoneking M, Fu
1050 Q, Tang K. 2015. A Chronological Atlas of Natural Selection in the Human Genome
1051 during the Past Half-million Years. *BioRxiv*. doi: [10.1101/018929](https://doi.org/10.1101/018929).

1052 **Zhu Y**, Yang H, Bi Y, Zhang Y, Zhen C, Xie S, Qin H, He J, Liu L, Liu Y. 2011. Positive
1053 association between *NTNG1* and schizophrenia in Chinese Han population. *Journal of*
1054 *Genetics* **90**:499-502.




Figure 1. NTNG paralogs complementation within neurological disorders and brain transcriptome. (A-1, A-2) Reported cognitive disorders associations for *NTNG1* and *NTNG2*. *denotes rather an indirect association via a direct interaction with the research target. (B-1, B-2) RNA-seq of the STG of healthy (circle) and SCZ (cross) human subjects. The original dataset was produced by (Wu et al., 2012), accession number E-MTAB-1030 on ArrayExpress (ST1a) and reprocessed as described in SM. Five *NTNG1* and four *NTNG2* transcripts, consistently expressed across all 16 human samples are shown. Two samples (one healthy and one SCZ) have been omitted due to unsatisfactory quality of reads and expression profiling (ST1b). For the SNPs calling by SAMtools see ST1c. Data are presented as a mean RPKM \pm SEM. (C) Total number of the assembled transcripts across all samples for both paralogs (see ST1d for the completely reconstructed transcriptome). Dash-outlined are co-spliced exon clusters.




Figure 2. *NTNG* paralogs expression dynamics classification (A-C) in the human brain across the life span. A (1-4): further subdivision of the classifier. RNA-seq data are from the BrainSpan (www.brainspan.org) presented as a mean \pm SEM. TCx = temporary neocortex; OCx = occipital neocortex; PCx = parietal neocortex; LGE = lateral ganglionic eminence; CB(C) = cerebellum(cortex); DTH = dorsal thalamus; URL = upper (rostral) rhombic limb; pcw = post-conception week. Two data points (MD, 12-13 pcw, and mPFC, 16-19 pcw) for *NTNG1* expression were omitted as they were 6-7 times different from the mean of other replicas. All processed brain samples are listed in ST2.




Figure 3. Human *NTNG* paralogs DNA and protein sequence comparisons and “structure-function” rule incongruity. (A) Identical gene structures with different sizes of introns. RNA-seq data from Figure 1B were used to precisely deduce the exon/intron junction boundaries. The sizes of exons 1, 10 and introns (1-2) are not indicated due to observed among the splice transcripts lengths variability (see ST1a for details). Arrows indicate location of CNS = conserved non-coding sequences underwent accelerated evolution in human compare to mice (mCNS) and chimpanzee (chCNS), as per Prabhakar et al., 2006; and ASC = anthropoid-specific constrained regions in human compare to marmoset (maASC), as per del Rosario et al., 2014. (B) Identical exonal composition of the longest *NTNG* encoded RNA paralog transcripts and corresponding proteins with relatively high percent of identity among them dependent on the included/excluded Ukd domain (B-2) encoded by the exons 6 and 7 (B-1). Notably, the protein sequence represents higher percent of the paralogs difference than encoded it DNA. The matrices were obtained by GeneJockey II (Biosoft). (C) Protein alignments for the longest human *NTNG* encoded proteins, Netrin-G1m and Netrin-G2b, with Loops I-III highlighting binding sites for their cognate post-synaptic binding partners NGL-1 (*Lrrc4c*) and NGL-2 (*Lrrc4*), respectively, as determined by Seiradake et al. (2011). Arrow indicates a putative secretory cleavage site location, as calculated by SignalIP (Petersen et al., 2011), the blue rectangle delineates the area of the lowest identity (3'-domain LE1+Ukd domain); ω – denotes a point of putative GPI-attachment, as predicted by Big-PI (Eisenhaber et al., 2000). PSD-95 interaction site via the SH3-binding domain (Arbuckle et al., 2010, as determined for mice Netrin-G2) overlaps with the Loop III NGL-2 binding surface. Two stars indicate a modern human (T346A) and a hominin-specific (S371A/V) amino acid substitutions (Prosselkov et al., 2015). (D) Identical structural motif of the Netrin-G1/NGL1 and Netrin-G2/NGL2 complexes as per Seiradake et al. (2011). The figure’s reproduction is covered by the Creative Commons license.