

Structure, inhibition, and regulation of two-pore channel TPC1 from *Arabidopsis thaliana*

Alexander F. Kintzer¹ and Robert M. Stroud¹

¹Department of Biochemistry and Biophysics, University of California, San Francisco, California 94158, USA.

Main text

Two-pore channels (TPCs) comprise a subfamily (TPC1-3) of eukaryotic voltage- and ligand-gated cation channels^{1,2} that contain two non-equivalent tandem pore-forming subunits that then dimerize to form quasi-tetramers. Found in vacuolar³ or endolysosomal⁴ membranes, they regulate the conductance of sodium⁵ and calcium^{3,6} ions, intravesicular pH⁵, trafficking⁷ and excitability^{8,9}. TPCs are activated by a decrease in transmembrane potential^{1,3,9,10}, an increase in cytosolic calcium concentrations^{1,10}, and inhibited by luminal low pH, and calcium¹¹, and regulated by phosphorylation^{12,13}. We report the crystal structure of TPC1 from *Arabidopsis thaliana* (TPC1) at 2.8x4.0x3.3Å resolution as a basis for understanding ion permeation^{3,4,10}, channel activation^{1,5,10}, the location of voltage-sensing domains^{1,9,10}, and regulatory ion-binding sites^{11,14}. We determined sites of phosphorylation^{3,4} in the N-terminal and C-terminal domains that are positioned to allosterically modulate cytoplasmic Ca²⁺-activation. One of the two voltage sensing domains (VSDII) encodes voltage sensitivity and inhibition by luminal Ca²⁺ and adopts a conformation distinct from the activated state observed in structures of other voltage-gated ion

channels^{15,16}. The structure shows that potent pharmacophore trans-NED19¹⁷ allosterically acts by clamping the pore domains to VSDII. In animals NED19 prevents infection by Ebola virus and Filoviruses presumably by altering their fusion with the endolysosome, and delivery of their contents into the cytoplasm⁷ (Supplementary Discussion).

Diffraction was optimized and the final conditions depended on relipidation, partial dehydration, NED19¹⁷, and deletion of residues 2-11. The structure was determined *de novo* to 2.87Å by 9 metal substitutions and derivatives, and refined to $R_{\text{cryst}}=29.7\%$, $R_{\text{free}}=33.9\%$ (Methods and Extended Data Table 1,2,3).

TPC1 consists of two non-identical Shaker-like^{18,19} pore-forming subunits, domain I and II, separated by an EF-hand domain (Fig. 1). The central pore domains PI (S5-S6) and PII (S11-S12) couple the voltage-sensing domains VSDI (S1-S4) and VSDII (S7-S10) and modulatory cytosolic N-terminal (NTD), the central EF-hand (EF), and C-terminal (CTD) domains, which extend 20Å into the cytoplasm (Fig. 2, Extended Data Fig. 1, Methods for detailed domain assignments). The experimentally determined electron density unbiased by any model (Extended Data Fig. 2), showed that TPC1 forms a rectangular structure, ~100Å x 70Å in the membrane plane (Fig. 1). During review of our manuscript and after referee reviews, we became aware of the preceding manuscript that reports the crystal structure of the same molecule, accompanied by extensive functional characterizations¹⁰. The structures are generally in excellent agreement.

We additionally observe the CTD, a lipid, the inhibitor NED19, and there is some variation in observed ion binding sites. We discuss our results in light of their observations.

Plant and human TPCs respond to phosphoregulation by specific kinases^{3,4}. We determined sites of phosphorylation in TPC1 using a combination of mass spectrometry and binding of ProDiamond-Q to N- and C-terminal truncations (Extended Data Fig. 3 and Supplementary Discussion). Residues S22, T26, and T29 in the NTD are sites of phosphorylation in TPC1. S22 is conserved in both human TPCs, whereas T26 is only found in hTPC1 (Extended Data Fig. 1). The ordered structure begins with NTD helix 2 (NTDh2; residues P30-G46). NTDh2 makes hydrophobic and polar contacts with the EF-hand domain helix 1 (EFh1), EFh2, EFh4, and VSDI (S2-S3), leading into a NTD loop 2 (NTDl2) that makes nearly a 180° turn to connect to the pre-S1 helix of VSDI. The proximity of the NTD phosphorylation sites to the Ca²⁺-activation site in the EF-hand^{14,10} EFh3 and loop EFh3, and VSDI suggests that phosphorylation in the NTD could modulate channel opening by influencing the local structure of the Ca²⁺ site or conformation of VSDI via a salt-bridge (E50-R200) between NTDl2 and VSDI (S4-S5).

Plant and human TPC1s form voltage-dependent inwardly-rectifying (towards the cytosol) ion channels, whereas hTPC2 is voltage insensitive^{1,5,9,10,12}. In contrast to Ca_v, Na_v K_v

channels, activation of TPC channels is characteristically much slower, and they do not inactivate hence the original naming for plant TPCs, as slow vacuolar channels^{1,5,9,10,12}.

Voltage-sensing is primarily encoded by VSDII and not by VSDI (Fig. 3)¹⁰. VSDII contains a classic voltage-sensing motif with four arginines R531, R537, R540, R543 in S10 (Fig. 3b, Extended Data Fig. 4a) and a conserved counter anion charge-transfer center in (Y475-E478-R537) in S8 (Extended Data Fig. 4b), as in NavAb¹⁶. Three other arginines form corresponding ion pairs with anions on S9 (E511-R531, D500-R540, E494-R543). The accompanying paper shows that the three arginines R537, R540, R543 are required for voltage-sensing in TPC1¹⁰. The second ‘voltage-sensing’ arginine R537 interacts with the conserved luminal charge transfer center near Y475^{10,16}, creating an opening in the cytosolic side of VSDII, and exposing R543 to the cytosol. VSDII is in a novel conformation with respect to other voltage-gated ion channel structures, which each place the last charge, equivalent to R543, at the charge transfer center, thought to represent activated VSDs^{15,16}. The luminal/extracellular side of activated VSDs adopts a more open conformation, while the cytosolic side is narrow and closed. This may mean that the TPC1 VSDII structure represents an inactive state. VSDII is indicated as the functional voltage sensor in human TPC1 also, since the R539I mutation (corresponding to R537 in plant TPC1 VSDII) eliminates most of the observed voltage-sensitivity⁹. In the human TPC2 channel the corresponding residue is I551, consistent with its lack of voltage-sensitivity.

VSDI is structurally divergent from VSDII, containing only two arginines R185 and R191 in S4, and has no conserved charge transfer center in S2 (Fig. 3a, Extended Data Fig. 4a,b) therefore not predicted to form a functional voltage sensor. These predictions are corroborated by site-directed mutagenesis and electrophysiology¹⁰. Intervening lateral amphipathic helices on the cytoplasmic leaflet S4-S5 and S10-S11 link VSDI and II to the pore domains PI and PII, respectively, making polar and hydrophobic contacts with the pore gates I (Gate I) and II (Gate II), and providing connections for coupling voltage-sensing, phosphorylation, and binding of ions, inhibitors, or proteins to the pore. S4-S5 is critical for coupling voltage to the pore gates⁹ in human TPC1.

Structural divergence between VSDI and II generates the rectangular shape of TPC1. Variations in the S4 angle are found in symmetrical tetrameric ion channels, though the functional relevance is not clear (Extended Data Fig. 5). In the structure of Ca_v1.1 asymmetry of the pore domains PI-PIV plays a role in ion permeation and selectivity while the VSDs remain close to symmetrical²⁰. TPCs and other tandem channels may use structural asymmetry to expand the repertoire of sensory responses and facilitate channel gating, as in the regulation of Ca_v channels by ions, toxins, pharmacophores, phosphorylation, or additional subunits²¹.

Luminal Ca²⁺ inhibits plant TPC1 conductance^{10,11}. The plant TPC1 mutant D454N/*fou2* in the S7-S8 loop abolishes luminal Ca²⁺ sensitivity and increases the rate of channel opening¹¹.

In our structure there are two Ca^{2+} binding sites on the luminal side of VSDII. Site I is coordinated by E457 from S7-S8 loop and E239 of PI, whereas Site II uses D454, D240 of PI, and E528 of S10. Based on mutagenesis and patch-clamp electrophysiology of TPC1 only Site 2 is critical for luminal inhibition by Ca^{2+} ions¹⁰. Heavy atom mimetics of Ca^{2+} (Ba^{2+} and Yb^{3+}) validate the placement of Ca^{2+} in Sites I and II, defined by isomorphous and anomalous diffraction difference maps (Fig. 3, Extended Data Fig. 2c). A single fully occupied Yb^{3+} ion (Yb2) and Ba^{2+} site (Ba1) overlaps with the Site 2. In hTPC2, E528 is an aspartate, suggesting that there may be mechanistic similarities in ion and pH sensitivities between hTPC2 and plant TPC1. VSDI binds a structural Ca^{2+} ion, confirmed by replacement with a Yb^{3+} ion (Yb1) coordinated by acidic residues E124 in S2 and D166 and D170 in S3 (Fig. 3a). Human and plant TPCs have an acidic residue E124 in S2 suggesting that this region could be a conserved Ca^{2+} -binding site. However, this ion binding site has not been studied functionally.

NED19^{7,17} and Ca_v channel agonists and antagonists inhibit opening of TPC channels. Mutagenesis in human TPC1 demonstrated that L273 is important for channel activation and Ca^{2+} -release²². The equivalent residues in TPC1 (M258 and L623) line PI and PII and form a hydrophobic surface that could bind hydrophobic ligands or stabilize the closed state in the absence of agonist (Extended Data Fig. 6). We observe density in TPC1 corresponding to the polycyclic TPC inhibitor NED19 (Fig. 4). Interactions with NED19 involve F229 in S5, both

hydrophobic pi-stacking and a hydrogen bond between the carboxylate of NED19 and the sidechain amino group of W232 in S5, L255 in PI, F444 in S10, and W647 in S12. NED19 effectively clamps the pore domains and VSDII together, allosterically blocking channel activation. The binding site is consistent with NED19 as a high-affinity TPC-specific inhibitor^{7,17}. TPC1 S6 and S12 are 25-50% conserved with S6 helices in Ca_v channels (Extended Data Fig. 4c). Dihydropyridines (DHP) and phenyl alkyl amines (PPA) pharmacophores bind to specific aromatic residues at a nearby, yet distinct, region in S6 of the third and fourth tandem pore forming domains in Ca_v channels²³. Homologous DHP-binding residues Y288, Y656, L663, L664 in TPC1 and conserved residues in hTPC1 and hTPC2 could bind Ca_v pharmacophores in a parallel mechanism of inhibition⁷.

Ion selectivity varies among TPC channels. Plant TPCs are non-selective^{3,10}, whereas human TPC1 is Na⁺-selective^{5,9}, and TPC2 conducts Na⁺, Ca²⁺, and possibly H⁺ ions^{5,9,24}. While the selectivity filter in TPCs is fairly conserved with Na_v, and Ca_v channels, the specific anionic residues known to impart Na⁺ or Ca²⁺ selectivity^{16,25} are replaced by hydrophobic side chains (Extended Data Fig. 4d). The largely hydrophobic selectivity filter of TPC1 agrees with the reported lack of ion selectivity¹⁰. A single anion in the selectivity filter of hTPC1 pore loop II E642 could confer Na⁺-selectivity as this residue, N624 faces the pore lumen in TPC1. Pore loop I forms a re-entrant helix-turn-helix motif that forms a wide lining of two sides of the pore mouth

(Fig. 5a). The high resolution data and crystallization conditions enabled visualization of electron density for lipid molecules, one bound to the backside of PI at the interface of VSDII, flanking the luminal Ca^{2+} -binding Site II. Modeled as a palmitic acid, the lipid is a hydrogen bond acceptor from T241 and M237 of PI and bound adjacent to D240 the site for luminal Ca^{2+} -inhibition. This could provide a basis for the reported modulation of TPCs by fatty acids. (Supplementary Discussion). By contrast, pore loop II forms an extensive asymmetric constriction site at the pore mouth that brings four negative charges D605, D606 to its lining of which D605 likely coordinates a Ca^{2+} ion (Fig. 5b), confirmed by Yb^{3+} (Yb^{3+}) observed in isomorphous and anomalous diffraction difference maps. The coordination distance of 2.5 Å suggests that this site could recognize ions from the lumen. However, this site is unlikely to function in selecting ions, as the wide pore mouth ~12-15 Å of PI does not provide a barrier to ion flow (Fig. 5c). D269 of PI faces the pore, but the distance to the lumen is much too far 12.6 Å to function in ion selectivity.

Two N631 residues converge to 4.9 Å to form a narrowed region that separates the upper vestibule from the putative selectivity filter. However, opposing Q633 residues from PI are 12.2 Å apart leave the gasket open to ion flow. This is the narrowest part of the selectivity filter.

Two T264 residues, conserved in Ca_v , Na_v , and K_v subunits, define the upper boundary of the central cavity in TPC1 (Extended Data Fig. 4d, Fig. 5c). The central cavity contains solvent

molecules and barium ions (Ba²⁺), defined by isomorphous and anomalous difference maps.

However, no ions are observed in the selectivity filter, unlike highly selective Na_v, Ca_v, or K_v channels^{16,20,15}. The central cavity is water filled and lined by hydrophobic residues. Two rings of hydrophobic residues L301, V668, Y305, and L672 seal the cavity toward the cytoplasm, and are part of the extensively studied pore gates of K_v channels^{26–28} and Gates I and II in TPC1. On the cytosolic face of the gates, a salt-bridge K309-E673 staples Gate I and II together. K⁺ ions have been observed in the gate and lower selectivity filter of MlotiK1²⁹ and KcsA³⁰ and mutation of similarly positioned hydrophobic residues in the gate can modulate the channel state or flux of ions through the channel. This supports the idea that the VSDs, selectivity filter, and gate undergo coordinated movements during channel opening and closing to regulate ion conductance.

In plant TPC1, two EF-hand domains between domains I and II, confer sensitivity to cytosolic Ca²⁺ ions¹⁴. These domains consist of two Ca²⁺ binding motifs EF11 and EF13, with 25-30% sequence conservation to human calmodulin (hCAM, Extended Data Fig. 4e). EFh1 forms a 70Å long helix continuous with the gate helix S6 (Fig. 2b, Fig. 6a) that is connected to the first Ca²⁺-binding site in EF11 formed by sidechain interactions with D335 (2.6Å), D337 (2.8Å), N339 (3.2Å), and the backbone of E341 (2.3Å). This site is probably structural, as it does not function in Ca²⁺-activation^{10,14}. The second site in EF13 has been shown to facilitate channel activation by

cytosolic Ca^{2+} , as a D376A mutant abolished all Ca^{2+} -dependent activation¹⁴. We observe an Ca^{2+} ion in EF13 coordinated by E374 (2.9Å) and D377 (5.2Å), but surprisingly not by D376 which is 8.5Å away. Thus the mechanism of Ca^{2+} -activation may involve a structural change in EF13 involving the Ca^{2+} -binding site and D376.

The CTD plays a critical role in plant TPCs (Supplementary Discussion), though least conserved across species (Extended Data Fig. 1). CTD helix 1 (CTDh1) continues from Gate II and S12 with a poly-anionic (poly-E) helix E682-E685 (Fig. 6b). Ca_v channels contain a similar conserved poly-anionic region of variable length (Extended Data Fig. 4c). Poly-E residues E682 and E685 hydrogen bond to R552 of S10-S11, directly linking the conformation of the CTDh1 to the voltage-sensor VSDII S10-S11. CTD loop 1 (CTDl1) from each monomer converges to form a charged constriction site D694, K695, and R696 (Fig. 6b, Fig. 5c), that bring six charged sidechains to stabilize the channel through inter-monomer contacts. The intermolecular constriction site partially includes a poly-R motif formed by R696, R698, R699, R700, and may play a structural role or regulate binding of second messenger lipids or proteins (Extended Data Fig. 7). Importantly, the CTD forms an intra-molecular complex with the Ca^{2+} -activation site EF13 (Fig. 6c). Surprising, R700 from CTDl1 makes a salt-bridge with the critical residue for Ca^{2+} -activation R700-D376. S701 coordinates the bound Ca^{2+} (4.2Å) in EF13 along with E374 and D377. These interactions include the potential site of phosphorylation S706 in CTDl1, not

observed in the structure. Thus, reversible phosphorylation in the CTD could disrupt the EFh3-CTD11 interaction and modulate Ca^{2+} -sensing or couple directly to the VSDII via the R552-E682-E685 interaction (Fig. 6c). The interactions between the regulatory sites and the central channel are schematized in Extended Data Fig. 7. The channel conductance of human and plant TPCs is multi-modal³. Our structure suggests a mechanism for channel opening, whereby Ca^{2+} concentrations, voltage, and phosphoregulation are integrated through conformational changes in the VSDs, selectivity filter, gate, NTDs, and CTDs to govern the conduction of ions (Extended Data Fig. 7).

Methods

Protein Production. AtTPC1 was expressed in codon-optimized form in *Saccharomyces cerevisiae* based on p423_GAL1³¹ (Uniprot accession number Q94KI8). A C-terminal Thrombin cleavage site was followed by a 12 residue glycine-serine linker and a 10 residue histidine tag. Transformed *S. cerevisiae* (strain DSY-5) were grown from overnight starter cultures in synthetic complete defined medium lacking histidine (CSM-HIS) in 1L shaker flasks at 30°C to OD_{600nm} of 12-15. Protein expression was induced by adding 8% Galactose dissolved in 4x YP medium, to a final concentration of 2% Galactose, at 30°C for 18-20hr. Cells were harvested by centrifugation, resuspended in lysis buffer (50 mM Tris pH 7.4, 500 mM NaCl, 1 mM EDTA, 1 mM PMSF), followed by bead beating using 0.5mm beads for 30 minutes at 1 minute intervals. The lysate was then centrifuged at 21,000g for 20 minutes, followed by collection of membranes by centrifugation at 186,010g for 150 minutes. Membranes were resuspended (50 mM Tris pH 7.4, 500 mM NaCl, 10% Glycerol), aliquoted, and frozen in liquid nitrogen for storage at -80°C. 7-12 g aliquots of membranes were solubilized in membrane buffer containing 1% (0.2 g/g of membrane) n-dodecyl-beta-D-maltopyranoside (DDM), 0.1 mM cholesteryl hemisuccinate (CHS), 0.03 mg/mL soy polar lipids, 1 mM CaCl₂, and 1 mM PMSF, for 1.5 hours at 4°C. The solubilized material was collected by centrifugation at 104,630g for 20 minutes, filtered through a 5µm filter, and incubated with 3 mL of NiNTA beads in the presence of 20 mM Imidazole pH

208 7.4 for 7.5 hours at 4°C. NiNTA beads were collected by draining the mixture through a column
 209 and the beads were washed successively with 25 mL of NiNTA wash buffer (50 mM Tris pH
 210 7.4, 500 mM NaCl, 5% Glycerol, 0.1% DDM, 0.1 mM CHS, 0.03 mg/mL soy polar lipids, 1 mM
 211 CaCl₂) plus 20 mM Imidazole pH 7.4 for 20 minutes at 4°C, then 25 mL of NiNTA wash buffer
 212 plus 75 mM Imidazole pH 7.4 for 10 minutes at 4°C, followed by 10 mL of NiNTA elution
 213 buffer (50 mM Tris pH 7.4, 200 mM NaCl, 5% Glycerol, 0.1% DDM, 0.1 mM CHS, 0.03
 214 mg/mL soy polar lipids, 2 mM CaCl₂). TPC1 protein was obtained by incubating NiNTA beads
 215 with 200 units of bovine thrombin in 25 mL of NiNTA elution buffer overnight at 4°C. The
 216 following day, NiNTA beads were washed with an additional 25 mL of NiNTA wash buffer plus
 217 20 mM Imidazole pH 7.4 for 20 minutes at 4°C, concentrated, and loaded on a Superose 6
 218 column equilibrated in Size exclusion buffer (20 mM Hepes pH 7.3, 200 mM NaCl, 5%
 219 Glycerol, 0.05% DDM, 0.1 mM CHS, 0.03 mg/mL soy polar lipids, and 1 mM CaCl₂). Peaks
 220 fractions were pooled and concentrated to 5 mg/mL for crystallization. TPC1 and mutants were
 221 prepared similarly, except for the S701C mutant where 1mM and 100μM TCEP were included
 222 during solubilization and final size exclusion chromatography, respectively.

223 **Crystallization.** Several eukaryotic TPCs, including human hTPC1, hTPC2, and AtTPC1
 224 (TPC1) were evaluated for expression in *S. cerevisiae* and SF9 cells, and homogeneity by
 225 fluorescence-based and conventional size-exclusion chromatography. TPC1 was selected based

on expression level, stability, homogeneity and crystallization trials. Nevertheless, detergent-solubilized TPC1 was unstable and resistant to crystallization. Thermostability screening³² identified cholesteryl hemisuccinate (CHS), lipids, and Ca²⁺ ions as improving TPC1 stability and homogeneity.

Wild-type TPC1 crystals were obtained in the presence of CaCl₂, CHS, and soy polar lipids. Initial crystals from 200nL drops in sparse matrix trays diffracted to ~10Å resolution, with few diffracting to 4Å. Reproducibility in 2μL drops was achieved through extensive screening trials. To improve diffraction, truncation constructs were made. A construct lacking the N-terminal residues 2-11 (TPC1) diffracted on average to 7Å, with 1% showing diffraction to 3.9Å resolution, but were too radiation sensitive and anisotropic to collect complete datasets. The TPC inhibitor trans-NED19 (NED19) was added to 1mM final, 1 hour before crystallization. These diffracted to ~5Å resolution, with 50% to 3.9Å resolution. Adding 100μL of protein and 1mg of DDM to a thin-film of 0.03 mg soy polar lipids with stirring overnight at 4°C led to the TPC1 high lipid-detergent (HiLiDe)³³ mixture following ultracentrifugation for 20 minutes in a TLA-55 rotor at 112,000g. Average diffraction was 4Å, with 10% diffracting to 3.5-3.7Å resolution, the best being 3.5 x 5.5 x 4.0Å. Anisotropic resolution was determined using the CCP4 program Truncate³⁴ and the UCLA Anisotropy Server³⁵. Large improvements were achieved through partial dehydration. HiLiDe crystals were incubated with additional amounts of polyethylene

glycol 300, 0-20 % (v/v), prior to harvesting. Incubating with additional 15% PEG300, increased the average diffraction to 4Å, best diffraction to 2.7Å resolution, and diminished the diffraction anisotropy. Derivative crystals were obtained by soaking in 0.1-50 mM of heavy-atom compounds in a 1:1 mixture of Crystal buffer (0.1M Glycine pH 9.3, 50 mM KCl, 1mM CaCl₂, 35% PEG300) for 24-48 hours at 20°C. Partial dehydration was performed after derivatization.

Data Collection and Reduction. X-ray diffraction was collected at the Advanced Light Source Beamlines (ALS) 8.3.1, 5.0.2 and the Stanford Synchrotron Radiation Laboratory Beamline (SSRL) 12-2. Data were reduced using XDS³⁶. Native data were collected at wavelength 1.000Å. Data from heavy atom derivative crystals were collected at the anomalous peak of the L-III edge for each element (Ba=1.750Å, Hg=1.006Å, Ta=1.2546Å, Yb=1.3854Å).

Structure Determination of TPC1. De novo structure determination was based on 9 heavy metal and cluster derivatives at 3.5Å (Extended Data Table 1). Dehydration improved diffraction to 2.7Å while diminishing the anisotropy (Extended Data Table 2, Extended Data Table 3). Experimental phases were calculated by multiple isomorphous replacement with anomalous scattering (MIRAS) from both non-dehydrated (Extended Data Table 1) and dehydrated (Extended Data Table 2) crystals derivatized with tantalum bromide clusters (Ta₆Br₁₂), barium(II) chloride, ytterbium(III) chloride, and a mutant of TPC1 derivatized with mercury(II) chloride.

A single tantalum atom was identified by Single Isomorphous Replacement (SIR) with Anomalous Scattering (SIRAS) using SHELXC/D³⁷. A Ta₆Br₁₂ cluster was refined and phases calculated using SHARP³⁸. These phases were used to place sites in other derivatives and calculate phases using SHARP³⁸. Initial MIRAS phases were determined with non-dehydrated crystals with 100μM Ta₆Br₁₂ clusters (Derivative 1; 1 site, Isomorphous phasing power (PP_{iso}; defined as $PP_{iso} = \text{RMS} (|F_H|_{\text{calc}} / \sum ||F_{PH} - F_P|_{\text{obs}} - |F_{PH} - F_P|_{\text{calc}}|)$ or $\text{RMS} (|F_H|_{\text{calc}} / \sum \epsilon_{iso})$, where $|F_H|$ is the heavy atom structure factor amplitude, $|F_{PH}|$ the heavy-atom derivative structure factor amplitude, $|F_P|$ the native structure factor amplitude, and $|F_H|$ is the heavy atom structure factor amplitude, and ϵ_{iso} the phase-integrated lack of closure for isomorphous differences) of 1.0/0.77 (acentric/centric); Anomalous phasing power (PP_{ano}; defined as $PP_{ano} = \text{RMS} (\sum |F_{PH+} - F_{PH-}|_{\text{calc}} / \sum \epsilon_{ano})$ or $\text{RMS} (\sum |F_{PH+} - F_{PH-}|_{\text{calc}} / \sum ||F_{PH+} - F_{PH-}|_{\text{obs}} - |F_{PH+} - F_{PH-}|_{\text{calc}}|)$, where $|F_{PH+} - F_{PH-}|$ is the mean anomalous difference between Friedel pairs, and ϵ_{ano} the phase-integrated lack of closure for anomalous differences) of 0.31; Isomorphous R_{cullis} (R_{cullis}_{iso}; defined as $R_{cullis_{iso}} = \sum \epsilon_{iso} / \sum |F_{PH} - F_P|_{\text{obs}}$) of 0.766/0.755 (acentric/centric); Anomalous R_{cullis} (R_{cullis}_{ano}; defined as $R_{cullis_{ano}} = \sum \epsilon_{ano} / \sum |F_{PH+} - F_{PH-}|_{\text{obs}}$) of 0.584), 1mM Ta₆Br₁₂ clusters (Derivative 2; 1 site, PP_{iso} of 2.32/1.94 (acentric/centric), PP_{anom} of 0.761, R_{cullis}_{iso} of 0.525/0.536 (acentric/centric), R_{cullis}_{ano} of 0.792), or 2mM Ta₆Br₁₂ clusters (Derivative 3; 1 site, PP_{iso} of 0.89/1.1 (acentric/centric), PP_{anom} of 1.36, R_{cullis}_{iso} of 0.850/0.765

(acentric/centric), $R_{\text{cullis_ano}}$ of 0.572), using merged data (Native 1) from three non-dehydrated crystals as a source of isomorphous native amplitudes (Extended Data Table 1).

This map allowed building of all transmembrane segments and partial building of the EF-hand domain in TPC1. Sidechain features and density for loop regions were largely absent from calculated maps, precluding subunit identification, and objective determination of the helical register. Initial MIRAS phases were further improved by including merged data from five non-dehydrated crystals derivatized with 50mM barium(II) chloride (Derivative 4; 3 sites, PP_{iso} of 0.75/0.58 (acentric/centric), PP_{anom} of 0.59, $R_{\text{cullis_iso}}$ of 0.905/0.888 (acentric/centric), $R_{\text{cullis_ano}}$ of 1.00) in the MIRAS calculation, resulting in an overall FOM of 0.262/0.333 (acentric/centric) (Extended Data Table 1). Combined tantalum and barium phases allowed fitting of some sidechains to density, placement of ions binding sites in the channel lumen and VSDs (Extended Data Fig. 2, Supplementary Data Table 1). The best source of phases came from derivatives obtained from crystals that were dehydrated after a 24-48 hour soak in heavy atom solution. MIRAS phases were calculated from native crystals derivatized with 1mM $\text{Ta}_6\text{Br}_{12}$ clusters (Derivative 5; 1 sites, PP_{iso} of 4.09/3.56 (acentric/centric), PP_{anom} of 2.42, $R_{\text{cullis_iso}}$ of 0.302/0.286 (acentric/centric), $R_{\text{cullis_ano}}$ of 0.495) and Derivative 6; 1 sites, PP_{iso} of 8.54/8.91 (acentric/centric), PP_{anom} of 2.29, $R_{\text{cullis_iso}}$ of 0.149/0.122 (acentric/centric), $R_{\text{cullis_ano}}$ of 0.517) or 2mM $\text{Ta}_6\text{Br}_{12}$ clusters (Derivative 7; 1 sites, PP_{iso} of 3.65/3.54

298 (ancentric/centric), PP_{anom} of 2.25, $R_{cullis_{iso}}$ of 0.372/0.323 (ancentric/centric), $R_{cullis_{ano}}$ of
299 0.535), 1mM ytterbium(III) chloride (Derivative 8; 4 sites, PP_{iso} of 0.851/0.795
300 (ancentric/centric), PP_{anom} of 0.699, $R_{cullis_{iso}}$ of 0.962/0.884 (ancentric/centric), $R_{cullis_{ano}}$ of
301 0.982)), and a TPC1 S701C mutant derivatized with 1mM mercury(II) chloride (Derivative 9; 5
302 sites, PP_{iso} of 0.792/0.822 (ancentric/centric), PP_{anom} of 0.736, $R_{cullis_{iso}}$ of 0.935/0.704
303 (ancentric/centric), $R_{cullis_{ano}}$ of 1.02), followed by dehydration resulting in a FOM of
304 0.267/0.341 (ancentric/centric) (Extended Data Table 2). Non-isomorphism errors were
305 minimized by using merged data from three non-dehydrated crystals extending to 3.5Å
306 resolution (Native 1) as a source of isomorphous native amplitudes for all phasing calculations,
307 density modification, and initial model building (Extended Data Table 1, Extended Data Table
308 2). MIRAS phases from dehydrated derivatives extend to higher resolution, contain less
309 diffraction anisotropy, and display much clearer electron density for mainchain and sidechain
310 features, and allow objective determination of helical register (Extended Data Fig. 2b). Mercury
311 heavy atom positions were used as restraints, along with sequence homology, for objective
312 placement of cysteine residues (Supplementary Data Table 1). Ytterbium and barium heavy atom
313 positions were used in conjunction with homology for placing likely (D, E) chelate residues
314 within the pore, VSDs, and pore (Extended Data Fig. 2c, Supplementary Data Table 1). MIRAS
315 phases from non-dehydrated and dehydrated derivatives were further improved through solvent

flipping in SOLOMON³⁹ with 75% and 70% solvent content and FOMs of 0.637 and 0.470.

Phases were then combined and extended to 3.5Å resolution through iterative cycles of density modification in DMMULTI⁴⁰, using histogram matching, solvent flattening, gradual phase extension⁴¹, and 3-fold cross-crystal averaging with un-phased, dehydrated, native amplitudes (Native 2) with 70% solvent content with a FOM of 0.493 (Extended Data Fig. 2b). Averaging masks were generated by NCSMASK³⁴ with radius of 4Å from atomic models from each iteration of building and refinement. Model phases were purposely left out of experimental density modification procedures to reduce bias.

Initial model building was guided by the position of heavy atoms determined from isomorphous difference maps (Extended Data Fig. 2c, Supplementary Data Table 1). Using anomalous differences yielded equivalent maps and structural restraints.

Refinement. The best native data (Native 2) extends to overall resolution 2.8 x 4.0 x 3.3Å (ref.³⁵), and enabled building of 85% of the TPC1 protein sequence. Structure interpretation was using COOT⁴², with refinement in PHENIX⁴³. Simulated-annealing was employed alongside refinement of coordinates, TLS restraints, and several cycles of automated building using ARP/wARP³⁴ in CCP4. Building of the NTD, CTD, and loop regions were guided by restraints based on heavy atom positions (Supplementary Data Table 1), omit maps, and gradual improvement of the map and crystallographic Rfactors. Refinement of the structure in triclinic

space group P1 led to a substantial reduction in R_{free} (39.9%). During review of our manuscript and after referee reviews, we became aware of the accompanying manuscript that reports the crystal structure of the same molecule, under review at the same time, at the same journal. The authors made their coordinates of atTPC1 (PDBID 5E1J) available¹⁰. The topology of the two structures were identical and their analysis had determined the registration of ~14 additional sites by mutation of residues to cysteine. Therefore we checked our structure for registration against theirs. This led us to several corrections of registration in the sequence and led to improvements in map quality and crystallographic Rfactors (Extended Data Table 3). Anisotropy corrected data ($2.8 \times 4.0 \times 3.3 \text{ \AA}$) were used during the final stages of refinement with an applied isotropic B-factor of -78.72 \AA^2 . Our structure of TPC1 was refined to 2.87 \AA resolution with final $R_{\text{work}}/R_{\text{free}}$ of 29.71% and 33.89%. Analysis by Molprobit shows Ramachandran geometries of 91.8%, 6.9%, and 1.3% for favored, allowed, and outliers. The structure contains 17.0% rotamer outliers.

Domain assignment. Based on experimental electron density (Extended Data Fig. 2b), heavy atom restraints (Extended Data Fig. 2c and Supplementary Data Table 1), and sequence homology (Extended Data Fig. 1), we have assigned the TPC1 domain structure as follows (Fig. 2a, Extended Data Fig. 1): N-terminal domain (NTD; residues 30-55), pre-S1 helix (pre-S1; residues 60-71), Voltage Sensing Domain I (VSDI; S1-S4, residues 72-215), pore domain I (PI;

352 S5-S6, residues 216-301), gate I (residues 302-321), EF-hand (EF; residues 322-399), Domain I-
353 II linker (residues 400-418), pre-S7 helix (pre-S7; residues 419-433), VSDII (S7-S10; residues
354 434-564), pore domain II (PII; S11-S12, residues 565-668), gate II (residues 669-685), C-
355 terminal domain (CTD; residues 686-703). Unstructured regions include part of the NTD
356 (residues 1-29,56-59), S3-S4 (residues 174-183), Domain I-II linker (residues 408-412), S9-S10
357 (residues 519-523), PII (residues 591-593), and CTD (residues 704-733) for a total of 84 missing
358 residues (11.5% unstructured). The structure of TPC1 comprises 649 residues (88.5% of total
359 sequence). Supplementary Data Table 1 summarizes the restraints used for structure
360 determination.

361 Sequence alignments were performed using MUSCLE⁴⁴ and ALINE⁴⁵. Structural figures were
362 made using CHIMERA⁴⁶ and PyMOL. Pore radius calculations using HOLE⁴⁷.

363

References

1. Hedrich, R. & Neher, E. Cytoplasmic calcium regulates voltage-dependent ion channels in plant vacuoles. *Nature* **329**, 833–836 (1987).
2. Calcraft, P. J. *et al.* NAADP mobilizes calcium from acidic organelles through two-pore channels. *Nature* **459**, 596–600 (2009).
3. Hedrich, R. & Marten, I. TPC1 – SV Channels Gain Shape. *Mol. Plant* **4**, 428–441 (2011).
4. Patel, S. Function and dysfunction of two-pore channels. *Sci Signal* **8**, re7–re7 (2015).
5. Wang, X. *et al.* TPC proteins are phosphoinositide- activated sodium-selective ion channels in endosomes and lysosomes. *Cell* **151**, 372–383 (2012).
6. Pitt, S. J. *et al.* TPC2 Is a Novel NAADP-sensitive Ca²⁺ Release Channel, Operating as a Dual Sensor of Luminal pH and Ca²⁺. *J. Biol. Chem.* **285**, 35039–35046 (2010).
7. Sakurai, Y. *et al.* Two-pore channels control Ebola virus host cell entry and are drug targets for disease treatment. *Science* **347**, 995–998 (2015).
8. Hedrich, R., Flügge, U. I. & Fernandez, J. M. Patch-clamp studies of ion transport in isolated plant vacuoles. *FEBS Lett.* **204**, 228–232 (1986).
9. Cang, C., Bekele, B. & Ren, D. The voltage-gated sodium channel TPC1 confers endolysosomal excitability. *Nat. Chem. Biol.* **10**, 463–469 (2014).

- 381 10. Guo, J. *et al.* Structure of the voltage-gated two-pore channel TPC1 from Arabidopsis
382 thaliana. *Nature advance online publication*, (2015).
- 383 11. Beyhl, D. *et al.* The fou2 mutation in the major vacuolar cation channel TPC1 confers
384 tolerance to inhibitory luminal calcium. *Plant J.* **58**, 715–723 (2009).
- 385 12. Cang, C. *et al.* mTOR Regulates Lysosomal ATP-Sensitive Two-Pore Na⁺ Channels to
386 Adapt to Metabolic State. *Cell* **152**, 778–790 (2013).
- 387 13. Morgan, A. J., Davis, L. C., Ruas, M. & Galione, A. TPC: the NAADP discovery channel?
388 *Biochem. Soc. Trans.* **43**, 384–389 (2015).
- 389 14. Schulze, C., Sticht, H., Meyerhoff, P. & Dietrich, P. Differential contribution of EF-
390 hands to the Ca²⁺-dependent activation in the plant two-pore channel TPC1. *Plant J.* **68**,
391 424–432 (2011).
- 392 15. Long, S. B., Campbell, E. B. & Mackinnon, R. Crystal structure of a mammalian
393 voltage-dependent Shaker family K⁺ channel. *Science* **309**, 897–903 (2005).
- 394 16. Payandeh, J., Scheuer, T., Zheng, N. & Catterall, W. A. The crystal structure of a
395 voltage-gated sodium channel. *Nature* **475**, 353–358 (2011).
- 396 17. Naylor, E. *et al.* Identification of a chemical probe for NAADP by virtual screening.
397 *Nat. Chem. Biol.* **5**, 220–226 (2009).

18. Kamb, A., Iverson, L. E. & Tanouye, M. A. Molecular characterization of Shaker, a *Drosophila* gene that encodes a potassium channel. *Cell* **50**, 405–413 (1987).
19. Papazian, D. M., Schwarz, T. L., Tempel, B. L., Jan, Y. N. & Jan, L. Y. Cloning of genomic and complementary DNA from Shaker, a putative potassium channel gene from *Drosophila*. *Science* **237**, 749–753 (1987).
20. Wu, J. *et al.* Structure of the voltage-gated calcium channel Cav1.1 complex. *Science* **350**, aad2395 (2015).
21. Catterall, W. A., Perez-Reyes, E., Snutch, T. P. & Striessnig, J. International Union of Pharmacology. XLVIII. Nomenclature and Structure-Function Relationships of Voltage-Gated Calcium Channels. *Pharmacol. Rev.* **57**, 411–425 (2005).
22. Brailoiu, E. *et al.* Essential requirement for two-pore channel 1 in NAADP-mediated calcium signaling. *J. Cell Biol.* **186**, 201–209 (2009).
23. Peterson, B. Z., Tanada, T. N. & Catterall, W. A. Molecular Determinants of High Affinity Dihydropyridine Binding in L-type Calcium Channels. *J. Biol. Chem.* **271**, 5293–5296 (1996).
24. Pitt, S. J., Lam, A. K. M., Rietdorf, K., Galione, A. & Sitsapesan, R. Reconstituted human TPC1 is a proton-permeable ion channel and is activated by NAADP or Ca²⁺. *Sci. Signal.* **7**, ra46 (2014).

25. Tang, L. *et al.* Structural basis for Ca²⁺ selectivity of a voltage-gated calcium channel.
Nature **505**, 56–61 (2014).
26. Armstrong, C. M. Time Course of TEA⁺-Induced Anomalous Rectification in Squid
Giant Axons. *J. Gen. Physiol.* **50**, 491–503 (1966).
27. Holmgren, M., Smith, P. L. & Yellen, G. Trapping of Organic Blockers by Closing of
Voltage-dependent K⁺ Channels Evidence for a Trap Door Mechanism of Activation
Gating. *J. Gen. Physiol.* **109**, 527–535 (1997).
28. Perozo, E., Cortes, D. M. & Cuello, L. G. Three-dimensional architecture and gating
mechanism of a K⁺ channel studied by EPR spectroscopy. *Nat. Struct. Mol. Biol.* **5**, 459–
469 (1998).
29. Clayton, G. M., Altieri, S., Heginbotham, L., Unger, V. M. & Morais-Cabral, J. H.
Structure of the transmembrane regions of a bacterial cyclic nucleotide-regulated
channel. *Proc. Natl. Acad. Sci.* **105**, 1511–1515 (2008).
30. Zhou, Y., Morais-Cabral, J. H., Kaufman, A. & MacKinnon, R. Chemistry of ion
coordination and hydration revealed by a K⁺ channel–Fab complex at 2.0 [[angst]]
resolution. *Nature* **414**, 43–48 (2001).

31. Mumberg, D., Müller, R. & Funk, M. Regulatable promoters of *Saccharomyces cerevisiae*: comparison of transcriptional activity and their use for heterologous expression. *Nucleic Acids Res.* **22**, 5767–5768 (1994).
32. Hattori, M., Hibbs, R. E. & Gouaux, E. A Fluorescence-Detection Size-Exclusion Chromatography-Based Thermostability Assay for Membrane Protein Precrystallization Screening. *Structure* **20**, 1293–1299 (2012).
33. Gourdon, P. *et al.* HiLiDe—Systematic Approach to Membrane Protein Crystallization in Lipid and Detergent. *Cryst. Growth Des.* **11**, 2098–2106 (2011).
34. Winn, M. D. *et al.* Overview of the CCP4 suite and current developments. *Acta Crystallogr. D Biol. Crystallogr.* **67**, 235–242 (2011).
35. Strong, M. *et al.* Toward the structural genomics of complexes: Crystal structure of a PE/PPE protein complex from *Mycobacterium tuberculosis*. *Proc. Natl. Acad. Sci.* **103**, 8060–8065 (2006).
36. Kabsch, W. Automatic processing of rotation diffraction data from crystals of initially unknown symmetry and cell constants. *J. Appl. Crystallogr.* **26**, 795–800 (1993).
37. Sheldrick, G. M. Experimental phasing with SHELXC/D/E : combining chain tracing with density modification. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 479–485 (2010).

38. Bricogne, G., Vonrhein, C., Flensburg, C., Schiltz, M. & Paciorek, W. Generation, representation and flow of phase information in structure determination: recent developments in and around SHARP 2.0. *Acta Crystallogr. D Biol. Crystallogr.* **59**, 2023–2030 (2003).
39. Abrahams, J. P. & Leslie, A. G. W. Methods used in the structure determination of bovine mitochondrial F1 ATPase. *Acta Crystallogr. D Biol. Crystallogr.* **52**, 30–42 (1996).
40. Cowtan, K. 'dm': An automated procedure for phase improvement by density modification. *Jt. CCP4 ESF-EACBM Newsl. Protein Crystallogr.* **31**, 34–38 (1994).
41. Pedersen, B. P., Morth, J. P. & Nissen, P. Structure determination using poorly diffracting membrane-protein crystals: the H⁺-ATPase and Na⁺,K⁺-ATPase case history. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 309–313 (2010).
42. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta Crystallogr. D Biol. Crystallogr.* **60**, 2126–2132 (2004).
43. Adams, P. D. *et al.* PHENIX : a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 213–221 (2010).
44. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* **32**, 1792–1797 (2004).

45. Bond, C. S. & Schüttelkopf, A. W. ALINE: a WYSIWYG protein-sequence alignment editor for publication-quality alignments. *Acta Crystallogr. D Biol. Crystallogr.* **65**, 510–512 (2009).
46. Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and analysis. *J. Comput. Chem.* **25**, 1605–1612 (2004).
47. Smart, O. S., Neduvélil, J. G., Wang, X., Wallace, B. A. & Sansom, M. S. P. HOLE: A program for the analysis of the pore dimensions of ion channel structural models. *J. Mol. Graph.* **14**, 354–360 (1996).
48. Jones, D. T. Protein secondary structure prediction based on position-specific scoring matrices1. *J. Mol. Biol.* **292**, 195–202 (1999).
49. Cuff, J. A., Clamp, M. E., Siddiqui, A. S., Finlay, M. & Barton, G. J. JPred: a consensus secondary structure prediction server. *Bioinformatics* **14**, 892–893 (1998).
50. Blom, N., Sicheritz-Pontén, T., Gupta, R., Gammeltoft, S. & Brunak, S. Prediction of post-translational glycosylation and phosphorylation of proteins from the amino acid sequence. *PROTEOMICS* **4**, 1633–1649 (2004).
51. Liao, M., Cao, E., Julius, D. & Cheng, Y. Structure of the TRPV1 ion channel determined by electron cryo-microscopy. *Nature* **504**, 107–112 (2013).

484 52. Paulsen, C. E., Armache, J.-P., Gao, Y., Cheng, Y. & Julius, D. Structure of the TRPA1 ion
 485 channel suggests regulatory mechanisms. *Nature* **520**, 511–517 (2015).

486

Acknowledgements. We are grateful to J. Holton and G. Meigs at ALS beamline 8.3.1 and A. Gonzalez at SSRL beamline 12-2 for help with data collection, B. P. Pedersen, B. H. Schmidt, and J. Finer-Moore for critical analysis of the manuscript. This work was supported by NIH grant GM24485 to R.M.S. Beamline 8.3.1 at the Advanced Light Source is operated by the University of California Office of the President, Multicampus Research Programs and Initiatives grant MR-15-328599 and Program for Breakthrough Biomedical Research, which is partially funded by the Sandler Foundation. The Berkeley Center for Structural Biology is supported in part by the National Institutes of Health, National Institute of General Medical Sciences, and the Howard Hughes Medical Institute. The Advanced Light Source is supported by the Director, Office of Science, Office of Basic Energy Sciences, of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. Use of the Stanford Synchrotron Radiation Lightsource, SLAC National Accelerator Laboratory, is supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences under Contract No. DE-AC02-76SF00515. The SSRL Structural Molecular Biology Program is supported by the DOE Office of Biological and Environmental Research, and by the National Institutes of Health, National Institute of General Medical Sciences (including P41GM103393) . The contents of this publication are solely the responsibility of the authors and do not necessarily represent the official views of NIGMS or NIH. Figures were made with Chimera developed by the Resource for Biocomputing,

Visualization, and Informatics at the University of California, San Francisco (supported by NIGMS P41-GM103311). And PyMOL developed by the late Warren DeLano. Mass spectrometry was carried out by the UCSF mass spectrometry facility (supported by NIGMS P41- RR001614). The authors declare no competing financial interests.

Author Contributions. A.F.K performed the experiments, collected and processed the data, determined, refined, and analyzed the structure. R.M.S. supervised the project, and aided in structure determination and analysis. A.F.K and R.M.S. wrote the manuscript.

Author Information. Coordinates and structure factors have been deposited in the Protein Data Bank with the accession number 5DQQ. Correspondence and requests for materials should be addressed to R.M.S. (stroud@msg.ucsf.edu).

516 **Figures**

517




Figure 1 Overview of the TPC1 Structure. **a**, Top view from the luminal side onto the membrane plane and **b**, side view from the right side with the perpendicular to the membrane plane vertical of TPC1 structure labeled by domain and transmembrane helices. E, M, and C denote approximate endolysosomal, membrane, and cytosolic boundaries. The positions of Ca ion binding sites, bound lipids, and NED19 are shown. Approximate geometric dimensions of TPC1 are indicated.

520

521




Figure 2 Structural details of the TPC1 monomer. Structure and domain boundaries in TPC1. **a**, A diagram describing the arrangement of structural domains, colored to match the structures below. Helices are depicted as cylinders and loops as lines. Structural details of TPC1 **b**, domain I and **c**, domain II from three perpendicular views. Phosphorylation sites are shown (*). Charges in S4 (++) and S10 (++++), poly-E (----) in CTDh1, poly-R (++++ in CTDI1, and CxxCxxC (CCC) are shown. Ca^{2+} ions are shown in green and NED19 binding site shown as a purple hexagon.

522

523




Figure 3 Voltage-dependence and its modulation by ions. Side (left) and top (right) views of Ca^{2+} -binding sites in **a**, VSDI and **b**, VSDII. Ca^{2+} ions are shown in green. Ion coordinating residues are labeled in VSDI and II. Voltage-sensing residues are shown in VSDII. Ba^{2+} (cyan) and Yb^{3+} (magenta) isomorphous difference density peaks and atom positions contoured at 10σ numbered according to Extended Data Fig. 2c.




Figure 4 Binding site for trans-NED19. Structural interactions between the TPC1 and NED19. (top) Surface rendering showing the location of the NED19 binding site. (bottom) Molecular details of the NED19 binding-pocket at the interface between PI, PII, and VSDII. Simulated-annealing omit density contoured at 1σ is shown for NED19 (red) and interacting side chains (blue).

524

525




Figure 5 The TPC1 ion channel. Cross-sectional views of the TPC1 ion channel separately through PI and PII. Domains removed for clarity. **a**, PI to PI (orange) and **b**, PII to PII (blue) with sharpened $2mF_o-DF_c$ electron density contoured at 1σ and Ba^{2+} (cyan) and Yb^{3+} (magenta) isomorphous difference density contoured at 10σ . Membrane boundaries are indicated by dashes. **c**, Pore radius calculations through separate pore domains, PI (orange) and PII (blue) using HOLE (See Methods). Approximate boundaries for the putative selectivity filter (SF), cavity (C), Gates I and II, and CTD are shown. Channel axis is vertical.




Figure 6 Cytosolic sensory domains. Structure of EF-hand and coupling with NTD, VSDII, and CTD. Sideviews through **a**, the EF-hand (green), CTD (chartreuse), and NTD (yellow), **b**, constriction site formed by Gate II and CTD, and **c**, interactions between EF-hand and CTD. Phosphorylation sites are indicated by stars (*). Transmembrane and adjacent domains removed for clarity.


528

529

530 **Extended Data Figures**

531

532




533


534

535


Extended Data Figure 1 Sequence alignment of TPCs with TPC1 experimental and predicted secondary structure. A sequence alignment based on seven human and plant TPC orthologs with observed TPC1 (black) and secondary structure predictions by Psipred⁴⁸ (red) and Jpred⁴⁹ (blue). Helices are shown as cylinders, beta sheets as planks, coil as solid lines, and unstructured regions as dashed lines. Level of conservation is indicated by color (>50% yellow; >80% red). Blue dots (•) mark arginines in S4. Red dots (•) mark charge-transfer anions. Green dots (•) mark Ca²⁺-binding sites in the EF-hand. Solid and dashed green lines represent observed and absent peptides from mass spectrometry experiments. Magenta stars (★), orange stars (★), cyan stars (★), blue stars (★), black stars (★), and grey stars (★) mark observed phosphorylation sites using ESI-MS (Extended Data Fig. 3a), potential phosphorylation sites identified by truncation constructs (Extended Data Fig. 3e), predicted phosphorylation sites by NetPhosK⁵⁰, non-phosphorylated sites, unlikely sites due to solvent inaccessibility, and mark unknown or unidentified regions, respectively.




Extended Data Figure 2 Crystal packing of TPC1 and experimental electron densities. **a**, Views of the TPC1 C222₁ crystal lattice viewed along 2-fold axes parallel to **a** (left) and **b** (right). Unit cell boundaries are shown. The asymmetric unit is shown in yellow. **b**, Transverse view of TPC1 is shown with overlaid FOM-weighted experimental electron density calculated using native amplitudes (Native 1) and heavy-atom phases contoured at 1 σ . (left to right) Density modified phases from non-dehydrated derivatives (Extended Data Table 1), dehydrated derivatives (Extended Data Table 2), and combined phases with solvent flattening, histogram matching, phase extension to high resolution native (Native 2), and cross-crystal averaging (See Methods, Extended Data Table 3). **c**, Transverse views of TPC1 with overlaid heavy atom electron densities calculated from isomorphous differences.




Extended Data Figure 3 Determination of phosphorylation sites in TPC1. **a**, Electrospray ionization mass spectrometry (ESI-MS) peptide sequence coverage from an in-gel digest of wild-type (WT) TPC1 by four enzyme combinations (Trypsin/Asp-N, Trypsin/Glu-C, Lys-C, or Chymotrypsin). Measured peptides (Top, red highlight), (Middle) predicted (*) and observed (*) phosphorylation sites are shown. (Bottom) Observed phosphopeptides are listed with the sites of phosphorylation colored red. **b-e**, Polyacrylamide gels of purified TPC1 (10µg) stained with phosphoprotein-specific probe ProDiamond-Q, SyproRuby, or Coomassie (left to right). Molecular weights of standards are indicated in kDa. units to the left. First two lanes are PrecisionPlus protein molecular weight standards and PeppermintStick phosphoprotein molecular weight standards. **b**, WT TPC1 and crystallographic TPC1. **c**, Untreated and treated WT TPC1 with Lambda phosphatase for 1 hour at 25°C. **d**, Schematic of TPC1 truncations (NΔ1; 2-11, NΔ2; 2-21, NΔ3; 2-30, CΔ1; 682-733, CΔ2; 693-733, CΔ3; 709-733, CΔ4; 724-733). **e**, Analysis of TPC1 N- and C-terminal truncations for binding to ProDiamond-Q. NΔ1CΔ1 was unstable during purification.




Extended Data Figure 4 Sequence alignment of TPC1 subdomains with hTPC1, hTPC2, human Ca_v 1.1-1.4, Na_v 1.1, Na_vAb, and hCAM. **a**, S4 voltage-sensing segments. Conserved arginines are highlighted in cyan. Stars (★) mark potential voltage-sensing residues. **b**, S2 segments. Stars (★) mark conserved charge-transfer residues. **c**, S6 segments, pore gate, and poly-E motif. Conserved hydrophobic residues are highlighted in cyan. Dihydropyridine-binding residues in the Ca_v S6 domain III, domain IV, and corresponding residues in TPC1 are highlighted in red and green, respectively. Stars (★) mark the position of phenylalkylamine drug-binding residues in Ca_vs. Conserved residues in the gate and poly-E motif are highlighted in magenta and purple, respectively. **d**, Pore loops. Conserved residues are highlighted in red. **e**, Alignment of hCAM EF-hand domains (EF-hand 1, residues 1-71; EF-hand 2, residues 72-149) and atTPC1 (residues 322-398). Stars (★) mark calcium binding motifs and orange dots (•) mark the interaction site with CTD.



Extended Data Figure 5 Comparison of VSDs between TPC1 and symmetrical ion channels. Structural alignment of S11-S12 segments of TPC1 domain II (blue) with S5-S6 of **a**, TPC1 domain I (red), **b**, NavAb¹⁶ (PDBID 3RVY, grey), **c**, Kv1.2¹⁵ (PDBID 2A79, green), **d**, TRPV1⁵¹ (PDBID 3J5P, orange), and **e**, TRPA1⁵² (PDBID 3J9P, yellow). Angles between S4 segments with respect to S10 of TPC1 domain II are shown.



Extended Data Figure 6 Electrostatic surfaces and Hydrophobic surfaces. (left to right) Top from the luminal side and bottom from the cytoplasmic side views of an **a**, electrostatic surface representation and **b**, surface representation colored according to Kyte-Doolittle hydrophobicity of TPC1. Important domains and residues are labeled. Electrostatic potential and Kyte-Doolittle hydrophobicity without any bound ions was generated using Chimera⁴⁶. The EF-hand and NTD domains are negatively charged and bind cations. The poly-R motif accounts for the positively charged region in the CTD.



Extended Data Figure 7 Mechanism for TPC gating. A schematic summarizing structural features of TPC1 that suggest mechanisms for voltage-sensing, ion permeation, inhibition by NED19, luminal Ca^{2+} -inhibition, and cytosolic Ca^{2+} -activation (EF-hand), and phosphoregulation (NTD/CTD). Ca^{2+} binding (green) and lanthanide (red) binding sites are shown. An ion permeation pathway through the putative selectivity filter, gate, and poly-E and poly-R motifs are summarized. M^{+} represents a general cation (Na^{+} , K^{+} , Ca^{2+}) and + signs are gating charges.

Supplementary Information

Structure, inhibition, and regulation of two-pore channel TPC1 from *Arabidopsis thaliana*

Alexander F. Kintzer and Robert M. Stroud

Supplementary Discussion

Introduction. Two pore channels (TPCs) comprise members of the superfamily of voltage-gated ion channels^{1,2}. Electrophysiological recordings from whole plant vacuoles identified a voltage-gated, slow-vacuolar (SV), calcium-activated calcium current³. The gene encoding the SV channel was later identified and named TPC1 in rats⁴ and plants⁵ based on sequence homology to human voltage-gated Na⁺ (Na_v) and Ca²⁺ (Ca_v) channels⁶. Analogous to Na_v and Ca_v channels, which encode four tandem pore-forming domains on a single chain, TPCs encode two tandem domains separated by a cytosolic domain. Plant TPC1 contains cytosolic EF-hand domains homologous to calmodulin (hCAM, ref.⁷), replaced with a domain of unknown function in human TPCs (Extended Data Figure 1). TPCs show pharmacological similarities to Na_v and Ca_v channels⁸. In plants TPC1 regulates stomatal movement, germination, and long-range stress-induced Ca²⁺ waves^{9–11}. They are non-selective ion channels that conduct Ca²⁺, Na⁺, and K⁺ ions, whereas human TPC1 is Na⁺-selective and human TPC2 conducts Ca²⁺, Na⁺, and H⁺ ions¹². TPC1 functions as the major plant vacuolar calcium-induced calcium release channel¹³. In

animals, however, Ca^{2+} -release by TPC2 from the endolysosome, is subsequently amplified by ryanodine and inositol triphosphate receptors embedded in the endoplasmic reticulum¹⁴. Cytosolic Ca^{2+} ion concentration uniquely potentiates plant TPCs mediated by distinct calmodulin-like domains⁷, not present in mammalian TPCs⁴. TPCs dimerize to form quasi-tetrameric channels composed of alternating copies of the two divergent tandem pore-forming domains (D1 and D2)¹⁵. They have two homologous Shaker-like voltage-sensing domains in transmembrane segments S1-S4, and S7-S10, pore loops in S5-S6 and S11-S12, and activation gates following S6 and S12. D1 and D2 share higher sequence identity with particular domains in Ca_v and Na_v channels, than with one another⁶, suggesting that they are evolutionarily derived from a common ancestor and may share functional mechanisms.

TPCs sense and respond to numerous luminal and cytosolic agents that regulate ion release from the vacuole, or the endolysosome^{3,12,16–24}. These include Ca^{2+} , pH, and cellular redox potential. Plant TPC1 is inhibited by polyunsaturated fatty acids²⁰, whereas the lysosome-specific second messenger lipid $\text{PI}(3,5)\text{P}_2$ increases the open probability of human TPC2, but not TPC1 channels²¹.

The N-terminal domain of plant TPC1 contains a vacuolar localization signal sequence that is conserved with the endolysosomal targeting signal sequence in animals^{25,26}. A binding site on the N-terminus of human TPCs for Rab GTPases serves to regulate trafficking²⁷. The C-

terminus may form a hub for binding of apoptotic machinery²⁸. However, it is not known yet whether plant TPCs participate in protein-protein interactions via the N- or C-termini.

Both plant and human TPCs are known to respond to phosphoregulation^{29–32}. CAM-like dependent kinase, for instance, activates plant TPC1, but phosphorylation at a distinct site by a PKA, PKC, or PKG-like kinase inhibits the channel. mTOR kinase³³ phosphorylates human TPCs and so tunes the release of nutrients and ions from the endolysosome in response to cellular growth signals, stress, or starvation³⁰. Human diseases, including diabetes, obesity, cancer, and age-related diseases result from defective signaling both upstream and downstream of mTOR³³. Targeting the TPC-mTOR complex is therefore an avenue for treatment of various human diseases. Hence phosphoregulation plays an important role in TPC function *in vivo*, as with other ion channels.

Human TPCs form a stable complex with mTOR kinase, facilitating direct phosphorylation of the channel and stabilizing the closed state³⁰. MAPK, JNK, and P38 kinases^{34,35} also inhibit the conductance of hTPC2 by phosphorylation²⁹. LRRK2 kinase has been shown to bind and regulate hTPC2³⁶, though effects on the channel conductance have not been measured. NTDh2 in hTPC2 contains a Rab GTPase-binding motif, not conserved in hTPC1, that facilitates interaction with Rabs and regulates TPC1 trafficking and channel function²⁷. The interaction of plant TPCs with Rab GTPases has not been tested, but it is conceivable that

binding to NTDh2 could affect channel opening. Thus the NTD plays important roles in channel function, phosphoregulation, and intracellular trafficking.

In most eukaryotic cells, though apparently not in plants or yeast, nicotinic acid adenine dinucleotide phosphate (NAADP) triggers the release of Ca^{2+} ions from acidic intracellular stores, causing temporary increases in the cytosolic Ca^{2+} ion concentration^{37,38}. Ryanodine receptors further amplify these Ca^{2+} transients to produce changes in important biological processes ranging from cell differentiation to cardiac function^{32,39,40}. NAADP triggers opening of hTPC2 channels indirectly^{21,22}, probably via a complex with an as yet unidentified NAADP-binding protein²⁷.

Filoviruses, such as Ebola virus require host-cell receptors, endocytosis, proteolytic cleavage, and fusion with the endolysosomal membrane for release of viral material into the cytoplasm^{23,41-43}. TPC antagonists disrupt the intracellular release of Na^+ or Ca^{2+} ions from the endolysosome and inhibit the trafficking of Filoviruses, preventing infection²³. Antagonists trans-NED19 (NED19; ref.⁴⁴), and approved medications of the benzothiazepine, dihydropyridine (DHP), and phenylalkylamine (PAA) classes of L-type Ca_v antagonists inhibit TPCs directly by blocking channel opening²². The most potent anti-viral TPC antagonist, tetraandrine, provides an effective treatment for Ebola disease in animals²³. Both DHP and PAA pharmacophores inhibit plant TPC channels, owing to conserved hydrophobic residues in

transmembrane segment S6⁴⁵⁻⁴⁷. Thus, understanding the basis for inhibition of plant and human TPCs by pharmacophores may help determine the molecular underpinnings and novel avenues for the treatment of critical human diseases.

Determination of Phosphorylation sites in TPC1. We determined that purified plant TPC1 is phosphorylated by binding of a phosphoprotein-specific probe (ProDiamond-Q) to both wild-type TPC1 and TPC1 (Extended Data Figure 3). Treatment with a phosphatase reduces binding of the probe, indicating that the sites of phosphorylation are mostly solvent accessible and reversible (Extended Data Figure 3b).

We determined sites of phosphorylation in plant TPC1 using a combination of mass spectrometry and binding of ProDiamond-Q to N- and C-terminal truncations. Electrospray-ionization mass spectrometry (ESI-MS) analysis of peptides derived from proteolytic digests of gel bands (trypsin/Asp-N, Chymotrypsin, and Lys-C) identified peptides for 87.2% of residues in non-transmembrane regions, 38.1% in transmembrane regions, and 72.6% overall (Extended Data Figure 3a). The ESI-MS data covers most of the predicted solvent exposed sites of phosphorylation, with the exception of several sites in the NTD, P1, S7-S8, CTD, and transmembrane segments. To detect phosphorylation sites within the gaps in ESI-MS coverage, N- (Δ N1-3) and C-terminal (Δ C1-4) truncations of TPC1 (Extended Data Figure 3e) were analyzed for binding to ProDiamond-Q. Deleting part of the C-terminus (Δ N1 Δ C2) abrogated

binding of the probe, suggesting that the CTD may contain sites of phosphorylation and localizes the likely sites to a patch of serine residues in the CTD.

Modulation of VSDs by ions. Voltage-gated ion channels present charged residues in S2 and S4 that respond to the transmembrane electric field. Movement of S4 is mechanically coupled to gating of the channel, requiring at least three regularly spaced arginine residues in S4, and a counter anion in S2^{48,49}. VSDs 1 and 2 in plant TPC1 share roughly 25% sequence identity with each other and with the VSDs in human TPCs and Ca_vs (Extended Data Fig. 4).

Ion selectivity and pore blockade. Lanthanide ions block plant TPCs^{10,16}. The Yb³⁺-bound TPC_{cryst} structure shows direct blockade of the upper pore loop 2 coordinated by four carboxyls (Fig. 5, Extended Data Figure 2c), suggesting that lanthanides may block TPCs by binding near the pore mouth.

Cytoplasmic sensory domains. The plant vacuole functions as the primary intracellular Ca²⁺ store and comprises 90% of the cell volume^{3,5}. Plant TPCs coordinate Ca²⁺-activated release of Ca²⁺ ions from the vacuole to the cytoplasm. The CTD plays a critical role in plant TPCs²⁶, though least conserved across species (Extended Data Fig. 1). Deletion of the CTD results in inactive plant TPC1 channels. Plant TPC channels are known to be sensitive to cytoplasmic redox potential¹⁹. A Hg²⁺ site (Hg1) binds to a cysteine (C347) in EFh1. This site could reasonably function as the redox sensor, which facilitates channel opening in response to

934 lowering of cytoplasmic pH and more reducing conditions. On the luminal side of the membrane
935 a CxxCxxC (C574, C577, C580) motif in S11 (Extended Data Fig. 1) is characteristic of an iron-
936 sulfur cluster-binding site and often linked to redox processes in the endoplasmic reticulum. The
937 function of this site in plant biology has not been described. And the CxxCxxC motif is not
938 conserved in human TPCs.

939

940

941

942 **Supplementary Tables**

943

944

Model	Domain	Secondary Structure	Restraint	Basis	Experimental Electron Density
S2, S3	Domain 1	Helical	E124, D170	Deriv. 8 Yb1 position	Y
S3	Domain 1	Helical	C159	Deriv. 9 Hg2 position	Y
P1	Domain 1	Helical	D269	Deriv. 8 Yb3 position	Y
P1	Domain 1	Helical	E239, D240	Deriv. 4 Ba1 position, Deriv. 8 Yb2 position	
S6, Gate 1, EFh1	Domain 2, EF-hand	Helical	D306, E310	Deriv. 4 Ba1 position	Y
EFh2	EF-hand	Helical	C347, C392	Deriv. 9 Hg1 position	Y
S7-S8	Domain 2	Helical	D454, D457	Deriv. 4 Ba1 position, Deriv. 8 Yb2 position	Y
S10	Domain 2	Helical	E528	Deriv. 4 Ba1 position, Deriv. 8 Yb2 position	Y
S11	Domain 2	Helical	C574, C577, C580, M620	Deriv. 9 Hg3 positions, Deriv. 9 Hg4 positions	Y
P2	Domain 2	Helical	E605, D606	Deriv. 8 Yb3 position	Y
S12, Gate 2, CTD	Domain 2, CTD	Helical	E673, D680	Deriv. 4 Ba1 position	Y

945 **Supplementary Table 1 Building restraints for TPC1.** Experimental restraints used for
946 objective assignment of the TPC1 sequence to electron density.

947

948

949

Supplementary Information References

1. Catterall, W. A. Structure and Function of Voltage-Gated Ion Channels. *Annu. Rev. Biochem.* **64**, 493–531 (1995).
2. Yu, F. H. & Catterall, W. A. The VGL-Chanome: A Protein Superfamily Specialized for Electrical Signaling and Ionic Homeostasis. *Sci STKE* **2004**, re15 (2004).
3. Hedrich, R. & Neher, E. Cytoplasmic calcium regulates voltage-dependent ion channels in plant vacuoles. *Nature* **329**, 833–836 (1987).
4. Ishibashi, K., Suzuki, M. & Imai, M. Molecular Cloning of a Novel Form (Two-Repeat) Protein Related to Voltage-Gated Sodium and Calcium Channels. *Biochem. Biophys. Res. Commun.* **270**, 370–376 (2000).
5. Furuichi, T., Cunningham, K. W. & Muto, S. A Putative Two Pore Channel AtTPC1 Mediates Ca²⁺ Flux in Arabidopsis Leaf Cells. *Plant Cell Physiol.* **42**, 900–905 (2001).
6. Rahman, T. *et al.* Two-pore channels provide insight into the evolution of voltage-gated Ca²⁺ and Na⁺ channels. *Sci Signal* **7**, ra109–ra109 (2014).
7. Schulze, C., Sticht, H., Meyerhoff, P. & Dietrich, P. Differential contribution of EF-hands to the Ca²⁺-dependent activation in the plant two-pore channel TPC1. *Plant J.* **68**, 424–432 (2011).

8. Hashimoto, K., Saito, M., Matsuoka, H., Iida, K. & Iida, H. Functional Analysis of a Rice Putative Voltage-Dependent Ca²⁺ Channel, OsTPC1, Expressed in Yeast Cells Lacking its Homologous Gene CCH1. *Plant Cell Physiol.* **45**, 496–500 (2004).
9. Peiter, E. *et al.* The vacuolar Ca²⁺-activated channel TPC1 regulates germination and stomatal movement. *Nature* **434**, 404–408 (2005).
10. Choi, W.-G., Toyota, M., Kim, S.-H., Hilleary, R. & Gilroy, S. Salt stress-induced Ca²⁺ waves are associated with rapid, long-distance root-to-shoot signaling in plants. *Proc. Natl. Acad. Sci.* **111**, 6497–6502 (2014).
11. Gilroy, S. *et al.* A tidal wave of signals: calcium and ROS at the forefront of rapid systemic signaling. *Trends Plant Sci.* **19**, 623–630 (2014).
12. Pitt, S. J. *et al.* TPC2 Is a Novel NAADP-sensitive Ca²⁺ Release Channel, Operating as a Dual Sensor of Luminal pH and Ca²⁺. *J. Biol. Chem.* **285**, 35039–35046 (2010).
13. Bewell, M. A., Maathuis, F. J. M., Allen, G. J. & Sanders, D. Calcium-induced calcium release mediated by a voltage-activated cation channel in vacuolar vesicles from red beet. *FEBS Lett.* **458**, 41–44 (1999).
14. Gerasimenko, J. V. *et al.* Both RyRs and TPCs are required for NAADP-induced intracellular Ca²⁺ release. *Cell Calcium* doi:10.1016/j.ceca.2015.05.005

- 984 15. Rietdorf, K. *et al.* Two-pore Channels Form Homo- and Heterodimers. *J. Biol. Chem.*
985 **286**, 37058–37062 (2011).
- 986 16. Kawano, T. *et al.* Aluminum as a specific inhibitor of plant TPC1 Ca²⁺ channels.
987 *Biochem. Biophys. Res. Commun.* **324**, 40–45 (2004).
- 988 17. Beyhl, D. *et al.* The fou2 mutation in the major vacuolar cation channel TPC1 confers
989 tolerance to inhibitory luminal calcium. *Plant J.* **58**, 715–723 (2009).
- 990 18. Carpaneto, A., Cantù, A. M. & Gambale, F. Redox agents regulate ion channel activity
991 in vacuoles from higher plant cells. *FEBS Lett.* **442**, 129–132 (1999).
- 992 19. Scholz-Starke, J. *et al.* Redox-dependent modulation of the carrot SV channel by
993 cytosolic pH. *FEBS Lett.* **576**, 449–454 (2004).
- 994 20. Gutla, P. V. K., Boccaccio, A., De Angeli, A., Gambale, F. & Carpaneto, A. Modulation of
995 plant TPC channels by polyunsaturated fatty acids. *J. Exp. Bot.* **63**, 6187–6197 (2012).
- 996 21. Wang, X. *et al.* TPC proteins are phosphoinositide- activated sodium-selective ion
997 channels in endosomes and lysosomes. *Cell* **151**, 372–383 (2012).
- 998 22. Ruas, M. *et al.* Expression of Ca²⁺-permeable two-pore channels rescues NAADP
999 signalling in TPC-deficient cells. *EMBO J.* **34**, 1743–1758 (2015).
- 1000 23. Sakurai, Y. *et al.* Two-pore channels control Ebola virus host cell entry and are drug
1001 targets for disease treatment. *Science* **347**, 995–998 (2015).

- 1002 24. Genazzani, A. A. *et al.* Pharmacological properties of the Ca²⁺-release mechanism
1003 sensitive to NAADP in the sea urchin egg. *Br. J. Pharmacol.* **121**, 1489–1495 (1997).
- 1004 25. Churamani, D., Hooper, R., Rahman, T., Brailoiu, E. & Patel, S. The N-terminal region
1005 of two-pore channel 1 regulates trafficking and activation by NAADP. *Biochem. J.* **453**,
1006 147–151 (2013).
- 1007 26. Larisch, N., Schulze, C., Galione, A. & Dietrich, P. An N-Terminal Dileucine Motif
1008 Directs Two-Pore Channels to the Tonoplast of Plant Cells. *Traffic* **13**, 1012–1022
1009 (2012).
- 1010 27. Lin-Moshier, Y. *et al.* The Two-pore channel (TPC) interactome unmasks isoform-
1011 specific roles for TPCs in endolysosomal morphology and cell pigmentation. *Proc. Natl.*
1012 *Acad. Sci.* **111**, 13087–13092 (2014).
- 1013 28. Lam, A. K. M., Galione, A., Lai, F. A. & Zissimopoulos, S. Hax-1 identified as a two-pore
1014 channel (TPC)-binding protein. *FEBS Lett.* **587**, 3782–3786 (2013).
- 1015 29. Jha, A., Ahuja, M., Patel, S., Brailoiu, E. & Muallem, S. Convergent regulation of the
1016 lysosomal two-pore channel-2 by Mg²⁺, NAADP, PI(3,5)P₂ and multiple protein kinases.
1017 *EMBO J.* **33**, 501–511 (2014).
- 1018 30. Cang, C. *et al.* mTOR Regulates Lysosomal ATP-Sensitive Two-Pore Na⁺ Channels to
1019 Adapt to Metabolic State. *Cell* **152**, 778–790 (2013).

- 1020 31. Bethke, P. C. & Jones, R. L. Reversible protein phosphorylation regulates the activity
1021 of the slow-vacuolar ion channel. *Plant J.* **11**, 1227–1235 (1997).
- 1022 32. Morgan, A. J., Davis, L. C., Ruas, M. & Galione, A. TPC: the NAADP discovery channel?
1023 *Biochem. Soc. Trans.* **43**, 384–389 (2015).
- 1024 33. Laplante, M. & Sabatini, D. M. mTOR Signaling in Growth Control and Disease. *Cell*
1025 **149**, 274–293 (2012).
- 1026 34. Wullschleger, S., Loewith, R. & Hall, M. N. TOR Signaling in Growth and Metabolism.
1027 *Cell* **124**, 471–484 (2006).
- 1028 35. Johnson, G. L. & Lapadat, R. Mitogen-Activated Protein Kinase Pathways Mediated by
1029 ERK, JNK, and p38 Protein Kinases. *Science* **298**, 1911–1912 (2002).
- 1030 36. Gómez-Suaga, P. *et al.* Leucine-rich repeat kinase 2 regulates autophagy through a
1031 calcium-dependent pathway involving NAADP. *Hum. Mol. Genet.* **21**, 511–525 (2012).
- 1032 37. Cancela, J. M., Churchill, G. C. & Galione, A. Coordination of agonist-induced Ca²⁺-
1033 signalling patterns by NAADP in pancreatic acinar cells. *Nature* **398**, 74–76 (1999).
- 1034 38. Calcraft, P. J. *et al.* NAADP mobilizes calcium from acidic organelles through two-
1035 pore channels. *Nature* **459**, 596–600 (2009).
- 1036 39. Galione, A. A primer of NAADP-mediated Ca²⁺ signalling: From sea urchin eggs to
1037 mammalian cells. *Cell Calcium* **58**, 27–47 (2015).

- 1038 40. Patel, S. Function and dysfunction of two-pore channels. *Sci Signal* **8**, re7–re7
1039 (2015).
- 1040 41. Lee, J. E. & Saphire, E. O. Ebolavirus glycoprotein structure and mechanism of entry.
1041 *Future Virol.* **4**, 621–635 (2009).
- 1042 42. Carette, J. E. *et al.* Ebola virus entry requires the cholesterol transporter Niemann-
1043 Pick C1. *Nature* **477**, 340–343 (2011).
- 1044 43. Chandran, K., Sullivan, N. J., Felbor, U., Whelan, S. P. & Cunningham, J. M. Endosomal
1045 Proteolysis of the Ebola Virus Glycoprotein Is Necessary for Infection. *Science* **308**,
1046 1643–1645 (2005).
- 1047 44. Naylor, E. *et al.* Identification of a chemical probe for NAADP by virtual screening.
1048 *Nat. Chem. Biol.* **5**, 220–226 (2009).
- 1049 45. Hockerman, G. H., Johnson, B. D., Abbott, M. R., Scheuer, T. & Catterall, W. A.
1050 Molecular Determinants of High Affinity Phenylalkylamine Block of l-type Calcium
1051 Channels in Transmembrane Segment IIIS6 and the Pore Region of the $\alpha 1$ Subunit. *J. Biol.*
1052 *Chem.* **272**, 18759–18765 (1997).
- 1053 46. Peterson, B. Z., Tanada, T. N. & Catterall, W. A. Molecular Determinants of High
1054 Affinity Dihydropyridine Binding in L-type Calcium Channels. *J. Biol. Chem.* **271**, 5293–
1055 5296 (1996).

- 1056 47. White, P. J. Calcium channels in higher plants. *Biochim. Biophys. Acta BBA* -
1057 *Biomembr.* **1465**, 171–189 (2000).
- 1058 48. Seoh, S.-A., Sigg, D., Papazian, D. M. & Bezanilla, F. Voltage-Sensing Residues in the S2
1059 and S4 Segments of the Shaker K⁺ Channel. *Neuron* **16**, 1159–1167 (1996).
- 1060 49. Aggarwal, S. K. & MacKinnon, R. Contribution of the S4 Segment to Gating Charge in
1061 the Shaker K⁺ Channel. *Neuron* **16**, 1169–1177 (1996).
- 1062