

1 Adaptation of the Arizona Cognitive Task Battery for use with the Ts65Dn Mouse Model
2 of Down syndrome

3 Michael R. Hunsaker*

4 Department of Psychology, University of Utah, Salt Lake City, UT

5 Genevieve K. Smith

6 Department of Psychology, University of Utah, Salt Lake City, UT

7 Raymond P. Kesner**

8 Department of Psychology, University of Utah, Salt Lake City, UT

9 Author Note

10 *Current address: Special Education Department, Granite School District, 2500 S
11 State Street, Salt Lake City, UT 84115

12 **Please send correspondence and requests for offprint copies to: Raymond P.
13 Kesner, Department of Psychology, University of Utah

14 The authors wish to acknowledge Dr. Julie Korenberg for providing access to
15 facilities where these experiments were conducted.

16 All authors declare they have no competing financial or professional interests.

17 This research was supported by NIH grant R01HD067731.

Abstract

18

19 We propose and validate a clear strategy to efficiently and comprehensively characterize
20 neurobehavioral deficits in the Ts65Dn mouse model of Down syndrome. This novel
21 approach uses neurocognitive theory to design and select behavioral tasks that test specific
22 hypotheses concerning Down syndrome. In this manuscript we model in Ts65Dn mice the
23 Arizona Cognitive Task Battery used to study human populations with Down syndrome.
24 We observed specific deficits for spatial memory, impaired long-term memory for visual
25 objects, acquisition and reversal of motor responses, reduced motor dexterity, and
26 impaired adaptive function as measured by nesting and anxiety tasks. The Ts65Dn mice
27 showed intact temporal ordering, novelty detection, and visual object recognition with
28 short delays. These results phenocopy the performance of participants with Down
29 syndrome on the Arizona Cognitive Task Battery. This approach extends the utility of
30 mouse models of Down syndrome by integrating the expertise of clinical neurology and
31 cognitive neuroscience into the mouse behavioral laboratory. Further, by directly
32 emphasizing the reciprocal translation of research between human disease states and the
33 associated mouse models, we demonstrate that it is possible for both groups to mutually
34 inform each others' research to more efficiently generate hypotheses and elucidate
35 treatment strategies.

36 *Keywords:* Down syndrome, Mouse Model, Ts65Dn, Attribute, Spatial Memory,
37 Spatial Processing, Temporal Processing, Sensory/Perceptual Processing, Executive
38 Function, Motor Function, Rule-Based Memory

39 Adaptation of the Arizona Cognitive Task Battery for use with the Ts65Dn Mouse Model
40 of Down syndrome

41 **Introduction**

42 In order to design a battery of behavioral/neurocognitive tasks that could be
43 presented to individuals with Down syndrome across a wide age range in a single testing
44 session, Edgin et al. (2010) developed and validated the Arizona Cognitive Task Battery
45 (ACTB). What makes this battery different than others that are available at present (*e.g.*,
46 Cambridge Neuropsychological Testing Automated Battery (CANTAB)) is that the ACTB
47 has been developed to keep the following issues in mind: 1) when one studies a population
48 with a neurodevelopmental disease, particularly a chromosomal aneuploidy, there is a very
49 real possibility of floor effects confounding analyses of behavioral or cognitive task
50 performance. 2) Additionally, individuals with Down syndrome show language deficits,
51 limiting the tasks that can be used to test cognitive function without a language confound.
52 3) Finally, and perhaps most importantly, the ACTB was developed with the goal of
53 maximizing the sensitivity to identify effects that are present in Down syndrome.

54 The IQ in Down syndrome is typically moderately to severely intellectually disabled
55 range (*i.e.*, IQ = 25-55) and mental age rarely moves beyond 8 years. Paradoxically, it has
56 been suggested that early on, Down syndrome only presents with a mild to moderate
57 intellectual disability (*i.e.*, 55-70), but with age the IQ drops as mental age no longer
58 increases with chronological age (Edgin et al., 2010; Virji-Babul, Kerns, Zhou, Kapur, &
59 Shiffar, 2006).

60 It has been hypothesized that visual-spatial abilities appear to be normal in Down
61 syndrome. However, this appears to be something of an artifact when visual-spatial
62 memory is directly compared to auditory and verbal performance. In tests specifically
63 assessing visual and spatial abilities in Down syndrome, there is a clear deficit relative to
64 typically developing or age matched control populations (Edgin et al., 2010, 2012;
65 Pennington, Moon, Edgin, Stedron, & Nadel, 2003).

66 Within the memory domain, Down syndrome results in deficits for digit or word span
67 as well as general memory deficits with long delays prior to recall. Working memory,
68 specifically verbal working memory, is disrupted in Down syndrome (Edgin, Spano, Kawa,
69 & Nadel, 2014; Pennington et al., 2003; Stedron, Sahni, & Munakata, 2005; Vicari,
70 Bellucci, & Carlesimo, 2005). For visual and spatial memory, it appears that Down
71 syndrome results in specific memory deficits when memory span is increased (Carretti &
72 Lanfranchi, 2010; S Lanfranchi, Carretti, Spano, & Cornoldi, 2009; Silvia Lanfranchi,
73 Cornoldi, Vianello, & Conners, 2004). Again, as suggested by the language deficits, it has
74 been shown that individuals with Down syndrome have greater impairments for verbal
75 than visual-spatial span. Down syndrome also results in long-term memory deficits
76 (Pennington et al., 2003; Vicari, 2006).

77 Despite these memory deficits, implicit memory and perceptual priming appear to be
78 normal (Pennington et al., 2003; Vicari, 2006). This pattern suggests that there is an
79 explicit memory deficit in Down syndrome, meaning that when memory requires temporal
80 or spatial processing, there is a deficit. This has implicated hippocampus and medial
81 temporal lobe function in Down syndrome pathology, as well as the prefrontal cortex for
82 working memory. Implicit memory, dependent upon different brain areas (*e.g.*, parietal
83 cortex), appears to be spared, if not slightly facilitated in Down syndrome compared to
84 other cognitive domains (*i.e.*, word stem or perceptual priming tasks).

85 It has been shown that motor development in Down syndrome is slower than age and
86 mental age matched peers. Intriguingly, early motor markers like rolling and sitting up
87 have been shown to be only very subtly slowed in Down syndrome, but crawling and
88 walking has been shown to be more dramatically delayed. Despite this delay, it does
89 appear that children with Down syndrome develop through the same milestones as
90 typically developing children, these milestones just occur dramatically later in
91 development. Motor skill development appear to show the same developmental delays as
92 these early markers of motor abilities (Connolly & Michael, 1986; U. Frith & Frith, 1974;

93 Gemus et al., 2002; Rast & Harris, 1985; Vicari, 2006; Virji-Babul et al., 2006).

94 To date, the majority of behavioral assays used to test the behavioral phenotype of
95 the mouse models of Down syndrome have focused on spatial memory. More specifically,
96 focus has been placed on the Morris water maze test of spatial memory (Escorihuela et al.,
97 1995; Reeves et al., 1995; Sago et al., 1998). Later experiments have focused on novel
98 object recognition at short and long delays as a proxy for general memory deficits observed
99 across wide range of mouse disease models (Faizi et al., 2011). As a measure of executive
100 function or rostral cortical function, spontaneous alternation has been used
101 (A. M. Kleschevnikov et al., 2012, 2004). The majority of motor tests use the rotarod or
102 locomotor behavior in an open field as the primary measure (Faizi et al., 2011).

103 In this study we propose and then evaluate a clear strategy to efficiently and
104 comprehensively characterize neurobehavioral deficits in the Ts65Dn mouse model of Down
105 syndrome by developing a mouse variant of the Arizona Cognitive Task Battery (Mouse
106 Cognitive Task Battery; mCTB). This approach uses neurocognitive theory to design and
107 select behavioral tasks that test specific hypotheses concerning the genetic disorder being
108 studied-specifically those proposed as part of the Arizona Cognitive Task Battery (ACTB)
109 used to study human populations with Down syndrome (Edgin et al., 2010; Hunsaker,
110 2012a).

111 This approach specifically relies on known anatomical data regarding human and
112 mouse model brain function as important considerations in task design and selection,
113 similar to the ACTB (Edgin et al., 2010). This approach extends the utility of mouse
114 models by integrating the expertise of clinical neurology and cognitive neuroscience into
115 the mouse behavioral laboratory. Further, by directly emphasizing the reciprocal
116 translation of research between human disease states and the associated mouse models, we
117 demonstrate that it is possible for both groups to mutually inform each others' research to
118 more efficiently generate hypotheses and elucidate treatment strategies (*cf.*, Hunsaker,
119 2012a, 2016).

Materials and Methods

Animals

In this study, 10 segmentally trisomic Ts(1716)65Dn (Ts65Dn) male mice and 10 age-matched wildtype littermates were obtained from Jackson Laboratories (Bar Harbor, ME) and tested at 5-7 months of age, weighing 33 +/- 3.8g (standard error). The Ts65Dn/DnJ stock, commercially available from Jackson Laboratory (B6EiC3Sn.BLiA-Ts(1716)65Dn/DnJ), is homozygous for the wildtype allele for retinal degeneration. The stock is maintained by repeated backcrossing of Ts65Dn females to B6EiC3H F1 hybrid males derived from a new congenic strain of C3H mice. This new congenic strain (C3Sn.BLiA-Pde6b+) lacks the blindness causing recessive mutant allele. Animals were kept on a 12-h light/dark cycle, in a temperature and humidity controlled environment with *ad libitum* access to food and water. During no point in experimentation was food deprivation used. Care was taken to assure mice showed motivation to seek sucrose pellet rewards. All behavioral tests were conducted during the light portion of the cycle (06:00-18:00). Mice were housed in same-genotype groups of 2-3 per cage. Animal care and experimental testing procedures conformed to NIH, IACUC, and AALAC standards and protocols.

Experimental Design for Behavioral Testing

The week prior to testing, all animals were handled daily for 15 min sessions and given an opportunity to habituate to a clear and red apparatus for at least 15 min each and acclimate to sucrose pellet rewards. It was verified that prior to the end of this training period that mice consumed sucrose pellets as soon as placed on the apparatus. Behavioral tasks emphasizing exploratory behaviors were presented in a pseudo-randomized order between mice (randomized within the Ts65Dn mice and a 2N wildtype littermate was yoked to a given Ts65Dn mouse to account for any potential task order effects), followed by spontaneous alternation and motor tasks, then response and

146 reversal learning tasks. The 2N wildtype mice were the same age (within 15 days of age)
147 as the Ts65Dn mice.

148 After these tasks, mice received training on the cheeseboard, and then finally were
149 presented with test designed to evaluate quality of life/adaptive functional measures to
150 reduce the influence of any anxiety measures on later task performance.

151 To specifically isolate the contribution of spatial and nonspatial cues to task
152 performance, behavioral tasks were run two times, once in a clear box and many extra
153 maze cues, and a second time in a red box without extra maze cues (Dees & Kesner,
154 2013). This was done because **Smith2014dentate** noticed that there was a pattern of
155 deficits in Ts65Dn mice that were better explained by the mice having access to the
156 extra-maze context than by any specific memory process. As such, they ran every
157 experiment twice, one time using a clear box that allowed access to extra-maze cues and
158 another time in a red box that blocked the view of the extra maze cues. They found that
159 visual object recognition deficits at a 1 hour delay were seen in the clear box experiment,
160 whereas experiments in the red box showed intact visual object memory at a 1 hour delay.
161 They attributed this effect to extra-maze or distal context interfering with the visual
162 object recognition due to interference. Experiments in rats exploring the same effect
163 revealed similar results, and further unpacked the neural correlates of this effect Dees and
164 Kesner (2013). The rationale for this procedure comes from work reported by Smith,
165 Kesner, and Korenberg (2014) in Ts65Dn mice and Edgin et al. (2014) in children with
166 Down syndrome showing that context is particularly influential during object recognition
167 tasks in children with Down syndrome relative to typically developing children. In other
168 words, children with Down syndrome are particularly susceptible to memory interference
169 during cognitive tasks.

170 For every experiment a novel set of objects were used, such that no mouse ever
171 encountered the same object during different experiments. At the end of every experiment,
172 95% ethanol was used to reduce and spread olfactory cues and prevent odor effects

173 impacting future task performance.

174 **Tests of Spatial Attribute**

175 **Spatial Navigation using Cheeseboard.** Apparatus: A white, circular Plexiglas
176 platform with a series of 2 cm diameter holes centered every 5 cm was used as the
177 cheeseboard apparatus. The apparatus was placed approximately 1.5 m off the ground in a
178 space surrounded by extra maze, distal cues to provide a rich spatial context to guide
179 mouse navigation. Paths taken by the mice were recorded by an overhead camera and
180 analyzed using Noldus EthoVision software.

181 Method: Each mouse was habituated to the cheeseboard for 30 min the day prior to
182 experimentation with banana flavored sucrose pellets distributed in each hole (Bio-Serv,
183 #F07257). All mice consumed sucrose pellets and showed a random foraging pattern prior
184 to beginning of training. At the beginning of each trial, a single sucrose reward pellet was
185 placed in one of the holes of the cheeseboard (located within the midpoint of the
186 North-East, North-West, South-East or South-West quadrant). A mouse was then released
187 at one of the cardinal points (*e.g.*, North, South, East, or West at the edge of the
188 cheeseboard) as latency in seconds and distance in centimeters traveled to locate and
189 consume the reward was recorded. Each day, the mouse received a trial from each of the
190 four cardinal directions (order randomized between mice and between days within mice).
191 There were 5 minutes separating each trial for each mouse. After the fourth day of
192 training, the mice were given a probe trial wherein there was no reward. The search
193 patterns of the mice were evaluated. This protocol was modified from the original rat
194 protocol (Kesner, Farnsworth, & DiMattia, 1989) for mice after experiments reported by
195 Lopez, Hauser, Feldon, Gargiulo, and Yee (2010).

196 **Metric/Coordinate Processing.** Apparatus: The apparatus for these
197 experiments consisted of a large Plexiglas box 40 cm wide by 40 cm deep with clear walls
198 40 cm in height and a dark gray floor. An inset made of translucent red Plexiglas 39 cm in

199 width x 39 cm in height was constructed for easy insertion and removal from the original
200 clear box, therefore enabling the experimenter to block distal cues in the testing
201 environment when desired. The box was placed on a circular white table 1 m in diameter.
202 Four distinct two-dimensional black and white cues were placed 30 cm away from each side
203 of the box (methods after Smith et al. (2014)). Exploration was recorded with an overhead
204 video camera and the duration of exploration was measured with a stopwatch. Proximal
205 objects were made from various washable, non-porous materials (plastic, metal, glass,
206 etc.), 2-7 cm in height and had various color, pattern, and textures to ensure each object
207 was visually distinct. New objects were used between experiments so mice were never
208 exposed to the same object during different experiments. To prevent use of olfactory cues
209 to guide behavior, the boxes and objects were disinfected and deodorized with a sterilizing
210 cleaning agent after each use. The mouse was presented with entirely novel object sets for
211 every experiment. All locomotor activity was collected by the Noldus EthoVision software
212 calibrated to measure to the nearest cm (Noldus USA, North Carolina).

213 Method: Each mouse had previously been habituated to clear and red experimental
214 boxes. For the metric/coordinate processing test (Hunsaker, 2012a, 2013; Hunsaker, Kim,
215 Willemsen, & Berman, 2012, 2009; Kesner et al., 2014; Smith et al., 2014), two objects
216 were placed in the box separated by 25 cm (from inner edges) and mice were allowed to
217 explore the objects for 15 minutes. After a 5 min interval during which the mice were
218 covered by an opaque, heavy cup, the objects were moved closer together to an 8 cm
219 separation and the mouse was allowed to explore for 5 min. This procedure was carried
220 out in the clear box that allowed the mouse to see the extra-maze, distal cues as well as in
221 the red box that blocked the ability of the mouse to see these cues (Dees & Kesner, 2013;
222 Smith et al., 2014). Exploration during the last 5 min of habituation and during the 5 min
223 test session were converted into a ratio value ranging [-1,1] to control for overall
224 exploration. As such, a ratio value approaching -1 is interpreted as the mouse showing
225 continued habituation and thus not noticing the change. A ratio value approaching 1

226 suggest the mouse dramatically explored the change.

227 **Topological/Categorical Processing.** Apparatus: This experiment used the
228 same apparatus as the Metric/Coordinate experiment. A similar ratio value was computed
229 as a dependent measure.

230 Method: Each mouse had previously been habituated to clear and red experimental
231 boxes. For the topological/categorical processing test (Hunsaker, 2012a, 2013; Hunsaker
232 et al., 2012, 2009; Kesner et al., 2014; Lee et al., 2009; Smith et al., 2014), four objects
233 were placed in a square in the box separated by 25 cm (from inner edges) and mice were
234 allowed to explore the objects for 15 minutes. After a 5 min interval during which the
235 mice were covered by a heavy cup, the front two objects were transposed, and the mouse
236 was allowed to explore for 5 min. This procedure was carried out in the clear box that
237 allowed the mouse to see the extra-maze, distal cues as well as in the red box that blocked
238 the ability of the mouse to see these cues. Exploration during the last 5 min of
239 habituation and during the 5 min test session were converted into a ratio value ranging
240 [-1,1] to control for overall exploration. As such, a ratio value approaching -1 is interpreted
241 as the mouse showing continued habituation and thus not noticing the change. A ratio
242 value approaching 1 suggest the mouse dramatically explored the change in the object's
243 spatial location and/or distance from each other.

244 **Spatial Location Recognition.** Apparatus: This experiment used the same
245 apparatus as the Metric/Coordinate experiment. A similar ratio value was computed as a
246 dependent measure.

247 Method: Each mouse had previously been habituated to clear and red experimental
248 boxes. For the location recognition test (Smith et al., 2014), two objects were placed in
249 the box separated by 25 cm (from inner edges) and mice were allowed to explore the
250 objects for 15 minutes. After a 5 min interval during which the mice were covered by a
251 heavy cup, one of the objects was moved at a diagonal to a new location (still 25 cm
252 separation between the two objects), and the mouse was allowed to explore for 5 min. This

253 procedure was carried out in the clear box that allowed the mouse to see the extra-maze,
254 distal cues as well as in the red box that blocked the ability of the mouse to see these cues.
255 Exploration during the last 5 min of habituation and during the 5 min test session were
256 converted into a ratio value ranging [-1,1] to control for overall exploration. As such, a
257 ratio value approaching -1 is interpreted as the mouse showing continued habituation and
258 thus not noticing the change. A ratio value approaching 1 suggest the mouse dramatically
259 explored the change in which object occupied which spatial location.

260 **Tests of Temporal Attribute**

261 **Temporal Ordering for Visual Objects.** Apparatus: This experiment used the
262 same apparatus as the Metric/Coordinate experiment. A similar ratio value was computed
263 as a dependent measure.

264 Method: During session 1, two identical copies of a first object (object 1) were placed
265 at the ends of the box 2.5 cm from the end walls and centered between the long walls
266 (Hunsaker, 2013; Hunsaker, Goodrich-Hunsaker, Willemsen, & Berman, 2010, 2012). The
267 mouse was placed in the center of the box facing away from both objects. The mouse was
268 given 5 min to freely explore the objects. After 5 min, the mouse was removed to a small
269 holding cup for 5 min. During this time, the first objects were replaced with two
270 duplicates of a second object (Object 2). For Session 2, the mouse was again placed in the
271 apparatus and allowed to explore. After 5 min, the mouse was removed to the holding cup
272 for 5 min and the objects were replaced with two duplicates of a third object (Object 3).
273 For Session 3, the mouse was given 5 min to explore. After 5 min, the mouse was removed
274 into a small cup for 5 min and an unused copy of the first and an unused copy of the third
275 object were placed into the box. The mouse was again placed into the box and allowed to
276 explore the two objects (*i.e.*, Objects 1 and 3) during a 5 min test session. This procedure
277 was carried out in the clear box that allowed the mouse to see the extra-maze, distal cues
278 as well as in the red box that blocked the ability of the mouse to see these cues.

279 Exploration of each object during the test session were converted into a ratio value ranging
280 [-1,1] to control for overall exploration. As such, a ratio value approaching -1 is interpreted
281 as the mouse showing an absolute preference for the third over the first object. A ratio
282 value approaching 1 suggest the mouse strongly explored the first over the third object.

283 **Temporal Order Control - Novelty Detection for Visual Objects.**

284 Apparatus: This experiment used the same apparatus as the Metric/Coordinate
285 experiment. A similar ratio value was computed as a dependent measure.

286 Method: In addition to reflecting impaired temporal ordering, increased exploration
287 of the first object over the third could also be interpreted as being due to difficulty in
288 remembering the first object prior to the test session (Hunsaker, 2012a, 2013; Hunsaker
289 et al., 2010). To minimize and control for such general memory deficits, a novelty
290 detection of visual objects task was performed. Briefly, on a different day mice received
291 three sessions during which they were allowed to explore three novel sets of objects
292 (Objects 4, 5, 6) similarly to the temporal ordering tasks. During the test session, the first
293 object and a novel fourth object (Object 7) were presented and the mice were allowed 5
294 min to explore. This procedure was carried out in the clear box that allowed the mouse to
295 see the extra-maze, distal cues as well as in the red box that blocked the ability of the
296 mouse to see these cues (*cf.*, Dees and Kesner, 2013; Smith et al., 2014). Exploration of
297 each object during the test session were converted into a ratio value ranging [-1,1] to
298 control for overall exploration. As such, a ratio value approaching -1 is interpreted as the
299 mouse showing an absolute preference for the familiar over the novel object. A ratio value
300 approaching 1 suggest the mouse strongly explored the novel over the familiar object.

301 **Sensory/Perceptual Attribute**

302 **Feature Ambiguity.** Apparatus: This experiment used the same apparatus as the
303 Metric/Coordinate experiment. A similar ratio value was computed as a dependent
304 measure.

305 Method: Each mouse had previously been habituated to clear and red experimental
306 boxes. For the configural recognition condition (Bartko, Winters, Cowell, Saksida, &
307 Bussey, 2007; Bussey, Saksida, & Murray, 2002, 2006; Smith et al., 2014), mice were
308 placed for 15 min in the red box containing two compound objects, A-B and C-D,
309 separated by 15 cm. Following a 5 min delay under a heavy cup, the mouse underwent a
310 5-min Test Phase in which one object from the Study Phase remained the same (A-B) and
311 the other compound object is created from one component of each of the previous familiar
312 objects, (*e.g.*, A-D). That is, the "novel" object (A-D) was composed of the same elements,
313 but rearranged into a novel configuration. Therefore, the object is "novel" by virtue of its
314 configuration, not by its elements, each of which was present in one of the original
315 compound stimuli. Exploration of each compound object was scored as a single unit.
316 Exploration during the last 5 min of habituation and during the 5 min test session were
317 converted into a ratio value ranging [-1,1] to control for overall exploration. As such, a
318 ratio value approaching -1 is interpreted as the mouse showing continued habituation and
319 thus not noticing the change. A ratio value approaching 1 suggest the mouse dramatically
320 explored the change.

321 **Feature Ambiguity Control - Novelty Detection for Configuration of**
322 **Objects.** Apparatus: This experiment used the same apparatus as the
323 Metric/Coordinate experiment. A similar ratio value was computed as a dependent
324 measure.

325 Method: Each mouse had previously been habituated to clear and red experimental
326 boxes. For the configural recognition condition (Bartko et al., 2007; Bussey et al., 2002,
327 2006; Smith et al., 2014), mice were placed for 15 min in the red box containing two
328 compound objects, A-B and C-D, separated by 15 cm. Following a 5 min delay under a
329 heavy cup, the mouse underwent a 5-min control task during which C-D was replaced by
330 two never before seen objects (E-F) was also performed. This procedure was carried out in
331 the clear box that allowed the mouse to see the extra-maze, distal cues as well as in the

332 red box that blocked the ability of the mouse to see these cues. Exploration during the
333 last 5 min of habituation and during the 5 min test session were converted into a ratio
334 value ranging [-1,1] to control for overall exploration. As such, a ratio value approaching -1
335 is interpreted as the mouse showing continued habituation and thus not noticing the
336 change. A ratio value approaching 1 suggest the mouse dramatically explored the change.

337 **Object Recognition at 1 and 24 Hour Delays.** Apparatus: This experiment
338 used the same apparatus as the Metric/Coordinate experiment. A similar ratio value was
339 computed as a dependent measure.

340 Method: Each mouse had previously been habituated to clear and red experimental
341 boxes. For the object recognition test (Moore, Deshpande, Stinnett, Seasholtz, & Murphy,
342 2013; Smith et al., 2014), two objects were placed in the box separated by 25 cm (from
343 inner edges) and mice were allowed to explore the objects for 15 minutes. After a 5 min
344 interval during which the mice were covered by a heavy cup, one of the objects was
345 replaced by a novel object that had never before been experienced by the mouse, and the
346 mouse was allowed to explore for 5 min. This procedure was carried out in the clear box
347 that allowed the mouse to see the extra-maze, distal cues as well as in the red box that
348 blocked the ability of the mouse to see these cues. This procedure was carried out in each
349 box separately for delays of 1 hour and 24 hours. Exploration during the last 5 min of
350 habituation and during the 5 min test session were converted into a ratio value ranging
351 [-1,1] to control for overall exploration. As such, a ratio value approaching -1 is interpreted
352 as the mouse showing continued habituation and thus not noticing the change. a ratio
353 value approaching 1 suggest the mouse dramatically explored the change.

354 **Tests of Executive Function**

355 **Spontaneous Alternation.** Apparatus: For this experiment, a Y maze with each
356 arm measuring 45 cm in length by 30 cm in height with a runway width of 6 cm was used.
357 It was made from opaque gray Plexiglas to prevent the use of any extra-maze cues to

358 guide behavioral performance. As this was a spontaneous alternation task, no rewards
359 were provided at the end of the arms of the Y maze.

360 Method: Mice were placed in the stem of a Y maze and allowed to explore (Faizi
361 et al., 2011; A. M. Kleschevnikov et al., 2012, 2004). Whenever the mouse entered one of
362 the arms of the Y maze with all four limbs their response was recorded. Upon reaching the
363 end of the arm, the mouse was gently picked up and replaced in the stem of the Y maze.
364 The number of times the mouse alternated (*i.e.*, did not repeat the previous turn), was
365 recorded as an alternation.

366 **Response Learning.** Apparatus: For this experiment, a plus maze with each arm
367 measuring 50 cm in length by 25 cm in height with a runway width of 8 cm was used.
368 There was a 2 cm diameter depression at the end of the arms wherein a sucrose pellet was
369 placed to reward a correct response. It was made from opaque gray Plexiglas to prevent
370 the use of any extra-maze cues to guide behavioral performance. At any time the mouse
371 was required to make a 90 degree turn to the right or left to make a choice. The remaining
372 arm was blocked off using a gray Plexiglas block that fit snugly into the arms of the plus
373 maze.

374 Method: Mice were placed in the stem of a plus maze with one of the arms blocked
375 off (forming a T maze). Mice were given five trials to determine if there was any
376 preference for one direction over the other. As no such preference was observed, mice were
377 randomly assigned the rule to turn right or turn left. Mice received 20 trials per day for 4
378 days (Bissonette et al., 2008; Ragozzino, Detrick, & Kesner, 1999, 2002). Entry into an
379 arm with all four limbs was recorded as a choice and mice were not allowed to self correct
380 when they made mistakes. Upon reaching the end of the arm, the mouse was gently picked
381 up and replaced in the stem of the plus maze.

382 **Reversal Learning.** Apparatus: This experiment is a continuation of the
383 Response acquisition experiment and used the same apparatus. For this experiment, the
384 previously rewarded arm was now unrewarded and the previously unrewarded arm was

385 now rewarded by a sucrose pellet.

386 Method: The day after mice finished training on response learning, they received 80
387 trials of reversal training (Bissonette et al., 2008; Ragozzino et al., 1999, 2002). This
388 means that the turn the mice had just learned to make for reward was now incorrect,
389 rather the mice had to make the opposite turn to receive reward. Upon reaching the end
390 of the arm, the mouse was gently picked up and replaced in the stem of the plus maze.
391 Number of previously correct choices made were recorded as errors and error type was
392 evaluated as perseverative or regressive based on the work of Aggleton and Ragozzino
393 (Ragozzino et al., 2002; E. C. Warburton, Baird, Morgan, Muir, & Aggleton, 2001;
394 E. Warburton, Baird, Morgan, Muir, & Aggleton, 2000). Briefly, errors during trials 1-20
395 were considered perseverative errors (perseverating or inflexibly following a previously
396 learned rule) and errors during trials 21-40 were considered regressive errors (regressing or
397 returning to a previously learned rule). Additionally, a behavioral change point algorithm
398 was used to define the point at which each mouse consistently switched their responses
399 from the previously learned rule to the new rule. This was done after the work reported by
400 Diep et al. (2012) by taking the derivative of the learning curve at each point and
401 evaluating when the derivative significantly changed slope (analysis code available at
402 http://www.github.com/mrhunsaker/Change_Point).

403 **Motor Function**

404 **Capellini Handling.** Apparatus: For this experiment, a 250 mL Nalgene beaker
405 was used as a testing environment to assist in video recording mouse behavior. A small
406 mirror was set up behind the beaker and the camera was placed to capture a front and
407 rear view of the mouse to record trials.

408 Method: Mice were habituated over a weekend with approximately 20-30 dried
409 capellini pasta presented in their cages (Tennant et al., 2010). Each mouse was placed in a
410 250 mL beaker and given a 5 cm piece of dried capellini. Their behaviors while eating

411 were recorded for an offline analysis of their motor behaviors. Their latency to finish each
412 piece of pasta was recorded, as were abnormal behaviors including the mouse having its
413 paws together while eating, losing contact with the pasta with one or both paws, and
414 using the mouth to pull the pasta rather than using the digits to feed the pasta into the
415 mouth.

416 **Parallel Rung Walking.** Apparatus: Mice were placed in a box measuring 15 cm
417 wide by 15 cm deep by 45 cm tall with 1.5 mm diameter parallel rungs making up the
418 floor. The rungs were designed with same spacing used by Hunsaker et al. (2011).
419 However, as this was a box rather than a runway, locomotor activity was collected using
420 the Noldus EthoVision software to evaluate any effects of locomotor activity on motor
421 coordination.

422 Method: The mice were allowed to freely explore the box for 5 minutes (Cummings,
423 Engesser-Cesar, Cadena, & Anderson, 2007; Farr, Liu, Colwell, Whishaw, & Metz, 2006;
424 Hunsaker et al., 2011). The number of times a paw slipped through the parallel rod floor
425 beyond the wrist or ankle, a "foot slip" error was recorded (protocol simplified after Farr
426 et al. (2006)). Total number of steps was also recorded to be used as an adjustment factor
427 in later analyses.

428 **Adaptive Function**

429 **Nesting Behaviors.** Apparatus: A 10 cm long piece of 5 cm diameter PVC pipe
430 capped at one end was used as the apparatus. Sawdust similar to that used as mouse
431 bedding was used as a nesting substrate.

432 Method: Sawdust was used to fill a 10 cm long piece of 5 cm diameter PVC pipe
433 that was capped at one end (dry fit, no glue was used). This pipe was placed in a cage
434 with each mouse and the latency to contact the sawdust in the pipe, the latency to start
435 digging in the sawdust, and the latency to finalize the nest were recorded (Filali &
436 Lalonde, 2009).

437 **Neophobia.** Apparatus: The home cage of the mouse, a 35 cm diameter metal
438 platter, and a novel white Plexiglas box measuring 15 cm in all dimension were used to
439 assess neophagia.

440 Method: Mice were given three neophobia tests (specifically hyperneophagia tests)
441 based on the work of Bannerman et al. (2002). The first test was in each mouse's home
442 cage. Each mouse was provided a food they had never encountered (Cheerios cereal) and
443 the latency for the mouse to take the first bite was recorded. The second test was each
444 mouse was placed on a large platter in a bright area in the testing room and the latency
445 for the mouse to take a bite from a reward pellet (familiar food) was recorded. The final
446 test consisted of each mouse being placed in a novel white box and fed a Cheerio that had
447 been stored in a sealed container filled with thyme overnight, resulting in a novel food
448 (Vale-Martínez, Baxter, & Eichenbaum, 2002). Again, latency for the mouse to take the
449 first bite was recorded.

450 **Statistical Methods**

451 **Dependent Measures and Data Visualization.** For the Dry Land Watermaze
452 on the cheeseboard, mean latency to reach the rewarded location as well as total path
453 length were collected using the EthoVision software. The learning curves were normalized
454 to percentage of 1st day latencies and distances to specifically ascertain if there were
455 differences in the shape of the learning curves.

456 For the probe trial, mean distance from the reward location as well as percent time
457 in the quadrant of the cheeseboard containing the previously rewarded location were
458 collected.

459 For all exploratory tasks (Spatial, Temporal, and Sensory/Perceptual tasks), ratio
460 values were computed after the following formula: Exploration of the object of interest (or
461 all objects in the 5 min session of interest) minus the exploration of the other objects or
462 last 5 min of the habituation session. This was divided by the sum of all exploration

463 across both sessions or of both objects. As a formula this is depicted as: $(A-B)/(A+B)$.

464 For the reversal learning, the number of perseverative errors (continuing old rule)
465 during the first 20 (1-20) trials were computed. The number of regressive errors (returning
466 to old rule) were calculated during trials 21-40. A frequentist change point algorithm
467 developed by Gallistel, Fairhurst, and Balsam (2004) and translated in the R programming
468 language by Diep et al. (2012) was used to compute the point at which each mouse showed
469 evidence for having learned to apply the new rule (analysis code available for download at
470 http://github.com/mrhunsaker/Change_Point). This code takes the derivative of the
471 learning curve at every point and determines when the slope has significantly changed.

472 The threshold for significant change was conservatively set at $p < .001$ for the current task.

473 Data were all plotted in DataGraph (4.01 beta, Visual Data Tools, Inc. Chapel Hill,
474 NC.). Ratio data and computed factors are plotted as bar graphs with standard error of
475 the mean (SEM) error bars. Repeated data/learning curves are presented as a line graph
476 at the mean of each block with SEM error bars.

477 **Tests for equal variance and heteroscedasticity.** Prior to statistical analyses,
478 the data were tested for normalcy (Shapiro-Wilk test) and homoscedacity
479 (Browne-Forsythe test) to determine if the data met the assumptions for parametric
480 analyses of variance (ANOVA). Repeated measures were evaluated for sphericity using
481 Mauchly's test of sphericity and necessary adjustments were made using the Huhn-Feldt
482 correction using R 3.2.4 (Team, 2014).

483 **Parametric Statistical Analysis.** Once deemed appropriate, further statistical
484 analyses were performed using parametric analyses of variance (ANOVA). For exploratory
485 task ratios and computed factors were compared using a one-way ANOVA with groups
486 (2N control, Ts65Dn). For acquisition tasks wherein learning was quantified across trials
487 as well as locomotor data, statistical analyses were performed using a mixed model
488 ANOVA with group (2N control, Ts65Dn) as a between groups factor and block of trials as
489 a repeated within factor. If locomotor activity was significantly different between the

490 groups during any trial, locomotor activity was included in the statistical analysis as a
491 covariate.

492 All results were considered significant at an $\alpha < .05$ and Power $(1-\beta) > .80$: Analyses
493 were performed to determine observed power and effect size for all reported effects.
494 Statistical analyses were performed in R 3.2.4 language and environment and observed
495 statistical power was calculated using both R and the statistical program G*Power 3
496 (Faul, Erdfelder, Buchner, & Lang, 2009, 2007). All reported p values were adjusted for
497 False Discovery Rate (Benjamini, Drai, Elmer, Kafkafi, & Golani, 2001; Hunsaker, 2013)
498 using a custom script written in R 3.2.4 (Team, 2014).

499 Results

500 Spatial Attribute

501 **Cheeseboard.** To evaluate spatial navigation and general spatial memory, mice
502 were tested on a dry land version of the Morris water maze (cheeseboard). The Ts65Dn
503 mice showed deficits relative to 2N control mice for raw latency to find reward (Figure 1a;
504 groups ($F(1,76)=185.645$, $p < .0001$), no interaction among group and trial block
505 ($F(1,76)=0.333$, $p = .566$)). These deficits are present as well when the data are adjusted
506 for total latency on trial 1 (groups ($F(1,76)=48.44$, $p < .0001$); Figure 1b) Ts65Dn mice have
507 impaired learning in the Ts65Dn mice in the adjusted data ($F(1,76)=14.74$, $p = .00025$).
508 The same pattern of effects was observed for the data when evaluated for raw distance
509 covered to find reward (Figure 1c; groups ($F(1,76)=88.406$, $p < .0001$) no interaction among
510 group and block ($F(1,76)=0.258$, $p = .613$). Similarly to the latency data, an interaction
511 emerges with Ts65Dn mice showing a shallower learning curve when the data are adjusted
512 for total distance on trial 1 (groups ($F(1,76)=25.194$, $p < .0001$), interaction
513 ($F(1,76)=3.887$, $p = .0523$); Figure 1d).

514 During the probe trial (Figure 1), Ts65Dn mice spent significantly less time in the
515 quadrant where the reward was previously located (Figure 1e, $F(1,18)=91.25$, $p < .0001$).

516 Ts65Dn mice also on average were a further distance away from the previously rewarded
517 spatial location ($F(1,18)=41.7$, $p<.0001$; Figure 1f).

518 **Metric/Coordinate processing.** To evaluate dentate gyrus dependent spatial
519 processing, mice were tested for detection of a metric change (Figure 2a), Ts65Dn mice
520 showed significant impairments relative to 2N control mice. There was a main effect for
521 groups for the clear box ($F(1,18)=39.38$, $p<.0001$) as well as the red box ($F(1,18)=29.94$,
522 $p<.0001$). Deficits in both the clear and red box suggest that metric/coordinate processing
523 is specifically impaired in Ts65Dn mice, supporting earlier reports of dentate gyrus
524 dysfunction in Ts65Dn mice.

525 **Topological/Categorical processing.** To evaluate parietal lobe dependent
526 spatial processing, mice were tested for detection of a topological change (Figure 2b),
527 Ts65Dn mice showed significant impairments relative to 2N control mice. There was a
528 main effect for groups for the clear box ($F(1,18)=78.52$, $p<.0001$) but not for the red box
529 ($F(1,18)=1.489$, $p=.238$). Deficits in only the clear box suggests that topological
530 processing is only impaired when extra-maze cues are present, suggesting a general spatial
531 memory deficit rather than one specific to topological/categorical processing.

532 **Location Recognition.** To test general spatial memory, mice were tested for
533 detection of a change in the spatial location of a visual object (Figure 2c), Ts65Dn mice
534 showed significant impairments relative to 2N control mice. There was a main effect for
535 groups for the clear box ($F(1,18)=36.39$, $p<.0001$) as well as in the red box ($F(1,18)=62.0$,
536 $p<.0001$), suggesting spatial novelty detection deficits in Ts65Dn mice.

537 **Temporal Attribute**

538 **Temporal Ordering of Visual Objects.** To test CA1 function in Ts65Dn mice,
539 mice were tested for a simple temporal ordering task (Figure 2d). Ts65Dn mice did not
540 show significant impairments relative to 2N control mice. There was a main effect for
541 groups for the clear box ($F(1,18)=68.24$, $p<.0001$) but not for the red box ($F(1,18)=2.267$,

542 p=.149). These data suggest that the presence of spatial cues, but not temporal ordering
543 resulted in deficits in the clear box. For the novelty detection task run as a control for
544 temporal ordering (Figure 2e), Ts65Dn mice did not show significant impairments relative
545 to 2N control mice. There was a main effect for groups for the clear box ($F(1,18)=82.78$,
546 $p<.0001$) but not for the red box ($F(1,18)=2.909$, $p=.105$). These data suggest that the
547 presence of spatial cues, but not temporal ordering or novelty detection resulted in deficits
548 in the clear box.

549 **Sensory/Perceptual Attribute**

550 **Feature Ambiguity.** To test perirhinal function in Ts65Dn mice, a configural
551 feature ambiguity test was given (Figure 3a). Ts65Dn mice did not show significant
552 impairments relative to 2N control mice. There was a main effect for groups for the clear
553 box ($F(1,18)=34.13$, $p<.0001$) but not for the red box ($F(1,18)=.021$, $p=.984$). These data
554 suggest that the presence of spatial cues, but not configural feature ambiguity resulted in
555 deficits in the clear box. Ts65Dn mice were not impaired in a configural ambiguity control
556 task (Figure 3b). There was a main effect for groups for the clear box ($F(1,18)=12.27$,
557 $p=.0025$) but not for the red box ($F(1,18)=.012$, $p=.916$). These data suggest that the
558 presence of spatial cues, but not configural feature novelty detection ordering resulted in
559 deficits in the clear box.

560 **Object Recognition after 1 and 24 delays.** Object memory was tested in
561 Ts65Dn mice using object recognition memory at 1 and 24 hours (Figure 3c), Ts65Dn mice
562 did not show significant impairments relative to 2N control mice. There was a main effect
563 for groups for the clear box ($F(1,18)=29.51$, $p<.0001$) but not for the red box
564 ($F(1,18)=.908$, $p=.353$). These data suggest that the presence of spatial cues, but not
565 object recognition resulted in deficits in the clear box. For object recognition memory at
566 24 hours (Figure 3d), there was a main effect for groups for the clear box ($F(1,18)=46.23$,
567 $p<.0001$) as well as for the red box ($F(1,18)=31.36$, $p<.0001$). These data suggest that at

568 24 hours, the Ts65Dn mice were unable to retrieve the memory for the object, whereas
569 they were able to do so at 1 hour.

570 **Executive Function**

571 **Spontaneous Alternation.** Spontaneous alternation was used to test working
572 memory in the Ts65Dn mice (Figure 4a). Ts65Dn mice showed fewer alternations than 2N
573 control mice ($F(1,18)=23.85$, $p=.0001$).

574 **Rule Learning on a Plus Maze.** To evaluate inhibitory control and the ability
575 to learn a turn response (Figure 4b), Ts65Dn mice took significantly longer to learn the
576 rule than 2N control mice. There was a main effect for groups ($F(1,76)=4.24$, $p=.013$), a
577 main effect for block of trials ($F(1,76)=502.86$, $p<.0001$). There was also an interaction
578 among group and block ($F(1,76)=7.82$, $p=.0065$). This interaction was the result of the
579 Ts65Dn mice taking longer to learn the rule. For the final block of 20 trials, there were no
580 differences in performance for Ts65Dn and 2N control mice.

581 **Rule Reversal Learning on a Plus Maze.** To evaluate rule reversal learning
582 (behavioral flexibility) in Ts65Dn mice, the reversal of a turn response was evaluated
583 (Figure 4c). Ts65Dn mice took a significantly greater number of trials to learn the rule
584 than 2N control mice. There was a main effect for groups ($F(1,76)=4.952$, $p=.029$), a main
585 effect for block of trials ($F(1,76)=24.62$, $p<.0001$). There was also a nonsignificant
586 interaction among group and block ($F(1,76)=3.21$, $p=.077$). Looking at Figure 4c, the
587 nonsignificant interaction was the result of the Ts65Dn mice taking longer to learn to
588 reverse the rule. In fact, the Ts65Dn mice were only impaired relative to the 2N control
589 mice for the first block of 20 trials. For the remaining blocks of trials there were no
590 differences in performance for Ts65Dn and 2N control mice. There was a main effect for
591 groups for the trial at which the mice changed preference from old rule to new rule
592 (change point; $F(1,18)=21.43$, $p=.0002$; Figure 4d). For the first 20 trials of reversal
593 learning, Ts65Dn mice showed a greater number of perseverative errors ($F(1,18)=11.98$,

594 $p=.0028$; Figure 4e). For trials 21-40, there was no difference between Ts65Dn mice and
595 2N control mice for regressive errors ($F(1,18)=.287$, $p=.599$; Figure 4f).

596 **Motor Function**

597 **Capellini Eating Task.** For the capellini task of manual dexterity (Figure 5),
598 Ts65Dn mice showed significant impairments relative to 2N control mice. There was a
599 main effect for latency, with Ts65Dn mice taking longer to eat the pasta on average
600 ($F(1,18)=14.74$, $p=.0012$; Figure 5a). Ts65Dn mice also made a greater number of pasta
601 handling errors ($F(1,18)=92.68$, $p<.0001$; Figure 5b). There was also a main effect for
602 groups for the number of times the paws came together ($F(1,18)=42.34$, $p<.0001$;
603 Figure 5c), for the number of times the mouse lost contact with the pasta ($F(1,18)=20.35$,
604 $p=.0003$; Figure 5d) and the number of times the mouse pulled the pasta with their mouth
605 rather than using the hands to move it ($F(1,18)=21.46$, $p=.0002$; Figure 5e).

606 **Parallel Rung Walking Task.** During a parallel rung walking task (Figure 5f),
607 Ts65Dn mice showed significant impairments relative to 2N control mice. There was a
608 main effect for the number of foot slips in a 1 minute session ($F(1,18)=27.32$, $p<.0001$).
609 When adjusted for number of steps, Ts65Dn mice still showed a greater number of foot
610 slip errors ($F(1,18)=11.70$, $p=.0031$; Figure 5g).

611 **Adaptive Function / Quality of Life**

612 **Nesting Behavior.** Ts65Dn mice showed significant impairments relative to 2N
613 control mice for measures of nesting (Figure 6). Ts65Dn mice took longer to make contact
614 with the nesting material ($F(1,18)=152.9$, $p<.0001$; Figure 6a), for the time it took for
615 them to dig in the media (measured from time of first contact) ($F(1,18)=318.6$, $p<.0001$;
616 Figure 6b), and the time it took from starting to dig to finish the nest ($F(1,18)=94.3$,
617 $p<.0001$; Figure 6c).

618 **Neophobia.** Ts65Dn mice showed significant impairments relative to 2N control
619 mice for neophobia (Figure 6). Ts65Dn mice took longer to eat a novel food in a familiar

620 environment ($F(1,18)=19.59$, $p=.0003$; Figure 6d), took longer to eat a familiar food in a
621 novel environment ($F(1,18)=40.87$, $p<.0001$; Figure 6e), and took longer to eat a novel
622 food in a novel environment ($F(1,18)=83.74$, $p<.0001$; Figure 6f).

623 Discussion

624 Briefly, Ts65Dn mice displayed specific deficits for spatial processing, long-term
625 memory, motor function, executive function, and adaptive function (Table 1). These
626 deficits phenocopy the results from the ACTB used in testing children with Down
627 syndrome, including the report that providing distracting contextual cues may impair
628 memory function in Down syndrome (Edgin et al., 2010, 2012, 2014).

629 Overall, these data clearly demonstrate that the Ts65Dn mouse do in fact show a
630 similar pattern of behavioral deficits on the mouse variant of the Arizona Cognitive Task
631 Battery (mCTB) as individuals with Down syndrome show on the human ACTB. The
632 task similarities between the mouse and human ACTB are outlined in Table 1. In cases
633 where Down syndrome participants show deficits on the ACTB (Edgin et al., 2010), the
634 mice in the present study phenocopy those effects (also *cf.*, Edgin et al. (2012)). Similarly,
635 the Ts65Dn mice showed the same pattern of strengths as Down syndrome individuals
636 showed on the ACTB.

637 The pattern of Ts65Dn performance on spatial and temporal processing tasks
638 support the hypothesis that Ts65Dn mice show clear deficits for spatial processing tasks
639 dependent upon the dentate gyrus with sparing of spatial and temporal processing
640 dependent upon the CA1 subregion (Goodrich-Hunsaker, Hunsaker, & Kesner, 2008;
641 Kesner, Lee, & Gilbert, 2004; Kesner & Rolls, 2015; Rolls & Kesner, 2006; Smith et al.,
642 2014). Similarly, it appears that spatial processing dependent on neocortical processing is
643 spared (*cf.*, Goodrich-Hunsaker, Hunsaker, and Kesner (2005)).

644 These findings were confirmed by verifying that any spatial or temporal processing
645 deficits observed in the presence of distal cues was confirmed in a task that removed these

646 cues (Dees & Kesner, 2013). The data show that metric/coordinate processing and
647 location recognition deficits are similar in the presence or absence of distal cues, suggesting
648 that these hippocampus (more specifically the dentate gyrus) dependent spatial processes
649 are disrupted. The topological/categorical deficits observed in the clear box are absent
650 when tested in the absence of extramaze cues in a red box. These data suggest that
651 CA1/parietal cortex related spatial memory processes are intact when tested without
652 extra-maze cues available (*cf.*, Kesner et al. (2004), Kesner and Rolls (2015)).

653 Similarly, the temporal ordering deficits present in the clear box were absent in the
654 red box, and the novelty detection control task showed the same pattern, suggesting
655 temporal processing is intact in the Ts65Dn mice, but object identification may be
656 impaired if extra-maze distal cues are present. This hypothesis was confirmed in the
657 sensory/perceptual tests wherein the Ts65Dn mice were able to correctly process feature
658 ambiguity and feature novelty in the red, but not clear boxes. And finally, object
659 recognition was impaired even at only 1 hour delays for Ts65Dn mice when extramaze cues
660 were available. In the red box, the Ts65Dn mice were able to identify previously
661 encountered objects until a 24 hour delay was imposed.

662 For response learning or executive function, Ts65Dn mice were impaired for
663 spontaneous alternation (they alternated on fewer trials than wildtype mice), as well as
664 response learning and reversal learning of a previously learned rule. However, it appeared
665 that the Ts65Dn mice just learned the tasks more slowly since the early trials show deficit,
666 but later blocks of trials do not. For reversal learning, it is clear the Ts65Dn mice take a
667 greater number of trials to learn the reversal based on the changepoint calculated for the
668 learning curves (Ts65Dn mean=50 compared to mean=30 for 2N wildtype mice) as well as
669 the greater number of perseverative errors during trials 1-20 of the reversal learning task.
670 Interestingly, once the Ts65Dn mice showed learning of the reversal, they did not make
671 any more regressive errors than the 2N control mice.

672 These data support earlier theories that suggested there were specific deficits to

673 spatial memory in Down syndrome (Carlesimo, Marotta, & Vicari, 1997; Carretti &
674 Lanfranchi, 2010; S Lanfranchi et al., 2009; Silvia Lanfranchi et al., 2004; Vicari et al.,
675 2005; Visu-Petra, Benga, & Miclea, 2007). What these data clarify are the neural
676 substrates and specific domains of medial temporal lobe function are impaired in Down
677 syndrome. There are specific deficits on tasks that test dentate gyrus function, but sparing
678 of function on tasks that test parietal and perirhinal cortices as well as CA1 function.
679 Similarly, there are specific deficits in the Ts65Dn mouse that are attributable to
680 cerebellar function and executive functional deficits attributable to the rostral cortices
681 (analogue of the human prefrontal cortex). For thorough descriptions of neuroanatomic
682 correlates of the behavioral tasks included in the mCTB the reader is referred to the
683 descriptions of the original tasks (*cf.*, Bartko et al. (2007), Bussey et al. (2002), Kesner
684 et al. (2004), Kesner and Rolls (2015), Ragozzino et al. (1999, 2002), Rolls and Kesner
685 (2006)

686 For the motor tasks, the Ts65Dn mice showed clear deficits for handling the capellini
687 and greater difficulties walking on parallel rungs. For adaptive function, the Ts65Dn mice
688 took longer to build nests and consume novel foods in novel locations, suggesting reduced
689 adaptive function or quality of life relative to 2N control mice.

690 An important consideration in adopting a behavioral screen like this mCTB is the
691 relative throughput for the tasks. All of the tasks used to test medial temporal lobe
692 function take 30 minutes per session of testing, and can be repeated numerous times on
693 any given mouse after 24 hours have passed since the first test. The motor and adaptive
694 function tests are similarly high throughput, as is the spontaneous alternation task. The
695 only tasks that require a significant time investment are the dry land watermaze (Lopez
696 et al., 2010) on the cheeseboard and the rule acquisition and rule reversal learning tasks
697 (Bissonette et al., 2008; Ragozzino et al., 1999, 2002). The dry land watermaze task on
698 the cheeseboard follows a standard water maze protocol that lasts 5 days, and the
699 response learning and reversal learning tasks together take an additional week.

700 A second consideration is adopting the mCTB is the advantage of the anatomical
701 specificity of known neural substrates underlying each behavioral task (Bartko et al., 2007;
702 Bussey et al., 2002, 2006; Farr et al., 2006; Goodrich-Hunsaker et al., 2005, 2008;
703 Hunsaker, 2012a; Kesner et al., 2004; Kesner & Rolls, 2015) and previous comparison of
704 rodent performance on many of the behavioral tasks to human cognitive function
705 (Baumann, Chan, & Mattingley, 2012; Baumann & Mattingley, 2013; Goodrich-Hunsaker
706 & Hopkins, 2010; Goodrich-Hunsaker et al., 2005; Kesner & Goodrich-Hunsaker, 2010). As
707 such, these tasks can be used to dissociate function of brain areas within the mouse
708 models being tested. The final consideration is the lack of negative reinforcement or
709 aversive stimulus. This means mouse models displaying depression, anxiety, or anhedonia
710 are theoretically testable using the mCTB (*cf.*, Hunsaker (2012a, 2012b)).

711 An interesting complication emerged in the data that the mCTB was solved by
712 nature of how it was designed. On a number of nonspatial tasks, there was a confound of
713 distal cues interfering with the processing of proximal objects that were of interest in the
714 task. For example, in the temporal ordering and novelty detection for novel objects tasks,
715 the Ts65Dn mice looked like they had deficits, but only in the clear box that allowed
716 access to distal cues (Dees & Kesner, 2013; Smith et al., 2014). The feature ambiguity
717 task and the control condition showed the same pattern. The addition of a distal cue-free
718 condition (the red box) was essential for separating the effects of proximal-distal cue
719 interactions from the memory processes being tested by the tasks. The disparate
720 performance across clear and red boxes (or in presence of absence of extra maze contextual
721 cues) allowed us to assess the role of context and distracting cues in memory function in
722 Ts65Dn mice, a conceptual replication of Edgin et al. (2014) in Down syndrome and rats
723 as shown by Dees and Kesner (2013).

724 **Limitations**

725 The primary limitation of the present study is the lack of tests for language or
726 language like attributes in the Ts65Dn mouse model. However, such assays exist and can
727 easily be added to the task battery without significantly increasing the amount of time
728 required to perform the mCTB (Zampieri, Fernandez, Pearson, Stasko, & Costa, 2014).
729 The present experiment also only assayed the Ts65Dn mouse model of Down syndrome as
730 a proof of concept. Further studies will be necessary to evaluate whether other mouse
731 models of Down syndrome (*e.g.*, Ts2Cje, Ts1Yah, and Dep(17)1Yey/+; Das and Reeves
732 (2011)) show the same pattern of results as the Ts65Dn mouse model.

733 **Conclusions**

734 That deficits in the mouse and human ACTB are comparable suggests that the
735 mCTB may be useful for guiding the development of treatment strategies by providing
736 reliable, valid behavioral endpoints and outcome measures. These outcome measures
737 reported in the mCTB appear to show high face, content, and predictive validity with the
738 ACTB, at least so far as Ts65Dn performance mimics the performance of Down syndrome
739 patient populations. As we were able to identify such a clear phenotype in Ts65Dn mice,
740 the mouse mCTB may well turn out to be a useful tool for studying behavioral prodrome
741 of early Alzheimer-like pathology and cognitive decline in mouse models related to Down
742 syndrome. Similarly, the mCTB may serve as a powerful and comprehensive screening tool
743 for preclinical tests of pharmacological interventions in Down syndrome.

744

References

745

746 Bannerman, D., Deacon, R., Offen, S., Friswell, J., Grubb, M., & Rawlins, J. (2002).

747 Double dissociation of function within the hippocampus: spatial memory and

748 hyponeophagia. *Behavioral neuroscience*, *116*(5), 884.

749 Bartko, S. J., Winters, B. D., Cowell, R. A., Saksida, L. M., & Bussey, T. J. (2007).

750 Perirhinal cortex resolves feature ambiguity in configural object recognition and

751 perceptual oddity tasks. *Learning & Memory*, *14*(12), 821–832.

752 Baumann, O., Chan, E., & Mattingley, J. B. (2012). Distinct neural networks underlie

753 encoding of categorical versus coordinate spatial relations during active navigation.

754 *Neuroimage*, *60*(3), 1630–1637.

755 Baumann, O. & Mattingley, J. B. (2013). Dissociable roles of the hippocampus and

756 parietal cortex in processing of coordinate and categorical spatial information.

757 *Frontiers in human neuroscience*, *8*, 73–73.

758 Benjamini, Y., Drai, D., Elmer, G., Kafkafi, N., & Golani, I. (2001). Controlling the false

759 discovery rate in behavior genetics research. *Behavioural brain research*, *125*(1),

760 279–284.

761 Bissonette, G. B., Martins, G. J., Franz, T. M., Harper, E. S., Schoenbaum, G., &

762 Powell, E. M. (2008). Double dissociation of the effects of medial and orbital

763 prefrontal cortical lesions on attentional and affective shifts in mice. *The Journal of*

764 *Neuroscience*, *28*(44), 11124–11130.

765 Bussey, T. J., Saksida, L. M., & Murray, E. A. (2002). Perirhinal cortex resolves feature

766 ambiguity in complex visual discriminations. *European Journal of Neuroscience*,

767 *15*(2), 365–374.

768 Bussey, T. J., Saksida, L. M., & Murray, E. A. (2006). Perirhinal cortex and

769 feature-ambiguous discriminations. *Learning & Memory*, *13*(2), 103–105.

- 770 Carlesimo, G. A., Marotta, L., & Vicari, S. (1997). Long-term memory in mental
771 retardation: evidence for a specific impairment in subjects with down's syndrome.
772 *Neuropsychologia*, *35*(1), 71–79.
- 773 Carretti, B. & Lanfranchi, S. [S]. (2010). The effect of configuration on vswm performance
774 of down syndrome individuals. *Journal of Intellectual Disability Research*, *54*(12),
775 1058–1066.
- 776 Connolly, B. H. & Michael, B. T. (1986). Performance of retarded children, with and
777 without down syndrome, on the bruininks oseretsky test of motor proficiency.
778 *Physical Therapy*, *66*(3), 344–348.
- 779 Cummings, B. J., Engesser-Cesar, C., Cadena, G., & Anderson, A. J. (2007). Adaptation
780 of a ladder beam walking task to assess locomotor recovery in mice following spinal
781 cord injury. *Behavioural brain research*, *177*(2), 232–241.
- 782 Das, I. & Reeves, R. H. (2011). The use of mouse models to understand and improve
783 cognitive deficits in down syndrome. *Disease Models and Mechanisms*, *4*(5), 596–606.
- 784 Dees, R. L. & Kesner, R. P. (2013). The role of the dorsal dentate gyrus in object and
785 object-context recognition. *Neurobiology of learning and memory*, *106*, 112–117.
- 786 Diep, A. A., Hunsaker, M. R., Kwock, R., Kim, K., Willemsen, R., & Berman, R. F.
787 (2012). Female cgg knock-in mice modeling the fragile x premutation are impaired
788 on a skilled forelimb reaching task. *Neurobiology of learning and memory*, *97*(2),
789 229–234.
- 790 Edgin, J. O., Mason, G. M., Allman, M. J., Capone, G. T., DeLeon, I., Maslen, C., . . .
791 Nadel, L. (2010). Development and validation of the arizona cognitive test battery
792 for down syndrome. *Journal of neurodevelopmental disorders*, *2*(3), 149.
- 793 Edgin, J. O., Mason, G. M., Spano, G., Fernández, A., & Nadel, L. (2012). 7 human and
794 mouse model cognitive phenotypes in down syndrome: implications for assessment.
795 *Progress in brain research*, *197*, 123.

- 796 Edgin, J. O., Spano, G., Kawa, K., & Nadel, L. (2014). Remembering things without
797 context: development matters. *Child development*, *85*(4), 1491–1502.
- 798 Escorihuela, R. M., Fernández-Teruel, A., Vallina, I. F., Baamonde, C., Lumbreras, M. A.,
799 Dierssen, M., . . . Flórez, J. (1995). A behavioral assessment of ts65dn mice: a
800 putative down syndrome model. *Neuroscience letters*, *199*(2), 143–146.
- 801 Faizi, M., Bader, P. L., Tun, C., Encarnacion, A., Kleschevnikov, A., Belichenko, P., . . .
802 Mobley, W. C., et al. (2011). Comprehensive behavioral phenotyping of ts65dn
803 mouse model of down syndrome: activation of beta 1-adrenergic receptor by
804 xamoterol as a potential cognitive enhancer. *Neurobiology of Disease*, *43*(2),
805 397–413.
- 806 Farr, T. D., Liu, L., Colwell, K. L., Whishaw, I. Q., & Metz, G. A. (2006). Bilateral
807 alteration in stepping pattern after unilateral motor cortex injury: a new test
808 strategy for analysis of skilled limb movements in neurological mouse models.
809 *Journal of neuroscience methods*, *153*(1), 104–113.
- 810 Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical power analyses
811 using g* power 3.1: tests for correlation and regression analyses. *Behavior research*
812 *methods*, *41*(4), 1149–1160.
- 813 Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G* power 3: a flexible
814 statistical power analysis program for the social, behavioral, and biomedical sciences.
815 *Behavior research methods*, *39*(2), 175–191.
- 816 Filali, M. & Lalonde, R. (2009). Age-related cognitive decline and nesting behavior in an
817 appswe/ps1 bigenic model of alzheimer’s disease. *Brain research*, *1292*, 93–99.
- 818 Frith, U. & Frith, C. D. (1974). Specific motor disabilities in downs syndrome. *Journal of*
819 *Child Psychology and Psychiatry*, *15*(4), 293–301.
- 820 Gallistel, C. R., Fairhurst, S., & Balsam, P. (2004). The learning curve: implications of a
821 quantitative analysis. *Proceedings of the national academy of Sciences of the united*
822 *States of america*, *101*(36), 13124–13131.

- 823 Gemus, M., Palisano, R., Russell, D., Rosenbaum, P., Walter, S. D., Galuppi, B., &
824 Lane, M. (2002). Using the gross motor function measure to evaluate motor
825 development in children with down syndrome. *Physical & Occupational Therapy in*
826 *Pediatrics*, *21*(2-3), 69–79.
- 827 Goodrich-Hunsaker, N. J. & Hopkins, R. O. (2010). Spatial memory deficits in a virtual
828 radial arm maze in amnesic participants with hippocampal damage. *Behavioral*
829 *neuroscience*, *124*(3), 405.
- 830 Goodrich-Hunsaker, N. J., Hunsaker, M. R., & Kesner, R. P. (2005). Dissociating the role
831 of the parietal cortex and dorsal hippocampus for spatial information processing.
832 *Behavioral neuroscience*, *119*(5), 1307.
- 833 Goodrich-Hunsaker, N. J., Hunsaker, M. R., & Kesner, R. P. (2008). The interactions and
834 dissociations of the dorsal hippocampus subregions: how the dentate gyrus, ca3, and
835 ca1 process spatial information. *Behavioral neuroscience*, *122*(1), 16.
- 836 Hunsaker, M. R. (2012a). Comprehensive neurocognitive endophenotyping strategies for
837 mouse models of genetic disorders. *Progress in neurobiology*, *96*(2), 220–241.
- 838 Hunsaker, M. R. (2012b). The importance of considering all attributes of memory in
839 behavioral endophenotyping of mouse models of genetic disease. *Behavioral*
840 *neuroscience*, *126*(3), 371.
- 841 Hunsaker, M. R. (2013). Neurocognitive endophenotypes in cgg ki and fmr1 ko mouse
842 models of fragile x-associated disorders: an analysis of the state of the field.
843 *F1000Research*, *2*.
- 844 Hunsaker, M. R. (2016). Applying the attribute model to develop behavioral tasks that
845 phenocopy human clinical phenotypes using mouse disease models: an
846 endophenotyping approach. In *The neurobiological basis of memory* (pp. 337–366).
847 Springer.

- 848 Hunsaker, M. R., Goodrich-Hunsaker, N. J., Willemsen, R., & Berman, R. F. (2010).
849 Temporal ordering deficits in female cgg ki mice heterozygous for the fragile x
850 premutation. *Behavioural brain research*, *213*(2), 263–268.
- 851 Hunsaker, M. R., Kim, K., Willemsen, R., & Berman, R. F. (2012). Cgg trinucleotide
852 repeat length modulates neural plasticity and spatiotemporal processing in a mouse
853 model of the fragile x premutation. *Hippocampus*, *22*(12), 2260–2275.
- 854 Hunsaker, M. R., von Leden, R. E., Ta, B. T., Goodrich-Hunsaker, N. J., Arque, G.,
855 Kim, K., . . . Berman, R. F. (2011). Motor deficits on a ladder rung task in male and
856 female adolescent and adult cgg knock-in mice. *Behavioural brain research*, *222*(1),
857 117–121.
- 858 Hunsaker, M. R., Wenzel, H. J., Willemsen, R., & Berman, R. F. (2009). Progressive
859 spatial processing deficits in a mouse model of the fragile x premutation. *Behavioral*
860 *neuroscience*, *123*(6), 1315.
- 861 Kesner, R. P., Farnsworth, G., & DiMattia, B. V. (1989). Double dissociation of egocentric
862 and allocentric space following medial prefrontal and parietal cortex lesions in the
863 rat. *Behavioral neuroscience*, *103*(5), 956.
- 864 Kesner, R. P. & Goodrich-Hunsaker, N. J. (2010). Developing an animal model of human
865 amnesia: the role of the hippocampus. *Neuropsychologia*, *48*(8), 2290–2302.
- 866 Kesner, R. P., Hui, X., Sommer, T., Wright, C., Barrera, V. R., & Fanselow, M. S. (2014).
867 The role of postnatal neurogenesis in supporting remote memory and spatial metric
868 processing. *Hippocampus*, *24*(12), 1663–1671.
- 869 Kesner, R. P., Lee, I., & Gilbert, P. (2004). A behavioral assessment of hippocampal
870 function based on a subregional analysis. *Reviews in the Neurosciences*, *15*(5),
871 333–352.
- 872 Kesner, R. P. & Rolls, E. T. (2015). A computational theory of hippocampal function, and
873 tests of the theory: new developments. *Neuroscience & Biobehavioral Reviews*, *48*,
874 92–147.

- 875 Kleschevnikov, A. M., Belichenko, P. V., Faizi, M., Jacobs, L. F., Htun, K., Shamloo, M.,
876 & Mobley, W. C. (2012). Deficits in cognition and synaptic plasticity in a mouse
877 model of down syndrome ameliorated by gabab receptor antagonists. *The Journal of*
878 *Neuroscience*, *32*(27), 9217–9227.
- 879 Kleschevnikov, A. M., Belichenko, P. V., Villar, A. J., Epstein, C. J., Malenka, R. C., &
880 Mobley, W. C. (2004). Hippocampal long-term potentiation suppressed by increased
881 inhibition in the ts65dn mouse, a genetic model of down syndrome. *The Journal of*
882 *neuroscience*, *24*(37), 8153–8160.
- 883 Lanfranchi, S. [S], Carretti, B., Spano, G., & Cornoldi, C. (2009). A specific deficit in
884 visuospatial simultaneous working memory in down syndrome. *Journal of Intellectual*
885 *Disability Research*, *53*(5), 474–483.
- 886 Lanfranchi, S. [Silvia], Cornoldi, C., Vianello, R., & Conners, F. (2004). Verbal and
887 visuospatial working memory deficits in children with down syndrome. *American*
888 *Journal on Mental Retardation*, *109*(6), 456–466.
- 889 Lee, J. Y., Huerta, P. T., Zhang, J., Kowal, C., Bertini, E., Volpe, B. T., & Diamond, B.
890 (2009). Neurotoxic autoantibodies mediate congenital cortical impairment of
891 offspring in maternal lupus. *Nature medicine*, *15*(1), 91–96.
- 892 Lopez, L. L., Hauser, J., Feldon, J., Gargiulo, P., & Yee, B. (2010). Evaluating spatial
893 memory function in mice: a within-subjects comparison between the water maze test
894 and its adaptation to dry land. *Behavioural brain research*, *209*(1), 85–92.
- 895 Moore, S. J., Deshpande, K., Stinnett, G. S., Seasholtz, A. F., & Murphy, G. G. (2013).
896 Conversion of short-term to long-term memory in the novel object recognition
897 paradigm. *Neurobiology of learning and memory*, *105*, 174–185.
- 898 Pennington, B. F., Moon, J., Edgin, J., Stedron, J., & Nadel, L. (2003). The
899 neuropsychology of down syndrome: evidence for hippocampal dysfunction. *Child*
900 *development*, *74*(1), 75–93.

- 901 Ragozzino, M. E., Detrick, S., & Kesner, R. P. (1999). Involvement of the
902 prelimbic–infralimbic areas of the rodent prefrontal cortex in behavioral flexibility for
903 place and response learning. *The Journal of Neuroscience*, *19*(11), 4585–4594.
- 904 Ragozzino, M. E., Ragozzino, K. E., Mizumori, S. J., & Kesner, R. P. (2002). Role of the
905 dorsomedial striatum in behavioral flexibility for response and visual cue
906 discrimination learning. *Behavioral neuroscience*, *116*(1), 105.
- 907 Rast, M. M. & Harris, S. R. (1985). Motor control in infants with down syndrome.
908 *Developmental Medicine & Child Neurology*, *27*(5), 682–685.
- 909 Reeves, R. H., Irving, N. G., Moran, T. H., Wohn, A., Kitt, C., Sisodia, S. S., ...
910 Davisson, M. T. (1995). A mouse model for down syndrome exhibits learning and
911 behaviour. *Nat. Genet*, *11*, 177–184.
- 912 Rolls, E. T. & Kesner, R. P. (2006). A computational theory of hippocampal function, and
913 empirical tests of the theory. *Progress in neurobiology*, *79*(1), 1–48.
- 914 Sago, H., Carlson, E. J., Smith, D. J., Kilbridge, J., Rubin, E. M., Mobley, W. C., ...
915 Huang, T.-T. (1998). Ts1cje, a partial trisomy 16 mouse model for down syndrome,
916 exhibits learning and behavioral abnormalities. *Proceedings of the National Academy
917 of Sciences*, *95*(11), 6256–6261.
- 918 Smith, G. K., Kesner, R. P., & Korenberg, J. R. (2014). Dentate gyrus mediates cognitive
919 function in the ts65dn/dnj mouse model of down syndrome. *Hippocampus*, *24*(3),
920 354–362.
- 921 Stedron, J. M., Sahni, S. D., & Munakata, Y. (2005). Common mechanisms for working
922 memory and attention: the case of perseveration with visible solutions. *Journal of
923 cognitive neuroscience*, *17*(4), 623–631.
- 924 Team, R. C. (2014). R: a language and environment for statistical computing. vienna,
925 austria: r foundation for statistical computing; 2013.
- 926 Tennant, K. A., Asay, A. L., Allred, R. P., Ozburn, A. R., Kleim, J. A., & Jones, T. A.
927 (2010). The vermicelli and capellini handling tests: simple quantitative measures of

- 928 dexterous forepaw function in rats and mice. *JoVE (Journal of Visualized*
929 *Experiments)*, (41), e2076–e2076.
- 930 Vale-Martínez, A., Baxter, M. G., & Eichenbaum, H. (2002). Selective lesions of basal
931 forebrain cholinergic neurons produce anterograde and retrograde deficits in a social
932 transmission of food preference task in rats. *European Journal of Neuroscience*,
933 *16*(6), 983–998.
- 934 Vicari, S. (2006). Motor development and neuropsychological patterns in persons with
935 down syndrome. *Behavior genetics*, *36*(3), 355–364.
- 936 Vicari, S., Bellucci, S., & Carlesimo, G. A. (2005). Visual and spatial long-term memory:
937 differential pattern of impairments in williams and down syndromes. *Developmental*
938 *Medicine & Child Neurology*, *47*(05), 305–311.
- 939 Virji-Babul, N., Kerns, K., Zhou, E., Kapur, A., & Shiffar, M. (2006). Perceptual-motor
940 deficits in children with down syndrome: implications for intervention. *Down*
941 *Syndrome Research and Practice*, *10*(2), 74–82.
- 942 Visu-Petra, L., Benga, O., Miclea, M., et al. (2007). Visual-spatial processing in children
943 and adolescents with down's syndrome: a computerized assessment of memory skills.
944 *Journal of Intellectual Disability Research*, *51*(12), 942–952.
- 945 Warburton, E. C., Baird, A., Morgan, A., Muir, J. L., & Aggleton, J. P. (2001). The
946 conjoint importance of the hippocampus and anterior thalamic nuclei for allocentric
947 spatial learning: evidence from a disconnection study in the rat. *The Journal of*
948 *Neuroscience*, *21*(18), 7323–7330.
- 949 Warburton, E., Baird, A. L., Morgan, A., Muir, J. L., & Aggleton, J. P. (2000).
950 Disconnecting hippocampal projections to the anterior thalamus produces deficits on
951 tests of spatial memory in rats. *European Journal of Neuroscience*, *12*(5), 1714–1726.
- 952 Zampieri, B. L., Fernandez, F., Pearson, J. N., Stasko, M. R., & Costa, A. C. (2014).
953 Ultrasonic vocalizations during male–female interaction in the mouse model of down
954 syndrome ts65dn. *Physiology & behavior*, *128*, 119–125.

Table 1

Comparison of Arizona Cognitive Task Battery (ACTB) and Mouse Variant Reported in this Manuscript (mCTB)

Domain/Test in ACTB	Abilities Assessed	Analogous Task in mCTB	Ts65Dn performance
<i>Benchmark, General Cognitive Ability</i>			
KBIT-II Verbal Subscale	Receptive and Productive Language	<i>not modeled</i>	n/a
KBIT-II Nonverbal Subscale	Problem Solving	<i>not modeled</i>	n/a
Scales of Independent Behavior-Revised (SIB-R)	Adaptive Function	Nesting, Neophobia	deficits for Adaptive Function
CANTAB Spatial Span	Immediate Memory for Spatiotemporal Information	Temporal Order for Visual Objects	no deficits for Temporal Order
<i>Prefrontal Cortex - Executive Function, Response Attribute</i>			
Modified Dots Task	Inhibitory Control and Working Memory	Spontaneous Alternation	deficits for Spontaneous Alternation
CANTAB IED	Set Shifting	Rule Response Learning, Rule Reversal Learning	deficits for Response Learning, deficits for Reversal Learning
<i>Medial Temporal Lobe - Spatial Attribute</i>			
CANTAB PALS	Spatial Associative Memory	Location Recognition	deficits for Location Recognition
Virtual Water Maze	Spatial Memory/Navigation	Dry Land Water Maze (Cheeseboard)	deficits for Acquisition and Retrieval of Spatial Navigation
<i>not evaluated</i>	Spatial Relationships	Coordinate, Categorical	deficits for Coordinate task, no deficits for Categorical task
<i>Medial Temporal Lobe - Temporal Attribute</i>			
<i>not evaluated</i>	Temporal Processing/Sequence Learning	Temporal Order for Visual Objects	no deficits for Temporal Order
<i>Medial Temporal Lobe - Sensory/Perceptual Attribute</i>			
<i>not evaluated</i>	Object Recognition	Feature Ambiguity, Object Recognition, Novel Object Detection	No deficits at 1 hour delay, deficits at 24 hour delay
<i>Cerebellum - Motor Function</i>			
Finger Sequencing Task	Motor Sequencing	Capellini Handling	deficits for Motor Sequencing
NEPSY Visuomotor Precision	Visuomotor Tracking/Hand-Eye Coordination	Parallel Rung Walk, Capellini Handling	deficits for Motor Coordination
CANTAB SRT	Motor Response Time/Attention	<i>not modeled</i>	n/a

The mCTB was designed to model as many of the functions as the ACTB was designed to tests in humans. Cognitive deficits summarized in the table phenocopy the effects seen in Down syndrome on the ACTB or subsequent follow-up studies (Edgin et al., 2010, 2012). Similarly, the performance of Ts65Dn mice on the mCTB recapitulates intact cognitive function seen in participants with Down syndrome when tested using the ACTB




Figure 1. Dry land water maze performance on a cheeseboard for *Ts65Dn* and *2N* wildtype control mice. *Ts65Dn* mice showed impaired spatial navigation abilities during the 4 days of acquisition, even when adjusted for initial performance. *Ts65Dn* mice also show spatial memory deficits during the probe trial relative to *2N* wildtype control mice, reflected in reduced time in the quadrant containing the reward location and greater average distance from the previously rewarded location compared to *2N* control mice.




Figure 2. Spatial and Temporal Attribute task battery. The data suggest Ts65Dn mice show deficits relative to 2N wildtype control mice for location recognition and metric/coordinate processing, but no deficits for topological/categorical processing. The Ts65Dn mice do not show deficits for temporal ordering for visual objects compared to 2N wildtype control mice.




Figure 3. Sensory/Perceptual Attribute task battery. Overall, *Ts65Dn* mice do not show impaired sensory/perceptual function relative to *2N* wildtype mice. *Ts65Dn* mice also do not show deficits for object recognition at a 1 hour delay, but do show deficits for object recognition at 24 hour delays.




Figure 4. Executive Function / Rule Based Memory Task Battery. *Ts65Dn* mice show fewer alternations on a spontaneous alternation task relative to *2N* control mice. *Ts65Dn* mice show mild deficits for acquisition and reversal of a rule based response on a plus maze. During reversal training, *Ts65Dn* mice learn to apply the new rule on later trials than control mice, reflected by an increased number of perseverative, but not regressive, errors.




Figure 5. Motor Function Task Battery. Ts65Dn mice showed reduced motor dexterity during a Capellini Handling task reflected as an increase in the number of abnormal behaviors and increased latency to consume the capellini as well a greater number of foot slips during a Parallel Rung Walking task, even when adjusted for total number of steps.




Figure 6. Adaptive Function / Quality of Life Task Battery. *Ts65Dn* mice take longer to make a nest out of preferred nesting material and show increased neophobia for both food and environments.