

1 HydDB: A web tool for hydrogenase 2 classification and analysis

3 Dan Søndergaard^a, Christian N. S. Pedersen^a, Chris Greening^{c, d *}

4

5 ^a Aarhus University, Bioinformatics Research Centre, C.F. Møllers Allé 8, Aarhus
6 DK-8000, Denmark

7 ^b Australian National University, Research School of Chemistry, Sullivans Creek
8 Road, Acton, ACT 2601, Australia

9 ^c The Commonwealth Scientific and Industrial Research Organisation, Land and
10 Water Flagship, Clunies Ross Street, Acton, ACT 2060, Australia

11 ^d Monash University, School of Biological Sciences, Clayton, VIC 2800, Australia

12

13 **Correspondence:**

14 Dr Chris Greening (chris.greening@monash.edu), Monash University, School of
15 Biological Sciences, Clayton, VIC 2800, Australia

16 Dan Søndergaard (das@birc.au.dk), Aarhus University, Bioinformatics Research
17 Centre, C.F. Møllers Allé 8, Aarhus DK-8000, Denmark

18

19 **Journal:** Scientific Reports

20 **Short Title:** HydDB, a hydrogenase database

21 **Figures:** Two colour figures, two tables

22

23 **Abstract**

24 H₂ metabolism is the most ancient and diverse mechanism of energy-generation.
25 The metalloenzymes mediating this metabolism, hydrogenases, are encoded by over
26 60 microbial phyla and are present in all major ecosystems. We developed a
27 classification system and web tool, HydDB, for the structural and functional analysis
28 of these enzymes. We show that hydrogenase function can be predicted by primary
29 sequence alone using an expanded classification scheme (comprising 29 [NiFe]-
30 hydrogenase subgroups, 8 [FeFe]-hydrogenase subtypes, [Fe]-hydrogenases).
31 Using this scheme, we built a web tool that rapidly and reliably classifies
32 hydrogenase primary sequences using a combination of *k*-nearest neighbors'
33 algorithms and CDD referencing. Demonstrating its capacity, the tool reliably
34 predicted hydrogenase content and function in 12 newly-sequenced bacteria,
35 archaea, and eukaryotes. HydDB also provides the capacity to browse 3248
36 annotated sequences and contains a detailed repository of physiological,
37 biochemical, and structural information about the 38 hydrogenase classes defined
38 here. The database and classifier are freely and publicly available at
39 <http://services.birc.au.dk/hyddb/>

40

41 **Introduction**

42 Microorganisms conserve energy by metabolizing H₂. Oxidation of this high-energy
43 fuel yields electrons that can be used for respiration and carbon-fixation. This
44 diffusible gas is also produced in diverse fermentation and anaerobic respiratory
45 processes ¹. H₂ metabolism contributes to the growth and survival of microorganisms
46 across the three domains of life: chemotrophs and phototrophs, lithotrophs and

47 heterotrophs, aerobes and anaerobes, mesophiles and extremophiles alike ^{1,2}. On
48 the ecosystem scale, H₂ supports microbial communities in most terrestrial, aquatic,
49 and host-associated ecosystems ^{1,3}. It is also generally accepted that H₂ was the
50 primordial electron donor ⁴. In biological systems, metalloenzymes known as
51 hydrogenases are responsible for oxidizing and evolving H₂ ^{1,5}. Our recent survey
52 showed there is a far greater number and diversity of hydrogenases than previously
53 thought ². It is predicted over 55 microbial phyla and up to half of all microorganisms
54 harbor hydrogenases ^{2,6}. Better understanding H₂ metabolism and the enzymes that
55 mediate it also has wider implications, particularly in relation to human health and
56 disease ^{3,7}, biogeochemical cycling ⁸, and renewable energy ^{9,10}.

57

58 There are three classes of hydrogenase, the [NiFe], [FeFe], and [Fe] hydrogenases,
59 that are distinguished by their metal composition. Whereas the [Fe]-hydrogenases
60 are a small methanogenic-specific family ¹¹, the [NiFe] and [FeFe] classes are widely
61 distributed and functionally diverse. They comprise numerous different groups and
62 subgroups/subtypes with distinct biochemical features (e.g. directionality, affinity,
63 redox partners, and localization) and physiological roles (i.e. respiration,
64 fermentation, bifurcation, sensing) ^{1,5}. For example, while Group 2a and 2b [NiFe]-
65 hydrogenases share > 35% sequence identity, they have distinct roles as respiratory
66 uptake hydrogenases and H₂ sensors respectively ^{12,13}. Building on previous work
67 ^{14,15}, we recently created a comprehensive hydrogenase classification scheme
68 predictive of biological function ². This scheme was primarily based on amino acid
69 sequence phylogeny, but also factored in genetic organization, metal-binding motifs,
70 and functional information. This analysis identified 22 subgroups (within four groups)

71 of [NiFe]-hydrogenases and six subtypes (within three groups) of [FeFe]-
72 hydrogenases, each with unique physiological roles ².

73


74 In this work, we build on these findings to develop the first web database for the
75 classification and analysis of hydrogenases. We developed an expanded
76 classification scheme that captures the full sequence diversity of hydrogenase
77 enzymes and predicts their biological function. Using this information, we developed
78 a classification tool based on the *k*-nearest neighbors' (*k*-NN) method. This tool is a
79 more reliable, efficient, and user-friendly method for hydrogenase classification than
80 standard approaches involved phylogenetic tree construction, with a precision of
81 more than 99.8%.

82

83 **Results and Discussion**

84 **A sequence-based classification scheme for hydrogenases**

85 We initially developed a classification scheme to enable prediction of hydrogenase
86 function by primary sequence alone. To do this, we visualized the relationships
87 between all hydrogenases in sequence similarity networks ¹⁶, in which nodes
88 represent individual proteins and the distances between them reflect BLAST *E*-
89 values. As reflected by our analysis of other protein superfamilies ^{17,18}, SSNs allow
90 robust inference of sequence-structure-function relationships for large datasets
91 without the problems associated with phylogenetic trees (e.g. long-branch attraction).
92 Consistent with previous phylogenetic analyses ^{2,14,15}, this analysis showed the
93 hydrogenase sequences clustered into eight major groups (Groups 1 to 4 [NiFe]-
94 hydrogenases, Groups A to C [FeFe]-hydrogenases, [Fe]-hydrogenases), six of
95 which separate into multiple functionally-distinct subgroups or subtypes at narrower



96

97 **Figure 1.** Sequence similarity network of hydrogenase sequences. Nodes represent
 98 individual proteins and the edges show the BLAST E-values between them at the
 99 logE filter defined at the bottom-left of each panel. The sequences are colored by
 100 class as defined in the legends. Figure S1 shows the further delineation of the
 101 encircled [NiFe] hydrogenase classes.

102 *logE* filters (**Figure 1; Figure S1**). The SSNs demonstrated that all [NiFe]-
103 hydrogenase subgroups defined through phylogenetic trees in our previous work ²
104 separated into distinct clusters, which is consistent with our evolutionary model that
105 such hydrogenases diverged from a common ancestor to adopt multiple distinct
106 functions ². The only exception were the Group A [FeFe]-hydrogenases, which as
107 previously-reported ^{2,15}, cannot be classified by sequence alone as they have
108 principally diversified through changes in domain architecture and quaternary
109 structure. It remains strictly necessary to analyze the organization of the genes
110 encoding these enzymes to determine their specific function, e.g. whether they serve
111 fermentative or electron-bifurcating roles.

112

113 The SSN analysis revealed that several groups and subgroups that clustered
114 together in the phylogenetic tree analysis ² separate into several subclades of
115 probable distinct function (**Figure 1**). On this basis, we refined and expanded the
116 hydrogenase classification scheme to reflect the sequence diversification observed
117 (**Table 1**). Three lineages originally classified as Group 1a [NiFe]-hydrogenases
118 were reclassified as new subgroups, the Coriobacteria (Group 1i), Archaeoglobi
119 (Group 1j), and Methanosarcinales (Group 1i). The previously-defined 4b and 4d
120 subgroups ² were dissolved, as the SSN analysis confirmed they were highly
121 polyphyletic. These sequences are reclassified here into five new subgroups: the
122 formate- and carbon monoxide-respiring Mrp-linked complexes (Group 4b) ¹⁹, the
123 ferredoxin-coupled Mrp-linked complexes (Group 4d) ²⁰, the well-described
124 methanogenic Eha (Group 4h) and Ehb (Group 4i) supercomplexes ²¹, and a more
125 loosely clustered class of unknown function (Group 4g). Three crenarchaeotal
126 hydrogenases were also classified as their own family (Group 2e); these enzymes

[NiFe] Group 1: Respiratory H₂-uptake [NiFe]-hydrogenases

1a	Periplasmic	Electron input for sulfate, metal, and organohalide respiration. [NiFeSe] variants.	2
1b	Prototypical	Electron input for sulfate, fumarate, metal, and nitrate respiration.	2
1c	Hyb-type	Electron input for fumarate, nitrate, and sulfate respiration. Physiologically reversible.	2
1d	Oxygen-tolerant	Electron input for aerobic respiration and oxygen-tolerant anaerobic respiration.	2
1e	Isp-type	Electron input primarily for sulfur respiration. Physiologically reversible.	2
1f	Oxygen-protecting	Unresolved role. May liberate electrons to reduce reactive oxygen species.	2
1g	Crenarchaeota-type	Electron input primarily for sulfur respiration.	2
1h	Actinobacteria-type	Electron input for aerobic respiration. Scavenges electrons from atmospheric H ₂ .	2,40
1i	Coriobacteria-type (putative)	Undetermined role. May liberate electrons for anaerobic respiration.	This work
1j	Archaeoglobi-type	Electron input for sulfate respiration ⁴¹ .	This work
1k	Methanophenazine-reducing	Electron input for methanogenic heterodisulfide respiration ⁴² .	This work

[NiFe] Group 2: Alternative and sensory uptake [NiFe]-hydrogenases

2a	Cyanobacteria-type	Electron input for aerobic respiration. Recycles H ₂ produced by other cellular processes.	14
2b	Histidine kinase-linked	H ₂ sensing. Activates two-component system controlling hydrogenase expression.	14
2c	Diguanylate cyclase-linked (putative)	Undetermined role. May sense H ₂ and regulate processes through cyclic di-GMP production.	2
2d	Aquificae-type	Unresolved role. May generate reductant for carbon fixation or have a regulatory role.	2
2e	Metallosphaera-type (putative)	Undetermined role. May liberate electrons primarily for aerobic respiration ²² .	This work

[NiFe] Group 3: Cofactor-coupled bidirectional [NiFe]-hydrogenases

3a	F ₄₂₀ -coupled	Couples oxidation of H ₂ to reduction of F ₄₂₀ during methanogenesis. Physiologically reversible. [NiFeSe] variants.	14
3b	NADP-coupled	Couples oxidation of NADPH to evolution of H ₂ . Physiologically reversible. May have sulfhydrogenase activity.	14
3c	Heterodisulfide reductase-linked	Bifurcates electrons from H ₂ to heterodisulfide and Fd _{ox} in methanogens. [NiFeSe] variants.	14
3d	NAD-coupled	Interconverts electrons between H ₂ and NAD depending on cellular redox state.	14

[NiFe] Group 4: Respiratory H₂-evolving [NiFe]-hydrogenases

4a	Formate hydrogenlyase	Couples formate oxidation to fermentative H ₂ evolution. May be H ⁺ -translocating.	2
4b	Formate-respiring	Respires formate or carbon monoxide using H ⁺ as electron acceptor. Na ⁺ -translocating via Mrp ¹⁹ .	This work
4c	Carbon monoxide-respiring	Respires carbon monoxide using H ⁺ as electron acceptor. H ⁺ -translocating.	2

4d	Ferredoxin-coupled, Mrp-linked	Couples Fd _{red} oxidation to H ⁺ reduction. Na ⁺ -translocating via Mrp complex ²⁰ .	This work
4e	Ferredoxin-coupled, Ech-type	Couples Fd _{red} oxidation to H ⁺ reduction. Physiologically reversible via H ⁺ /Na ⁺ translocation.	2
4f	Formate-coupled (putative)	Undetermined role. May couple formate oxidation to H ₂ evolution and H ⁺ translocation.	2
4g	Ferredoxin-coupled (putative)	Undetermined role. May couple Fd _{red} oxidation to proton reduction and H ⁺ /Na ⁺ translocation.	This work
4h	Ferredoxin-coupled, Eha-type	Couples Fd _{red} oxidation to H ⁺ reduction in anaplerotic processes. H ⁺ /Na ⁺ -translocating ²¹ .	This work
4i	Ferredoxin-coupled, Ehb-type	Couples Fd _{red} oxidation to H ⁺ reduction in anabolic processes. H ⁺ /Na ⁺ -translocating ²¹ .	This work
[FeFe] Hydrogenases			
A1	Prototypical	Couples ferredoxin oxidation to fermentative or photobiological H ₂ evolution.	2,15
A2	Glutamate synthase-linked (putative)	Undetermined role. May couple H ₂ oxidation to NAD reduction, generating reductant for glutamate synthase.	2,15
A3	Bifurcating	Reversibly bifurcates electrons from H ₂ to NAD and Fd _{ox} in anaerobic bacteria.	2,15
A4	Formate dehydrogenase-linked	Couples formate oxidation to H ₂ evolution. Some bifurcate electrons from H ₂ to ferredoxin and NADP.	2,15
B	Colonic-type (putative)	Undetermined role. May couple Fd _{red} oxidation to fermentative H ₂ evolution.	15
C1	Histidine kinase-linked (putative)	Undetermined role. May sense H ₂ and regulate processes via histidine kinases ² .	This work
C2	Chemotactic (putative)	Undetermined role. May sense H ₂ and regulate processes via methyl-accepting chemotaxis proteins ² .	This work
C3	Phosphatase-linked (putative)	Undetermined role. May sense H ₂ and regulate processes via serine/threonine phosphatases ² .	This work
[Fe] Hydrogenases			
All	Methenyl-H ₄ MPT dehydrogenase	Reversibly couples H ₂ oxidation to 5,10-methenyltetrahydromethanopterin reduction.	14

127

128 **Table 1.** Expanded classification scheme for hydrogenase enzymes. The majority of the classes were defined in previous work
129 ^{2,14,15,40}. The [NiFe] Group 1i, 1j, 1j, 2e, 4d, 4g, 4h, and 4i enzymes and [FeFe] Groups C1, C2, and C3 enzymes were defined in
130 this work based on their separation into distinct clusters in the SSN analysis (**Figure 1**). HydDB contains detailed information on
131 each of these classes, including their taxonomic distribution, genetic organization, biochemistry, and structures, as well a list of
132 primary references.

133 enable certain crenarchaeotes to grow aerobically on O₂^{22,23} and hence may
134 represent a unique lineage of aerobic uptake hydrogenases currently
135 underrepresented in genome databases. The Group C [FeFe]-hydrogenases were
136 also separated into three main subtypes given they separate into distinct clusters
137 even at relatively broad logE values (**Figure 1**); these enzymes likely have a sensory
138 role^{2,15} and are each co-transcribed with different regulatory elements (**Table 1**).

139


140 **HydDB reliably predicts hydrogenase class using the *k*-NN method and CDD** 141 **referencing**

142 Using this information, we built a web tool to classify hydrogenases. Hydrogenase
143 classification is determined through a two-step process following input of the catalytic
144 subunit sequence. In the first, the Conserved Domain Database (CDD)²⁴ is
145 referenced to confirm that the inputted sequence has a hydrogenase catalytic
146 domain, i.e. “Complex1_49kDa superfamily” (cl21493) (for NiFe-hydrogenases),
147 “Fe_hyd_lg_C superfamily” (cl14953) (for FeFe-hydrogenases), and “HMD”
148 (pfam03201) (for Fe-hydrogenases). The sequence is subsequently classified
149 through the *k*-NN method that determines the most similar sequences listed in the
150 HydDB reference database. To determine the optimal *k* for the dataset, we
151 performed a 5-fold cross-validation for *k* = 1...10 and computed the accuracy for
152 each *k*. The results are shown in **Figure 2**. The classifier predicted the classes of the
153 3248 hydrogenase sequences with 99.8% accuracy and high robustness when
154 performing a 5-fold cross-validation (as described in the Methods section) for *k* = 4.
155 The six sequences where there were discrepancies between the SSN and *k*-NN
156 predictions are shown in **Table S1**. The classifier has also been trained to detect and

157 exclude protein families that are homologous to hydrogenases but do not metabolize
158 H₂ (Nuo, Ehr, NARF, HmdII^{1,2}) using reference sequences of these proteins.

159

160 **Figure 2.** Evaluation of the k -NN classifier for $k = 1 \dots 10$. For each k , a 5-fold cross-
161 validation was performed. The mean accuracy \pm two standard deviations of the folds
162 is shown in the figure (note the y-axis). $k = 1$ provides the most accurate classifier.
163 However, $k = 4$ provides almost the same accuracy and is more robust to errors in
164 the training set (reflected by the lower standard deviation). In general, the standard
165 deviation is very small, indicating that the predictions are robust to changes in the
166 training data.



167

168 Sequences of the [FeFe] Group A can be classified into functionally-distinct subtypes
169 (A1, A2, A3, A4) based on genetic organization². The classifier can classify such
170 hydrogenases if the protein sequence immediately downstream from the catalytic
171 subunit sequence is provided. The classifier references the CDD to search for
172 conserved domains in the downstream protein sequence. A sequence is classified

173 as [FeFe] Group A2 if one of the domains “GltA”, “GltD”, “glutamate synthase small
174 subunit” or “putative oxidoreductase”, but not “NuoF”, is found in the sequence.
175 Sequences are classified as [FeFe] Group A3 if the domain “NuoF” is found and
176 [FeFe] Group A4 if the domain “HycB” is present. If none of the domains are found,
177 the sequence is classified as A1. These classification rules were determined by
178 collecting 69 downstream protein sequences. The sequences were then submitted to
179 the CDD and the domains which most often occurred in each subtype were
180 extracted.

181

182 In addition to its accuracy, the classifier is superior to other approaches due to its
183 usability (**Figure S2**). It is accessible as a free web service at
184 <http://services.birc.au.dk/hyddb/> HydDB allows the users to paste or upload
185 sequences of hydrogenase catalytic subunit sequences in FASTA format and run the
186 classification. When analysis has completed, results are presented in a table that
187 can be downloaded as a CSV file. This provides an efficient and user-friendly way to
188 classify hydrogenases, in contrast to the previous standard which requires
189 visualization of multiple sequence alignments in phylogenetic trees ²⁵.

190

191 **HydDB infers the physiological roles of H₂ metabolism**

192 As summarized in **Table 1**, hydrogenase class is strongly correlated with
193 physiological role. As a result, the classifier is capable of predicting both the class
194 and function of a sequenced hydrogenase. To demonstrate this capacity, we used
195 HydDB to analyze the hydrogenases present in 12 newly-sequenced bacteria,
196 archaea, and eukaryotes of major ecological significance. The classifier correctly
197 classified all 24 hydrogenases identified in the sequenced genomes, as validated

198 with SSNs (**Table 2**). On the basis of these classifications, the physiological roles of
199 H₂ metabolism were predicted (**Table 2**). For five of the organisms, these predictions
200 are confirmed or supported by previously published data ^{23,26–29}. Other predictions
201 are in line with metabolic models derived from metagenome surveying ^{30–32}. In some
202 cases, the capacity for organisms to metabolize H₂ was not tested or inferred in
203 previous studies despite the presence of hydrogenases in the sequenced genomes
204 ^{27,33–35}.

205

206 While HydDB serves as a reliable initial predictor of hydrogenase class and function,
207 further analysis is recommended to verify predictions. Hydrogenase sequences only
208 provide organism with the genetic capacity to metabolise H₂; their function is
209 ultimately modulated by their expression and integration within the cell ^{1,36}. In
210 addition, some classifications are likely to be overgeneralized due to lack of
211 functional and biochemical characterization of certain lineages and sublineages. For
212 example, it is not clear if two distant members of the Group 1h [NiFe]-hydrogenases
213 (*Robiginitalea biformata*, *Sulfolobus islandicus*) perform the same H₂-scavenging
214 functions as the core group ⁸. Likewise, it seems probable that the Group 3a [NiFe]-
215 hydrogenases of *Thermococci* and *Aquificae* use a distinct electron donor to the
216 main class ³⁷. Prominent cautions are included in the enzyme pages in cases such
217 as these. HydDB will be updated when literature is published that influences
218 functional assignments.

219

220 **HydDB contains interfaces for hydrogenase browsing and analyzing**

221 In addition to its classification function, HydDB is designed to be a definitive
222 repository for hydrogenase retrieval and analysis. The database presently contains

Organism	Phylum	Hydrogenase accession no.	HydDB classification	SSN classification	Predicted H ₂ metabolism	Confirmed H ₂ metabolism
<i>Pyrinomonas methylaliphatogenes</i>	Acidobacteria	WP_041979300.1	[NiFe] Group 1h	[NiFe] Group 1h	Persistence by aerobic respiration of atmospheric H ₂	Confirmed experimentally ²⁶
<i>Phaeodactylibacter xiamenensis</i>	Bacteroidetes	WP_044227713.1 WP_044216927.1 WP_044227053.1	[NiFe] Group 1d [NiFe] Group 2a [NiFe] Group 3d	[NiFe] Group 1d [NiFe] Group 2a [NiFe] Group 3d	Chemolithoautotrophic growth by aerobic H ₂ oxidation	Bacterium grows aerobically, but H ₂ oxidation untested ²⁷
Bathyarchaeota archaeon BA1	Bathyarchaeota	KPV62434.1 KPV62673.1 KPV62298.1	[NiFe] Group 3c [NiFe] Group 3c [NiFe] Group 4g	[NiFe] Group 3c [NiFe] Group 3c [NiFe] Group 4g	Couples Fd _{red} oxidation to H ₂ evolution in energy-conserving and bifurcating processes	Unconfirmed but consistent with metagenome-based models ³¹
<i>Lenisia limosa</i>	Proteobacteria (Epsilon class)	LenisMan28	[FeFe] Group A1	[FeFe] Group A	Fermentative evolution of H ₂	Confirmed experimentally ⁴³
<i>Acidianus copahuensis</i>	Crenarchaeota	WP_048100721.1 WP_048100713.1 WP_048100378.1 WP_048100359.1	[NiFe] Group 1g [NiFe] Group 1g [NiFe] Group 1h [NiFe] Group 2e	[NiFe] Group 1g [NiFe] Group 1g [NiFe] Group 1h [NiFe] Group 2e	Chemolithoautotrophic growth by H ₂ oxidation using O ₂ or S ₀ as electron acceptors	Partially confirmed experimentally ²³
<i>Arcobacter</i> sp. E1/2/3	Proteobacteria (Epsilon class)	Arc.peg.2312	[NiFe] Group 1b	[NiFe] Group 1b	Chemolithoautotrophic growth by anaerobic H ₂ oxidation	Confirmed experimentally ⁴³
<i>Methanoperedens nitroreducens</i>	Euryarchaeota (ANME)	WP_048088262.1 WP_048090768.1	[NiFe] Group 3b [NiFe] Group 3b	[NiFe] Group 3b [NiFe] Group 3b	Secondary role for H ₂ metabolism limited to fermentative evolution of H ₂	Unconfirmed but consistent with metagenome-based models ³⁰
<i>Kryptonium thompsoni</i>	Kryptonia	CUU03002.1 CUU06124.1	[NiFe] Group 1d [NiFe] Group 3b	[NiFe] Group 1d [NiFe] Group 3b	Chemolithoautotrophic growth by aerobic H ₂ oxidation, fermentative evolution of H ₂ .	Untested, candidate phylum identified by metagenomics ³⁴
<i>Lokiarchaeum</i> sp. GC14_75	Lokiarchaeota	KKK40681.1	[NiFe] Group 3c	[NiFe] Group 3c	Bifurcates electrons between H ₂ , heterodisulfide, and ferredoxin	Unconfirmed but consistent with metagenome-based models ⁴⁴

Nitrospira moscoviensis	Nitrospirae	WP_053379275.1	[NiFe] Group 2a	[NiFe] Group 2a	Chemolithoautotrophic growth by aerobic H ₂ oxidation	Confirmed experimentally ²⁸
Bacterium GW2011_GWE1_35_17	Moranbacteria	KKQ46070.1 KKQ45273.1	[NiFe] Group 1a [NiFe] Group 3b	[NiFe] Group 1a [NiFe] Group 3b	Chemolithoautotrophic growth by anaerobic H ₂ oxidation, fermentative evolution of H ₂ .	Unconfirmed but consistent with metagenome-based models ³²
Bacterium GW2011_GWA2_33_10	Peregrinibacteria	KKP36897.1	[FeFe] Group A3	[FeFe] Group A	Bifurcates electrons between H ₂ , NADH, and ferredoxin	Unconfirmed but consistent with metagenome-based models ³²
Entotheonella sp. TSY1	Tectomicrobia	ETW97737.1 ETW94065.1	[NiFe] Group 1h [NiFe] Group 3b	[NiFe] Group 1h [NiFe] Group 3b	Persistence by aerobic respiration of atmospheric H ₂ , fermentative evolution of H ₂	Untested, candidate phylum identified by metagenomics ³⁵

223

224 **Table 2.** Predictive capacity of the HydDB. HydDB accurately determined hydrogenase content and predicted the physiological roles of H₂
225 metabolism in 12 newly-sequenced archaeal and bacterial species.

entries for 3248 hydrogenases, including their NCBI accession numbers, amino acid sequence, hydrogenase class, taxonomic affiliation, and predicted behavior (**Figure S2**). To enable easy exploration of the data set, the database also provides access to an interface for searching, filtering, and sorting the data, as well as the capacity to download the results in CSV or FASTA format. There are individual pages for the 38 hydrogenase classes defined here (**Table 1**), including descriptions of their physiological role, genetic organization, taxonomic distribution, and biochemical features. This is supplemented with a compendium of structural information about the hydrogenases, which is integrated with the Protein Databank (PDB), as well as a library of over 1000 literature references (**Figure S5**).

Conclusions

To summarize, HydDB is a definitive resource for hydrogenase classification and analysis. The classifier described here provides a reliable, efficient, and convenient tool for hydrogenase classification and functional prediction. HydDB also provides browsing tools for the rapid analysis and retrieval of hydrogenase sequences. Finally, the manually-curated repository of class descriptions, hydrogenase structures, and literature references provide a deep but accessible resource for understanding hydrogenases.

Materials and Methods

Sequence datasets

The database was constructed using the amino acid sequences of all curated non-redundant 3248 hydrogenase catalytic subunits represented in the NCBI RefSeq

database in August 2014 ² (**Dataset S1**). In order to test the classification tool, additional sequences from newly-sequenced archaeal and bacteria phyla were retrieved from the Joint Genome Institute's Integrated Microbial Genomes database ³⁸.

Sequence similarity networks

Sequence similarity networks (SSNs) ¹⁶ were used to visualize the distribution and diversity of the 3248 retrieved hydrogenase sequences. In this analysis, nodes represent individual proteins and edges represent the all-versus-all BLAST E -values. Three networks were constructed using Cytoscape, namely for the [NiFe]-hydrogenase large subunit sequences, [FeFe]-hydrogenase catalytic domain sequences, and [Fe]-hydrogenase sequences. The relationships between them were viewed at different $\log E$ cutoffs using different subsets of sequences.

Classification method

The k -NN method is a well-known machine learning method for classification ³⁹. Given a set of data points x_1, x_2, \dots, x_N (e.g. sequences) with known labels y_1, y_2, \dots, y_N (e.g. type annotations), the label of a point, x , is predicted by computing the distance from x to x_1, x_2, \dots, x_N and extracting the k labeled points closest to x , i.e. the neighbors. The predicted label is then determined by majority vote of the labels of the neighbors. The distance measure applied here is that of a BLAST search. Thus, the classifier corresponds to a homology search where the types of the top k results are considered. However, formulating the classification method as a machine learning problem allows the use of common evaluation methods to estimate the accuracy of the method and perform model selection. The classifier was evaluated

using k -fold cross-validation. The dataset is first split in to k parts of equal size. $k - 1$ parts (the *training set*) are then used for training the classifier and the labels of the data points in the remaining part (the *test set*) are then predicted. This process, called a *fold*, is repeated k times. The predicted labels of each fold are then compared to the known labels and an accuracy can be computed.

Acknowledgements

We thank A/Prof Colin J. Jackson, Dr Hafna Ahmed, Dr Andrew Warden, and Dr Stephen Pearce for their helpful advice and comments regarding this manuscript. This work was supported by a PUMPKin Centre of Excellence PhD Scholarship awarded to DS, an Australian National University PhD Scholarship awarded to FHA, and a CSIRO Office of the Chief Executive Postdoctoral Fellowship awarded to CG.

Author Contributions

CG and DS designed experiments. DS and CG performed experiments. CG, DS, and CNSP analysed data. CNSP supervised students. CG and DS wrote the paper.

The authors declare no conflict of interest.

References

1. Schwartz, E., Fritsch, J. & Friedrich, B. *H₂-metabolizing prokaryotes*. (Springer Berlin Heidelberg, 2013).
2. Greening, C. *et al.* Genome and metagenome surveys of hydrogenase diversity indicate H₂ is a widely-utilised energy source for microbial growth and survival. *ISME J.* **10**, 761–777 (2016).

3. Cook, G. M., Greening, C., Hards, K. & Berney, M. in *Advances in Bacterial Pathogen Biology* (ed. Poole, R. K.) **65**, 1–62 (Academic Press, 2014).
4. Lane, N., Allen, J. F. & Martin, W. How did LUCA make a living? Chemiosmosis in the origin of life. *BioEssays* **32**, 271–280 (2010).
5. Lubitz, W., Ogata, H., Rüdiger, O. & Reijerse, E. Hydrogenases. *Chem. Rev.* **114**, 4081–148 (2014).
6. Peters, J. W. *et al.* [FeFe]- and [NiFe]-hydrogenase diversity, mechanism, and maturation. *Biochim. Biophys. Acta - Mol. Cell Res.* (2014). doi:10.1016/j.bbamcr.2014.11.021
7. Carbonero, F., Benefiel, A. C. & Gaskins, H. R. Contributions of the microbial hydrogen economy to colonic homeostasis. *Nat Rev Gastroenterol Hepatol* **9**, 504–518 (2012).
8. Greening, C. *et al.* Atmospheric hydrogen scavenging: from enzymes to ecosystems. *Appl. Environ. Microbiol.* **81**, 1190–1199 (2015).
9. Levin, D. B., Pitt, L. & Love, M. Biohydrogen production: prospects and limitations to practical application. *Int. J. Hydrogen Energy* **29**, 173–185 (2004).
10. Cracknell, J. A., Vincent, K. A. & Armstrong, F. A. Enzymes as working or inspirational catalysts for fuel cells and electrolysis. *Chem. Rev.* **108**, 2439–2461 (2008).
11. Shima, S. *et al.* The crystal structure of [Fe]-Hydrogenase reveals the geometry of the active site. *Science* **321**, 572–575 (2008).
12. Lenz, O. & Friedrich, B. A novel multicomponent regulatory system mediates H₂ sensing in *Alcaligenes eutrophus*. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 12474–12479 (1998).
13. Greening, C., Berney, M., Hards, K., Cook, G. M. & Conrad, R. A soil actinobacterium scavenges atmospheric H₂ using two membrane-associated, oxygen-dependent [NiFe] hydrogenases. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 4257–4261 (2014).
14. Vignais, P. M., Billoud, B. & Meyer, J. Classification and phylogeny of hydrogenases. *FEMS Microbiol. Rev.* **25**, 455–501 (2001).
15. Calusinska, M., Happe, T., Joris, B. & Wilmotte, A. The surprising diversity of clostridial hydrogenases: a comparative genomic perspective. *Microbiology* **156**, 1575–1588 (2010).

16. Atkinson, H. J., Morris, J. H., Ferrin, T. E. & Babbitt, P. C. Using sequence similarity networks for visualization of relationships across diverse protein superfamilies. *PLoS One* **4**, e4345 (2009).
17. Ahmed, F. H. *et al.* Sequence-structure-function classification of a catalytically diverse oxidoreductase superfamily in mycobacteria. *J. Mol. Biol.* **427**, 3554–3571 (2015).
18. Ney, B. *et al.* The methanogenic redox cofactor F₄₂₀ is widely synthesized by aerobic soil bacteria. *ISME J.* In press (2016).
19. Kim, Y. J. *et al.* Formate-driven growth coupled with H₂ production. *Nature* **467**, 352–5 (2010).
20. McTernan, P. M. *et al.* Intact functional fourteen-subunit respiratory membrane-bound [NiFe]-hydrogenase complex of the hyperthermophilic archaeon *Pyrococcus furiosus*. *J. Biol. Chem.* **289**, 19364–19372 (2014).
21. Lie, T. J. *et al.* Essential anaplerotic role for the energy-converting hydrogenase Eha in hydrogenotrophic methanogenesis. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 15473–8 (2012).
22. Auernik, K. S. & Kelly, R. M. Physiological versatility of the extremely thermoacidophilic archaeon *Metallosphaera sedula* supported by transcriptomic analysis of heterotrophic, autotrophic, and mixotrophic growth. *Appl. Environ. Microbiol.* **76**, 931–935 (2010).
23. Giaveno, M. A., Urbietta, M. S., Ulloa, J. R., González Toril, E. & Donati, E. R. Physiologic versatility and growth flexibility as the Main characteristics of a novel thermoacidophilic *Acidianus* strain isolated from Copahue geothermal area in Argentina. *Microb. Ecol.* **65**, 336–346 (2012).
24. Marchler-Bauer, A. & Bryant, S. H. CD-Search: protein domain annotations on the fly. *Nucleic Acids Res.* **32**, W327–31 (2004).
25. Berney, M., Greening, C., Hards, K., Collins, D. & Cook, G. M. Three different [NiFe] hydrogenases confer metabolic flexibility in the obligate aerobe *Mycobacterium smegmatis*. *Environ. Microbiol.* **16**, 318–330 (2014).
26. Greening, C. *et al.* Persistence of the dominant soil phylum *Acidobacteria* by trace gas scavenging. *Proc. Natl. Acad. Sci.* **112**, 10497–10502 (2015).

27. Chen, Z. *et al.* *Phaeodactylibacter xiamenensis* gen. nov., sp. nov., a member of the family *Saprospiraceae* isolated from the marine alga *Phaeodactylum tricornutum*. *Int. J. Syst. Evol. Microbiol.* **64**, 3496–3502 (2014).
28. Koch, H. *et al.* Growth of nitrite-oxidizing bacteria by aerobic hydrogen oxidation. *Science* **345**, 1052–1054 (2014).
29. Carere, C. R. *et al.* Growth and persistence of methanotrophic bacteria by aerobic hydrogen respiration. *Proc. Natl. Acad. Sci. U. S. A.* (2016).
30. Haroon, M. F. *et al.* Anaerobic oxidation of methane coupled to nitrate reduction in a novel archaeal lineage. *Nature* **500**, 567–70 (2013).
31. Evans, P. N. *et al.* Methane metabolism in the archaeal phylum *Bathyarchaeota* revealed by genome-centric metagenomics. *Science* **350**, 434–438 (2015).
32. Brown, C. T. *et al.* Unusual biology across a group comprising more than 15% of domain *Bacteria*. *Nature* **523**, 208–211 (2015).
33. Spang, A. *et al.* Complex archaea that bridge the gap between prokaryotes and eukaryotes. *Nature* **521**, 173–179 (2015).
34. Eloe-Fadrosh, E. A. *et al.* Global metagenomic survey reveals a new bacterial candidate phylum in geothermal springs. *Nat Commun* **7**, (2016).
35. Wilson, M. C. *et al.* An environmental bacterial taxon with a large and distinct metabolic repertoire. *Nature* **506**, 58–62 (2014).
36. Greening, C. & Cook, G. M. Integration of hydrogenase expression and hydrogen sensing in bacterial cell physiology. *Curr. Opin. Microbiol.* **18**, 30–8 (2014).
37. Greening, C. *et al.* Physiology, biochemistry, and applications of F₄₂₀- and F_o-dependent redox reactions. *Microbiol. Mol. Biol. Rev.* **80**, 451–493 (2016).
38. Markowitz, V. M. *et al.* IMG: the integrated microbial genomes database and comparative analysis system. *Nucleic Acids Research* **40**, D115–22 (2012).
39. Cover, T. & Hart, P. Nearest neighbor pattern classification. *IEEE Trans. Inf. Theory* **13**, (1967).

40. Constant, P., Chowdhury, S. P., Pratscher, J. & Conrad, R. Streptomyces contributing to atmospheric molecular hydrogen soil uptake are widespread and encode a putative high-affinity [NiFe]-hydrogenase. *Environ. Microbiol.* **12**, 821–829 (2010).
41. Stetter, K. O. *Archaeoglobus fulgidus* gen. nov., sp. nov.: a new taxon of extremely thermophilic archaeobacteria. *Syst. Appl. Microbiol.* **10**, 172–173 (1988).
42. Deppenmeier, U. & Blaut, M. Analysis of the *vhoGAC* and *vhtGAC* operons from *Methanosarcina mazei* strain Gö1, both encoding a membrane-bound hydrogenase and a cytochrome b. *Eur. J. Biochem.* **269**, 261–269 (1995).
43. Hamann, E. *et al.* Environmental *Breviatea* harbour mutualistic *Arcobacter* epibionts. *Nature* **534**, 254–258 (2016).
44. Sousa, F. L., Neukirchen, S., Allen, J. F., Lane, N. & Martin, W. F. Lokiarchaeon is hydrogen dependent. *Nat. Microbiol.* **1**, 16034 (2016).

Supporting Information

Figure S1. Sequence similarity networks showing the relationships between closely related subgroups of [NiFe]-hydrogenases as narrow logE filters.




Figure S2. Screenshot showing interface of HydDB classification page.

HydDB Classify Browse Information Pages

Classify

HydDB provides access to an accurate classifier for hydrogenase sequences and a curated database of hydrogenases by known type. The service is provided by the School of Biological Sciences, Monash University and the Bioinformatics Research Centre, Aarhus University.

Classify

Sequences

Sequences File

No file chosen

Check sequences using CDD?
If enabled, HydDB will use CDD to check whether the submitted sequences encode catalytic subunits of putative before classification. Since this step is time-consuming, you may want to uncheck this option if you are certain your sequences encode hydrogenase catalytic subunits.

Mail

If an e-mail address is provided, a mail will be sent when the job succeeds or fails.

Instructions

To use the classifier to predict the type of one or more hydrogenases from sequence, either:

- paste your FASTA-formatted protein sequences into the text area, or
- upload a FASTA-formatted file with your protein sequences.

Press the "Submit" button to upload the sequences and begin the classification.

If you provided an e-mail address you will receive an e-mail when your job finishes or fails including a link to the results. You will also be able to download the results as a CSV file.

Only sequences encoding the catalytic subunits of hydrogenases will be classified, i.e. those binding the [NiFe]-centre (NiFe-hydrogenases), [FeFe]-centre (FeFe-hydrogenases), or [Fe]-centre (Fe-hydrogenases). Electron-transfer subunits, accessory proteins, and maturation factors cannot be classified by this service.

Limits

A job can at most run for 2 hours. This should be enough for about 2500 sequences to be classified. Results will be stored for 2 weeks. However, we recommend to download the results as they may be deleted due to the rare event of a power outage or server crash.

Statistics

Jobs completed in total	40
Sequences classified in total	232
Jobs completed in the last 24 hours	0
Sequences classified in the last 24 hours	0

Figure S3. Screenshot showing the information provided in the data entry pages for 3248 individual hydrogenases in HydDB.

HydDB
Classify
Browse
Information Pages ▾

Entry WP_004030875.1

Phylum	Euryarchaeota
Order	Methanobacteriales
Organism	Methanobacterium formicicum
Hydrogenase	[Fe]
Activity (Predicted)	Bidirectional
Oxygen Tolerance (Predicted)	Tolerant
Subunits (Predicted)	1
Metal Centres (Predicted)	Fe ion
Accessory Subunits (Predicted)	None

```

MKLAILGAGCYRTHAASGITNFSRACEVAEQVGKPEIAMTHSTIAMGAEKELAGIDEIVVSDPVFDNDFTVIDDFEYEAVIEAHKDPESIMPQIREKVNVAKDLKPKPPKG
AIHFTHPEDLGFVTTDDNEAVQDADWMTWFPKGDMMQMGIIKEFADNLKEGAILTHACTVPTTFQKIFEDLSSDEMNIAPKVNVSYPHGAVPEMKGGVYIAEGYASEDAI
CKLVDWGVAAARGDAFKLPAELLPVCDMCSALTAITYAGILSYRDSVMNIIILGAPAGFAQWIAKESLTQVTDLMNSVGDHMEKLDPGALLGTADSMNFGAAADVLPVLEVL
ENRKGKGP TCNI
            
```


Figure S4. Screenshot showing the capacity for browsing hydrogenase data entries in HydDB.

NCBI Accession	Organism	Hydrogenase Class	Phylum	Order	Activity (Predicted)	Oxygen Tolerance (Predicted)	Subunits (Predicted)	Metal Centres (Predicted)	Accessory Subunits (Predicted)
WP_004030875.1	Methanobacterium formicum	[Fe]	Euryarchaeota	Methanobacteriales	Bidirectional	Tolerant	1	Fe ion	None
WP_012955328.1	Methanobrevibacter ruminantium	[Fe]	Euryarchaeota	Methanobacteriales	Bidirectional	Tolerant	1	Fe ion	None
WP_019263574.1	Methanobrevibacter smithii	[Fe]	Euryarchaeota	Methanobacteriales	Bidirectional	Tolerant	1	Fe ion	None
WP_016357634.1	Methanobrevibacter sp. AbM4	[Fe]	Euryarchaeota	Methanobacteriales	Bidirectional	Tolerant	1	Fe ion	None
WP_013296316.1	Methanothermobacter marburgensis	[Fe]	Euryarchaeota	Methanobacteriales	Bidirectional	Tolerant	1	Fe ion	None
WP_010876766.1	Methanothermobacter thermautotrophicus	[Fe]	Euryarchaeota	Methanobacteriales	Bidirectional	Tolerant	1	Fe ion	None
WP_013413799.1	Methanothermus fervidus	[Fe]	Euryarchaeota	Methanobacteriales	Bidirectional	Tolerant	1	Fe ion	None



Figure S5. Screenshot showing the detailed content of the information pages about each hydrogenase class on HydDB. Equivalent information pages are available for all 38 hydrogenase classes defined in this work (**Table 1**).

[NiFe] Group 1h-hydrogenase

This entry was last updated at: June 13, 2016, 11:11 a.m.

Properties	
Group	[NiFe] Group 1: Respiratory H ₂ -uptake [NiFe] hydrogenases
Subgroup	[NiFe] Group 1h: Actinobacteria-type
Function	Hydrogenotrophic respiration using O ₂ as terminal electron acceptor. Enzyme scavenges electrons from atmospheric H ₂ to fuel respiratory chain during carbon-starvation. Route of electron transfer unresolved.
Activity	H ₂ -uptake (unidirectional, high-affinity)
Oxygen tolerance	O ₂ -tolerant or O ₂ -insensitive
Localisation	Membrane-associated?

Distribution	
Ecosystem distribution	Upland soils, plant tissues, possibly surface waters
Taxonomic distribution	Widespread among obligately aerobic soil bacteria, especially Actinobacteria, Acidobacteria, and Chloroflexi



Architecture	
Structures	5AA5 (<i>Ralstonia eutropha</i> , 2.5 Å resolution, active)
Subunits	3?
Subunit description	HhyL (hydrogenase large subunit) HhyS (hydrogenase small subunit) HhyE (putative iron-sulfur protein and proposed physiological electron acceptor)
Catalytic site	[NiFe]-centre
FeS clusters	Proximal: 3Cys1Asp[4Fe4S] Medial: 4Cys[4Fe4S] Distal: 3Cys1His[4Fe4S]

Important Notes

The *Robiginitalea biformata* and *Sulfolobus islandicus* enzymes are relatively to distantly related to the main group. No studies have yet tested whether these enzymes have a H₂-scavenging role like other Group 1h [NiFe]-hydrogenases. They may instead represent founding members of a functionally-distinct lineage.

Sequences in this class									
NCBI Accession	Organism	Hydrogenase Class	Phylum	Order	Activity (Predicted)	Oxygen Tolerance (Predicted)	Subunits (Predicted)	Metal Centres (Predicted)	Accessory Subunits (Predicted)
WP_014267363.1	<i>Granulicella mallensis</i>	[NiFe] Group 1h	Acidobacteria	Acidobacteriales	Aerobic Uptake	Tolerant	3	[NiFe]-centre, 3 x [4Fe4S] clusters	[FeS] protein
WP_011688202.1	<i>Soilbacter usitatus</i>	[NiFe] Group 1h	Acidobacteria	Soilbacteriales	Aerobic Uptake	Tolerant	3	[NiFe]-centre, 3 x [4Fe4S] clusters	[FeS] protein
WP_021597135.1	<i>Actinomadura madurae</i>	[NiFe] Group 1h	Actinobacteria	Actinomycetales	Aerobic Uptake	Tolerant	3	[NiFe]-centre, 3 x [4Fe4S] clusters	[FeS] protein
WP_026402909.1	<i>Actinomadura rifamycinii</i>	[NiFe] Group 1h	Actinobacteria	Actinomycetales	Aerobic Uptake	Tolerant	3	[NiFe]-centre, 3 x [4Fe4S] clusters	[FeS] protein
WP_018330638.1	<i>Actinomycetospora chiangmaiensis</i>	[NiFe] Group 1h	Actinobacteria	Actinomycetales	Aerobic Uptake	Tolerant	3	[NiFe]-centre, 3 x [4Fe4S] clusters	[FeS] protein
WP_007735075.1	<i>Rhodococcus qingshengii</i>	[NiFe] Group 1h	Actinobacteria	Actinomycetales	Aerobic Uptake	Tolerant	3	[NiFe]-centre, 3 x [4Fe4S] clusters	[FeS] protein
WP_003935326.1	<i>Rhodococcus ruber</i>	[NiFe] Group 1h	Actinobacteria	Actinomycetales	Aerobic Uptake	Tolerant	3	[NiFe]-centre, 3 x [4Fe4S] clusters	[FeS] protein
WP_005443931.1	<i>Saccharomonospora azurea</i>	[NiFe] Group 1h	Actinobacteria	Actinomycetales	Aerobic Uptake	Tolerant	3	[NiFe]-centre, 3 x [4Fe4S] clusters	[FeS] protein

« 1 2 3 »

Literature

Genetics:

- Berney, M., Greening, C., Hards, K., Collins, D., and Cook, G.M. (2014) Three different [NiFe] hydrogenases confer metabolic flexibility in the obligate aerobic *Mycobacterium smegmatis*. *Environ. Microbiol.* **16**: 318-330.
- Constant, P., Chowdhury, S.P., Hesse, L., and Conrad, R. (2011) Co-localization of atmospheric H₂ oxidation activity and high affinity H₂-oxidizing bacteria in non-axenic soil and sterile soil amended with *Streptomyces* sp. PCB7. *Soil Biol. Biochem.* **43**: 1888-1893.
- Constant, P., Chowdhury, S.P., Hesse, L., Pratscher, J., and Conrad, R. (2011) Genome data mining and soil survey for the novel group 5 [NiFe]-hydrogenase to explore the diversity and ecological importance of presumptive high-affinity H₂-oxidizing bacteria. *Appl. Environ. Microbiol.* **77**: 6027-6035.
- Greening, C., Biswas, A., Carere, C.R., Jackson, C.J., Taylor, M.C., Stott, M.B., Cook, G.M., and Morales, S.E. (2016) Genomic and metagenomic surveys of hydrogenase distribution indicate H₂ is a widely utilised energy source for microbial growth and survival. *ISME J.* **10**: 761-777.
- Khdir, M., Hesse, L., Popa, M.E., Quiza, L., Lalonde, I., Meredith, L.K., Röckmann, T., and Constant, P. (2015) Soil carbon content and relative abundance of high affinity H₂-oxidizing bacteria predict atmospheric H₂ soil uptake activity better than soil microbial community composition. *Soil Biol. Biochem.* **85**: 1-9.

Physiology:

- Berney, M., Greening, C., Conrad, R., Jacobs, W.R., and Cook, G.M. (2014) An obligately aerobic soil bacterium activates fermentative hydrogen production to survive reductive stress during hypoxia. *Proc. Natl. Acad. Sci. U. S. A.* **111**: 11479-11484.
- Constant, P., Chowdhury, S.P., Pratscher, J., and Conrad, R. (2010) Streptomyces contributing to atmospheric molecular hydrogen soil uptake are widespread and encode a putative high-affinity [NiFe]-hydrogenase. *Environ. Microbiol.* **12**: 821-829.

1 **Table S1.** Hydrogenase sequences where there is disagreement between
2 classification by SSN and *k*-NN methods. These sequences represent six out of the
3 total 3248 sequences analyzed, i.e. 0.0018%.

4

NCBI Accession	Organism	<i>k</i> -NN Classification	SSN Classification
WP_027414715.1	Aneurinibacillus terranovensis	[NiFe] Group 1e	[NiFe] Group 1d
WP_027358538.1	Desulforegula conservatrix	[NiFe] Group 3d	[NiFe] Group 3c
WP_012532312.1	Geobacter bemidjiensis	[NiFe] Group 3d	[NiFe] Group 3c
WP_012469611.1	Geobacter lovleyi	[NiFe] Group 3d	[NiFe] Group 3c
WP_004512544.1	Geobacter metallireducens	[NiFe] Group 3d	[NiFe] Group 3c
WP_015839165.1	Geobacter sp. M21	[NiFe] Group 3d	[NiFe] Group 3c

5

6

7 **Dataset S1.** Excel spreadsheet listing the sequence, taxonomy, and hydrogenase
8 class of all 3248 hydrogenase catalytic subunit sequences listed in HydDB

9