

Modifications to a *LATE MERISTEM IDENTITY-1* gene are responsible for the major leaf shapes of Upland cotton (*Gossypium hirsutum* L.)

Authors

Ryan Andres^{1,6}, Viktoriya Coneva², Margaret H. Frank², John R. Tuttle¹, Sang-Won Han³, Luis Fernando Samayoa^{1,4}, Baljinder Kaur¹, Linglong Zhu¹, Hui Fang¹, Daryl Bowman¹, Marcela Rojas-Pierce³, Candace H. Haigler^{1,3}, Don C. Jones⁵, James B. Holland^{1,4}, Daniel H. Chitwood^{2,6}, Vasu Kuraparthi^{1,6}

Author Affiliations

¹ Department of Crop Science, North Carolina State University, Raleigh, North Carolina, USA.

² Danforth Plant Sciences Center, St Louis, Missouri, USA.

³ Department of Plant and Microbial Biology, North Carolina State University, Raleigh, North Carolina, USA.

⁴ USDA-ARS, Plant Science Research Unit, Raleigh, North Carolina, USA.

⁵ Cotton Incorporated, Cary, North Carolina, USA

⁶ These authors contributed equally to this work.

Correspondence should be addressed to VK (vasu_kuraparthi@ncsu.edu)

Abstract

Manipulation of cotton canopy architecture offers the potential to maximize yield while minimizing inputs. Here we show that the major leaf shapes of cotton at the *L-D₁* locus are controlled by a *Late Meristem Identity1-D1b* (*GhLMII-D1b*) gene which encodes a HD-Zip transcription factor. *Okra* leaf *GhLMII-D1b* has a 133-bp tandem duplication in the promoter, correlated with elevated expression, while an 8-bp deletion in the third exon of *normal* leaf *GhLMII-D1b* causes a frame-shifted and truncated coding sequence. Virus-induced gene silencing (VIGS) of *GhLMII-D1b* in an *okra* variety was sufficient to induce normal leaf formation. An intermediate leaf shape allele, *sub-okra*, lacks both the promoter duplication and the exonic deletion. Our results indicate that *sub-okra* is the ancestral leaf shape of tetraploid cotton and *normal* is a derived mutant allele that came to predominate and define the leaf shape of cultivated cotton.

Introduction

Cotton (*Gossypium* spp.) is the world's most important source of natural fiber as well as a leading oilseed crop. The cultivated cottons (*G. hirsutum* and *G. barbadense*) are allotetraploid species ($2n=4x=52$, AADD) formed from the hybridization of diploids *G. arboreum* ($2n=2x=26$, AA) and *G. raimondii* ($2n=2x=26$, DD)¹. Remarkable phenotypic diversity exists for leaf shape in cotton, widely ranging from entire (lacking dissection) to deeply lobed across both diploids and polyploids²⁻⁴. Although the role of leaf shape variation in the evolution and adaptation of *Gossypium* to diverse environments is not yet clearly established; leaf shape is an important agronomic trait that affects plant and canopy architecture, yield, stress tolerance, and other production attributes⁵. The four major leaf shapes of cotton: *normal*, *okra*, *sub-okra* and *super-okra* (**Fig. 1a**) are semi-dominant and allelomorphic at the *L-D₁* (*okra*) locus⁶⁻¹¹ while *laciniate*, similar in morphology to *okra*, maps to the orthologous diploid A-genome locus (*L-A₁*)¹².

The *okra* allelic series confers increasingly lobed leaf shapes from *sub-okra* to *okra* with the proximal lobes in mature *super-okra* reduced to a single linear blade (**Fig. 1a**). The characteristic shape of these four leaf morphs can be used both qualitatively and quantitatively^{2,13} to distinguish among their alleles (**Fig. 1b-e**). Classical development studies involving *okra* revealed the underlying factor acted early in leaf development in all tissue layers (L1, L2, L3) and cell autonomously^{2, 14}. Scanning Electron Micrographs (SEMs) of the shoot apical meristem show that the deeply lobed phenotype of *super-okra* is apparent by the P2 stage of leaf development (**Fig. 1f**), while the less severe *okra* manifests by the P4 stage (**Fig. 1g**).

The *L-D_I* locus was placed on the short arm of chromosome 15-D₁ (Chr15) using cytogenetics^{15, 16} and confirmed by QTL mapping¹⁷⁻²⁰. Recently, the *L-D_I* locus, was localized to a 5.4cM interval near the telomere of Chr15²¹. Shuttle mapping utilizing the *laciniate* gene (*L-A₂^L*) from *G. arboreum* further reduced the candidate region to 112kb and ten genes¹². Here, we report the identification of a *LATE MERISTEM IDENTITY1* (*GhLMI1-D1b*) gene, encoding an HD-Zip transcription factor, as the major determinant of leaf shape variation at the *L-D_I* locus in cotton.

Results

The *okra* locus explains a majority of leaf shape variation in cultivated cotton

To determine the quantitative extent that the *okra* locus is responsible for controlling leaf shape in cultivated cotton, we morphometrically analyzed 1504 leaves from 420 cultivated cotton lines (**Supplementary Table 1**). The eigenleaves resulting from a Principal Component Analysis (PCA) of the harmonic series of an Elliptical Fourier Descriptors (EFDs) analysis describe shape variance associated with linear vs. palmately-lobed leaf types (PC1) and pronounced distal lobing (PC3), in addition to fluctuating asymmetry (PC2) (**Fig. 1b**). PC1 and PC3 (in addition to other PCs not shown) separate *normal* from *sub-okra*, *okra*, and *super-okra* leaf types and explain the majority of shape variance in the cotton accessions analyzed (**Fig. 1c**). A Linear Discriminant Analysis (LDA) performed to distinguish cotton accessions by leaf shape led to the following correct percentage of assignments to phenotypic class by EFDs alone: 99.8% *normal* (1501 of 1504 cases); 70.5% *sub-okra* (31 of 44 cases), 91.6% *okra* (87 of 95 cases) and 79.2% *super-okra* (19 of 24 cases) (**Fig. 1d, e**). The classification became 99% correct (1651 of 1667 cases) if only two classes were formed: *normal* and non-*normal* (inclusive of *sub-okra*, *okra*, and *super-okra*) (**Fig. 1e**).

Our results indicate that *okra* ($L-D_1^O$), a monogenic locus, is quantitatively responsible for the majority of leaf shape variance in cotton, the alleles of which can be discriminated from each other at high correct classification rates using shape information alone. A strongly monogenic basis for leaf shape in cotton is in contrast to a polygenic basis for leaf shape described in other crops^{22,23}. Our results are consistent with classical morphometric work describing the profound role the $L-D_1^O$ locus plays in determining cotton leaf shape^{2,13}.

Fine mapping of the *L-D₁* locus in a large F₂ population

1,027 F₂ plants from the cross NC05AZ21 x NC11-2100 showed the expected phenotypic ratio for single gene inheritance of *okra* leaf shape (**Supplementary Table 2**). Genotyping with the co-dominant flanking SSRs of *L-D₁* (Gh565 and DPL0402) identified 122 recombinants and produced a genetic map (**Fig. 2a**) similar to the preliminary mapping²¹. Genotyping of these recombinants showed marker 13-LS-195 continued to co-segregate with leaf shape phenotype, confirming that 13-LS-195 is tightly linked to the leaf shape locus (*L-D₁*) in cotton²¹. The resulting 337 kb, 34 gene candidate interval (**Fig. 2b**)²¹ was further resolved to 112 kb and ten genes using orthologous mapping of the homeologous *laciniata* gene¹² (**Fig. 2c**).

Fine mapping using association mapping and isogenic lines

Novel markers: Using *G. raimondii* sequence information²⁴, four novel markers were developed within the ten gene candidate region (**Supplementary Fig. 1**): 1) a large (>100 bp) STS marker (GhLS-STS1) in the promoter region of *Gorai.002G244000*, 2) an 8-bp STS marker (13-LS-195) within *Gorai.002G244000*, 3) a C→A SNP (GhLS-SNP2) in the intron of *Gorai.002G244100* and 4) a 13-bp STS marker (GhLS-STS2) <300 bp downstream of the *Gorai.002G244200* stop codon. An additional SNP marker, (GhLS-SNP1) in the intron of *Gorai.002G243800* was also designed despite the fact that this gene had already been excluded by orthologous mapping (**Fig. 2c**).

Association mapping: Of the four markers used to genotype the diversity panel of 538 accessions, three markers (GhLS-STS1, 13-LS-195, and GhLS-SNP2) showed complete association with leaf shape (**Supplementary Table 3**) while marker GhLS-STS2 showed no association between the two most common leaf shapes, *normal* and *okra* (**Supplementary**

Table 3). The lack of association of this STS marker with leaf shape was sufficient to reduce the ten gene, 112-kb candidate region (**Fig. 2c**) to a four gene, 52 kb region between *Gorai.002G243900* and *Gorai.002G244200* (**Fig. 2d** and **Supplementary Table 3**).

Population structure was estimated in a subset of 404 lines of the 544 member panel using SSR markers distributed throughout the genome. Association tests of variants in the candidate gene region and of SSRs throughout the genome confirmed that the four remaining candidate genes showed very strong and significant ($p < 10^{-7}$) association with leaf shape after correcting for population structure (**Fig. 2e** and **Supplementary Table 4**). A few SSR markers also had significant associations, which may have occurred by chance when rare allele classes were observed in okra types (**Supplementary Fig. 2a** and **Supplementary Table 4**). After fitting the most significant candidate gene marker as a covariate and re-testing the background markers for associations, no markers outside the candidate gene interval were significant (**Supplementary Fig. 2b** and **Supplementary Table 4**).

Mapping using isogenic lines: We confirmed that genes conferring the *okra* phenotype in the mapping parent NC05AZ21 and *okra* isogenic line (LA213-*okra*) are allelic by phenotyping their F₁ (**Supplementary Fig. 3**) and an F₂ population. Genotyping of two sets of isolines (BC₈ in Stoneville 213 and BC₃ in Stoneville 7A backgrounds) with the four novel markers showed a similar marker pattern as observed in the association mapping (**Supplementary Table 3** and **Supplementary Table 5**), confirming the resolution of the candidate region to four genes and 52 kb.

Thus, characterizing the association-mapping panel and isolines with novel markers reduced the candidate genomic region to 52kb containing four genes. Of the four genes, *Gorai.002G244200* (here after *GhLMII-D1a*) and *Gorai.002G244000* (here after *GhLMII-*

D1b) are paralogues coding for HD-Zip transcription factors with 71.2% protein similarity. Their homologs were implicated in flowering time and leaf complexity in *Arabidopsis*^{25, 26, 27}. Of the remaining two putative genes, *Gorai002G244100* (hereafter *GhRLK1*), is a serine/threonine protein kinase while *Gorai.002G243900* (hereafter *GhHRA1*) is a trihelix transcription factor.

Expression analysis of leaf shape gene candidates

Expression analysis was performed to help identify the causal gene among the four remaining candidates. Semi-quantitative expression analysis revealed that neither *GhHRA1* nor *GhRLK1* were expressed in young leaf tissue (**Supplementary Fig. 4**). Based on this observation together with their lack of sequence polymorphisms from novel STS marker development, we eliminated these two genes from consideration. *GhLMII-D1a* was similarly expressed among leaf shapes while *GhLMII-D1b* expression was detected only in *okra* and *super-okra* (**Supplementary Fig. 4**). qRT-PCR confirmed the equivalent expression of *GhLMII-D1a* among leaf shapes while *GhLMII-D1b* was upregulated in *okra* and *super-okra* compared to *normal* and *sub-okra* (**Fig. 3a, b**). This indicated that up-regulation and/or ectopic expression of *GhLMII-D1b* in *okra* and *super-okra* may be responsible for these leaf shapes.

Sequencing of *GhLMII*-like genes

To identify DNA polymorphisms among leaf shapes, *GhLMII-D1a* and *GhLMII-D1b* were sequenced in 20 varieties (**Supplementary Table 6**) and comparisons were made relative to *sub-okra* (**Fig. 3c** and **Supplementary Fig. 5**). Sequence analysis of *GhLMII-D1a* identified two variants (**Supplementary Fig. 5**). Variant 1 was found in two *normal* and all five *okra* and *sub-okra* lines. Variant 2 was found in the remaining three *normal* and all five *super-*

okra lines. The 13-bp InDel (GhLS-STS2), which showed no association with leaf shape in the association analysis (**Supplementary Table 3**), is placed ~100 bp from the end of the 3' UTR. This polymorphism lies outside *GhLMII-D1a*, the proximity, lack of major polymorphisms, and identical expression among leaf shapes provide evidence against *GhLMII-D1a* as the candidate gene underlying *L-D₁*.

Sequence analysis of *GhLMII-D1b* identified two prominent polymorphisms among leaf shapes. First, a 133-bp tandem duplication located ~800 bp upstream of the transcription start site was unique to *okra* and *super-okra* (**Figure 3c**) and may explain the altered expression of *GhLMII-D1b* seen earlier (**Fig. 3b**). The second notable polymorphism was an 8-bp deletion in the 3rd exon of *GhLMII-D1b* found only in *normal* (**Fig. 3c**). The exonic location of this deletion was confirmed through Sanger sequencing of *GhLMII-D1b okra* cDNA. Translation of the resulting *normal* coding sequence produces a frameshift 156 amino acids (aa) into the protein and a premature stop codon truncating the protein from 228 aa to 178 aa (**Fig. 5a**). Neither of the conserved functional domains of an HD-Zip transcription factor appear directly impacted by the deletion. However, the frameshift introduces multiple leucines that may disturb the characteristic spacing of the leucine zipper and alter protein-protein interactions (**Fig. 5a**).

Both major polymorphisms (GhLS-STS1 for promoter duplication and 13-LS-195 for 3rd exon InDel) were completely associated with leaf shape in the association panel and isogenic lines (**Supplementary Table 3** and **5**). Taken together with the gene expression differences, these results provided strong evidence that modifications to *GhLMII-D1b* are responsible for the various leaf shapes of cotton.

Functional characterization of *GhLMII-D1b* using virus-induced gene silencing (VIGS)

As expression analysis indicated the elevated or ectopic expression of *GhLMII-D1b* could be responsible for okra leaf shape (**Fig. 3b**), we hypothesized that silencing of *GhLMII-D1b* in *okra* would reduce transcript levels and confer a simpler leaf shape. A 461bp fragment of *GhLMII-D1b*^{*Okra*} encompassing 268bp of the coding sequence (CDS) and 193bp of the 3' untranslated region (UTR) was used in VIGS. Including the 3' UTR sequence was expected to minimize off-target silencing of other *GhLMII*-like genes, especially *GhLMII-D1a*. Silencing of *GhLMII-D1b* in an *okra* isoline led to a pronounced reduction in leaf lobing compared with uninfected and negative controls and a brief period of normal leaf production (**Fig. 4a** and **Supplementary Fig. 6**). Silencing of *GhLMII-D1b* was eventually overcome, leading to a reversion to *okra* in a timeframe similar to the loss of silencing seen in the TRV: *CHLI* positive control (**Supplementary Fig. 6**).

The level of *GhLMII-D1b* transcript was substantially reduced in the TRV:*GhLMII-D1b* treatment compared to the negative controls (**Fig. 4b**). This proved that knocking down the *GhLMII-D1b* transcript through VIGS was sufficient to induce *normal* leaf formation in an *okra* variety. Furthermore, expression of *GhLMII-D1a* was unaffected by VIGS (**Fig. 4c**) demonstrating specificity to *GhLMII-D1b*. Thus, phenotyping and transcript profiling of TRV: *GhLMII-D1b* leaves confirmed that altered expression of *GhLMII-D1b* was responsible for the *Okra* leaf shape of cotton.

Phylogenetic analysis of *LMII*-like genes in diploid and tetraploid cotton

Outside of the promoter duplication *sub-okra*, *okra*, and *super-okra* *GhLMII-D1b* were identical except for one promoter SNP unique to both *okra* and *super-okra* (**Fig. 3c**).

Phylogenetic analysis confirmed the close sequence relatedness of the *sub-okra*, *okra*, and *super-okra* *GhLMII-D1b* alleles (**Fig. 5b**). However, *normal GhLMII-D1b* was considerably

different from the other alleles with six unique promoter SNPs and two SNPs in the second exon, both of which cause amino acid changes. *Normal* also carries a 1bp deletion in the 3rd exon, an additional third exon SNP, and a SNP in the 3' region (**Fig. 3c**).

The 1-bp deletion would also cause a frameshift and premature stop codon if not for the preceding 8-bp deletion. Interestingly, this 1-bp deletion is identical to one carried by the D genome donor *G. raimondii* in *Gorai.002G244000*. While it is unclear if the 1-bp deletion in *Gorai.002G244000* plays any role in the entire leaf shape of *G. raimondii* (**Fig. 5a**) it would be interesting to determine the leaf shape of a tetraploid carrying just the 1-bp deletion without the preceding 8-bp deletion. *Normal* leaf *GhLMII-D1b* also shares many of its other polymorphisms with *Gorai.002G244000* including four out of the six promoter SNPs, the G→A SNP in the 2nd exon, the T→A SNP in the third exon, and the 3' UTR SNP.

Phylogenetic analysis shows that *normal* leaf *GhLMII-D1b* CDS is more closely related to the *G. raimondii* *LMII-D1b* gene than to the alleles in other tetraploids with variable leaf shapes (**Fig. 5b**). *GaLMII-A1b* from moderately lobed *G. arboreum* is predicted to produce a full-length *LMII*-like protein that is identical in length to those produced by the non-*normal* leaf shapes of tetraploid cotton.

In order to further assess the variability of *LMII-D1b* genes in *Gossypium*, *GhLMII-D1b* was Sanger sequenced in two additional D genome diploids, *G. thurberi* and *G. trilobum*. Both of these species have highly lobed leaves similar to *G. hirsutum* *okra* (**Fig. 5a**). Both of the *G. thurberi* and *G. trilobum* *LMII-D1b* genes were full-length and similar to those found in the *G. hirsutum* leaf shape mutants and *G. arboreum*. Thus, while only a handful of species from the large *Gossypium* genus have been analyzed, there exists a trend that species with lobed leaves carry full-length *LMII-D1b* genes while those with entire

leaves produce truncated *LMII-1b* genes (**Fig. 5a**). Extending beyond *G. hirsutum*, this provides evidence for a broad role of *LMII-1b* genes in controlling the diversity of leaf shapes seen throughout *Gossypium*.

Transcriptome-wide changes accompanying *GhLMII-D1b*

The morphological effects of *okra* leaf shape are evident by the P4 leaf primordia stage (**Fig. 1g**). To analyze gene expression changes preceding observable changes in leaf morphology, we used RNA-Seq to determine transcriptional changes at the P2 stage of leaf development (**Fig. 6a**). 453 genes are differentially expressed between *okra* and *normal* BC₈ isolines (**Fig. 6b** and **Supplementary Table 7**). GO enrichment analysis for genes down regulated in *okra* indicated an association with microtubule and kinesin processes, suggesting a cellular basis for the observed changes in leaf morphology (**Supplementary Table 8**). The most GO terms for genes up regulated in *okra* relative to *normal* are related to photosynthesis (**Supplementary Table 9**). Previous transcriptomic analyses in tomato apices revealed photosynthetic-related gene expression and genetic changes in leaf complexity are positively correlated²³. The direction of this correlation is the same in *okra* (i.e., positively correlated with more dissected leaf morphology) implying possible connections between leaf development and generation of capacity for photosynthesis.

Subcellular/Nuclear localization of *GhLMII-D1b*

Transient expression of *35S::GhLMII-D1b^{Okra}-GFP* and *pUBQ10::GhLMII-D1b^{Okra}-GFP* in *Nicotiana benthamiana* was used to determine the subcellular localization of *GhLMII-D1b^{Okra}*. *GhLMII-D1b^{Okra}-GFP* was detected in the nucleus of transformed tobacco plants (**Supplementary Fig. 7**) in a pattern that co-localized with the nuclear marker RFP-

Histone2B (**Supplementary Fig. 7**). The nuclear localization of *GhLMII-D1b^{Okra}-GFP* is consistent with the predicted function of *LMII*-like genes as transcription factors²⁵⁻²⁷.

Discussion

Leaf shape varies dramatically across plant evolution and in response to the environment.

Understanding the genetic architecture of leaf shape is critical for harnessing its variation to modify plant physiology and improve agronomic profitability^{28, 29}. Using a diverse array of genomic and molecular tools, we established that the multi-allelic, major leaf shape locus *L-D₁* of cotton is governed by *GhLMII-D1b* encoding an HD-Zip transcription factor.

LMII-like genes, in particular their duplication and regulatory diversification, have recently been proposed as evolutionary hotspots for leaf shape diversity in model plants^{25, 26}. The evidence presented here indicates that the major leaf shapes of cultivated cotton are controlled by the same pathway. The near-tandem duplication of *LMII*-like genes in *Gossypium* (**Fig. 2d**) are unique from those previously described, as the Malvales and Brassicales diverged prior to duplications observed in the Brassicaceae^{26,27}. Therefore, the separate *LMII*-like duplication event in *Gossypium* indicates convergent evolution and strengthens the evidence that *LMII*-like genes are evolutionary hot spots for modifying leaf shape^{26,27}.

Sequence analyses of *GhLMII-D1b* in a set of twenty cultivars elucidated two major polymorphisms among different leaf shapes at the *L-D₁* locus (**Fig. 3c** and **Supplementary Table 5, 6**). First, an 8-bp deletion near the beginning of the third exon was found only in *normal GhLMII-D1b*. This deletion results in a frameshift and premature stop codon in the predicted *normal GhLMII-D1b* protein that may interfere with the function of the leucine zipper motif (**Fig. 5a**). Additionally, the C-terminal truncation may impact protein stability by removing residues necessary for proper folding. The second co-segregating polymorphism was a 133bp tandem duplication ~800bp upstream of the *GhLMII-D1b* transcription start site

(**Fig. 3c** and **Supplementary Table 5**). This duplication was present in all *super-okra* and *okra* alleles but absent from all *normal* and *sub-okra* alleles. Furthermore, expression of *GhLMII-D1b* was upregulated in plants with *okra* and *super-okra* leaf shapes (**Fig. 3b**). This finding is consistent with previous reports that promoter modifications to *LMII*-like genes alter expression and leaf shape in model plants^{26, 27}. Finally, the down-regulation of *GhLMII-D1b* by VIGS in a cotton accession with the *okra* leaf phenotype strongly reduced leaf lobing and resulted in the production of *normal* leaves (**Fig. 4a**).

Based on phylogenetic analysis, it appears that *normal* originated in *G. raimondii* *LMII-D1b* (**Fig. 5b**). Owing to their shared 1-bp deletion, frameshift, and premature stop codon, this protein may already have been non- or partially functional. The subsequent 8-bp deletion, which causes an earlier frameshift and premature stop codon, would be expected to even further compromise the protein function. Sequence analysis indicated the other three leaf shape alleles are not derivatives of *G. raimondii* *LMII-D1b*. This ancestral allele may have derived from a lobed D genome diploid, although likely not *G. thurberi* or *G. trilobum* (**Fig. 5b**). Once the *sub-okra* allele developed within the D genome, it likely gave rise to *okra* via a single duplication of 133bp in the promoter region. The origin of this promoter duplication is currently unclear but it may have arisen via unequal crossing over or gene conversion. Consistent with the expression results and gene silencing results obtained here, this promoter duplication led to ectopic and/or over-expression of *GhLMII-D1b* and an increase in the degree of leaf lobing and complexity. The finding of only a single additional promoter SNP in the entire *GhLMII-D1b* genic region between *sub-okra* and *okra* indicates this event happened relatively recently. Only two promoter SNPs differentiate *super-okra* from *okra* (**Fig. 3c**) consistent with the knowledge that *super-okra* originated from *okra*

within the last 90 years⁸. Phylogenetic analysis involving sequences from all *Gossypium* species would establish the evolution and adaptive significance of *LMII* genes in cotton.

Okra is an exceptional mutation affecting leaf development. It inspired an early quantitative framework for leaf shape across evolution and development^{2, 13} and through chimeric studies, revealed some of the first insights into the morphogenetic and cellular basis of leaf morphology^{14, 30}. In two separate instances within the Brassicaceae, the homologs of *GhLMII-D1b* have been implicated in evolutionary shifts in leaf morphology^{26, 27}. That the *okra* locus confers a monogenic basis to most of the leaf shape variance in cotton, the mechanisms for which have been studied in other model organisms, and that it is implicated in the productivity of a major crop, demonstrates a unique intersection between agriculture and the evolutionary and developmental basis of leaf morphology.

URLs

Clustal Omega Multiple Sequence Alignment, <http://www.ebi.ac.uk/Tools/msa/>; *G. raimondii* genome, www.phytozome.net; mRNA Translation to Protein Sequence, <http://web.expasy.org/translate/>; FancyGene <http://bio.ieo.eu/fancygene/>; Helix-turn-helix Prediction, https://npsa-prabi.ibcp.fr/cgi-bin/primanal_hth.pl ; Leucine Zipper Prediction, <http://2zip.molgen.mpg.de/>; Cotton Database, www.cottongen.org; <https://gtac.wustl.edu/>; Sequence Alignment, <http://samtools.sourceforge.net>; Read mapping <http://bedtools.readthedocs.org/en/latest/content/tools/multicov.html>; Pairwise gene comparison <https://bioconductor.org/packages/3.0/bioc/html/edgeR.html>.

METHODS

Methods and any associated references are available in the online version of the paper.

Note: Supplementary information is available in the online version of the paper.

ACKNOWLEDGEMENTS: Cotton Incorporated supported this work through its core-funding and PhD Fellowship program. For additional support of this research, we thank the NC Cotton Growers Association Inc. and the National Science Foundation: grants IOS-1238014 and IOS-1025947 supported the work of LFS and JRT, respectively. We appreciate the technical help by Jared Smith and Sharon Williamson USDA-ARS and Cathy Herring of NCDA. We appreciate Drs. Richard Percy and James Frelichowski of the USDA-NCGC for supplying the cotton germplasm.

AUTHOR CONTRIBUTIONS

VK, RA and DC designed the research and wrote the manuscript, with contributions and final review by the other authors. RA and VK performed fine mapping, gene cloning and functional genomic analyses. VC, MF and DC performed morphometric, SEM and RNA-Seq

analyses. LFS and JBH performed association analyses. LZ, BK, HF genotyped the diversity panel. RA, JRT, CHH and VK performed VIGS experiments. MRP and SWH performed the tobacco transient expression assay.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

References

1. Wendel, J.F., Brubaker, C., Alvarez, I., Cronn, R., Stewart, J.M. Evolution and natural history of the cotton genus. In: *Genetics and Genomics of Cotton* A.H. Paterson (ed.) 3-19 (2009).
2. Hammond, D. The expression of genes for leaf shape in *Gossypium hirsutum* L. and *Gossypium arboreum* L. I. The expression of gene for leaf shape in *Gossypium hirsutum* L. *Am. J. Bot.* **28**, 124-137 (1941).
3. Hutchinson, J.B., Silow, R.A., Stephens S.G., Evolution of *Gossypium*: Oxford Univ. Press, London. (1947).
4. Saunders, J.H. The wild species of *Gossypium*. Oxford Univ. Press, London (1961).
5. Andres, R.J., Bowman, D.T., Kuraparthi, V. Agronomic effects of major leaf shapes in cotton production *J. Cot. Sci.* Submitted (2016).
6. Shoemaker, D.N. A study of leaf characters in cotton hybrids. *J. Hered.* **5**, 116-119 (1909).
7. McClendon, C.A. Mendelian inheritance in cotton hybrids. *Geo. Agr. Exp. St. Bull.* **99**, 139-228 (1912).
8. Harland, S.C. The genetics of *Gossypium*. *Bibliographica Genetica* **9**, 107-182 (1932).
9. Hutchinson, J.B., Silow, R.A. Gene symbols for use in cotton genetics. *J. Hered.* **30**, 461-464 (1939).
10. Stephens, S.G. A genetic survey of leaf shape in new world cottons – a problem in critical identification of alleles. *J. Genet.* **46**, 313-330 (1945).
11. Green, J.M. Sub-okra, a new leaf shape in Upland cotton. *J. Hered.* **44**, 229-232 (1953).
12. Kaur, B., Andres, R.J., Kuraparthi, V. Major leaf shape genes, laciniate in diploid cotton and okra in polyploid Upland cotton, map to an orthologous genomic region. *Crop Sci.* **56**, 1-11 (2016).

13. Hammond, D. The expression of genes for leaf shape in *Gossypium hirsutum* L. and *Gossypium arboreum* L. II. The expression of gene for leaf shape in *Gossypium arboreum* L. *Am. J. Bot.* **28**, 138-150 (1941).
14. Dolan, L., Poethig, R.S. Genetic analysis of leaf development in cotton. *Dev. Suppl.* **1**, 39-46 (1991).
15. Endrizzi, J.E., Brown, M.S. Identification of monosomes for six chromosomes in *Gossypium hirsutum*. *Am. J. Bot.* **51**, 117-120 (1964).
16. Endrizzi, J.E., Kohel, R.J. Use of telosomes in mapping three chromosomes in cotton. *Genetics* **54**, 535-550 (1966).
17. Jiang, C., Wright, R.J., Woo, S.S., DelMonte, T.A., Paterson, A.H. QTL analysis of leaf morphology in tetraploid *Gossypium* (cotton). *Theor. Appl. Genet.* **100**, 409-418 (2000).
18. Song, X.L., Guo, W.Z., Han, Z.G., Zhang, T.Z. Quantitative trait loci mapping of leaf morphological traits and chlorophyll content in cultivated tetraploid cotton. *J. Integr. Plant Biol.* **47**, 1382-1390 (2005).
19. Lacape, J.M. *et al.* Mapping QTLs for traits related to phenology, morphology, and yield components in an inter-specific *Gossypium hirsutum* x *G. barbadense* cotton RIL population. *Field Crop Res.* **144**, 256-267 (2013).
20. Zhu, Q.H., Spriggs, A., Taylor, J.M., Llewellyn, D., Wilson, I. Transcriptome and complexity-reduced, DNA-based identification of intraspecies single-nucleotide polymorphisms in the polyploid *Gossypium hirsutum* L. *Genes, Genomes, Genetics* **4**, 1893-1905 (2014).
21. Andres, R.J., Bowman, D.T., Kaur, B., Kuraparthi, V. Mapping and genomic targeting of the major leaf shape gene (*L*) in Upland cotton (*Gossypium hirsutum* L.). *Theor. Appl. Genet.* **127**, 167-177 (2014).
22. Tian F., *et al.* Genome-wide association study of leaf architecture in the maize nested association mapping population. *Nat. Genet.* **43**, 159-162 (2011).
23. Chitwood, D.H. *et al.* A quantitative genetic basis for leaf morphology in a set of precisely defined tomato introgression lines. *Plant Cell* **25**, 2465-2481 (2013).
24. Paterson, A.H. *et al.* Repeated polyploidization of *Gossypium* genomes and the evolution of spinnable fibers. *Nature* **492**, 423-427 (2012).

25. Saddic, L.A. *et al.* The LEAFY target LMI1 is a meristem identity regulator and acts together with LEAFY to regulate expression of CAULIFLOWER. *Development* **133**, 1673-1682 (2006).
26. Vlad, D. *et al.* Leaf shape evolution through duplication, regulatory diversification, and loss of a homeobox gene. *Science* **343**, 780-783 (2014).
27. Sicard, A. *et al.* Repeated evolutionary changes of leaf morphology caused by a mutation to a homeobox gene. *Curr. Biol.* **24**, 1880-1886 (2014).
28. Nicotra, A.B., Leigh, A., Boyce, C.K., Jones, C.S., Niklas, K.J., Royer, D.L., Tsukaya, H. The evolution and functional significance of leaf shape in the angiosperms. *Functional Plant Biol.* **38**, 535-552 (2011).
29. Chitwood, D.H., Sinha, N.R. Evolutionary and environmental forces sculpting leaf development. *Curr. Biol.* **26**, R297-R306 (2016).
30. Dolan L., Poethig, S.R. The OKRA leaf shape mutation in cotton is active in all cell layers of the leaf. *Am. J. Bot.* **85**, 322-327 (1998).

Figure legends

Figure 1 Morphometric analysis of leaf shapes conferred by the *sub-okra*, *okra*, and *super-okra* mutations. (a) Left to right, leaves representative of *normal*, *sub-okra*, *okra*, and *super-okra* leaf shape phenotypes. (b) Eigenleaves representative of leaf shapes found ± 3.5 standard deviations along each Principal Component (PC) axis calculated from the harmonic series of an Elliptical Fourier Descriptor (EFD) analysis. Percent variance explained by each PC is provided. (c) PC1 and PC3 (PC2 not shown, as it explains asymmetric shape variance) separate *normal* and various *okra* leaf shape classes. 95% confidence ellipses provided. (d) Linear Discriminant Analysis (LDA) maximizes the discrimination of *normal* and *okra* leaf shape classes. LD1 and LD2 are shown, and the percent separation between phenotypic classes of leaves indicated. 95% confidence ellipses provided. (e) A confusion matrix, showing actual vs. predicted leaf shapes, constructed using linear discriminants. Leaf shape alone discriminates a majority of *normal* from *sub-okra*, *okra*, and *super-okra* leaf types. (f) Scanning Electron Micrographs (SEMs) of the Shoot Apical Meristem (SAM), P1, and P2 leaf primordia for *normal*, *okra*, and *super-okra* shoot apices. Note the displaced lobe in the P2 of *super-okra* relative to *normal*. (g) SEM of *normal* and *okra* shoot apices showing a more pronounced leaf lobe present by the P4 stage of leaf primordium development in *okra* relative to *normal*. Colors: *normal*, red; *sub-okra*, green; *okra*, blue; *super-okra*, purple.

Figure 2 Genetically resolving the *L-D₁* locus in Upland cotton. (a) Genetic mapping of *L-D₁* locus based on bi-parental mapping. (b) Orthologous mapping to the sequenced D genome donor (*G. raimondii*) chromosome 2 (337kb, 34 putative genes)²¹. (c) Shuttle mapping utilizing the orthologous *laciniate* (*L-A₂*) locus from donor diploid *G. arboreum* (112kb, 10 putative genes)¹². (d) Fine mapping combining an association mapping panel and two sets (BC₈ and BC₃) of isogenic lines (52kb, putative 4 genes). (e) Association analysis statistics, adjusting for population structure for variants within candidate gene region.

Figure 3 Nucleotide polymorphisms of *GhLMII-D1b* gene and expression analysis of candidate genes using qRT-PCR among different leaf shapes. (a, b) Relative transcript levels of leaf shape candidate genes (*GhLMII-D1a* and *GhLMII-D1b*) among isolines at ~90 days after planting. There is no difference in the expression of *GhLMII-D1a* among leaf shapes but a significant increase of *GhLMII-D1b* expression in *okra* and *super-okra*. Error bars represent the standard deviation of the fold change. (c) Polymorphisms among *GhLMII-D1b* sequences from the four major leaf shapes. The 133bp tandem promoter duplication was found only in *okra* and *super-okra* and in parallel with elevated gene expression. The 8bp deletion was found only in *normal* and causes a frameshift mutation and premature stop codon. All other polymorphisms are SNPs with unknown effect on the gene expression and protein function. *Sub-okra* is set as the standard to which the other three leaf shapes are compared.

Figure 4 Functional characterization of *GhLMII-D1b* using virus-induced gene silencing (VIGS). (a) Representative 6th true leaf and 4 week old plants from VIGS experiment showing the reversion to *normal* leaf shape in the *GhLMII-D1b* silencing treatment. (b) Relative transcript levels of candidate genes in the *GhLMII-D1b* silenced and control LA213-Okra plants confirmed the effective knockdown of *GhLMII-D1b*. (c) Transcript levels of *GhLMII-D1a* were unaffected by VIGS treatment, confirming silencing was specific to *GhLMII-D1b*.

Figure 5 Functional prediction and phylogenetic analysis of *LMII*-like genes among different cotton leaf shapes. (a) Amino acid translations in the *Gossypium LMII-D1b* alleles. Normal in *G. hirsutum* and simple leaf in *G. raimondii* show truncated proteins while *G. hirsutum-okra*, *G. trilobum* and *G. thurberii* with dissected leaves code for functional *LMII-D1b* protein. Frameshift mutation resulting from 8bp deletion in *normal* introduces additional leucines (L) at 7 amino acid intervals (highlighted in red) that may interfere with the functionality of this domain. (b) Phylogenetic analysis showing the close relationship between *okra*, *sub-okra*, and *super-okra* but not to *G. thurberi* or *G. trilobum*. Conversely, *normal* CDS appears more closely related *G. raimondii*.

Figure 6 Transcriptomic comparison of *okra* versus *normal* P2 stage primordia. (a) SEMs of *normal* and *okra* shoot apices. P2 stage leaf primordia (highlighted in green) were hand dissected and processed for Illumina RNA-sequencing. (b) Heatmap visualization of the 453 genes that are significantly differentially expressed (false discovery rate ≤ 0.05) between *normal* and *okra* P2 samples (**Supplementary Table 7-9**). Scaled and centered reads per million values for *normal* and *okra* biological replicates are plotted in the left three columns and right three columns, respectively. Yellow indicates up regulated and blue indicates down regulated expression values. Scale bars = 100 micrometers.

Supplementary figures

Supplementary Figure 1 Markers used in association mapping and isogenic line studies. Four novel markers: GhLS-STS1, GhLS-STS2, GhLS-SNP1 and GhLS SNP2 were developed and run in both studies along with 13-LS-195 that had previously co-segregated with leaf shape phenotype²¹. Four markers: GhLS-STS1, 13-LS-195, GhLS-SNP1 and GhLS-SNP2 showed complete association with leaf shape phenotype. GhLS-STS2 showed no association with leaf shape.

Supplementary Figure 2 (a) Results of genome-wide association scan for leaf shape in diverse set of 406 cotton inbred lines. Blue circles represent the *P*-value of each marker tested in a logistic regression model that also included the first three principal components of

the population structure analysis. Three size-based markers from candidate genes are indicated by red arrows while two SNP markers are indicated by green arrows. The green and red dashed lines represent the 0.05 and 0.01 significance levels. **(b)** Results of second genome-wide association scan for leaf shape in diverse set of 406 cotton inbred lines, adjusting for the effects of the candidate gene. Blue circles represent the *P*-value of each marker tested in a logistic regression model that also included the most significant candidate gene marker from the initial scan plus the first three principal components of the population structure analysis. The green and red dashed lines represent the 0.05 and 0.01 significance levels.








Supplementary Figure 3 Allelism between *okra* leaf shapes of parental accession (NC05AZ21) and isoline (LA213-*okra*). *Okra* leaf shape genes in isoline LA213-*okra* and NC05AZ21 are allelic. Leaf shape phenotypes of greenhouse grown parents and their F₁ hybrid at approximately 40 days after planting.


Supplementary Figure 4 Semi-quantitative expression analysis of leaf shape candidate genes. Neither *GhHRA1* (*Gorai.002G243900*) nor *GhRLK1* (*Gorai.002G244100*) were expressed in critical young leaf tissue, eliminating these two genes from consideration. *GhLMII-D1a* (*Gorai.002G244200*) appeared equally expressed across leaf shapes. *GhLMII-D1b* (*Gorai.002G244000*) was expressed only in leaf shapes (*okra* and *super-okra*) with the larger promoter indicating differential expression of this gene could play a major role in leaf shape. Glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) was the reference gene.


Supplementary Figure 5 Nucleotide polymorphisms of the *GhLMII-D1a* gene among different leaf shapes. Two variants of *GhLMII-D1a* were found, neither of which perfectly associated with leaf shape. Both variants encode full-length proteins and have no obvious polymorphisms that would alter function.




Supplementary Figure 6 Leaves from representative VIGS plants. True leaves five through ten from representative plants of all VIGS treatments are shown. Severe reductions in leaf lobing and sinus depth was seen in the LA213-*okra* TRV: *LMII-D1b* that briefly produced normal leaves. Abolishment of viral silencing proceeded similar to that seen in the TRV: CHLI positive control treatment.

Supplementary Figure 7 *GhLMII-D1b*^{*Okra*} localizes to the nucleus. A *GhLMII-D1b*^{*Okra*}-*GFP* fusion construct driven by the 35S (a,b,c) or pUBQ10 (d,e,f) promoter was transiently expressed in *Nicotiana benthamiana* leaves. Signal from *GhLMII-D1b-Okra-GFP* (a,d), the nuclear marker *RFP-Histone2B* (b,e) or the merged images including brightfield (c,f) are shown. Scale bar: 20µm.







a**b****c**

a

LA213-Okra: Uninfected





LA213-Okra: TRV-GhLM1-D1b



LA213-Okra: TRV-Mock



LA213-Normal: Uninfected

**b****c**

a



Gh-normal



Gh-okra



G. raimondii



G. thurberi



G. trilobum

GGGGLL-CL NormalPectin
GGGGLL-CL Carbohydrate
GGGGLL-CL Protein
GGGGLL-CL Protein
GGGGLL-CL Protein

GGGGLL-CL NormalPectin
GGGGLL-CL Carbohydrate
GGGGLL-CL Protein
GGGGLL-CL Protein
GGGGLL-CL Protein

GGGGLL-CL NormalPectin
GGGGLL-CL Carbohydrate
GGGGLL-CL Protein
GGGGLL-CL Protein
GGGGLL-CL Protein

GGGGLL-CL NormalPectin
GGGGLL-CL Carbohydrate
GGGGLL-CL Protein
GGGGLL-CL Protein
GGGGLL-CL Protein

GGGGLL-CL NormalPectin
GGGGLL-CL Carbohydrate
GGGGLL-CL Protein
GGGGLL-CL Protein
GGGGLL-CL Protein

Multiple sequence alignment of Gh-normal, Gh-okra, G. raimondii, G. thurberi, and G. trilobum.

GGGGLL-CL NormalPectin
GGGGLL-CL Carbohydrate
GGGGLL-CL Protein
GGGGLL-CL Protein
GGGGLL-CL Protein

Multiple sequence alignment of Gh-normal, Gh-okra, G. raimondii, G. thurberi, and G. trilobum.

GGGGLL-CL NormalPectin
GGGGLL-CL Carbohydrate
GGGGLL-CL Protein
GGGGLL-CL Protein
GGGGLL-CL Protein

GGGGLL-CL NormalPectin
GGGGLL-CL Carbohydrate
GGGGLL-CL Protein
GGGGLL-CL Protein
GGGGLL-CL Protein

b



G. arboreum



G. raimondii



G. hirsutum (Normal)



G. hirsutum (Okra)



G. hirsutum (Super-Okra)




G. hirsutum (Sub-Okra)




G. thurberi



G. trilobum





a*normal**okra***b**


G. raimondii

Chr 02

Gene 24...










LA213-Okra



F₁: LA213-Okra x NC05AZ21



NC05AZ21






Variant 1



2-normals, 5-okras,
5-sub-okra lines

Variant 2



3-normals, 5-okras,
5-super-okras

Intron 
UTR 
Exon 



LA213-Okra (uninfected)



LA213-Okra (Mock)




LA213-Okra (TRV:GFP)



LA213-Okra (TRV:CHLI)



LA213-Okra (TRV:LMI1-D1b)





LA213-Normal

35S::GhLMI1-D1b-GFP

RFP-Histone2B



Merged



UBQ10::GhLMI1-D1b-GFP

RFP-Histone2B

Merged



Scale Bar = 20 μ m