

Flow environment and matrix structure interact to determine spatial competition in *Pseudomonas aeruginosa* biofilms

Carey D. Nadell^{1*}, Deirdre Ricaurte^{2*}, Jing Yan², Knut Drescher¹, and Bonnie L. Bassler^{2,3†}

5

¹ Max Planck Institute for Terrestrial Microbiology, Marburg, HE, Germany

² Department of Molecular Biology, Princeton University, Princeton, NJ, USA

³ Howard Hughes Medical Institute, Chevy Chase, MD, USA

10 * Co-first authorship

† Correspondence to: bbassler@princeton.edu

Abstract

Bacteria often live in biofilms, which are dense microbial communities surrounded by a secreted extracellular matrix. Here, we demonstrate that hydrodynamic flow and matrix organization interact to shape competitive dynamics in *Pseudomonas aeruginosa* biofilms. Irrespective of initial frequency, in competition with matrix mutants, wild type cells always increase in relative abundance in flat-surfaced, straight-tunnel microfluidic devices under simple flow regimes. By contrast, in microenvironments with complex, irregular flow profiles – which are common in natural environments – wild type matrix-producing and isogenic non-producing strains can coexist. This result stems from local obstruction of flow by wild-type matrix producers, which generates regions of near-zero flow speed that favor occupation by cells that do not invest in matrix secretion. Our findings connect the evolutionary stability of matrix production with the hydrodynamics and spatial structure of the surrounding environment, providing a potential explanation for the variation in biofilm matrix secretion observed among bacteria in natural environments.

25 Introduction

In nature, bacteria predominantly exist in biofilms, which are surface-attached or free-floating communities of cells held together by a secreted matrix [1-3]. The extracellular matrix defines biofilm structure, promoting cell-cell and cell-surface adhesion and conferring resistance to chemical and physical insults [4-6]. The matrix also plays a role in the population dynamics of biofilm-dwelling bacteria [2, 7, 8]; simulations and experiments using *Vibrio cholerae*, *Pseudomonas fluorescens*, and *Pseudomonas aeruginosa* show that matrix-secreting cell lineages can physically smother or laterally displace other cell lineages and, in so doing, outcompete neighboring matrix non-producing cells [9-16]. However, not all wild and clinical isolates of these species produce biofilm matrices, despite the clear ecological and competitive benefits of possessing a matrix [17-19]. Fitness trade-offs between the benefits of being adhered to surfaces and the ability to disperse to new locations can cause variability in matrix production [10, 18]. However, it is not known how selective forces within the biofilm environment itself might drive the coexistence of strains that make matrix with strains that do not. Here we explore how selection for matrix production occurs in biofilms on different surface geometries and flow regimes, including those that are relevant inside host organisms and in abiotic environments such as soil.

The local hydrodynamics associated with natural environments can have dramatic effects on biofilm matrix organization. This phenomenon has been particularly well established for *P. aeruginosa*, a common soil bacterium [20] and opportunistic pathogen that thrives in open wounds [21, 22], on sub-epithelial medical devices [23], and in the lungs of cystic fibrosis patients [24-26]. Under laminar flow in simple microfluidic channels, *P. aeruginosa* forms biofilms with intermittent mushroom-shaped tower structures [24, 27-29]. Under irregular flow regimes in more complex environments, however, *P. aeruginosa* also produces sieve-like biofilm streamers that protrude into the liquid phase above the substratum [30-33]. These streamers – whose structure depends on the secreted matrix – are proficient at catching cells, nutrients, and debris that pass by, leading to clogging and termination of local flow [34]. The spatial and temporal characteristics of flow thus combine to alter matrix morphology, which, in turn, feeds back to alter local hydrodynamics and nutrient advection and diffusion. By modifying community structure and solute transport in and around biofilms [35], this feedback could have a significant influence on the evolutionary dynamics of matrix secretion in natural environments.

Here, we examined the relationship between flow regime and matrix-mediated biofilm competition using strains of *P. aeruginosa* PA14 that differ only in their production of Pel, a visco-elastic matrix polysaccharide that is central to biofilm and streamer formation [27, 34, 36, 37]. Using a combination of fluid flow visualization and population dynamics analyses, we reveal a novel interaction between hydrodynamic conditions, biofilm architecture, and competition within bacterial communities.

60 Results / Discussion

We performed competition experiments with wild type PA14 and an otherwise isogenic strain deleted for *pelA*, which is required for synthesis of Pel [38]. Deletion of *pelA* significantly impairs biofilm formation in PA14, which does not naturally produce Psl, an additional matrix polysaccharide secreted by other *P. aeruginosa* isolates [39, 40]. Wild type cells produced GFP, and $\Delta pelA$ mutants produced mCherry. Experiments in shaken liquid culture using genetically identical cells producing GFP or

mCherry confirmed that fluorescent protein expression constructs had no measurable effect on growth rate (Figure S1). Our first goal was to compare the population dynamics of the wild type and $\Delta pelA$ strains in typical straight-chamber microfluidic devices, which have simple parabolic flow regimes, and in porous environments containing turns and corners, which have irregular flow profiles and better reflect the packed soil environments that *P. aeruginosa* often occupies [20, 41, 42]. To approximate the latter environment, we used microfluidic chambers containing column obstacles; the size and spacing distributions of the column obstacles were designed specifically to simulate soil or sand. Analogous methods have been used previously to study bacterial growth [43] and the behavior of *Caenorhabditis elegans* [44] in realistic environments while maintaining accessibility to microscopy. In our setup, flow was maintained through these chambers at rates comparable to those experienced by *P. aeruginosa* in a soil environment [45].

The flow regime alters selection for matrix production in biofilms

Several approaches are available to study how competitive dynamics differ in particular flow environments. Most commonly, one would monitor biofilm co-cultures of wild type and $\Delta pelA$ PA14 cells over time until their strain compositions reached steady state. Performing such time-series experiments was not possible here due to a combination of low fluorescence output in the early phases of biofilm growth coupled with phototoxicity incurred by cells during epifluorescence imaging. To circumvent this issue, we measured the change in frequency of wild type and $\Delta pelA$ cells as a function of their initial ratio in both straight-tunnel chambers and soil-mimicking chambers containing column obstacles. From these measurements, we could infer the final stable states of Pel-producing and non-producing cells as a function of surface topography and flow conditions. This method is commonly used to measure the behavior of dynamical systems, and it has been employed successfully in a variety of analogous experimental applications [10, 11, 16, 46-48].

In simple flows, wild type PA14 increased in relative abundance regardless of initial population composition, indicating uniform positive selection for Pel secretion. This result is consistent with recent studies of *V. cholerae* and *Pseudomonas* spp. demonstrating that the secreted matrix cannot be readily exploited by non-producing mutants (Figure 1A,B) [10, 12, 14, 16]. Wild type cells obtained their competitive advantage in biofilms despite a reduction in maximum growth rate relative to $\Delta pelA$ mutants in shaken liquid culture (Figure S1). Confocal microscopy revealed that Pel-producers mostly excluded non-Pel-producing cells from biofilm clusters in straight tunnel chambers (Figure 2A,B), although some $\Delta pelA$ mutants resided on the periphery of wild type biofilms. The liquid effluent from these chambers contained an over-representation of the $\Delta pelA$ mutant relative to wild type, consistent with the interpretation that $\Delta pelA$ strain was displaced from the substratum over time (Figure 2C). When wild type and $\Delta pelA$ cells competed in microfluidic devices simulating porous microenvironments, there was a pronounced shift to negative frequency-dependent selection for Pel production (Figure 1A,C). Wild type PA14 was selectively favored at initial frequencies below ~0.6. Above this critical frequency, the $\Delta pelA$ mutant was favored. From this result, we can infer that in this porous environment, $\Delta pelA$ null mutants can grow and stably coexist with wild type Pel-producers.

Wild type P. aeruginosa obstructs porous environments, generating shear-free regions that Pel-deficient mutants can occupy

Previous work has shown that in environments with corners, wild type *P. aeruginosa* produces biofilm streamers that catch cells and other debris in the surrounding flow [34]. Such streamers were produced by the wild type in our system. Although $\Delta pelA$ cells could be found in these structures, they were not abundant, so cell capture by streamers cannot account for the observed coexistence of the two strains. This result suggests that streamers do indeed catch debris, consistent with prior studies [31, 34], but that in our system, $\Delta pelA$ cells are not abundant enough in the passing liquid phase to accumulate substantial population sizes by this manner (Figure S2).

Our microscopy-based observation of soil-mimicking chambers suggested that wild type biofilms gradually obstructed some of the regions located between columns over time. We hypothesized that partial clogging could render those portions of the chambers more suitable for growth of the $\Delta pelA$ strain, which was previously shown to be sensitive to removal by shear [39, 40]. This hypothesis predicts that $\Delta pelA$ cells should be found predominantly in regions of the chamber that have been clogged by wild type biofilms, or immediately downstream of such regions. We tested this prediction by repeating our co-culture competition experiment with wild type and $\Delta pelA$ cells in chambers containing columns, measuring the distribution of each strain as above, and then re-inserting new influent tubing through which fluorescent beads were introduced to the chambers (Figure S3). By tracking the beads with high frame-rate microscopy, we could trace the presence and absence of fluid flow and then superimpose this information onto the spatial distributions of wild type and $\Delta pelA$ mutant cells. Wild type biofilms accumulated primarily between columns parallel to the long axis of the chamber, with clusters of $\Delta pelA$ cells in close proximity. Importantly, in support of our prediction, $\Delta pelA$ cells grew most successfully in regions that were locally obstructed by wild type biofilms (Figure 3A,B). $\Delta pelA$ cells did not clog chambers when grown in isolation, supporting the interpretation that $\Delta pelA$ accumulation relies on and only occurs after clogging by the wild type. This result supports prior indications that Pel-producers do indeed have a higher tolerance for shear than Pel-deficient cells [39, 40]. We directly confirmed this prediction in our system by independently growing each strain in isolation under varying shear stress in simple straight-tunnel chambers (Figure 3C).

Collectively, our findings suggest that when *P. aeruginosa* wild type and $\Delta pelA$ mutants experience irregular flow in heterogeneous environments, wild type biofilm formation causes partial clogging, regionally reducing local flow speed. The lack of flow generates favorable conditions for the $\Delta pelA$ strain, whose biofilm would otherwise be removed by shear forces, enabling it to proliferate locally using nutrients diffusing from the bulk liquid phase [49].

Conclusions

Our results show that a feedback process operates between hydrodynamic flow conditions, biofilm spatial architecture, and competition in *P. aeruginosa* biofilms. Pel non-producers do not appear to directly exploit the matrix material produced by wild type cells, but they can take advantage of the low-shear conditions that wild type matrix producers generate in complex flow environments. *P. aeruginosa* is notorious as an opportunistic pathogen of plants and animals, including humans. It also thrives outside of hosts, for example, in porous niches such as soil. Despite the well-documented ecological benefits of matrix secretion during biofilm formation, environmental and clinical isolates of *P. aeruginosa* exhibit considerable variation in their production of matrix components, including loss

or overexpression of Pel [17]. Our results offer an explanation for natural variation in the ability of *P. aeruginosa* to produce extracellular matrix, particularly among bacteria in porous environments: the evolutionary stable states of extracellular matrix secretion vary with the topographical complexity of the flow environment in which bacteria reside.

Materials and Methods

Strains

All strains are derivatives of *Pseudomonas aeruginosa* PA14. Wild type PA14 strains constitutively producing fluorescent proteins [34] were provided by Albert Siryaporn (UC Irvine), and they harbor genes encoding either EGFP or mCherry under the control of the $P_{A1/04/03}$ promoter in single copy on the chromosome [50]. The $\Delta pelA$ strain was constructed using the lambda red system modified for *P. aeruginosa* [51].

Liquid growth rate experiments

To determine maximum growth rates and the potential for fluorescent protein production to cause fitness differences, bacterial strains were grown overnight in Luria Bertani (LB) medium at 37° C. To acclimatize the cultures to the metabolic shift that occurs in minimal medium, cultures were back-diluted into minimal M9 medium with 0.5% glucose and monitored until their optical densities at 600 nm were ~0.6, corresponding to mid-logarithmic phase. Cultures were back diluted again into minimal M9 medium with 0.5% glucose and transferred to 96 well plates. Hourly measurements of culture optical density were taken until saturation, corresponding to stationary phase. MatLab curve fitting software was used to calculate the maximum growth rate of each strain. These experiments confirmed that the fluorescent protein markers had no measurable effect on growth rates and thus did not contribute to competitive outcomes in our experiments.

Microfluidics and competition experiments

Microfluidic devices consisting of poly(dimethylsiloxane) (PDMS) bonded to 36 mm x 60 mm glass slides were constructed using standard soft photolithography techniques [52]. We used flat, straight-tunnel microfluidic devices with no obstacles to simulate environments with simple parabolic flow profiles, and we used devices with PDMS pillars interspersed throughout the chamber volume to simulate environments with complex (i.e. irregular, non-parabolic) flow profiles. The size and spatial distributions of these column obstacles were determined by taking a cross-section through a simulated volume of packed beads mimicking a simple soil environment. Flow rates through these two chamber types were adjusted to equalize the average initial flow velocities, although the local flow velocity varied as biofilms grew during experiments (see main text).

For all competition experiments, bacterial strains were grown overnight. The following morning, aliquots of the overnight cultures were added to Eppendorf tubes, and their optical densities were equalized prior to preparation of defined mixtures of wild type and $\Delta pelA$ cells. 100 μ L volumes of the wild type strain alone, the $\Delta pelA$ strain alone, or mixtures of the two strains (for competition experiments), were introduced into microfluidic chambers using 1 mL syringes and Cole-Parmer polytetrafluoroethylene tubing (inner diameter = 0.30 mm; outer diameter = 0.76 mm). After 3 h, fresh tubing connected to syringes containing minimal M9 medium with 0.5% glucose were inserted into the

inlet channels. Syringes (3 mL BD Syringe, 27G) were mounted onto high precision syringe pumps (Harvard Apparatus), which were used to tune flow rate according to empirical measurements of flow speeds in soil [45]. It should be noted that microfluidic experiments in the chambers containing columns experience a high failure rate, in which no biofilms appear to grow after the 72 h period of the experiment. This problem was overcome by performing the experiment at high replication in parallel. Sufficient data were thus collected to populate the relevant panels in Figures 1 and 3 of the main text.

Microscopy and image analysis

Mature biofilms were imaged using a Nikon Ti-E inverted microscope via a widefield epifluorescence light path (using a 10x objective) or a Borealis-modified Yokogawa CSU-X1 spinning disk confocal scanner (using a 60x TIRF objective). A 488 nm laser line was used to excite EGFP, and a 594 nm laser line was used to excite mCherry. For all competition experiments, 3-5 biological replicates were prepared on different days in different microfluidic chambers. Within each chamber, 4-6 non-overlapping, randomly selected areas (technical replicates) were selected and imaged for quantification. Quantification of biofilm composition was performed using custom MatLab scripts for image analysis [48]. Imaging of biofilms could only be performed once for each experiment, precluding time-series analyses, due to phototoxicity effects after multiple rounds of imaging. Phototoxicity was a particularly notable issue here due to dimness of the fluorescent proteins in *P. aeruginosa*, which made long exposures necessary to capture images of sufficient quality for later analysis. For this reason we opted for inferential population dynamics analysis as described in the main text.

Effluent Measurements

To measure strain frequencies in the biofilm effluent of straight-tunnel chambers (Figure S2), 1:1 strain mixtures of wild type and $\Delta pelA$ cells were prepared and inoculated into simple flow chambers according to the procedure outlined above for competition experiments. At 0, 24, and 48 h, 5 μ L samples were collected from the microfluidic chamber outlet tubing, mixed vigorously by vortex, and plated onto agar in serial dilution. After overnight growth at 37° C, plates were imaged with an Image Quant LAS 4000. Cy3 and Cy5 fluorescence settings were used for EGFP and mCherry excitation, respectively. Image Quant TL Colony Counting software was used to measure the relative abundance of each strain.

Flow Tracking Experiments

1:1 mixtures of the wild type and the $\Delta pelA$ mutant were prepared and introduced into obstacle-containing flow chambers according to the procedure described above. Minimal M9 medium with 0.5% glucose was introduced into the chambers for 72 h as described above. The entire chamber was then imaged using widefield epifluorescence microscopy to document the locations of wild type and $\Delta pelA$ cell clusters. Subsequently, the influent syringes were replaced with syringes containing yellow-green fluorescent beads (Invitrogen; sulfate-modified, diameter = 2 μ m) at a concentration of 0.3%. Bead suspensions were flowed into the microfluidic chambers. To determine the presence or absence of flow with respect to the spatial distributions of wild type and $\Delta pelA$ cells, and to obtain large images for statistics, the entire chamber was imaged with a 1 s exposure time, over which traveling beads were captured as streaks. It should be noted that this experiment also has a high failure rate due to the

235 sensitivity of the microfluidic chambers to removal and re-insertion of tubing, and required
optimization to execute successfully. Custom MatLab code was written to correlate the presence or
absence of fluid flow with the accumulation of wild type and *ΔpelA* cells. In brief, the positions of the
columns were first identified and used to divide the chamber into triangular sampling areas using a
240 network structure in which columns served as nodes and straight lines between column centers served
as vertices. Within each sampling triangle, the area covered by columns was first removed, and
subsequently, the averaged *ΔpelA* and wild type fluorescence intensities in the remaining area were
used to determine if a particular region had wild type and/or *ΔpelA* accumulation. In parallel, each
sampling area was scored for the presence of flow in the corresponding bead tracking images Figure
S3).

245

Author Contributions

CDN and BB conceived the project. CDN, DR, JY, and KD performed experiments and analyzed data.
KD and JY contributed new analytical tools. All authors wrote the manuscript.

250 Conflict of Interest

The authors declare no conflict of interest.

Acknowledgments

We thank members of the B.L.B. laboratory, as well as Thomas Bartlett and Alvaro Banderas for
255 helpful discussions. This work was supported by the Alexander von Humboldt Foundation (CDN), the
Human Frontier Science Program Grant CDA00084/2015-C (KD), the Max Planck Society (KD), the
Howard Hughes Medical Institute (BLB), NIH Grant 2R37GM065859 (BLB), National Science
Foundation Grant MCB-0948112 (BLB), and National Science Foundation Grant MCB-1344191
(BLB).

260

References

1. Hall-Stoodley, L., Costerton, J.W., and Stoodley, P. (2004). Bacterial biofilms: from the
natural environment to infectious diseases. *Nat Rev Microbiol* 2, 95-108.
2. Nadell, C.D., Xavier, J.B., and Foster, K.R. (2009). The sociobiology of biofilms. *Fems*
265 *Microbiol Rev* 33, 206-224.
3. Flemming, H.-C., Wingender, J., Szewzyk, U., Steinberg, P., Rice, S.A., and Kjelleberg, S.
(2016). Biofilms: an emergent form of bacterial life. *Nat Rev Microbiol* 14, 563-575.
4. Flemming, H.-C., and Wingender, J. (2010). The biofilm matrix. *Nat Rev Microbiol* 8, 623-
633.
- 270 5. Hobley, L., Harkins, C., MacPhee, C.E., and Stanley-Wall, N.R. (2015). Giving structure to the
biofilm matrix: an overview of individual strategies and emerging common themes. *Fems*
Microbiol Rev.
6. Teschler, J.K., Zamorano-Sanchez, D., Utada, A.S., Warner, C.J.A., Wong, G.C.L., Linington,
R.G., and Yildiz, F.H. (2015). Living in the matrix: assembly and control of *Vibrio cholerae*
275 biofilms. *Nat Rev Micro* 13, 255-268.
7. Steenackers, H.P., Parijs, I., Foster, K.R., and Vanderleyden, J. (2016). Experimental
evolution in biofilm populations. *Fems Microbiol Rev*, fuw002.

8. Nadell, C.D., Drescher, K., and Foster, K.R. (2016). Spatial structure, cooperation, and competition in bacterial biofilms. *Nat Rev Microbiol* **14**, 589-600.
- 280 9. Xavier, J.B., and Foster, K.R. (2007). Cooperation and conflict in microbial biofilms. *Proc Natl Acad Sci U S A* **104**, 876-881.
10. Nadell, C.D., and Bassler, B.L. (2011). A fitness trade-off between local competition and dispersal in *Vibrio cholerae* biofilms. *Proc Natl Acad Sci USA* **108**, 14181-14185.
11. Nadell, C.D., Drescher, K., Wingreen, N.S., and Bassler, B.L. (2015). Extracellular matrix structure governs invasion resistance in bacterial biofilms. *ISME J* **9**, 1700-1709.
- 285 12. Kim, W., Racimo, F., Schluter, J., Levy, S.B., and Foster, K.R. (2014). Importance of positioning for microbial evolution. *Proc Natl Acad Sci U S A* **111**, E1639-1647.
13. Schluter, J., Nadell, C.D., Bassler, B.L., and Foster, K.R. (2015). Adhesion as a weapon in microbial competition. *ISME J* **9**, 139-149.
- 290 14. Irie, Y., Roberts, A., Kragh, K.N., Gordon, V.D., Hutchison, J., Allen, R.J., Melaugh, G., Bjarnsholt, T., West, S.A., and Diggle, S.P. (2016). The *Pseudomonas aeruginosa* PSL polysaccharide is a social but non-cheatable trait in biofilms. *bioRxiv*.
15. Yan, J., Sharo, A.G., Stone, H.A., Wingreen, N.S., and Bassler, B.L. (2016). *Vibrio cholerae* biofilm growth program and architecture revealed by single-cell live imaging. *Proceedings of the National Academy of Sciences*, 201611494.
- 295 16. Madsen, J.S., Lin, Y.-C., Squyres, G.R., Price-Whelan, A., de Santiago Torio, A., Song, A., Cornell, W.C., Sørensen, S.J., Xavier, J.B., and Dietrich, L.E. (2015). Facultative control of matrix production optimizes competitive fitness in *Pseudomonas aeruginosa* PA14 biofilm models. *Appl Environ Microb* **81**, 8414-8426.
- 300 17. Mann, E.E., and Wozniak, D.J. (2012). *Pseudomonas* biofilm matrix composition and niche biology. *Fems Microbiol Rev* **36**, 893-916.
18. Yawata, Y., Cordero, O.X., Menolascina, F., Hehemann, J.-H., Polz, M.F., and Stocker, R. (2014). Competition-dispersal tradeoff ecologically differentiates recently speciated marine bacterioplankton populations. *Proc Natl Acad Sci U S A* **111**.
- 305 19. Chowdhury, G., Bhadra, R.K., Bag, S., Pazhani, G.P., Das, B., Basu, P., Nagamani, K., Nandy, R.K., Mukhopadhyay, A.K., and Ramamurthy, T. (2016). Rugose atypical *Vibrio cholerae* O1 El Tor responsible for 2009 cholera outbreak in India. *J Med Microbiol*.
20. Green, S.K., Schroth, M.N., Cho, J.J., Kominos, S.D., and Vitanza-Jack, V.B. (1974). Agricultural plants and soil as a reservoir for *Pseudomonas aeruginosa*. *Applied microbiology* **28**, 987-991.
- 310 21. Fazli, M., Bjarnsholt, T., Kirketerp-Møller, K., Jørgensen, B., Andersen, A.S., Kroghfelt, K.A., Givskov, M., and Tolker-Nielsen, T. (2009). Nonrandom Distribution of *Pseudomonas aeruginosa* and *Staphylococcus aureus* in Chronic Wounds. *Journal of Clinical Microbiology* **47**, 4084-4089.
- 315 22. Burmølle, M., Thomsen, T.R., Fazli, M., Dige, I., Christensen, L., Homøe, P., Tvede, M., Nyvad, B., Tolker-Nielsen, T., Givskov, M., et al. (2010). Biofilms in chronic infections - a matter of opportunity - monospecies biofilms in multispecies infections. *Fems Immunol Med Mic* **59**, 324-336.
23. Guaglianone, E., Cardines, R., Vuotto, C., Di Rosa, R., Babini, V., Mastrantonio, P., and Donelli, G. (2010). Microbial biofilms associated with biliary stent clogging. *FEMS Immunology & Medical Microbiology* **59**, 410-420.
- 320 24. Harmsen, M., Yang, L., Pamp, S.J., and Tolker-Nielsen, T. (2010). An update on *Pseudomonas aeruginosa* biofilm formation, tolerance, and dispersal. *Fems Immunol Med Mic* **59**, 253-268.

- 325 25. Ciofu, O., Hansen, C.R., and Hoiby, N. (2013). Respiratory bacterial infections in cystic
fibrosis. *Curr Opin Pulm Med* 19, 251-258.
26. Folkesson, A., Jelsbak, L., Yang, L., Johansen, H.K., Ciofu, O., Hoiby, N., and Molin, S. (2012).
Adaptation of *Pseudomonas aeruginosa* to the cystic fibrosis airway: an evolutionary
perspective. *Nat Rev Microbiol* 10, 841-851.
- 330 27. Friedman, L., and Kolter, R. (2004). Genes involved in matrix formation in *Pseudomonas*
aeruginosa PA14 biofilms. *Mol Microbiol* 51, 675-690.
28. Miller, J.K., Badawy, H.T., Clemons, C., Kreider, K.L., Wilber, P., Milsted, A., and Young, G.
(2012). Development of the *Pseudomonas aeruginosa* mushroom morphology and cavity
formation by iron-starvation: A mathematical modeling study. *J. Theo. Biol.* 308, 68-78.
- 335 29. Parsek, M.R., and Tolker-Nielsen, T. (2008). Pattern formation in *Pseudomonas aeruginosa*
biofilms. *Current Opinion in Microbiology* 11, 560-566.
30. Persat, A., Nadell, C D., Kim, M K., Ingremeau, F., Siryaporn, A., Drescher, K., Wingreen, N S.,
Bassler, B L., Gitai, Z., and Stone, H A. (2015). The Mechanical World of Bacteria. *Cell* 161,
988-997.
- 340 31. Kim, M.Y., Drescher, K., Bassler, B.L., and Stone, H.A. (2014). Rapid Formation and Flow
Around *Staphylococcus Aureus* Biofilm Streamers. *Biophysical Journal* 106, 422A-422A.
32. Kim, M.K., Drescher, K., Pak, O.S., Bassler, B.L., and Stone, H.A. (2014). Filaments in curved
streamlines: rapid formation of *Staphylococcus aureus* biofilm streamers. *New Journal of*
Physics 16.
- 345 33. Rusconi, R., Lecuyer, S., Guglielmini, L., and Stone, H.A. (2010). Laminar flow around
corners triggers the formation of biofilm streamers. *Journal of The Royal Society Interface*,
rsif20100096.
34. Drescher, K., Shen, Y., Bassler, B.L., and Stone, H.A. (2013). Biofilm streamers cause
catastrophic disruption of flow with consequences for environmental and medical
350 systems. *Proceedings of the National Academy of Sciences* 110, 4345-4350.
35. Stewart, P.S. (2012). Mini-review: Convection around biofilms. *Biofouling* 28, 187-198.
36. Chew, S.C., Kundukad, B., Seviour, T., van der Maarel, J.R.C., Yang, L., Rice, S.A., Doyle, P.,
and Kjelleberg, S. (2014). Dynamic Remodeling of Microbial Biofilms by Functionally
Distinct Exopolysaccharides. *Mbio* 5.
- 355 37. Jennings, L.K., Storek, K.M., Ledvina, H.E., Coulon, C., Marmont, L.S., Sadovskaya, I., Secor,
P.R., Tseng, B.S., Scian, M., and Filloux, A. (2015). Pel is a cationic exopolysaccharide that
cross-links extracellular DNA in the *Pseudomonas aeruginosa* biofilm matrix. *Proceedings*
of the National Academy of Sciences 112, 11353-11358.
- 360 38. Franklin, M.J., Nivens, D.E., Weadge, J.T., and Howell, P.L. (2011). Biosynthesis of the
Pseudomonas aeruginosa extracellular polysaccharides, alginate, Pel, and Psl. *Frontiers in*
microbiology 2.
39. Colvin, K.M., Irie, Y., Tart, C.S., Urbano, R., Whitney, J.C., Ryder, C., Howell, P.L., Wozniak, D.J.,
and Parsek, M.R. (2012). The Pel and Psl polysaccharides provide *Pseudomonas*
aeruginosa structural redundancy within the biofilm matrix. *Environ Microbiol* 14, 1913-
365 1928.
40. Colvin, K.M., Gordon, V.D., Murakami, K., Borlee, B.R., Wozniak, D.J., Wong, G.C.L., and
Parsek, M.R. (2011). The Pel Polysaccharide Can Serve a Structural and Protective Role in
the Biofilm Matrix of *Pseudomonas aeruginosa*. *PLoS Pathog* 7, e1001264.
41. Das, K., and Mukherjee, A.K. (2007). Crude petroleum-oil biodegradation efficiency of
370 *Bacillus subtilis* and *Pseudomonas aeruginosa* strains isolated from a petroleum-oil
contaminated soil from North-East India. *Bioresource technology* 98, 1339-1345.

42. Stover, C.K., Pham, X.Q., Erwin, A., Mizoguchi, S., Warrenner, P., Hickey, M., Brinkman, F., Hufnagle, W., Kowalik, D., and Lagrou, M. (2000). Complete genome sequence of *Pseudomonas aeruginosa* PAO1, an opportunistic pathogen. *Nature* 406, 959-964.
- 375 43. Vos, M., Wolf, A.B., Jennings, S.J., and Kowalchuk, G.A. (2013). Micro-scale determinants of bacterial diversity in soil. *Fems Microbiol Rev* 37, 936-954.
44. Lockery, S.R., Lawton, K.J., Doll, J.C., Faumont, S., Coulthard, S.M., Thiele, T.R., Chronis, N., McCormick, K.E., Goodman, M.B., and Pruitt, B.L. (2008). Artificial dirt: microfluidic substrates for nematode neurobiology and behavior. *Journal of neurophysiology* 99, 3136-3143.
- 380 45. Heath, R.C. (1983). Basic ground-water hydrology, Volume 2220, (US Geological Survey).
46. Chuang, J.S., Rivoire, O., and Leibler, S. (2009). Simpson's paradox in a synthetic microbial system. *Science* 323, 272-275.
47. Sanchez, A., and Gore, J. (2013). Feedback between population and evolutionary dynamics determines the fate of social microbial populations. *PLoS Biol* 11, e1001547.
- 385 48. Drescher, K., Nadell, C., Stone, H., Wingreen, N., and Bassler, B. (2014). Solutions to the Public Goods Dilemma in Bacterial Biofilms. *Curr Biol* 24, 50-55.
49. Bottero, S., Storck, T., Heimovaara, T.J., van Loosdrecht, M.C.M., Enzien, M.V., and Picioreanu, C. (2013). Biofilm development and the dynamics of preferential flow paths in porous media. *Biofouling* 29, 1069-1086.
- 390 50. Choi, K.-H., and Schweizer, H.P. (2006). mini-Tn7 insertion in bacteria with single attTn7 sites: example *Pseudomonas aeruginosa*. *Nature protocols* 1, 153-161.
51. Lesic, B., and Rahme, L.G. (2008). Use of the lambda Red recombinase system to rapidly generate mutants in *Pseudomonas aeruginosa*. *BMC molecular biology* 9, 20.
- 395 52. Sia, S.K., and Whitesides, G.M. (2003). Microfluidic devices fabricated in poly (dimethylsiloxane) for biological studies. *Electrophoresis* 24, 3563-3576.

Figures & Legends




Figure 1. Wild type *P. aeruginosa* PA14 outcompetes the $\Delta pelA$ mutant under simple flow conditions, but the two strains coexist under complex flow conditions. (A) Wild type and $\Delta pelA$ strains were co-cultured at a range of initial frequencies in the simple straight-tunnel (black data) or soil-mimicking (blue data) microfluidic chambers. In both cases, fresh minimal M9 medium with 0.5% glucose was introduced at flow rates adjusted to the cross-sectional area of each chamber type to equalize the average initial flow velocity in both environments ($\sim 180 \mu m/s$). The diagonal gray line denotes the maximum possible increase in wild type frequency for a given initial condition. Each data point is an independent biological replicate. (B) A maximum intensity projection (top-down view) of a confocal z-stack of wild type (green) and $\Delta pelA$ (red) biofilms in simple flow chambers. (C) An epifluorescence micrograph (top-down view) of wild type (green) and $\Delta pelA$ (red) biofilms after 72 h growth in a flow chamber containing column obstacles to simulate a porous environment with irregular flows. Images in (B) and (C) were taken from chambers in which the wild type was inoculated at a frequency of 0.7, indicated by the arrow in panel (A). In (B) and (C), the global flow direction is indicated by the blue arrows.





Figure 2. Matrix production confers a competitive advantage to wild type *P. aeruginosa* PA14 in biofilms under simple flow conditions. (A) Absolute abundances of wild type and $\Delta pelA$ strains in monoculture and co-culture in simple flow microfluidic chambers (Error bars denote means \pm S.E. for $n = 4-6$). The two strains were inoculated alone (left two bars) or together at a 1:1 ratio (right two bars). Flow speed was $\sim 180 \mu m/s$. (B) Single optical plane $3 \mu m$ from the surface, and yz - and xz -projections at right and bottom respectively, of the wild-type and $\Delta pelA$ strains growing in co-culture after inoculation at a 1:1 ratio. (C) Relative abundance of wild type and $\Delta pelA$ strains in the liquid effluent of straight-chamber microfluidic devices. Wild type and $\Delta pelA$ cells were combined at a 1:1 initial ratio and co-inoculated on the glass substratum of simple flow chambers. At 0, 24, and 48 h, 5 μL of effluent was collected from chamber outlets and plated on LB agar plates in serial dilution to obtain colony counts.



435 **Figure 3.** Pel-deficient mutants occupy locations protected from flow due to local clogging by wild
type *P. aeruginosa* biofilms. (A) Wild type (green) and $\Delta pelA$ (red) *P. aeruginosa* strain mixtures were
inoculated into complex flow chambers with irregularly-spaced column obstacles. Biofilms were
imaged using confocal microscopy, after which fluorescent beads were flowed through the chamber.
The presence or absence of flow was monitored through averaging successive exposures of bead tracks
440 (white lines are bead tracks; blue arrows highlight flow trajectories). (B) Analysis of co-occurrence of
flow and wild type or $\Delta pelA$ cell growth at the end of 1:1 competition experiments in complex flow
chambers with column obstacles, as illustrated by the micrograph in (A). The occurrence of wild type
(gray) and $\Delta pelA$ (white) cell clusters are shown as a function of whether local flow has been blocked
or remains open after 72 h of competition (Bars denote means \pm S.E. for $n = 3$). (C) Biofilm growth
445 of wild type *P. aeruginosa* PA14 (gray) and the $\Delta pelA$ mutant (white) in monoculture in flat, straight-
tunnel flow chambers under different shear stress exposure treatments (Bars denote means \pm S.E. for
 $n = 4$).

Supplemental Information

Flow environment and matrix structure interact to determine spatial competition in *Pseudomonas aeruginosa* bacterial biofilms

Carey D. Nadell^{1*}, Deirdre Ricaurte^{2*}, Jing Yan², Knut Drescher¹, and Bonnie L. Bassler^{2,3†}

¹ Max Planck Institute for Terrestrial Microbiology, Marburg, HE, Germany

² Department of Molecular Biology, Princeton University, Princeton, NJ, USA

³ Howard Hughes Medical Institute, Chevy Chase, MD, USA

* Co-first authorship

† Correspondence to: bbassler@princeton.edu

Supplemental Materials and Methods





Figure S1. The maximum growth rates of *P. aeruginosa* PA14 wild type and $\Delta pelA$ cells in mixed liquid culture (M9 + 0.5% glucose). $n = 3$ biological x 10 technical replicates. Growth rates were measured as the maximum slope of growth curves measured using cultures inoculated into 96-well plates and measured for optical density at 600nm every 20 minutes until stationary phase.

26



27

28

29 **Figure S2.** Streamer structures produced by wild type *P. aeruginosa* PA14 (green) in microfluidic
30 chambers with complex flow profiles do not capture large numbers of co-cultured $\Delta pelA$ mutants
31 (red) over 72 h of biofilm growth (black circles are column obstacles).




Figure S3. Analysis procedure for correlating local flow and accumulation of wild type and $\Delta pelA$ cells (only $\Delta pelA$ cells are shown in this figure for simplicity). (A) First, the positions of the columns were identified from the fluorescence image. The centers of the columns served as nodes to divide the spaces into triangular regions. Within each region, the area covered by the column was removed. (B) Averaged intensity within each sampling region was calculated, and a threshold was set to determine if biofilm accumulation occurred in each region. White triangles correspond to regions identified as containing biofilms using this method. (C) The corresponding image with fluorescent beads (green) to track local flows. Typically, 10 to 15 images were taken and integrated to cover the flow regions sampled by the beads. Overlaid are triangular regions showing $\Delta pelA$ cell accumulation (from (B)). $\Delta pelA$ cells accumulated primarily in regions lacking flow due to upstream clogging (bottom) or in regions immediately downstream of obstructed areas, indicated by accumulation of non-moving beads (left, (C)).