

ARTICLE - DISCOVERY

Experimental evidence that thermal selection has shaped the latitudinal distribution of mitochondrial haplotypes in Australian fruit flies

5

M. Florencia Camus^{1,2*}, Jonci N. Wolff¹, Carla M. Sgrò¹, and Damian K. Dowling^{1*}

10 ¹ School of Biological Sciences, Monash University, Clayton, Victoria, 3800,
Australia

² Department of Genetics, Evolution & Environment, University College London, London
WC1E 6BT, UK

15

* To whom correspondence should be addressed.

Email: f.camus@ucl.ac.uk, damian.dowling@monash.edu

20

Abstract

Cellular metabolism is regulated by enzyme complexes within the mitochondrion, the function of which are sensitive to the prevailing temperature. Such thermal sensitivity, coupled with the observation that population frequencies of mitochondrial haplotypes tend to associate with latitude, altitude or climatic regions across species distributions, led to the hypothesis that thermal selection has played a role in shaping standing variation in the mitochondrial DNA (mtDNA) sequence. This hypothesis, however, remains controversial, and requires evidence the distribution of haplotypes observed in nature corresponds with the capacity of these haplotypes to confer differences in thermal tolerance. Specifically, haplotypes predominating in tropical climates are predicted to encode increased tolerance to heat stress, but decreased tolerance to cold stress, than temperate counterparts. We present direct evidence for these predictions, using mtDNA haplotypes sampled from the Australian distribution of *Drosophila melanogaster*. We show that the ability of flies to tolerate extreme thermal challenges is affected by sequence variation across mtDNA haplotypes, and that the thermal performance associated with each haplotype corresponds with its latitudinal prevalence. The haplotype that predominates at low (subtropical) latitudes confers greater resilience to heat stress, but lower resilience to cold stress, than counterparts predominating at higher (temperate) latitudes. We explore molecular mechanisms that might underlie these responses, presenting evidence that the effects are in part regulated by SNPs that do not change the protein sequence. Our findings indicate that standing genetic variation in the mitochondrial genome can be shaped by thermal selection, and could therefore contribute to evolutionary adaptation under climatic stress.

Keywords: Mitochondrial DNA; evolution; ecology; thermal adaptation; natural selection;

genome evolution

Introduction

The mitochondria are essential for eukaryote evolution, taking centre-stage in the process of cellular respiration. This process is regulated via a series of finely coordinated interactions between the genes of two obligate genomes – nuclear and mitochondrial (Rand, et al. 2004; Wolff, et al. 2014). Indeed, because of the strong dependence of cellular respiration on mitochondrial-encoded gene products, biologists traditionally assumed that strong purifying selection would efficiently prevent “function-encoding” genetic variation from accumulating within the mitochondrial DNA (mtDNA) sequence (Ballard and Kreitman 1994; Rand 2001; Dowling, et al. 2008). The assumption of selective neutrality has, however, been challenged over the past decade via analyses of mutational patterns within the mtDNA sequences of metazoans, which have uncovered the signature of recurrent adaptive evolution (Bazin, et al. 2006; James, et al. 2016). These analyses have been complemented by studies using experimental approaches with the power to partition mitochondrial from nuclear genetic effects, which have demonstrated that the intra-specific genetic variation that exists within the mitochondrial genome commonly affects the expression of core phenotypic traits, from morphological, to metabolic, to life-history (Rand 2001; Dowling, et al. 2008; Burton, et al. 2013; Dobler, et al. 2014).

Indeed, several lines of empirical evidence have emerged that support a novel hypothesis that posits that the standing genetic variation that delineates the mtDNA haplotypes of spatially-disjunct populations has been shaped by natural selection imposed by the prevailing thermal climate (Mishmar, et al. 2003; Ballard and Whitlock 2004; Ruiz-Pesini, et al. 2004; Wallace 2007; Dowling 2014). The first support for this *mitochondrial climatic adaptation* hypothesis was provided by studies of mtDNA variation in humans, where patterns of amino acid variation were observed to align closely to particular climatic regions (Mishmar, et al. 2003;

Ruiz-Pesini, et al. 2004), and where levels of genetic divergence between mtDNA haplotypes of different populations were shown to correlate with temperature differences between these populations (Balloux, et al. 2009). These studies on human mtDNA sequences are intriguing, but have in some cases proven difficult to replicate with different or larger datasets (Kivisild, 75 et al. 2006; Sun, et al. 2007).

Additional support for the hypothesis has been provided from studies of other metazoans, some of which have reported signatures of positive selection on mtDNA genes sampled from populations inhabiting particular thermal environments (Foote, et al. 2011; Silva, et al. 2014; 80 Ma, et al. 2015; Morales, et al. 2015), and others which have documented variation in mitochondrial gene or haplotype frequencies along clinal gradients associated with climate, such as latitude (Silva, et al. 2014; Consuegra, et al. 2015), altitude (Fontanillas, et al. 2005; Cheviron and Brumfield 2009), or with temperature itself (Quintela, et al. 2014). Inferences from these studies, however, are based only on correlations between haplotype frequencies 85 and environmental gradients, and could plausibly be explained by neutral demographic processes, such as by multiple colonisations from different origins into different locations followed by admixture, sex-specific patterns of dispersal and introgression (given that the mtDNA sequence is maternally-inherited) and by recurrent occurrences of secondary contact (Endler 1977; Toews and Brelsford 2012; Adrion, et al. 2015; Bergland, et al. 2016).

90

Finally, support has been provided through laboratory experiments in invertebrates, which have reported that the expression of life-history phenotypes (Dowling, et al. 2007; Arnqvist, et al. 2010; Hoekstra, et al. 2013; Wolff, et al. 2016), as well as the transmission dynamics (Matsuura, et al. 1997; Doi, et al. 1999), associated with particular mtDNA haplotypes, or

95 combinations of mitochondrial and nuclear (mito-nuclear) genotype, often depend on the thermal environment in which the study subjects are assayed . These studies have thus indicated that mitochondrial genetic variation is sensitive to thermal selection, at least when measured in the laboratory. However, these studies likewise have caveats. Invariably, the mtDNA haplotypes used were either sourced from disjunct global populations (Dowling, et
100 al. 2007; Arnqvist, et al. 2010; Wolff, et al. 2016), or alternatively levels of mtDNA divergence were artificially inflated by creating strains in which mtDNA haplotypes of one species were pitted against mtDNA haplotypes of a congener species, inside the nuclear backgrounds of one of the two species (Matsuura, et al. 1997; Doi, et al. 1999; Hoekstra, et al. 2013). Thus, inferences from these studies have been drawn from a ‘between population’
105 or ‘between species’ level, but such results are difficult to interpret within the ecological context of thermal adaptation, in which natural selection will act on standing variation in mtDNA haplotypes within a population of a given species.

Accordingly, in this study we sought to directly test the *mitochondrial climatic adaptation*
110 hypothesis, within the Australian distribution of the fruit fly, *Drosophila melanogaster*. This species invaded Australia over a century ago (Hoffmann and Weeks 2007), and it is thought the Australian population was established from multiple introductions of flies from two origins; Africa and Eurasia (David and Capi 1988; Singh and Long 1992). Indeed, a recent study of nuclear genome-wide allele frequencies from Australian populations concurs with
115 this conclusion, with flies sampled from high latitudes closely related to cold-adapted European populations, and those of low latitudes more closely related to African populations (Bergland, et al. 2016). These studies therefore provide a cautionary note, by indicating that colonisation history might well contribute to the existence of any latitudinal patterns in

mtDNA haplotype frequencies that occur within Australia, rather than thermal selection on
 120 the standing variation of mtDNA haplotypes (Adrion, et al. 2015; Bergland, et al. 2016).

Direct experimental evidence for the mitochondrial climatic adaptation hypothesis therefore
 requires a two-step approach; firstly evidence of shifts in the frequencies of mtDNA
 haplotypes along an gradient that aligns closely to the environment (e.g. latitude); and
 125 secondly experimental evidence that links thermal sensitivities of these haplotypes, when
 measured in the lab, to their latitudinal distributions in the field. This has never previously
 been achieved for the genetic variation that resides within the mitochondrial genome. Indeed,
 when it comes to the evolutionary significance of clinal variation in general, there are
 surprisingly few examples in which latitudinal variation in allele frequencies has been linked
 130 clearly to variation in fitness (Adrion, et al. 2015).

Results and Discussion

We first collected field-inseminated female flies from 11 populations along an eastern
 Australian latitudinal cline (Table S1), and used these flies to initiate isofemale lines (lines
 135 initiated by a solitary gravid female), and ultimately mass-bred populations per latitudinal
 location (with each population kept in independent duplicate). Previous research has shown
 strong linear associations between the expression of thermal tolerance phenotypes, and allele
 frequencies of underlying candidate nuclear genes, along this latitudinal cline (Hoffmann, et
 al. 2002; Weeks, et al. 2002; Hoffmann and Weeks 2007), uncovering signatures of thermal
 140 adaptation. To gauge levels of mtDNA sequence variation across these populations, we
 examined full mtDNA sequences across pooled samples of flies from each mass-bred
 population duplicate. Using this approach, we identified 15 SNPs in the mitochondrial
 genome, whilst the rest of the genome was highly conserved (Table 1). To probe levels of

haplotype diversity and estimate the frequencies of each haplotype within each of the source
145 populations, we designed a custom genotyping assay based on these 15 SNPs, and used this
assay to genotype the field-collected isofemale lines ($N = 312$). We identified a total of 10
unique haplotypes. All haplotypes fell into one of two main haplogroups, with a total of 12
SNPs delineating the two groups (Figure 1). Both haplogroups were found to segregate across
most of the 11 populations, but as a whole one haplogroup (haplogroup A) predominated in
150 the northern sub-tropical populations while the other (haplogroup B) predominated in
southern temperate populations (Figure S1). Furthermore, each of the A and B haplogroups
was dominated by one major haplotype (A1 accounting for 93.3% of A haplotypes; and B1
accounting for 77.2% of B haplotypes, Figure 1). The A1 haplotype appears more closely
related to other haplotypes of African origin. The ancestral origin of B1 haplotypes is,
155 however, less clear given they are most closely related to haplotypes other New World
populations, but also a haplotype from Japan (Figure 2). The frequency of the A1 haplotype
was negatively associated with the latitude of its source population ($R^2 = 0.4847$, $\beta = -$
 0.02881 , $p < 0.05$, Figure 3A), while the frequency of B1 exhibited a positive association (R^2
 $= 0.5137$, $\beta = 0.02718$ $p < 0.05$, Figure 3B). Thus, a latitudinal cline exists for the frequencies
160 of the A1 and B1 haplotypes along the east coast of Australia.

We next sought to experimentally assess whether the clinal associations of the A1 and B1
haplotypes are likely to have been shaped by thermal selection. To address this question, it
was necessary to disentangle effects attributable to mitochondrial genetic variation from
165 those caused by segregating nuclear allelic variation, or other sources of environmental
variance (Dowling, et al. 2008). We thus created eight genetic strains of flies, in which four
of the strains harboured the A1 haplotype, and the other four the B1 haplotype, in an

otherwise isogenic nuclear background derived from a distinct southern latitudinal population [Puerto Montt (PUE), Chile, South America]. We also ensured that all strains were free of *Wolbachia* infection, a maternally-inherited endosymbiotic bacterium, because variation in infection with different strains of *Wolbachia* would confound our capacity to map phenotypic effects to the mtDNA sequence. Furthermore, we created these eight strains such that each haplotype was replicated across two levels (intra- and inter-latitudinal replication per haplotype), which therefore enabled us to statistically partition effects attributable to the mitochondrial haplotype from those attributable to other sources of variation. Specifically, the haplotypes that sourced the strains were collected from each of two geographically-disjunct populations – Melbourne (37.99°S) and Brisbane (27.61°S). Because each mass-bred population was kept in independent duplicate, we ensured each duplicate contributed one A1 haplotype and one B1 haplotype to the strains (4 duplicates \times 2 haplotypes = 8 strains, thus creating replication within and between latitudes, Figure S2A & B).

Once the strains were created, full protein-coding resequencing of the mitochondrial genomes of each strain revealed that those harbouring the A1 haplotype were indeed all homogeneous; characterised by a single haplotype. The strains harbouring the B1 haplotype were, however, heterogeneous (Figure 1), and could be further partitioned into four unique “sub-haplotypes” (B-1, B-2, B-3, & B-4). Each B sub-haplotype was delineated by 1 to 4 SNPs, but all shared the same pool of 12 SNPs that delineate them from the A haplogroup (Figure 1, Table 2, Figure S3). This enabled us to partition mitochondrial genetic effects over two levels – at the level of the haplotype, and the sub-haplotype. The genetic variation differentiating the A1 and B1 haplotypes was comprised of 15 synonymous SNPs in the protein-coding genes. Synonymous SNPs have traditionally been considered to be functionally silent because they do not change the amino acid sequence. However, a growing body of empirical evidence

suggests that synonymous polymorphisms might routinely modify the phenotype and thus be of functional and evolutionary significance (Kimchi-Sarfaty, et al. 2007; Hurst 2011). On the other hand, the SNPs delineating the “sub-haplotypes” hubbed within the B1 haplotype consisted of a mixture of synonymous and non-synonymous SNPs (Table 2).

Flies harbouring the A1 haplotype, which predominates in the sub-tropics exhibited greater tolerance to an extreme heat challenge than flies harbouring the B1 haplotype (haplotype, $\chi^2=6.04$, $p = 0.014$, Table S2), but the magnitude of these effects changed across the sexes (haplotype \times sex, $\chi^2=24.7$, $p < 0.001$, Figure 4A-B, Table S2). We also uncovered sex-specific effects that mapped specifically to the level of the mtDNA sub-haplotype (sex \times sub-haplotype[haplotype], $\chi^2=25.04$, $p = <0.001$, Figure 4C). This interaction was primarily attributable to the B1-D sub-haplotype, which conferred inferior heat tolerance in males, but high heat tolerance in females, relative to the other sub-haplotypes. Only one synonymous SNP, located in the *mt:ND4* gene, delineates the protein-coding region of this sub-haplotype from the other B1 sub-haplotypes (Table 2). This polymorphism is therefore a candidate SNP in conferring sex-specific outcomes in heat tolerance, although we cannot rule out the possibility that further variation within the non-coding region of the mtDNA sequence and regulatory elements (which we did not sequence) contributed to this effect. Nonetheless, the observed pattern associated with this sub-haplotype is striking in the context of a hypothesis proposed by Frank and Hurst (1996), often called *Mother's Curse*, which proposes that maternal inheritance of the mitochondria will facilitate the accumulation of mtDNA mutations that are deleterious to males, but benign or only slightly deleterious to females (Frank and Hurst 1996; Gemmell, et al. 2004; Beekman, et al. 2014). However, while this haplotype harbours variation that causes a male-limited reduction in heat tolerance, it did not however confer a detrimental effect on male capacity to tolerate cold stress (Table S3, Figure

4D). Thus, further studies are required to determine whether the male-specificity of the B1-D sub-haplotype on heat tolerance effect extends to pleiotropic effects on other key life-history traits such as reproduction and longevity (Camus, et al. 2015), or whether this effect is sensitive to genotype-by-environment interactions that mediate the severity of effect in males (Mossman, et al. 2016; Wolff, et al. 2016).

Flies harbouring the B1 haplotype were superior at withstanding an extreme cold challenge, relative to their A1 counterparts ($\chi^2=34.31$, $p < 0.001$, Figure 4D-E, Table S3), but there was no significant effect traceable to the level of the sub-haplotype (Table S3). Importantly, the effects of mitochondrial haplotype on both thermal tolerance phenotypes was robust to the source of origin of the haplotypes (*i.e.*, whether they were sourced from Brisbane or Melbourne), providing clear evidence that the phenotypic effects are directly tied to the mtDNA sequence (Figure 4). Furthermore, all B1 sub-haplotypes exhibited decreased heat tolerance and increased cold tolerance when compared to the A1 haplotype, providing support for the contention that the differences in thermal tolerance observed between northern and southern haplogroups are mapped to the 15 SNPs that delineate the A1 and B1 haplotypes (or to cryptic variation in the non-coding region that we did not genotype), rather than the 4 SNPs that delineate the different sub-haplotypes hubbed within B1 (Table S2, Table S3, and Figure 4F). Alternatively, it is possible that the unique SNPs that delineate the sub-haplotypes drive the bulk of the differences in thermal response between the A1 and B1 haplotypes, and represent a case of parallel evolution for thermal tolerance brought about by different underlying SNPs (Arendt and Reznick 2008). While we cannot definitively disentangle these possibilities, we note that the polymorphisms that delineate the A1 from the B1 sub-haplotypes only include non-synonymous SNPs in two of four cases. Thus, while we are unable to definitively ascertain whether differences in the A1 and B1 thermal responses

are underpinned primarily by the 15 shared SNPs that separate all B1 from A1 haplotypes, by other cryptic regulatory variation in the non-coding region, or by the unique SNPs that delineate the B1 sub-haplotypes, our results suggest that SNPs that do not change the amino acid sequence are likely to be responsible for this thermal divergence in at least two cases.

We then examined whether the thermal tolerance phenotypes might be mediated by patterns of differential gene expression of protein-coding mtDNA genes, copy number variation in mtDNA, or codon usage bias across the A1 and B1 haplotypes. For the gene expression analyses, we examined five genes involved in complex I and complex IV of the electron transport chain (complex I: *mt:ND4*, *mt:ND5*, complex IV: *mt:COXI*, *mt:COXII*, *mt:COXIII*). Emerging evidence suggests that genetic variation within complex I genes (both mitochondrial and nuclear) might contribute disproportionately to trajectories of mitonuclear, and ultimately, life history evolution (Camus, et al. 2015; Garvin, et al. 2015; Morales, et al. 2015). Complex IV, on the other hand, harbours genes with the lowest levels of *dN/dS*, indicative of greater selective constraints on these mitochondrial genes across taxonomically diverse organisms (Nabholz, et al. 2013). Accordingly, we found that strains harbouring the B1 haplotype exhibited higher gene expression for the complex I genes *mt:ND4* and *mt:ND5*, which belong to the same transcriptional unit (Torres, et al. 2009), than strains with the A1 haplotype (haplotype \times gene < 0.001 , Figure 5, Table S4). All of the SNPs located in the *mt:ND4* and *mt:ND5* genes, which delineate northern from southern haplotypes, are synonymous. This observation is interesting in light of a recent report that found that patterns of expression of *mt:ND5* and *mt:CYTB* genes in *D. melanogaster* mapped to candidate SNPs that lay directly within the affected genes, and which presumably exerted their effects via post-transcriptional modification of RNA, potentially altering the stability of the transcripts

(Camus, et al. 2015). We did not, however, detect differences in mtDNA copy number between A1 and B1 haplotypes (Table S5).

270 Evidence is also mounting that variation in patterns of genomic DNA base composition (GC content (Šmarda, et al. 2014), as well as variation in codon usage bias across DNA sequences (Sharp, et al. 1995) can be shaped under natural selection. For example, in bacteria and metazoans, higher levels of GC base pairs have been associated with the thermal environment, with the GC base pair associated with higher thermal stability (Bernardi 2007).

275 In bacteria, this correlates with greater tolerance of higher temperatures (Musto, et al. 2004). In the green alga *Chlamydomonas*, experimental alteration of mitochondrial codons drastically changes translational efficiency, suggesting that mitochondrial codon usage has been optimised for translation of mitochondrial products (Salinas, et al. 2012). In our study, the A1 and B1 haplotypes differ by 15 synonymous SNPs that are evenly distributed across

280 the mitochondrial protein-coding region, with most protein-coding genes harbouring at least one SNP site. SNPs of the A1 haplotype show a high GC bias, with 80% of the SNPs represented by a guanine or cytosine, and conversely those of B1 reveal a GC content of only 20% (Table S6A, Fishers exact test, $p = 0.001$). Thus, the A1 haplotype, which confers higher tolerance to an extreme heat challenge, has a higher GC content; concordant with previous

285 observations in bacteria and metazoans suggesting higher thermal stability of the GC base pair relative to AT (Bernardi 2007). Additionally, the SNPs delineating the A1 haplotype change the codon bias and produce rarer codons (Table S6B, Fishers exact test $p = 0.002$). These findings suggest that GC content and codon bias may play a role in the observed haplotype effects on gene expression of *mt:ND4* and *mt:ND5*, with ultimate upstream effects

290 on thermal tolerance.

By harnessing an experimental genomic approach applied to the mitochondrial genome, within a robust ecological framework, we have documented clinal patterns in standing mtDNA haplotypes, and provided experimental evidence that these patterns are linked to the capacity of these haplotypes to tolerate thermal stress. While these results suggest the documented clinal patterns of mtDNA variation are likely to have been shaped at least in part by thermal selection, it is difficult to fully resolve the relative influence of thermal selection from history of colonisation and other demographic factors, given that the Australian east coast is thought to have been subjected to recurrent colonisation events from flies of two disparate origins, Africa and Eurasia, over the past 150 years. This, of course, is a caveat that is not unique to our study on mtDNA variation, but extends to all studies of clinal variation of New World populations. *D. melanogaster* is thought to have expanded out of equatorial Africa and into Eurasia around 15 000 years ago (Li and Stephan 2006). The Australian population consists of flies harbouring two haplotypes; the A1 haplotype appears more related to African than European mtDNA sequences, while the ancestral origin of the B1 haplotype is less clear, but it aligns most closely to other Asian haplotypes and haplotypes from New World populations (Figure 2). We thus note there is a possibility that these haplotypes might have been pre-adapted to tropical and temperate conditions prior to their introductions into Australia, and that the relevant mitochondrial variation under selection along the Australian cline is likely to have already existed upon the arrival of these haplotypes into Australia. Accordingly, the Australian distribution of mtDNA haplotypes is likely to have been shaped both by the history of colonization, followed by the subsequent and ongoing action of thermal selection in spatially sorting the haplotypes along the latitudinal cline. Our study suggests that further research into the mitochondrial climatic adaptation hypothesis is warranted. In particular, we suggest our conclusions can be tested by future studies that utilise other established latitudinal clines, in *D. melanogaster* and in other

species, to determine whether mtDNA haplotypes show similar associations to latitude as revealed in the Australian cline, and to determine whether the mtDNA haplotypes involved exhibit thermal sensitivities that concord with the clinal patterns.

320

In conclusion, our results provide support for the hypothesis that standing genetic variation within the mitochondrial genome has been shaped by natural selection imposed by thermal stress. We also presented evidence that phenotypic responses to thermal stress have been shaped in part by SNPs that do not change the amino acid sequence. This thus suggests a role
325 for a set of SNPs that were traditionally thought to evolve under neutrality (Kimchi-Sarfaty, et al. 2007; Hurst 2011), within a genome that was likewise traditionally thought to be devoid of phenotype-changing genetic variation (Ballard and Rand 2005; Dowling, et al. 2008), in the dynamics of thermal adaptation. Secondly, and more broadly, our results add to an emerging body of research in *Drosophila* (Sorensen, et al. 2007; Chen, et al. 2012; Lavington, et al. 2014; Cogni, et al. 2015), and other metazoans (Porcelli, et al. 2015), which
330 highlights metabolic genes (including those targeted to the mitochondria) act as important substrates on which thermal selection is likely to act to shape adaptive evolutionary responses. Several studies have now reported variation in allele frequencies, or expression patterns, of nuclear-encoded metabolic genes along latitudinal clines (Chen, et al. 2012; Lavington, et al. 2014; Cogni, et al. 2015), or across replicated laboratory populations that
335 have evolved under differing thermal regimes (Sorensen, et al. 2007), in *Drosophila melanogaster*. These studies, however, did not screen for involvement of the evolutionary-conserved mitochondrial genes. The function of key metabolic enzymes, however, relies on close coordination between mitochondrial and nuclear genomes (Rand, et al. 2004; Levin, et al. 2014; Wolff, et al. 2014; Quiros, et al. 2016). This point, when reconciled with the
340 emerging studies, would suggest that genetic interactions between the mitochondrial and

nuclear genomes could represent key mediators of evolutionary adaptation of the metabolic machinery under thermal stress.

Materials and Methods

Field Collection, Isofemale Line Establishment and Maintenance

Populations of *Drosophila melanogaster* were sampled from the east coast of Australia during March-April 2012 from 11 locations. The population names (latitude and longitude) are: Townsville (19.26,146.79), Rockhampton (23.15, 150.72), Brisbane (27.61, 153.30), Ballina (28.87,153.44), Coffs Harbour (30.23, 153.15), Port Macquarie (30.93, 152.90), Wollongong (34.34, 150.91), Narooma (36.25, 150.14), Gosford (33.31, 151.20), Bermagui (36.40, 150.06), Melbourne (37.99, 145.27). Samples were collected as close to sea level as possible to avoid altitudinal differences between the populations (Collinge, et al. 2006). Individual field-inseminated females were isolated into individual vials in the laboratory to initiate independent isofemale lines. At least twenty isofemale lines were generated for each population. Each line was treated with tetracycline to eliminate cytoplasmic endosymbionts, such as *Wolbachia* (Clancy and Hoffmann 1998), and tested using *Wolbachia*-specific primers. We further verified infection status when analysing next-generation sequencing data by seeing if any of our obtained reads mapped to the *Wolbachia* genome.

Three generations after the isofemale lines were established in the lab, one mass-bred population was created from the isofemale lines of each latitudinal location (11 locations). Specifically, the populations were established by combining 25 virgin males and 25 virgin females from each of 20 randomly-selected isofemale lines per latitudinal location. The following generation, each population was divided into two duplicates (11 populations \times 2 duplicates), which were kept separately from this point onward. A small sample of flies (~20-50 individuals) from each isofemale line was also collected at this time, and placed at -20°C for sequencing and genotyping. Mass-bred populations were kept at 25°C under a 12:12h

light:dark cycle. Genetic variation was maintained within each duplicate population by rearing flies across two bottles on potato-dextrose-agar food medium, with densities of approximately 300 flies per bottle. Every generation, newly-emerged flies from each duplicate were collected from both bottles and then randomly redistributed into two new fresh bottles.

Next Generation Sequencing and SNP Genotyping

To identify regions of variation between the 11 populations, we first used pooled samples of 100 individuals (both males and females) from each population and used next generation sequencing to obtain full mitochondrial genomes. DNA samples were enriched for mitochondrial DNA to obtain the best coverage possible. This process was achieved by using *Wizard SV* Miniprep Purification Kit (Promega, Madison, WI) for DNA extraction, which captures circular DNA. Enriched DNA samples were made into 200bp paired-end libraries and sequenced using the Illumina GAIIx platform (Micromon, Monash University, Australia). Reads were aligned to the *Drosophila melanogaster* mitochondrial reference genome (NCBI reference sequence: NC_001709.1) using *Geneious* (Kearse, et al. 2012), generating mitochondrial genomes for each of the 11 latitudinal locations.

To obtain allele frequencies from each population, variable sites obtained from the mass sequencing were used as markers. DNA from each isofemale line was extracted using the *GenEra Puregene Cell and Tissue Kit* (Qiagen, Hilden, Germany). Even though each mass-bred population was created using 20 randomly-chosen isofemale lines, we genotyped all isofemale lines collected from each latitudinal location. A custom SNP genotyping assay was

developed (Geneworks, Thebarton, Australia) for the 15 SNPs identified via mass sequencing, and genotyping was performed by Geneworks (Thebarton, Australia) on a SEQUENOM MassARRAY platform (Agena Bioscience, San Diego, CA). This genotyping
395 revealed the presence of northern-predominant (*i.e.*, predominating in northern latitude populations) and southern-predominant (*i.e.*, predominating in southern latitude populations) haplogroups, with each haplogroup characterised by one major haplotype (A1 and B1).

Creation of Mitochondrial Strains from Mass-bred Populations

400 We created “introgression strains” from each of the population duplicates (11 latitudes \times 2 population duplicates = 11 introgression strains \times 2 duplicates), by introgressing the pool of mtDNA variants of each population duplicate into a standard and isogenic nuclear background originally sourced from Puerto Montt (PUE), Chile (41.46°S, 72.93°W) (Calboli, et al. 2003), which had been created via 20 generations of full-sibling matings. To initiate
405 each strain, 100 virgin female flies were sampled from each population duplicate and crossed to 120 males from the PUE strain. Then, for 20 sequential generations, 100 daughters were collected per strain and backcrossed to 120 PUE males. This crossing scheme aimed to maintain the pool of segregating mitochondrial haplotypes within each population, while translocating them alongside that of an isogenic nuclear background, to enable partitioning of
410 mitochondrial genetic effects from cryptic variance tied to the nuclear genome (Figure S2A). In order to prevent mitochondrial contamination from the Puerto Montt (PUE) line, all lines were tested every 5 generations during the introgression regime, to ensure there were no instances of contamination of the lines (by rogue females of the PUE strain) by using qRT-PCR melt curve analysis that would detect PUE-specific mtDNA SNPs.

We then created a new set of isofemale lines from each of the introgression strain duplicates, and re-genotyped females of each line using the custom SNP genotyping assay described above (Geneworks, Thebarton, Australia). From the genotyping results, we were able to
420 identify female lineages that carried individual haplotypes (A1 [northern] or B1 [southern]), and using this information we then selected one isofemale line carrying the A1 haplotype and one isofemale line carrying the B1 haplotype, from each of the two independent population duplicates from two (Brisbane, Melbourne) of the 11 latitudinal locations (Figure S2B). We continued to backcross virgin females of each isofemale line to males of the isogenic PUE
425 line for a further seven generations. Prior to this step, the PUE line had been propagated via a protocol of mating between one full-sibling pair for five generations, to remove any genetic variation that had accumulated within this nuclear background during the course of the introgressions described above. We chose to use isofemale lines from Brisbane (latitude: 27.61°S) and Melbourne (latitude: 37.99°S) because they are geographically-disjunct, and
430 because re-genotyping confirmed that both A1 and B1 haplotypes were segregating in each of the introgression strain duplicates following the 20 generations of introgression. Following this process, each of the A1 and B1 haplotypes was represented across four independent genetic strains each, at two levels of replication; an intra-latitudinal (between the two population duplicates of a given latitude) and an inter-latitudinal (between two latitudes,
435 Brisbane and Melbourne) replicate (Figure S2B).

We then re-sequenced these strains, and obtained full complete mitochondrial genomes for all eight mitochondrial strains, again using the next generation sequencing approach described above. Resequencing results revealed that haplotype A1 was isogenic across all four A1

strains, while we found that the B1 strains could be delineated into four unique sub-haplotypes that were nested within the B1 haplotype. These four southern sub-haplotypes all shared the known SNPs that delineate the north and south haplogroups (and the A1 and B1 haplotypes), however they each carried between one and three additional SNPs (Table 2).

Extreme Heat Challenge

Tolerance to an extreme heat challenge was measured for 120 flies of each sex from each mitochondrial strain (Hoffmann, et al. 2002). Flies were placed in individual 5mL water-tight glass vials and subsequently exposed to a 39°C heat challenge, by immersion of the glass vials in a preheated circulating water bath. Heat “knock-down” time was recorded as the time taken for each individual fly to become immobilized (in a coma-like state) at 39°C (Williams, et al. 2012). This experiment was conducted over two trials within the same generation. Each trial of the experiment consisted of a fully-balanced replicate of the experimental units (*i.e.*, equal numbers of flies of each sex × mitochondrial strain), separated in time by 2 hours within the same day. The position of flies of each experimental unit was randomized within each trial of the experiment. The assay was conducted blind to the genotype or sex of the fly.

Extreme Cold Challenge

This assay measures the amount of time it takes a fly to regain consciousness and stand on all legs after succumbing to a cold-induced coma (Hoffmann, et al. 2002). In each trial of the assay, 40 flies from each experimental unit (N = 640) were placed individually in 1.7mL microtubes. These tubes were then submerged in a water bath set to 0°C (comprised of water and engine coolant) for 4 h, to place flies into coma. At 4 h, all microtubes were removed

from the bath, and laid out on a bench at 25°C, and the time taken (seconds) for each fly to regain consciousness and stand upright was recorded. The assay was conducted blind to the genotype or sex of the fly.

Statistical Analyses of Thermotolerance Data

We used separate multilevel linear mixed models to test the effects of mtDNA haplotype and sub-haplotype on responses to each of the heat and cold challenges. The response variable for the heat challenge assay was the time taken to fall into coma, while the response variable for the cold challenge assay was the time taken to wake from coma. Fixed effects were the identity of the mtDNA haplotype (A1, B1), the sub-haplotype nested within haplotype (A1, B1-A, B1-B, B1-C, B1-D), sex and their interactions. This analysis assumes the effect of the SNPs separating the A1 and the B1 haplotypes, and those that separate the B1 sub-haplotypes, are hierarchical and can be statistically partitioned (*i.e.*, any significant ‘haplotype’ effects in the model can be mapped to the 15 SNPs that separate the A1 and B1 haplotypes, while significant ‘sub-haplotype’ effects are mapped to the unique SNPs that separate the four B1 sub-haplotypes). We, however, acknowledge the alternative possibility that the unique SNPs separating B1 sub-haplotypes could in theory underpin the differences between the A1 and B1 thermal responses, if such SNPs have accumulated under parallel evolution (Arendt and Reznick 2008). Random effects described the biological structure of the mitochondrial strains; there were two tiers of replication – with each haplotype replicated across two “duplicates” within each of two latitudinal “populations”. Thus, duplicate nested within population was included as a random effect, as well as other known and random environmental sources of variance (the trial identity, and the identity of the person scoring the response variable [2 people]).

Parameter estimates were calculated using restricted maximum likelihood algorithm in the *lme4* package of R (Bates 2012). The fitted model was evaluated by simplifying a full model, by sequentially removing terms that did not change (at $\alpha = 0.05$) the deviance of the model, starting with the highest order interactions, using log-likelihood ratio tests to assess the change in deviance in the reduced model relative to the previous model (Fox 2002).

Haplotype Network, Divergence and Codon Bias

Relationships among haplotypes were visualized on a median-joining network (Bandelt, et al. 1999) and constructed in the software NETWORK version 4.6.1.2 (www.fluxus-engineering.com).

We obtained divergence estimates between A1 and B1 haplotypes using *Geneious* (Kearse, et al. 2012) and *MEGA6* (Tamura, et al. 2013). Using *Geneious*, divergence was calculated by examining the %identity function and subtracted that value from 100 to derive the percentage divergence. In *MEGA6*, we performed a pairwise distance comparison using a maximum composite likelihood model. Both methods gave concordant estimates of divergence (divergence = 0.001%)

We obtained *Drosophila melanogaster* mitochondrial codon usage bias values from the Codon Usage Database (Nakamura, et al. 2000). For both haplotypes, each SNP site was given the title “preferred” or “unpreferred” based on the codon usage bias score. Results were then analysed as a 2×2 contingency table using Fishers exact tests (Table S6A & B).

Total RNA/DNA Extraction and cDNA Synthesis

510 For RNA extractions, we placed single flies from each A1 and B1 strain into a microtube. We thus combined source population and duplicate into one sample. Each extraction was performed in triplicate, thus resulting in three microtubes with flies possessing the A1 haplotype and three microtubes with flies harbouring the B1 haplotype. In the case of the A1 haplotype all flies harboured the same haplotype (although originating from different rearing
515 vials), whereas for the B1 haplotype each biological replicate was formed by combining a single fly from each sub-haplogroup into a microtube.

We then performed a coupled RNA and DNA extraction as per the supplier's protocols using *TRIzol® Reagent* (Thermo Fisher Scientific, Waltham, MA) to first create a phase separation
520 of RNA and DNA from which the total RNA was then purified using a *HighPure RNA extraction kit* (Roche Applied Science, Penzberg, Germany). In this manner, both the DNA and RNA was independently separated and stored from the one sample. The separated nucleic acids (~100µL of each sample extracted) were quantified by Nanodrop UV/Vis spectrophotometry (Thermo Fisher Scientific, Waltham, MA) and *the* purity of total RNA
525 was confirmed using the A_{260}/A_{280} ratio with expected values between 1.8 and 2.0. The integrity of both the RNA and DNA was assessed by electrophoresis (1% TBE agarose gel).

The cDNA was synthesized from 1µg of RNA using the Transcriptor First Strand cDNA Synthesis Kit (Roche Applied Science, Penzberg, Germany) and a mixture of random
530 hexamers and oligodT primers to capture mitochondrial transcripts both in the transitory polycistronic stage and as individual polyadenylated single transcripts (Clayton 2000).

mtDNA Copy Number Quantification

mtDNA copy number was measured for each DNA extraction performed (see *Total*
 535 *RNA/DNA Extraction and cDNA Synthesis*) because it is known to differ across sexes (Yin, et al. 2004; Ballard, et al. 2007) and may be expected to covary with transcript abundance (Thomas 1992). mtDNA copy number was calculated relative to a single copy gene in the nuclear genome (Correa, et al. 2012). Copy number was determined using quantitative real-time PCR of a 113 bp region of the large ribosomal subunit (CR34094, FBgn0013686). No
 540 nuclear copies of this gene are found in the *Drosophila melanogaster* genome. Similarly, nuclear DNA was quantified by amplifying a 135 bp region of the single-copy (Aoyagi and Wassarman 2000) subunit of the RNA polymerase II gene (CG1554, FBgn0003277). The copy number was then determined as the relative abundance of the mtDNA to nuclear DNA ratio and thus reflects the average number of mtDNA copies per cell.

545

Gene expression quantification

Five of the thirteen total mitochondrial protein-coding genes were amplified to quantify gene expression levels. Quantified genes were: *mt:COI*, *mt:COII*, *mt:COIII*, *mt:ND4*, and *mt:ND5*. Gene expression of each biological replicate (three biological replicates per haplotype) was
 550 measured using quantitative real time (qRT)-PCR (Lightcycler 480 – Roche Applied Science, Penzberg, Germany). Reactions were performed in duplicate (technical duplicates) using a *SYBRGreen I Mastermix* (Roche Applied Science, Penzberg, Germany), whereby each well contained 5µl of SYBR buffer, 4µl of 2.5µM primer mix and 1µl of diluted cDNA. The

following amplification regime used was: 90°C (10s), 60°C (10s), 72°C (10s) for 45 cycles,

555 followed by a melt curve analysis to verify the specificity of the primer pair.

The *Bestkeeper*[®] software (Pfaffl, et al. 2004) was used to select nuclear housekeeping genes (HKGs) for quality assessment. Three suitable HKGs were chosen: succinate dehydrogenase A (CG17246), 14-3-3ε (CG31196), and an unknown protein-coding gene (CG7277). All
560 three genes had similar expression levels with high correlation coefficients (>0.8) against each other. For each experimental sample, the expression values of the mitochondrial target genes were standardized as follows:

The cycle threshold was calculated using the gene of interest (GOI) and the geometric mean
565 of the three housekeeping genes (GEOM):

$$\Delta Ct = Ct_{GOI} - Ct_{GEOM}$$

The cycle thresholds were then used to calculate the relative gene expression for each experimental sample in relation to the housekeeping genes.

$$\text{Relative gene expression} = 2^{-\Delta Ct}$$

570 Gene expression levels of all five mitochondrial genes were obtained by determining the ΔCt per sample, measured at the maximum acceleration of fluorescence, using the Second Derivative Maximum Method (Rasmussen 2001) in the *Lightcycler Software V1.5.0* (Roche Applied Science, Penzberg, Germany). When the ΔCt values between two technical duplicates for each sample fell within 0.5 units of each other, then the mean gene expression
575 estimates were pooled to form a single data point (Bustin, et al. 2009).

Statistical Analysis of Gene Expression Data and Copy Number Variation

We fitted linear models, in which mitochondrial copy number and gene expression data were modelled separately as response variables. Mitochondrial haplotype (A1, B1), and gene identity were modelled as fixed effects. Mitochondrial copy number values were added as a fixed covariate to the analysis of gene expression, and F statistics and associated probabilities estimated using a Type III sums-of-squares tests in the *car* package (Fox 2002) in *R* (R Development Core Team 2009). Mitochondrial copy number variation was modelled with haplotype (A1 and B1) as a factor.

References

- Adrion JR, Hahn MW, Cooper BS. 2015. Revisiting classic clines in *Drosophila melanogaster* in the age of genomics. Trends in Genetics 31:434-444.
- 590 Aoyagi N, Wassarman DA. 2000. Genes encoding *Drosophila melanogaster* RNA polymerase II general transcription factors: diversity in TFIIA and TFIID components contributes to gene-specific transcriptional regulation. Journal of Cell Biology 150:F45-50.
- Arendt J, Reznick D. 2008. Convergence and parallelism reconsidered: what have we learned about the genetics of adaptation? Trends in Ecology & Evolution 23:26-32.
- 595 Arnqvist G, Dowling DK, Eady P, Gay L, Tregenza T, Tuda M, Hosken DJ. 2010. Genetic Architecture of Metabolic Rate: Environment Specific Epistasis between Mitochondrial and Nuclear Genes in an Insect. Evolution 64:3354-3363.
- Ballard JW, Kreitman M. 1994. Unraveling selection in the mitochondrial genome of *Drosophila*. Genetics 138:757-772.
- 600 Ballard JW, Melvin RG, Miller JT, Katewa SD. 2007. Sex differences in survival and mitochondrial bioenergetics during aging in *Drosophila*. Aging Cell 6:699-708.
- Ballard JWO, Rand DM. 2005. The population biology of mitochondrial DNA and its phylogenetic implications. Annual Review of Ecology Evolution and Systematics 36:621-642.
- Ballard JWO, Whitlock MC. 2004. The incomplete natural history of mitochondria. Molecular Ecology 13:729-744.
- 605 Balloux F, Handley LJJ, Jombart T, Liu H, Manica A. 2009. Climate shaped the worldwide distribution of human mitochondrial DNA sequence variation. Proceedings of the Royal Society Biological Sciences Series B 276:3447-3455.
- Bandelt HJ, Forster P, Rohl A. 1999. Median-joining networks for inferring intraspecific phylogenies. Molecular Biology and Evolution 16:37-48.
- 610 Bates D, Maechler, M., Bolker, B. 2012. *lme4*: Linear mixed-effects models using Eigen and R syntax. R package version 0.999999-0. <http://cran.r-project.org/package=lme4>.
- Bazin E, Glémin S, Galtier N. 2006. Population Size Does Not Influence Mitochondrial Genetic Diversity in Animals. Science 312:570-572.
- 615 Beekman M, Dowling DK, Aanen DK. 2014. The costs of being male: are there sex-specific effects of uniparental mitochondrial inheritance? Philos Trans R Soc Lond B Biol Sci 369:20130440.
- Bergland AO, Tobler R, Gonzalez J, Schmidt P, Petrov D. 2016. Secondary contact and local adaptation contribute to genome-wide patterns of clinal variation in *Drosophila melanogaster*. Molecular Ecology 25:1157-1174.
- 620 Bernardi G. 2007. The neoselectionist theory of genome evolution. Proceedings of the National Academy of Sciences of the United States of America 104:8385-8390.
- Burton RS, Pereira RJ, Barreto FS. 2013. Cytonuclear Genomic Interactions and Hybrid Breakdown. Annual Review of Ecology, Evolution, and Systematics, Vol 44 44:281-302.
- Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, et al. 2009. The MIQE Guidelines: Minimum Information for Publication of Quantitative Real-Time PCR Experiments. Clinical Chemistry 55:611-622.
- 625 Calboli FC, Kennington WJ, Partridge L. 2003. QTL mapping reveals a striking coincidence in the positions of genomic regions associated with adaptive variation in body size in parallel clines of *Drosophila melanogaster* on different continents. Evolution 57:2653-2658.

- 630 Camus MF, Wolf JB, Morrow EH, Dowling DK. 2015. Single Nucleotides in the mtDNA Sequence Modify Mitochondrial Molecular Function and Are Associated with Sex-Specific Effects on Fertility and Aging. *Current Biology* 25:2717-2722.
- Chen Y, Lee SF, Blanc E, Reuter C, Wertheim B, Martinez-Diaz P, Hoffmann AA, Partridge L. 2012. Genome-wide transcription analysis of clinal genetic variation in *Drosophila*. *PLoS One* 7:13.
- 635 Cheviron ZA, Brumfield RT. 2009. Migration-Selection Balance and Local Adaptation of Mitochondrial Haplotypes in Rufous-Collared Sparrows (*Zonotrichia Capensis*) Along an Elevational Gradient. *Evolution* 63:1593-1605.
- Clancy DJ, Hoffmann AA. 1998. Environmental effects on cytoplasmic incompatibility and bacterial load in *Wolbachia*-infected *Drosophila simulans*. *Entomologia Experimentalis et Applicata* 86:13-24.
- 640 Clayton DA. 2000. Transcription and replication of mitochondrial DNA. *Human Reproduction* 15 Suppl 2:11-17.
- Cogni R, Kuczynski K, Lavington E, Koury S, Behrman EL, O'Brien KR, Schmidt PS, Eanes WF. 2015. Variation in *Drosophila melanogaster* central metabolic genes appears driven by natural selection both within and between populations. *Proceedings of the Royal Society Biological Sciences Series B* 282.
- 645 Collinge J, Hoffmann A, McKechnie S. 2006. Altitudinal patterns for latitudinally varying traits and polymorphic markers in *Drosophila melanogaster* from eastern Australia. *Journal of Evolutionary Biology* 19:473-482.
- Consuegra S, John E, Verspoor E, de Leaniz CG. (Consuegra2015 co-authors). 2015. Patterns of natural selection acting on the mitochondrial genome of a locally adapted fish species. *Genetics Selection Evolution (Paris)* 47:1-10.
- 650 Correa CC, Aw WC, Melvin RG, Pichaud N, Ballard JW. 2012. Mitochondrial DNA variants influence mitochondrial bioenergetics in *Drosophila melanogaster*. *Mitochondrion* 12:459-464.
- David JR, Capy P. 1988. Genetic variation of *Drosophila melanogaster* natural populations. *Trends in Genetics* 4:106-111.
- 655 Dobler R, Rogell B, Budar F, Dowling DK. 2014. A meta-analysis of the strength and nature of cytoplasmic genetic effects. *Journal of Evolutionary Biology* 27:2021-2034.
- Doi A, Suzuki H, Matsuura ET. 1999. Genetic analysis of temperature-dependent transmission of mitochondrial DNA in *Drosophila*. *Heredity* 82:555-560.
- 660 Dowling DK. 2014. Evolutionary perspectives on the links between mitochondrial genotype and disease phenotype. *Biochimica et Biophysica Acta* 1840:1393-1403.
- Dowling DK, Abiega KC, Arnqvist G. 2007. Temperature-specific outcomes of cytoplasmic-nuclear interactions on egg-to-adult development time in seed beetles. *Evolution* 61:194-201.
- 665 Dowling DK, Friberg U, Lindell J. 2008. Evolutionary implications of non neutral mitochondrial genetic variation. *Trends in Ecology & Evolution* 23:546-554.
- Endler JA. 1977. Geographic variation, speciation, and clines. *Monographs in Population Biology* 10:1-246.
- Fontanillas P, Dépraz A, Giorgi MS, Perrin N. (pdf co-authors). 2005. Nonshivering thermogenesis capacity associated to mitochondrial DNA haplotypes and gender in the greater white-toothed shrew, *Crocidura russula*. *Molecular Ecology* 14:661-670.
- 670 Foote AD, Morin PA, Durban JW, Pitman RL, Wade P, Willerslev E, Gilbert MTP, da Fonseca RR. 2011. Positive selection on the killer whale mitogenome. *Biology Letters* 7:116-118.
- Fox J. 2002. An R and S-Plus companion to applied regression. Thousand Oaks, Calif.: Sage Publications.
- 675 Frank SA, Hurst LD. 1996. Mitochondria and male disease. *Nature* 383:224.

- Garvin MR, Bielawski JP, Sazanov LA, Gharrett AJ. 2015. Review and meta-analysis of natural selection in mitochondrial complex I in metazoans. *Journal of Zoological Systematics and Evolutionary Research* 53:1-17.
- 680 Gemmell NJ, Metcalf VJ, Allendorf FW. 2004. Mother's curse: the effect of mtDNA on individual fitness and population viability. *Trends in Ecology & Evolution* 19:238-244.
- Hoekstra LA, Siddiq MA, Montooth KL. 2013. Pleiotropic effects of a mitochondrial-nuclear incompatibility depend upon the accelerating effect of temperature in *Drosophila*. *Genetics* 195:1129-1139.
- 685 Hoffmann AA, Anderson A, Hallas R. 2002. Opposing clines for high and low temperature resistance in *Drosophila melanogaster*. *Ecology Letters* 5:614-618.
- Hoffmann AA, Weeks AR. 2007. Climatic selection on genes and traits after a 100 year-old invasion: a critical look at the temperate-tropical clines in *Drosophila melanogaster* from eastern Australia. *Genetica* 129:133-147.
- Hurst LD. 2011. Molecular genetics: The sound of silence. *Nature* 471:582-583.
- 690 James JE, Piganeau G, Eyre-Walker A. 2016. The rate of adaptive evolution in animal mitochondria. *Molecular Ecology* 25:67-78.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, et al. 2012. Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647-1649.
- 695 Kimchi-Sarfaty C, Oh JM, Kim IW, Sauna ZE, Calcagno AM, Ambudkar SV, Gottesman MM. 2007. A "silent" polymorphism in the MDR1 gene changes substrate specificity. *Science* 315:525-528.
- Kivisild T, Shen PD, Wall DP, Do B, Sung R, Davis K, Passarino G, Underhill PA, Scharfe C, Torroni A, et al. 2006. The role of selection in the evolution of human mitochondrial genomes. *Genetics* 172:373-387.
- 700 Lavington E, Cogni R, Kuczynski C, Koury S, Behrman EL, O'Brien KR, Schmidt PS, Eanes WF. 2014. A Small System-High-Resolution Study of Metabolic Adaptation in the Central Metabolic Pathway to Temperate Climates in *Drosophila melanogaster*. *Molecular Biology and Evolution* 31:2032-2041.
- Levin L, Blumberg A, Barshad G, Mishmar D. 2014. Mito-nuclear co-evolution: the positive and negative sides of functional ancient mutations. *Frontiers in Genetics* 5:448.
- 705 Li H, Stephan W. 2006. Inferring the Demographic History and Rate of Adaptive Substitution in *Drosophila*. *PLOS Genetics* 2:e166.
- Ma X, Kang J, Chen W, Zhou C, He S. 2015. Biogeographic history and high-elevation adaptations inferred from the mitochondrial genome of Glyptosternoid fishes (Sisoridae, Siluriformes) from the southeastern Tibetan Plateau. *BMC Ecology and Evolution* 15:233.
- 710 Matsuura ET, Tanaka YT, Yamamoto N. 1997. Effects of the nuclear genome on selective transmission of mitochondrial DNA in *Drosophila*. *Genes & Genetic Systems* 72:119-123.
- Mishmar D, Ruiz-Pesini E, Golik P, Macaulay V, Clark AG, Hosseini S, Brandon M, Easley K, Chen E, Brown MD, et al. 2003. Natural selection shaped regional mtDNA variation in humans. *Proceedings of the National Academy of Sciences of the United States of America* 100:171-176.
- 715 Morales HE, Pavlova A, Joseph L, Sunnucks P. 2015. Positive and purifying selection in mitochondrial genomes of a bird with mitonuclear discordance. *Molecular Ecology* 24:2820-2837.
- Mossman JA, Biancani LM, Zhu CT, Rand DM. 2016. Mitonuclear Epistasis for Development Time and Its Modification by Diet in *Drosophila*. *Genetics* 203:463-484.
- 720 Musto H, Naya H, Zavala A, Romero H, Alvarez-Valín F, Bernardi G. 2004. Correlations between genomic GC levels and optimal growth temperatures in prokaryotes. *FEBS Letters* 573:73-77.

- Nabholz B, Ellegren H, Wolf JBW. 2013. High Levels of Gene Expression Explain the Strong Evolutionary Constraint of Mitochondrial Protein-Coding Genes. *Molecular Biology and Evolution* 30:272-284.
- 725 Nakamura Y, Gojobori T, Ikemura T. 2000. Codon usage tabulated from international DNA sequence databases: status for the year 2000. *Nucleic Acids Research* 28:292.
- Pfaffl MW, Tichopad A, Prgomet C, Neuvians TP. 2004. Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper--Excel-based tool using pair-wise correlations. *Biotechnology Letters* 26:509-515.
- 730 Porcelli D, Butlin RK, Gaston KJ, Joly D, Snook RR. 2015. The environmental genomics of metazoan thermal adaptation. *Heredity* 114:502-514.
- Quintela M, Johansson MP, Kristjansson BK, Barreiro R, Laurila A. 2014. AFLPs and mitochondrial haplotypes reveal local adaptation to extreme thermal environments in a freshwater gastropod. *PLoS One* 9:e101821.
- 735 Quiros PM, Mottis A, Auwerx J. 2016. Mitonuclear communication in homeostasis and stress. *Nature Reviews Molecular Cell Biology* 17:213-226.
- Rand DM. 2001. The Units of Selection on Mitochondrial DNA. *Annual Review of Ecology and Systematics* 32:415-448.
- 740 Rand DM, Haney RA, Fry AJ. (JLEBL, pdf co-authors). 2004. Cytonuclear coevolution: the genomics of cooperation. *Trends in Ecology & Evolution* 19:645-653.
- Rasmussen R. 2001. Quantification on the LightCycler. In: Meuer S, Wittwer C, Nakagawara K-I, editors. *Rapid Cycle Real-Time PCR*: Springer Berlin Heidelberg. p. 21-34.
- Ruiz-Pesini E, Mishmar D, Brandon M, Procaccio V, Wallace DC. 2004. Effects of purifying and adaptive selection on regional variation in human mtDNA. *Science* 303:223-226.
- 745 Salinas T, Duby F, Larosa V, Coosemans N, Bonnefoy N, Motte P, Marechal-Drouard L, Remacle C. 2012. Co-evolution of mitochondrial tRNA import and codon usage determines translational efficiency in the green alga *Chlamydomonas*. *PLOS Genetics* 8:e1002946.
- Sharp PM, Averof M, Lloyd AT, Matassi G, Peden JF. 1995. DNA sequence evolution: the sounds of silence. *Philosophical Transactions of the Royal Society of London B Biological Sciences* 349:241-247.
- 750 Silva G, Lima FP, Martel P, Castilho R. 2014. Thermal adaptation and clinal mitochondrial DNA variation of European anchovy. *Proceedings of the Royal Society Biological Sciences Series B* 281.
- Singh RS, Long AD. 1992. Geographic variation in *Drosophila*: From molecules to morphology and back. *Trends in Ecology & Evolution* 7:340-345.
- 755 Šmarda P, Bureš P, Horová L, Leitch IJ, Mucina L, Pacini E, Tichý L, Grulich V, Rotreklová O. 2014. Ecological and evolutionary significance of genomic GC content diversity in monocots. *Proceedings of the National Academy of Sciences of the United States of America* 111:E4096-E4102.
- Sorensen JG, Nielsen MM, Loeschcke V. 2007. Gene expression profile analysis of *Drosophila melanogaster* selected for resistance to environmental stressors. *Journal of Evolutionary Biology* 20:1624-1636.
- 760 Sun C, Kong QP, Zhang YP. 2007. The role of climate in human mitochondrial DNA evolution: A reappraisal. *Genomics* 89:338-342.
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution* 30:2725-2729.
- 765 Thomas C. 1992. Sugarbeet minicircular mitochondrial DNAs: high-resolution transcript mapping, transcript abundance and copy number determination. *Molecular and General Genetics MGG* 234:457-465.

Toews DP, Brelsford A. 2012. The biogeography of mitochondrial and nuclear discordance in animals. *Molecular Ecology* 21:3907-3930.

770 Torres TT, Dolezal M, Schlotterer C, Ottenwalder B. 2009. Expression profiling of *Drosophila* mitochondrial genes via deep mRNA sequencing. *Nucleic Acids Research* 37:7509-7518.

Wallace DC. 2007. Why Do We Still Have a Maternally Inherited Mitochondrial DNA? Insights from Evolutionary Medicine. *Annual Review of Biochemistry* 76:781-821.

775 Weeks AR, McKechnie SW, Hoffmann AA. 2002. Dissecting adaptive clinal variation: markers, inversions and size/stress associations in *Drosophila melanogaster* from a central field population. *Ecology Letters* 5:756-763.

Williams BR, Van Heerwaarden B, Dowling DK, SgrÒ CM. 2012. A multivariate test of evolutionary constraints for thermal tolerance in *Drosophila melanogaster*. *Journal of Evolutionary Biology* 25:1415-1426.

780 Wolff JN, Ladoukakis ED, Enríquez JA, Dowling DK. 2014. Mitonuclear interactions: evolutionary consequences over multiple biological scales. *Philosophical Transactions of the Royal Society of London B Biological Sciences* 369.

785 Wolff JN, Tompkins DM, Gemmell NJ, Dowling DK. 2016. Mitonuclear interactions, mtDNA-mediated thermal plasticity, and implications for the Trojan Female Technique for pest control. *Scientific Reports* 6:30016.

Yin PH, Lee HC, Chau GY, Wu YT, Li SH, Lui WY, Wei YH, Liu TY, Chi CW. 2004. Alteration of the copy number and deletion of mitochondrial DNA in human hepatocellular carcinoma. *British Journal of Cancer* 90:2390-2396.

790 **Acknowledgements**

We would like to thank Fiona Cockerell, Allannah Clemson, Winston Yee and Belinda Williams for assistance with the heat and cold tolerance assays. We thank Vanessa Kellerman and Winston Yee for fly collection. This research was supported by funding from the Hermon Slade Foundation (HSF 15/2), Australian Research Council (DP1092897, FT160100022), and
795 Monash University Research Fellowship to DKD. CMS was funded by an ARC fellowship and the Science and Industry Endowment Fund.

Tables and Figures

Table 1: Location of all SNPs identified via next-generation sequencing of the 11 mass-bred populations. For each SNP site, we identified nucleotides that were diagnostic of the northern and southern major haplotypes. Here we list the location (Site) of the SNP, the affected gene (Gene), and the codon position (Position). Additionally, for each north and south polymorphism, we list the nucleotide (nt), the codon, amino acid (AA), and the usage bias for the specific codon.

Site	Gene	Position	A Haplogroup				B Haplogroup			
			nt	Codon	AA	Usage Bias	nt	Codon	AA	Usage Bias
1154	<i>mt:ND2</i>	3	C	AAC	N	2.6	T	AAU	N	48.2
2661	<i>mt:COX1</i>	3	C	CCC	P	1.7	T	CCU	P	23.1
3583	<i>mt:COX2</i>	3	T	GCU	A	3	A	GCA	A	13
4247	<i>mt:ATPase6</i>	3	C	GGC	G	0.1	T	GGU	G	12.7
5396	<i>mt:COX3</i>	1	C	CUA	L	7.1	T	UUA	L	134.5
6299	<i>mt:tRNAE</i>		C				A			
6980	<i>mt:ND5</i>	3	G	UAC	Y	7.1	A	UAU	Y	35.6
7862	<i>mt:ND5</i>	3	G	UUC	F	5.6	A	UUU	F	85
8866	<i>mt:ND4</i>	1	A	UUA	L	134.5	G	CUA	L	7.1
8972	<i>mt:ND4</i>	3	C	UUG	L	4.3	T	UUA	L	134.5
10215	<i>mt:ND6</i>	1	C	CUA	L	7.1	T	UUA	L	134.5
10671	<i>mt:CYTB</i>	1	T	UUA	L	134.5	C	CUA	L	7.1
12121	<i>mt:ND1</i>	3	C	AUG	M	2.8	T	AUA	M	51.2
12334	<i>mt:ND1</i>	1	C	GGG	G	4.1	A	GGU	G	12.7
14665	<i>mt:srRNA</i>		C				T			

810 **Table 2:** Location of all SNPs identified via next-generation resequencing of the
mitochondrial genomes of each genetic strain, revealing that the S1 haplotype can be further
partitioned into four unique “sub-haplotypes”. Below is the list comprising the origin from
which each genetic strain was originally sourced (Origin), the identity of the duplicate of
each population (Dup), the haplotype associated with each strain (h.type), the sub-haplotype
815 (sub-h.type), the affected gene (Gene), whether the SNP is synonymous (S) or non-
synonymous (N), the nucleotide change (nt change), the location (Site) of the SNP, and
amino acid change (AA change).

Origin	dup	h.type	sub- h.type	Gene	SNP	nt change	Site	AA change
MEL	1	A1	A1					
MEL	2	A1	A1					
BRIS	1	A1	A1					
BRIS	2	A1	A1					
MEL	1	B1	B1-C	<i>mt:COXII</i>	N	C → T	3359	P → S
MEL	2	B1	B1-D	<i>mt:ND4</i>	S	C → T	8033	
BRIS	1	B1	B1-A	<i>tRNA-ASP</i>		A → C	3892	
				<i>mt:COXIII</i>	S	T → C	4954	
				<i>mt:ND5</i>	S	A → G	7877	V → I/M
BRIS	2	B1	B1-B	<i>mt:COXI</i>	S	G → A	2262	
				<i>mt:COXII</i>	S	C → T	3385	
				<i>mt:COXIII</i>	N	G → A	5162	
				<i>mt:ND4-L</i>	S	A → T	9341	




Figure 1: Haplotype network for mitochondrial protein-coding regions derived from genotyping of 15 SNPs. Circles indicate unique haplotypes with circle size proportional to haplotype frequency. The two different colours correspond to the two haplogroups; with red corresponding to northern haplotypes and grey to southern haplotypes. Further resequencing of A1 and B1 haplotypes revealed that the B1 haplotype is comprised of at least 4 sub-haplotypes (B1-A, B1-B, B1-C, B1-D). Sub-haplotypes all share the same diagnostic 15 SNPs that delineate the B1 from the A1 haplotype, however contain 1 to 4 additional SNPs scattered throughout the genome (see Table 2).




Figure 2: HKY neighbor-joining tree of 19 *Drosophila melanogaster* mitochondrial protein-coding regions. Protein-coding regions from the haplotypes obtained in this study are in red (A1, B1), whereas the rest of the haplotypes were obtained from Camus et al. (2012) and represent haplotypes derived from worldwide locations: Alstonville, New South Wales, Australia (**ALS**); Barcelona, Spain (**BAR**); Brownsville, Texas, USA (**BRO**); Dahomey -now Benin- West Africa (**DAH**); Hawaii, USA (**HAW**); Israel (**ISR**); Japan (**JAP**); Madang, Papua New Guinea (**MAD**); Mysore, India (**MYS**); Oregon, USA (**ORE**); Puerto Montt, Chile (**PUE**); Sweden (**SWE**); and Zimbabwe (**ZIM**).




Figure 3: Haplotype abundance along the Australian eastern coast. **A)** Haplotype A1 (red) is predominantly found in the north of Australia, decreasing in frequency as latitude increases ($R^2 = 0.4847$). **B)** Haplotype B1 (grey) is more common in the south, decreasing in frequency as latitude decreases ($R^2 = 0.5137$).




Figure 4: **A)** Heat tolerance (mean heat “knockdown” time \pm 1 S.E) of males carrying the A1 (red) and B1 (grey) haplotypes / sub-haplotypes. Means for each haplotype are shown separately according to population of origin; Bri refers to Brisbane, Mel refers to Melbourne. **B)** Heat tolerance (mean heat “knockdown” time \pm 1 S.E) of females carrying the A1 (red) and B1 (grey) haplotypes / sub-haplotypes. **C)** Differences in male and female heat tolerance means across mitochondrial haplotypes. **D)** Cold tolerance (100 - mean chill-coma recovery time \pm 1S.E) of males carrying the A1 (red) and B1 (grey) haplotypes / sub-haplotypes. **E)** Cold tolerance (100 - mean chill-coma recovery time \pm 1S.E) of females carrying the A1 (red) and B1 (grey) haplotypes / sub-haplotypes. **F)** Heat and cold tolerance (centred on a mean of zero and standard deviation of 1) across mitochondrial haplotypes.




Figure 5: Least-squares means (± 1 S.E.) of gene expression across A1 (red) and B1 (grey, B1A-D combined) haplotypes for the *mt:ND4* and *mt:ND5* genes (OXPHOS complex 1) and *mt:COI*, *mt:COII*, *mt:COIII* genes (OXPHOS complex IV). *mt:COI*, *mt:COII*, *mt:COIII* all belong to one transcriptional unit and encode subunits of complex IV, whilst *mt:ND4* and *mt:ND5* are members of a second transcriptional unit and encode subunits of complex 1 of the mitochondrial electron transport chain. Least-square means for all plots were derived from the multilevel models, which take into account mtDNA copy number as a covariate (Table S4)