

1 Space use and movement of jaguar (*Panthera onca*) in western Paraguay

2 Roy T McBride, Jr. ^{a,1}, Jeffrey J Thompson ^{*b,c,1}

3 ^a*Faro Moro Eco Research, Departamento de Boquerón, Paraguay*

4 ^b*Asociación Guyra Paraguay, Parque Ecológico Asunción Verde. Avda. Carlos Bóveda,*
5 *Asunción, Paraguay C.C:132*

6 ^c*Consejo Nacional de Ciencia y Tecnología (CONACYT), Asunción, Paraguay*

7 ¹These authors contributed equally to this paper.

8 *Corresponding author: jthompson.py@gmail.com

9 Guyra Paraguay
10 Parque Ecológico Asunción Verde. Avda. Carlos Bóveda
11 Asunción, Paraguay C.C:132
12 595 975 928079
13 jthompson.py@gmail.com

14
15 Faro Moro Eco Research
16 Departamento de Boquerón, Paraguay
17 rocktmcbride@yahoo.com
18

19 Abstract

20 We present the first estimates of home range and core area size for jaguar (*Panthera onca*) in
21 western Paraguay in the Dry Chaco, Humid Chaco and Pantanal. Using an autocorrelated kernel
22 density estimator we estimated mean home range sizes of 818 km² (95% CI:425-1981) in the Dry
23 Chaco and 237 km² (95% CI:90-427) in the Humid Chaco/Pantanal. The home range estimates
24 for the Dry Chaco are the largest recorded for jaguar. Core areas, defined as the home range area
25 where use was equal to expected use, was consistent across sexes and systems on average
26 represented by the 59% utility distribution isopleth (range:56-64%). Males had a higher

probability of larger home ranges and more directional and greater daily movements than females collectively and within systems. The large home ranges in the Dry Chaco are attributable to the relatively low productivity of that semi-arid ecosystem and high heterogeneity in resource distribution while the larger than expected home ranges in the Humid Chaco/Pantanal compared to home range estimates from the Brazilian Pantanal may be due to differences in geomorphology and hydrological cycle. Known mortality of study jaguars from persecution was high (~75%) and consequently we do not rule out the possibility that the observed large home ranges are partially a function of reduced densities. The large home ranges of jaguars in western Paraguay and a low proportional area of protected areas in the region demonstrate the importance of private ranchland for the long-term conservation of the species which, given the high mortality from persecution, highlight a crucial need to incorporate the livestock production sector in conservation decision making, including region-wide efforts to mitigate jaguar-livestock conflict.

Keywords: *Panthera onca*, home range, Adaptive Kernel Density Estimate, Chaco, Paraguay

Introduction

Globally, apex predators, and the maintenance of their functional roles, are severely threatened due to anthropogenic pressures, particularly associated with large spatial needs to access sufficient prey to meet metabolic requirements and persecution (Ripple et al. 2014). Habitat conversion and degradation and over hunting of prey species increase spatial requirements of apex predators, increasing conflict with humans and affecting social behavior,

dispersal and habitat use (Crooks 2002; Cardillo et al. 2004; McDonald 1983; Ripple et al. 2014). Consequently, an understanding of the space use and movement ecology of apex predators is key to effective conservation decision making for these species.

The jaguar (*Panthera onca*) is the largest feline in the Americas, distributed from the southwestern United States to northern Argentina, although it presently occupies < 50% of its original range, and <80% of the range outside of Amazonia, due to habitat loss and persecution (de la Torre et al. 2017; Sanderson et al. 2002; Zeller 2007). Given the contraction of the species' distribution, range-wide conservation efforts have focused upon maintaining connectivity among key populations throughout the species range (Sanderson et al. 2002; Rabinowitz and Zeller 2010), however, an effective implementation of this management approach is partly dependent upon a thorough understanding of the spatial and movement ecology of jaguars.

For a big cat the jaguar is relatively understudied (Brodie 2009), and although multiple studies have estimated jaguar home range size (Azevedo and Murray 2007; Cavalcanti and Gese 2009; Crawshaw and Quigley 1991; Crawshaw 1995; Crawshaw et al. 2004; Cullen 2006; Morato et al. 2016; Rabinowitz and Nottingham 1986; Schaller and Crawshaw 1980; Scognamillo et al. 2002; Silveira 2004; Tobler et al. 2013) and movements (Colchero et al. 2011, Conde et al. 2010; Sollman et al. 2011), there is still relatively little known about the species' spatial and movement ecology. Since anthropogenic factors drive jaguar occurrence throughout its range by determining habitat availability and quality (Petracca et al. 2014a,b; Thompson and Martinez 2015; Zeller et al. 2012) this conspicuous knowledge gap on how jaguars perceive and use the landscape is of concern as it limits managers' ability to quantifiably design and manage conservation landscapes for the jaguar. .

Of further concern is that until recently jaguar home range estimates likely underestimated space use as VHF-based estimates were based upon small number of locations, while GPS-based estimates failed to account for autocorrelation inherent in GPS telemetry data (Morato et al. 2016). Furthermore, only recently have movement parameters and quantitative assessment of home range residency been estimated for jaguar (Morato et al. 2016). Consequently, there is an important need for research that incorporates developing methodologies that account for and take advantage of autocorrelation in telemetry data to better quantify jaguar spatial and movement ecology.

Range-wide, the jaguar is considered near threatened (Caso et al. 2008), however, at the austral limit of its distribution the species is considered critically endangered in Argentina and endangered in Brazil and Paraguay. Although multiple studies have investigated space use by jaguar in Brazil and Argentina (Azevedo and Murray 2007; Cavalcanti and Gese 2009; Crawshaw and Quigley 1991; Crawshaw 1995; Crawshaw et al. 2004; Cullen 2006; Morato et al. 2016; Schaller and Crawshaw 1980; Silveira 2004) there has been no such research on the species in Paraguay despite a recognized need in the face of a rapid constriction in the species' distribution in relation to a country-wide expansion of the agricultural sector (Secretaría del Ambiente et al. 2016) which has resulted in some of the highest rates of deforestation in the world (Hansen et al. 2013).

Given the status of the jaguar in Paraguay, the lack of information on the spatial and movement ecology of the species is of concern within the context of continued habitat loss, the maintenance of in-country and trans-boundary connectivity of populations, and their implications for the range-wide conservation of the jaguar. Consequently, we used GPS-based telemetry to study space use and movements of jaguars in western Paraguay in the Dry Chaco, Humid Chaco

and Pantanal, the region with the largest jaguar population in the country. Moreover, we employed developing methodologies which allowed us to determine home range residency and account for autocorrelation in the data (Calabrese et al. 2016; Fleming et al. 2015, 2016), which in turn allowed for rigorous comparisons with estimates from other research employing the same methodologies (Morato et al. 2016).

Based upon carnivore ecology in general, and jaguar ecology specifically, we expected male home range and movement rates to be higher than females (Cavalcanti and Geese 2009; Conde et al. 2010; Mikael 1989; Morato et al. 2016; Sollmann et al. 2011) and that jaguars in the Dry Chaco would exhibit larger home ranges, higher movement rates, and more directional movement compared to those in the more productive habitats of the Humid Chaco and Pantanal (Mikael 1989; Fahrig 2007; Gutierrez-Gonzalez et al. 2012). Also, when comparing to other sites (Morato et al. 2016) we expected estimated from the Humid Chaco and Pantanal to be similar to those from the Brazilian Pantanal, while estimates from the Dry Chaco would be larger than those from more humid systems but possibly similar to jaguars from the Brazilian Cerrado due to biotic and abiotic similarities between systems. Apart from constituting an important contribution towards the conservation of jaguars within Paraguay, placing our results into a comparative context with research from neighboring countries will facilitate the efficacy of trans-boundary conservation efforts, with important implications for range-wide conservation strategies for jaguar.

Methods

Study area

We conducted our study in three ecosystems in western Paraguay; Dry Chaco, Humid Chaco and Pantanal, (Figure 1). The Dry Chaco is comprised of xeric forest, savannas, and grasslands and the Humid Chaco and Pantanal are a mosaic of seasonally flooded grasslands, palm savanna and xerophilic woodlands on higher ground (Olson et al. 2001; Mereles et al. 2013). We note that delineations between the Humid Chaco and Pantanal differ (Olson et al. 2001; Mereles et al. 2013), however, for our purposes the similarities between systems and among our study sites in those systems make this discrepancy moot and consequently we treat the Humid Chaco and Pantanal as a single system in our analysis.

The western half of Paraguay is generally semi-arid with a pronounced east–west gradient in precipitation and humidity which divides the Chaco into the Humid Chaco with precipitation approximately > 1000 mm/year and the Dry Chaco with precipitation < 1000 mm/year (Olson et al. 2001). The Pantanal is also subjected to this east-west precipitation gradient; however, it and the Humid Chaco are also strongly effect by the hydrological cycles of the Rio Paraguay (Mereles et al. 2013).

In the Humid Chaco our study area was Estancia Aurora, a 30,000 ha cattle ranch in the north of the department of Villa Hayes and in the Pantanal on the 65,000 ha ranch Estancia Fortín Patria and on the 80,000 ha ranch Estancia Leda. In the central Dry Chaco, we worked on the 40,000 ha Faro Moro ranch and more northerly in the 7,200 km² Defensores del Chaco National Park and the neighboring 269,000 ha of ranchland of the consortium *Grupo Chavoreca*.

Jaguar captures

Jaguars were captured using trained hounds to tree or bay jaguars which were then anesthetized using a weight-dependent dose of a mix of ketamine hydrochloride and xylazine hydrochloride injected by a dart shot from a tranquilizer gun (McBride and McBride 2007). Capture methods followed ASM protocols (Sikes 2016) and in > 60 captures and recaptures of jaguar and puma over the study period there were no deaths or noticeable injury to animals.

From 2002–2009 jaguars were fitted with Telonics Generation II, data stored-on-board, GPS collars (Telonics, Mesa, Arizona, USA) which were set to record locations at 4 hour intervals. Starting in 2009 we used Northstar GPS collars (D-cell, Northstar, King George, Virginia, USA) programmed to record locations at three or four hour intervals and in 2012 we switched to Telonics Generation III GPS collars (Telonics, Mesa, Arizona, USA) which were set to record locations daily every two hours from 1800 to 0600 hours.

Home range estimation

Semi-variogram analysis, model selection and AKDE estimates were undertaken using the *ctmm* package (Calabrese et al. 2016) in R 3.3.2 (R Development Core Team 2010). Starting values derived from semi-variograms were used for maximum likelihood model fitting with model selection based upon Akaike Information Criteria, adjusted for small sample size (AICc), and model weights (Calabrese et al. 2016; Fleming et al. 2014, 2015). We accounted for data collected with an irregular sampling schedule from collars used starting in 2012 with the *dt* argument within the *variogram* function in the *ctmm* package (Calabrese et al. 2016).

Movement models tested were an independent identically distributed (IID) model which ignores autocorrelation in the data and is equivalent to kernel density estimation (KDE) (Worton

et al. 1989), a random search model (Brownian motion) with no home range, Brownian motion within a home range (Ornstein–Uhlenbeck, OU), and Ornstein–Uhlenbeck motion with foraging (OUF) (Calabrese et al. 2016; Fleming et al. 2014). Both the OU and OUF models produce estimates of home range size and home range crossing time, while the OUF model additionally estimates the velocity autocorrelation time scale (a measure of path sinuosity) and mean distance traveled (Calabrese et al. 2016; Fleming et al. 2014).

Home ranges were estimated using the best fit model for each individual using AKDE (Calabrese et al. 2016; Fleming et al. 2015). For comparison with home range estimates from previous research we estimated 95% KDE home ranges using the IID model and 95% Minimum Convex Polygons (MCP) home ranges using the adehabitatHR package in R (Calenge 2006) (Supplementary Table S1).

Core area estimation

We estimated core areas of AKDE home ranges as the area encompassed within the isopleth where the proportional use of the estimated home range is equal to the predicted probability of use (Bingham and Noon 1997; Seaman and Powell 1990; Vander Wal and Rodgers 2012). We determined this by fitting an exponential curve to the isopleths of the utility distribution of each individual at 10% increments from 10% to 100% of the AKDE home range and the proportional area of the home range that each of those isopleths encompassed. We then determined the threshold where proportional home range size begins to increase at a rate greater than the probability of use (slope=1; Bingham and Noon 1997; Seaman and Powell 1990; Vander Wal and Rodgers 2012) to define the isopleth that represented the core area boundary.

182

183 *Statistical analyses*

184 For our statistical analysis we combined jaguars from the Humid Chaco and the Pantanal
185 into a single group as the characteristics of the system are highly similar , the delineation
186 between the two systems is debatable (Olson et al. 2001, Mereles et al. 2013), and consequently
187 jaguars from those systems are subjected to similar ecological and anthropogenic drivers.
188 Additionally, only individuals that exhibited residency in their movement behavior through semi-
189 variogram analysis and space use best explained by the OUF model were included in our
190 comparative analysis of differences between sexes and ecosystems.

191 We used a fixed-effect one-way analysis of variance (ANOVA) in a Bayesian modeling
192 framework to test for differences in estimates of home range size, home range crossing time,
193 directionality in movement (velocity autocorrelation time scale) and mean daily distance traveled
194 between sexes across systems, between systems (sexes combined), between sexes within a
195 system, and between same sexes between systems. We tested normality using the Shapiro-Wilk
196 test and log-transforming the data when its distribution did not meet assumptions of normality.

197 All analyses were undertaken in R 3.2.2. (R Development Core Team 2010) using
198 WinBUGS (Lunn et al. 2000) and the *R2bugs* package (Sturtz et al. 2005) for the Bayesian
199 analysis. We ran 3 chains in WinBUGS with 100,000 iterations and a 20,000 iteration burn-in
200 period; confirming convergence by a scale reduction factor ≤ 1.01 and visual inspection of trace
201 plots for lack of autocorrelation. We tested differences between groups by taking 10,000 random
202 samples from posterior distributions for each group of interest, comparing the proportional
203 frequency that posterior estimates parameters were greater for males than females overall and

within systems, greater for all individuals, and between same sexes, in the Dry Chaco compared to the Humid Chaco/Pantanal.

Results

Jaguar captures and data collection

We captured and collared 35 jaguars from June 2002 to June 2014 of which 19 individuals provided sufficient data for analysis; 7 in the Dry Chaco (5 males, 2 females), 9 in the Humid Chaco (3 males, 6 females) and 3 in the Pantanal (1 male, 2 females) with estimated ages between 2 and 10 years (Table 1). Collars collected data between 52 and 439 days, obtaining from 148 to 3462 locations (Table 1). The length of the study period and the annual frequency of captures were dependent upon resource availability and logistical restraints that dictated captures and collar recovery.

Home range, core area and movement parameter estimates.

Best fitting models for the movement of jaguars were either the OU or OUF models with 16 individuals demonstrating residency (Table 1). Estimated home range sizes varied between 86 and 2,909 km² and core areas between 21-509 km². Core areas were represented by a consistent proportion of the utility distribution; ranging between 56%-64% isopleths (Table 2).

Male and female mean home range size were 727 km² (95% CI:355-1954) and 255 km² (95% CI:90-578), respectively and 818 km² (95% CI:425-1981) and 237 km² (95% CI:90-427) for jaguars in the Dry Chaco and Humid Chaco/Pantanal, respectively (Fig. 2, Fig.3). In the Dry

Chaco mean home range size for males was 925 km² (95% CI:424-2035) and 551 km² (95% CI:513-590) for females, while in the Humid Chaco/Pantanal the mean home range was 398 km² (95% CI:345-427) and 156 km² (95% CI:90-267) for males and females, respectively (Fig. 4).

228

Table 1. Sex, age, sample characteristics and estimated movement parameters, AKDE home range, core area and core area utility distribution isopleths for study jaguars in the Paraguayan Dry Chaco, Humid Chaco and Pantanal.

ID	Sex/age (yr)	Number of fixes/days	Velocity autocorrelation timescale (h)	Home range crossing time (days)	Average distance traveled (km/day)	Home range (km ²) (95% CI)	Core (km ²)	Core area isopleths (%)
<i>Dry Chaco</i>								
DC1	M/5	1094/376	1.1	8.0	28.8	2909 (2115-3892)	504	59
DC3	M/2	722/363	1.8	11.5	7.9	501 (342-689)	107	63
DC4	M/5	620/82	1.9	3.5	15.0	597 (379-865)	182	58
DC6	M/7	1387/393	2.2	4.8	19.3	1266 (979-1589)	329	57
DC7	M/5	3462/439	1.4	2.7	17.1	409 (350-473)	85	64
DC5	F/6	1610/386	1.1	11.5	11.8	674 (468-917)	178	59
DC2	F/8	921/379	1.7	9.5	9.7	589 (419-788)	176	56
<i>Humid Chaco/Pantanal</i>								

Pan2	F/2	1694/375	1.1	4.3	7.9	113 (89-139)	24	60
HC5	F/4	593/242	0.5	1.2	20.9	96 (79-115)	23	61
HC4	F/3	288/266	1.5	6.2	9.3	308 (213-420)	86	57
HC7	F/1	980/170	0.2	10.2	13.7	125 (74-190)	32	58
HC8	F/6	1668/324	0.1	9.2	22.3	264 (185-356)	73	57
HC9	F/6	928/362	0.2	9.7	13.9	124 (87-168)	33	59
Pan3	M/6	727/192	1.4	3.4	16.6	514 (385-662)	134	57
HC2	M/4	983/143	1.4	4.4	15.0	525 (359-723)	138	56
HC6	M/10	660/133	0.9	5.5	13.4	338 (214-490)	91	60
Pan1	F/4	1695/366	NA	3.5	NA	95 (76-115)	21	60
HC1	M/6	148/88	NA	5.9	NA	1061 (591-1667)	283	58
HC3	F/6	280/54	NA	5.7	NA	86 (41-147)	22	57

232

233 Males demonstrated larger home ranges ($P=0.99$), higher daily movement ($P=0.84$),
234 greater directionality in movement (velocity autocorrelation time scale) ($P=0.84$) and lower
235 home range crossing times ($P=0.9$) (Table 2, Figure 2). Between systems, home ranges were
236 larger ($P=1$), movements more directional ($P=0.99$) and home range crossing times greater
237 ($P=0.77$) in the Dry Chaco, while daily travel distance was similar between systems but with a
238 slightly higher probability of being larger in the Dry Chaco ($P=0.61$, Figure 3).

239 Between systems males in the Dry Chaco had higher probabilities to have larger home
240 ranges ($P=0.91$), higher home range crossing time ($P=0.75$), greater directionality in movement
241 ($P=0.86$), and greater daily travel distances ($P=0.72$) (Table 2), although values for all
242 parameters were more variable in males from the Dry Chaco (Figure 4). A similar pattern was

243 evident between females in both systems for home range size ($P=0.99$), home range crossing
244 time ($P=0.89$) and directionality in movement ($P=0.96$) which were greater for females in the
245 Dry Chaco, however, females in the Dry Chaco had lower daily movements ($P=0.23$) than those
246 in the Humid Chaco/Pantanal (Table 4).

247

248 Table 2. Probabilities, based upon posterior distributions, that home range and movement
249 parameters are different between sex and ecosystem, between sexes within systems, and between
250 same sexes between systems.

	Home range (km ²)	Home range crossing time (days)	Velocity autocorrelation timescale (h)	Average distance traveled (km/day)
Dry Chaco male > Dry Chaco female	0.71	0.08	0.71	0.89
Humid Chaco/Pantanal male > Humid Chaco/Pantanal female	0.99	0.17	0.95	0.52
Dry Chaco male > Humid Chaco/Pantanal male	0.91	0.75	0.86	0.72
Dry Chaco female > Humid	0.99	0.89	0.96	0.23

Chaco/Pantanal female				
All Dry Chaco > All Humid Chaco/Pantanal	1	0.77	0.99	0.61
Male > Female	0.99	0.10	0.99	0.84

Discussion

We present the first estimates of movement parameters and home range and core area for jaguar in the Dry Chaco, Humid Chaco, and Paraguayan Pantanal, which furthermore take advantage of developing methods to empirically test for home range residency and account for autocorrelation in telemetry data when estimating space use (Calabrese et al. 2016; Fleming et al. 2014, 2015). Our results include the largest home range estimates recorded for jaguar (Dry Chaco) and, as expected, jaguars in the more productive Humid Chaco/Pantanal had smaller home ranges, lower movement rates and had less directionality in movements compared to jaguars in the Dry Chaco. Also, consistent with previous research males had larger home ranges, higher movement rates and more directional movements than females overall and within systems.

Overall and between systems male home ranges were larger than females which was expected (Calvalcanti and Gese 2009; Sollmann et al. 2011; Morato et al. 2016) as smaller home ranges of females are driven by food availability in relation to reproductive and offspring rearing needs which in-turn drives larger male home ranges towards optimizing reproductive

opportunities (Mikael 1989; Sunquist and Sunquist 1989). This relationship is further supported by our estimated movement parameters which showed that males traveled farther, faster, and more directionally than females in utilizing home ranges.

Consistent with our expectations home range sizes of jaguars in the Dry Chaco were larger than in the Humid Chaco and Pantanal, overall and between sexes within systems where male home ranges were greater than females. The larger home ranges in the Dry Chaco are attributable to the lower productivity of that semi-arid ecosystem, more heterogeneously distributed prey and water, and negative effects of anthropogenic factors (i.e., deforestation; Fahrig 2007; Gutierrez-Gonzalez et al. 2012). The difference between sexes within systems is attributable to differences in territorial organization stemming from aforementioned reproductive and social needs (Mikael 1989; Sunquist and Sunquist 1989).

Home range estimates from the Dry Chaco for both males and females are considerably larger than other estimates from this study and Morato et al. (2016), although our estimates of male home range size from the Dry Chaco (mean: 925 km², 95% CI: 424-2035) are consistent with the estimate for a single male from the Brazilian Cerrado (1269 km²), a semi-arid ecosystem with environmental and land use similarities to the Gran Chaco. Morato et al. (2016) demonstrated that increasing home range size of jaguars was associated with lower habitat quality, which is consistent with the very large home ranges from the Dry Chaco which were closest in size to Morato et al.'s (2016) home ranges in the Atlantic forest which they considered to be of the lowest habitat quality of their study areas.

We expected home range sizes from the Humid Chaco/Pantanal to be similar to estimates from the Brazilian Pantanal, however, our estimates were 59% and 112% larger for males and

females, respectively than home ranges reported for the Brazilian Pantanal; falling between estimates from the Amazon and Atlantic forest, although most similar to jaguars from the Amazon (Morato et al. 2016; Fig. 5). These differences may be related to differences in the geomorphology of the two regions and its interaction with the local hydrological cycles.

The Paraguayan Pantanal and our study area in the Humid Chaco have less forest area and a relatively greater area of inundated land during a large portion of the year compared to the Pantanal study areas of Morato et al. (2016) in Brazil. Consequently, the reduced forest area, with smaller and more isolated forest patches during annual flooding, could drive the comparatively larger home ranges observed in the Paraguayan Pantanal and Humid Chaco, although reduced jaguar densities resulting from persecution may also play a role in liberating available space and permitting greater space use.

Differences in the mean movement parameters were evident between jaguars in the Humid Chaco/Pantanal and in the Brazilian Pantanal whereby movements were more directional in the Humid Chaco/Pantanal, although still relatively sinuous but most similar to jaguars in the Atlantic forest, while daily movements were very similar to those in the Amazon (Fig. 6). Jaguars in the Dry Chaco had high movement rates and directionality in movement, similar to individuals in the Amazon from seasonally flooded forests (Morato et al. 2016).

We believe that these similarities are responses to movements among sporadically distributed critical resources despite the large differences in ecosystem characteristics. Conversely, although daily movement rate of jaguars in the Humid Chaco/Pantanal were similar to those in the Dry Chaco and Amazon, the relatively low directionality demonstrated by jaguars

in the Humid Chaco/Pantanal suggests that, although jaguars are covering relatively large areas, movements are in response to more homogenously distributed resources within home ranges.

Core areas, as measured by the utility distribution isopleth were highly similar across systems and sexes, encompassed on average by the 59% isopleth (95% CI:56-64%), which represented on average 29% (95% CI:21-34%) of total home range area. This indicates that despite home range size, sex, or system jaguars are most intensively using about a third of their home range area. Additionally, our results suggests a cautious interpretation of arbitrarily defined core area delimitations, typically assigned to the 50% utility distribution isopleths which falls outside of the 95% confidence limits of our estimates (Powell 2012).

In light of the extensive deforestation that is occurring in the Dry Chaco of western Paraguay, the large home ranges that we observed in this system, which are consistent with the estimated low density of jaguar in the Bolivian Dry Chaco (Noss et al. 2012), are of concern as they demonstrate the large forested area that jaguars in the Dry Chaco require. In the Humid Chaco/Pantanal spatial requirement of jaguars were greater than expected based on estimates from the Brazilian Pantanal, which suggests lower than expected densities in these systems in Paraguay and cautions against extrapolating population parameter estimates from other regions within the Pantanal to the Rio Paraguay flood plain in Paraguay.

In both the Dry Chaco and the Humid Chaco/Pantanal we recognize that there may be an important effect on space use caused by reduced jaguar densities from persecution which is pervasive throughout western Paraguay, illustrated by our confirmation, or high probability, of ~75% of our study animals being killed due to persecution. Persecution is common throughout the range of the jaguar, however, its practice and magnitude is not equivocal geographically and

consequently how the removal of individuals may impact space use, and subsequently comparisons among ecosystems and regions, needs to be considered and is of interest for future research.

The large spatial requirements of jaguars in western Paraguay, particularly in the Dry Chaco, indicate that the protected areas of the region which, represent <5% of the total regional area are likely insufficient to maintain a viable regional population, especially in light of the level of persecution on private lands. This highlights an urgent need to mitigate jaguar-human conflict in the region by actively including the livestock production sector in the conservation decision making process. Furthermore, given continuing deforestation, conservation initiatives need to take into account the large spatial needs of jaguar in western Paraguay by recognizing and incorporating the role of private lands in the long-term conservation of the species in Paraguay and in maintaining trans-boundary connectivity among populations.

Acknowledgements

We thank DVM Sybil Zavala, Cougar McBride and Caleb McBride for assistance in the field and the many sportsman and conservationist who contributed to supporting this work. This research was conducted under the permission of the Secretariat of the Environment (SEAM) of Paraguay. JJT was supported by the Consejo Nacional de Ciencia y Tecnología of Paraguay (CONACYT).

Appendix A

Table S1. Home range estimates of study jaguars using autocorrelated kernel density estimation (AKDE), kernel density estimation (KDE) and minimum convex polygon (MCP).

Literature Cited

- Azevedo, F. C. C., Murray, D. L., 2007. Spatial organization and food habits of jaguars (*Panthera onca*) in a floodplain forest. *Biol. Conserv.* 137, 391–401.
- Bingham, B.B., Noon, B.R., 1997. Mitigation of habitat “take”: Application to habitat conservation planning. *Conserv Biol.* 11, 127-139.
- Brodie, J.F. 2009. Is research effort allocated efficiently for conservation? Felidae as a global case study. *Biodivers Conserv.* 18, 2927–2939.
- Calabrese, J.M., Fleming, C.H., Gurarie, E., 2016. Ctm: an R package for analyzing animal relocation data as a continuous-time stochastic process. *Methods Ecol Evol.* 7, 1124-1132.
- Calenge, C., 2006. The package adehabitat for the R software: a tool for the analysis of space and habitat use by animals. *Ecol Model.* 197, 516-519.
- Cardillo, M., Purvis, A., Sechrest, W., Gittleman, J.L., Bielby, J., Mace, G.M., 2004. Human population density and extinction risk in the world's carnivores. *PLOS Biology* 2(7): e197. doi: 10.1371/journal.pbio.0020197.
- Caso, A., Lopez-Gonzalez, C., Payan, Eizirik, E., de Oliveira, T., Leite-Pitman, R., Kelly, M., Valderrama, C, 2008. *Panthera onca*. The IUCN Red List of Threatened Species 2008: e.T15953A5327466. <http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T15953A5327466.en>.
- Cavalcanti, S.M.C., E.M. Gese. 2009. Spatial ecology and social interactions of jaguars (*Panthera onca*) in the southern Pantanal, Brazil. *J. Mammal.* 90, 935–945.

Colchero, F., Conde, D., Manterola, C., Chávez, C., Rivera, A., Ceballos, G., 2011. Jaguars on the move: Modeling movement to mitigate fragmentation from road expansion in the Mayan Forest. *Anim Conserv.* 14, 158-166.

Conde, D.A., Colchero, F., Zarza, H., Christensen, N.L., Sexton, J.O., Manterola, C., Chávez, C., Rivera, A., Azuara, D. and Ceballos, G., 2010. Sex matters: Modeling male and female habitat differences for jaguar conservation. *Biol Conserv.* 143, 1980-1988.

Crawshaw, Jr., P.G., 1995. Comparative ecology of ocelot (*Felis pardalis*) and jaguar (*Panthera onca*) in a protected subtropical forest in Brazil and Argentina. Ph.D. dissertation, University of Florida. Gainesville, Florida, USA.

Crawshaw Jr., P.G., Quigley, H.B., 1991. Jaguar spacing, activity, and habitat use in a seasonally flooded environment in Brazil. *J. Zool.* 223, 357–370.

Crawshaw, Jr., P. G., Mahler, J. K., Indrusiak, C., Cavalcanti, S. M. C., Leite-Pitman, M. R. P., Silvius, K. M., 2004. Ecology and conservation of the jaguar (*Panthera onca*) in Iguazu National Park, Brazil, in: Silvius, K.M., Bodmer, R. E., Fragoso, J. M. V. (Eds.). *People in nature: wildlife conservation in South and Central America*. Columbia University Press, New York, pp. 286–296.

Crooks, K.R., 2002. Relative sensitivities of mammalian carnivores to habitat fragmentation. *Conserv Biol.* 16, 488-502.

Cullen Jr., L., 2006. Jaguar as landscape detectives for the conservation in the Atlantic Forest of Brazil. Ph.D. dissertation. University of Kent, UK.

de la Torre, J.A., González-Maya, J.F., Zarza, H., Ceballos, G., Medellín, R.A., 2017. The jaguar's spots are darker than they appear: assessing the global conservation status of the jaguar *Panthera onca*. *Oryx*. 1-16. doi:10.1017/S0030605316001046

Fahrig, L., 2007. Non-optimal animal movement in human-altered landscapes. *Funct Ecol.* 21,

399 1003-1015.

400 Fleming, C.H., Calabrese, J.M., Mueller, T., Olson, K.A., Leimgruber, P., Fagan, W.F., 2014.

401 From fine-scale foraging to home ranges: a semivariance approach to identifying movement

402 modes across spatiotemporal scales. *Am Nat.* 183: E154-E167.

403 Fleming, C.H., Fagan, W.F., Mueller, T., Olson, K.A., Leimgruber, P., Calabrese, J.M., 2015.

404 Rigorous home range estimation with movement data: a new autocorrelated kernel density

405 estimator. *Ecology.* 96, 1182-1188.

406 Gutiérrez-González, C.E., Gómez-Ramírez, M.Á., López-González, C.A., 2012. Estimation of

407 the density of the near threatened jaguar *Panthera onca* in Sonora, Mexico, using camera trapping

408 and an open population model. *Oryx*, 46, 431-437.

409 Hansen, M.C., Potapov, P.V., Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A., Thau,

410 D., Stehman, S.V., Goetz, S.J., Loveland, T.R., Kommareddy, A., 2013. High-resolution global

411 maps of 21st-century forest cover change. *Science*, 342, 850-853.

412 Lunn, D.J., Thomas, A., Best, N., and Spiegelhalter, D., 2000. WinBUGS – a Bayesian

413 modelling framework: concepts, structure, and extensibility. *Stat Comput*, 10, 325-337.

414 McBride, Jr., R.T., McBride, R.T., 2007. Safe and selective capture technique for jaguars in the

415 Paraguayan Chaco. *Southwest Nat.* 52, 570-577.

416 Macdonald, D.W., 1983 The ecology of carnivore social behaviour. *Nature.* 301, 379-384.

417 Mereles, F., Cartes, J.L., Clay, R.P., Cacciali, P., Paradedá, C., Rodas, O., A. Yanosky, 2013.

418 Análisis cualitativo para la definición de las ecorregiones de Paraguay occidental. *Paraquaria*

419 *Natural.* 1, 12-20.

420 Mikael, S. 1989. The mating tactics and spacing patterns of solitary carnivores, in: Gittleman,
421 J.L., (Ed.). Carnivore behavior, ecology, and evolution. Cornell University Press, New York. pp.
422 164-182.

423 Morato, R.G., Stabach, J.A., Fleming, C.H., Calabrese, J.M., De Paula, R.C., Ferraz, K.M.,
424 Kantek, D.L., Miyazaki, S.S., Pereira, T.D., Araujo, G.R. and Paviolo, A., 2016. Space Use and
425 Movement of a Neotropical Top Predator: The Endangered Jaguar. PloS one, 11(12),
426 p.e0168176.

427 Noss, A.J., Gardner, B., Maffei, L., Cuéllar, E., Montaña, R., Romero-Muñoz, A., Sollman, R.,
428 O'Connell, A.F., 2012. Comparison of density estimation methods for mammal populations with
429 camera traps in the Kaa-Iya del Gran Chaco landscape. Anim Conserv. 15, 527-535.

430 Olson, D.M., Dinerstein, E., Wikramanayake, E.D., Burgess, N.D., Powell, G.V., Underwood,
431 E.C., D'amico, J.A., Itoua, I., Strand, H.E., Morrison, J.C., Loucks, C.J., 2001. Terrestrial
432 Ecoregions of the World: A New Map of Life on Earth: A new global map of terrestrial
433 ecoregions provides an innovative tool for conserving biodiversity. BioScience, 51, 933-938.

434 Petracca, L.S., Hernández-Potosme, S., Obando-Sampson, L., Salom-Pérez, R., Quigley, H.,
435 Robinson, H.S., 2014a. Agricultural encroachment and lack of enforcement threaten connectivity
436 of range-wide jaguar (*Panthera onca*) corridor. J Nat Conserv. 22, 436-444.

437 Petracca, L.S., Ramírez-Bravo, O.E. Hernández-Santín, L., 2014b. Occupancy estimation of
438 jaguar *Panthera onca* to assess the value of east-central Mexico as a jaguar corridor. Oryx, 48,
439 133-140.

440 Powell, R. A., 2012. Movements, home ranges, activity, and dispersal, in: Boitani, L., Powell,
441 R.A., (Eds.), Carnivore ecology and conservation: a handbook of techniques. Oxford University
442 Press, London, United Kingdom. pp. 188-217.

443

444 R Development Core Team, 2010. R: A language and environment for statistical computing. R
445 Foundation for Statistical Computing, Vienna, Austria.

446 Rabinowitz, A. R., Nottingham, B. G., 1986. Ecology and behavior of the jaguar (*Panthera*
447 *onca*) in Belize, Central America. J Zool. 210, 149–159.

448 Rabinowitz, A., Zeller, K. A., 2010. A range-wide model of landscape connectivity and
449 conservation for the jaguar, *Panthera onca*. Biol. Conserv. 143, 939–945.

450 Ripple, W.J., Estes, J.A., Beschta, R.L., Wilmers, C.C., Ritchie, E.G., Hebblewhite, M., Berger,
451 J., Elmhagen, B., Letnic, M., Nelson, M.P. and Schmitz, O.J., 2014. Status and ecological effects
452 of the world's largest carnivores. Science. 343(6167), 1241484.

453 Sanderson, E.W., Redford, K. H., Chetkiewicz, C. B., Medellin, R.A., Rabinowitz, A.R.,
454 Robinson, J.G., Taber, A.B., 2002. Planning to save a species: the jaguar as a model. Conserv.
455 Biol. 16, 58–71.

456 Schaller, G.B., Crawshaw, P.G., Jr., 1980. Movement patterns of jaguar. Biotropica 12, 161–168.

457 Secretaría del Ambiente, Wildlife Conservation Society Paraguay, Itaipu Binacional, 2016. Plan
458 de Manejo de la *Panthera onca*, Paraguay 2017-2026. First ed., Asunción, Paraguay.

459 Seaman, D.E., Powell, R.A., 1990. Identifying patterns and intensity of home range use. Bears:
460 their biology and management Vol. 8, A Selection of Papers from the Eighth International
461 Conference on Bear Research and Management. Victoria, British Columbia, Canada, February
462 1989. International Association of Bear Research and Management. pp.243-249.

463 Scognamillo, D., Maxit, I., Sunquist, M., Farrell, L., 2002. Ecología del jaguar y el problema de
464 la depredación de ganado en un hato de los llanos venezolanos. in: Medellin, R., Equihua, C.,
465 Chetkiewicz, C. L.B., Crawshaw, Jr., P. G, Rabinowitz, A., Redford, K.H., Robinson, J.G.,

466 Sanderson, E.W., Taber, A.B. (Eds.), El jaguar en el nuevo milenio. Universidad Nacional
467 Autónoma de México and Wildlife Conservation Society, Distrito Federal, México. pp. 139–150

468 Sikes, R.S., 2016. Guidelines of the American Society of Mammalogists for the use of wild
469 mammals in research and education. J Mammal. 97, 663-688.

470 Silveira, L., 2004. Ecologia Comparada e Conservação da Onça-pintada (*Panthera onca*) e
471 Onça-parda (*Puma concolor*), no Cerrado e Pantanal. Ph.D. Dissertation, University of Brasilia,
472 Brasilia, Brasil.

473 Sollmann, R., Furtado, M.M., Gardner, B., Hofer, H., Jácomo, A.T., Tôrres, N.M. , Silveira, L.,
474 2011. Improving density estimates for elusive carnivores: accounting for sex-specific detection
475 and movements using spatial capture–recapture models for jaguars in central Brazil. Biol
476 Conserv., 144, 1017-1024.


477 Sturtz, S., Ligges, U., Gelman, A., 2005. R2WinBUGS: A Package for Running WinBUGS from
478 R. J Stat Soft., 12, 1-16.

479 Sunquist, M.E., Sunquist, F.C. 1989. Ecological constraints on predation by large felids, in:
480 Gittleman, J.L. (ed.), Carnivore behavior, ecology, and evolution. Cornell University Press,
481 Ithaca, New York. pp. 283–301

482 Thompson, J.J., Martínez Martí, C., 2015. Patterns and determinants of jaguar (*Panthera onca*)
483 occurrence in habitat corridors at the southwestern extent of the species range, in: Martínez Martí,
484 C., (Ed.), Cats, Cores and Corridors: A survey to assess the status of Jaguars and their habitat in
485 the southernmost part of their range. Panthera. New York. pp. 26-40.

486 Tobler, M.W., Carrillo-Percegué, S.E., Hartley, A.Z. and Powell, G.V., 2013. High jaguar
487 densities and large population sizes in the core habitat of the southwestern Amazon. Biol
488 Conserv, 159, 375-381.

- 489 Vander Wal, E., Rodgers, A.R., 2012. An individual-based quantitative approach for delineating
490 core areas of animal space use. *Ecol Model.* 224, 48–53.
- 491 Zeller, K.A., 2007. *Jaguars in the New Millennium Data Set Update: The State of the*
492 *Jaguar in 2006.* Wildlife Conservation Society, Bronx, New York.
- 493 Zeller, K.A., Nijhawan, S., Salom-Pérez, R., Potosme, S.H., Hines, J.E., 2011. Integrating
494 occupancy modeling and interview data for corridor identification: a case study for jaguars in
495 Nicaragua. *Biol Conserv.* 144, 892–901.



496
497 Figure 1. Map showing the distribution of the Dry and Humid Chaco and Pantanal in western
498 Paraguay (Olson et al. 2001) and study areas where jaguar movements were monitored.
499




Figure 2. Home range and movement parameters of male and female jaguars across all study sites.




Figure 3. Home range and movement parameters for study jaguars in the Dry Chaco and Humid Chaco/Pantanal.




Figure 4. Home range and movement parameters of male and female jaguars in the Dry Chaco and Humid Chaco/Pantanal.




Figure 5. Mean male and female home ranges (error bars represent 95% confidence interval) from this study and AKDE estimates from Morato et al. (2016) by ecosystem. The numbers next to points represent sample size.




Figure 6. Mean of the mean distance traveled and the velocity autocorrelation timescale (error bars represent SE) of jaguars from this study and mean estimates from Morato et al. (2016) by ecosystem.

530 Table S1. Comparative home range sizes for study jaguars based upon autocorrelated kernel
531 density estimator (AKDE), 95% kernel density estimator (KDE), and 95% minimum convex
532 polygon (MCP).

ID	Sex/age (yr)	AKDE Home range (km ²) (95% CI)	95% KDE (km ²)	95% MCP (km ²)
<i>Dry Chaco</i>				
DC1	M/5	2909 (2115-3892)	1674	1813
DC3	M/2	501 (342-689)	193	152
DC4	M/5	597 (379-865)	453	389
DC6	M/7	1266 (979-1589)	877	950
DC7	M/5	409 (350-473)	418	587
DC5	F/6	674 (468-917)	479	475
DC2	F/8	589 (419-788)	402	298
<i>Humid Chaco/Pantanal</i>				
Pan2	F/2	113 (89-139)	71	87
HC5	F/4	96 (79-115)	102	89
HC4	F/3	308 (213-420)	176	151
HC7	F/1	125 (74-190)	87	88
HC8	F/6	264 (185-356)	205	184
HC9	F/6	124 (87-168)	87	96
Pan3	M/6	514 (385-662)	350	288
HC2	M/4	525 (359-723)	302	255
HC6	M/10	338 (214-490)	280	256
Pan1	F/4	95 (76-115)	59	67
HC1	M/6	1061 (591-1667)	644	491
HC3	F/6	86 (41-147)	52	36

533