

## How, when, and where relic DNA biases estimates of microbial diversity

2

JT Lennon<sup>1\*</sup>, ME Muscarella<sup>1,3</sup>, SA Placella<sup>2</sup>, BK Lehmkuhl<sup>1</sup>

4

<sup>1</sup>Indiana University, Department of Biology

6

<sup>2</sup>Michigan State University, W.K. Kellogg Biological Station

<sup>3</sup>Current address: University of Illinois, Department of Plant Biology

8

\* corresponding author: lennonj@indiana.edu

10

12           **Extracellular or "relic" DNA is one of the largest pools of nucleic acids in the**  
13           **biosphere<sup>1,2</sup>. Relic DNA can influence a number of important ecological and evolutionary**  
14           **processes, but it may also bias estimates of microbial abundance and diversity, which has**  
15           **implications for understanding environmental, engineered, and host-associated ecosystems.**  
16           **We developed models capturing the fundamental processes that regulate the size and**  
17           **composition of the relic DNA pools to identify scenarios leading to biased estimates of**  
18           **biodiversity. Our models predict that bias increases with relic DNA pool size, but only**  
19           **when the species abundance distributions (SAD) of relic and intact DNA are distinct from**  
20           **one another. We evaluated our model predictions by quantifying relic DNA and assessing**  
21           **its contribution to bacterial diversity using 16S rRNA gene sequences collected from**  
22           **different ecosystem types, including soil, sediment, water, and the mammalian gut. On**  
23           **average, relic DNA made up 33 % of the total bacterial DNA pool, but exceeded 80 % in**  
24           **some samples. Despite its abundance, relic DNA had no effect on estimates of taxonomic**  
25           **and phylogenetic diversity, even in ecosystems where processes such as the physical**  
26           **protection of relic DNA are common and predicted by our models to generate bias. Rather,**  
27           **our findings are consistent with the expectation that relic DNA sequences degrade in**  
28           **proportion to their abundance and therefore may contribute minimally to estimates of**  
29           **microbial diversity.**

30           When microorganisms die, their DNA leaks into the surrounding environment. The fate  
31           of this relic DNA has important implications for evolutionary and ecological processes. For  
32           example, relic DNA can be taken up and incorporated into the genomes of some microorganisms  
33           via transformation, thereby serving as a reservoir of genetic information that can confer new  
34           traits and fitness benefits to distantly related taxa<sup>3</sup>. In addition, relic DNA is a high-quality

resource containing carbon, nitrogen, and phosphorus that is consumed by a diverse array of  
36 bacteria with consequences for microbial community structure and ecosystem processes<sup>1,4</sup>.

Relic DNA also has the potential to bias cultivation-independent estimates of diversity,  
38 which are widely used for addressing questions concerning the assembly, biogeography, and  
functioning of microbial communities. Microbial DNA extracted from environmental and host-  
40 associated samples is not solely derived from metabolically active organisms<sup>5</sup>. A large portion  
of the individuals in a microbial community is dormant or dead<sup>6,7</sup>. Although nucleic acids can be  
42 temporarily retained in non-viable cells, DNA is ultimately released into the environment when  
individuals die from autolysis, senescence, viral infection, or predation<sup>8</sup>. Together, these sources  
44 of mortality can create large pools of relic DNA<sup>2,9</sup>. For example, there is an estimated 0.45  
petagrams of relic DNA in global ocean sediments, which is 70-fold greater than the amount of  
46 DNA contained in intact cells from the same environments<sup>1</sup>. If included in cultivation-  
independent approaches, relic DNA could distort our understanding of the ecological and  
48 evolutionary processes that regulate the distribution, abundance, and function of microbial taxa.

The processes that regulate relic DNA dynamics vary among ecosystems. In well-mixed  
50 microbial habitats, the residence time of relic DNA can be short owing to rapid rates of  
hydrolysis, oxidation, and UV-mediated damage<sup>2</sup>. For example, in surface waters of freshwater  
52 and marine environments, the extracellular DNA pool turns over in less than one day<sup>10,11</sup> while  
plasmid DNA begins to degrade in just minutes<sup>12</sup>. As a result, the size distribution of relic DNA  
54 is skewed towards small fragments ranging between 100 and 500 base pairs<sup>13,14</sup>. In more  
structured microbial habitats, other factors contribute to the size and turnover of the relic DNA  
56 pool. For example, binding with inorganic and organic substances in soils and sediments reduces  
the rate of relic DNA degradation<sup>9</sup>. Likewise, biofilms, aggregates, and outer membrane vesicles

58 can protect relic DNA from hydrolytic enzymes<sup>8,15,16</sup>. Alternatively, the heterogeneous  
distribution of microorganisms in structured habitats creates "hot spots" of metabolic activity<sup>17</sup>  
60 that may increase relic DNA turnover. Collectively, these processes may help explain the  
accumulation of relic DNA in a range of ecosystems including ocean sediments<sup>1</sup>, the built  
62 environment<sup>18</sup>, and the mammalian gastrointestinal tract<sup>19</sup>.

To date, the documented effects of relic DNA on estimates of diversity are idiosyncratic.  
64 Even in samples with large amounts of relic DNA, bias can be non-existent or substantial, and  
can lead to the overestimation or underestimation of diversity<sup>20,21</sup>. Such observations reflect the  
66 need for a more mechanistic understanding of relic DNA dynamics so that microbial  
communities from different ecosystems can be consistently and accurately characterized. We  
68 address this issue by developing a theoretical framework that considers the processes regulating  
the size and turnover of the relic DNA pool. We emphasize that the direction and magnitude of  
70 bias is influenced by sampling from a joint species abundance distribution (SAD) consisting of  
sequences from the relic and intact DNA pools. We evaluated our models by quantifying the  
72 contribution of relic DNA to the abundance and diversity of bacterial communities in ecosystem  
types that are known to have contrasting relic DNA dynamics.

74

## RESULTS

76 We developed an interrelated set of models to identify conditions where relic DNA leads  
to biased estimates of microbial diversity. We began with a conceptual model representing the  
78 dynamics of intact and relic DNA (Fig. 1a). The size and composition of the intact DNA pool  
reflects sequences contained in viable bacteria belonging to different species, which are  
80 influenced by births, deaths, and immigration. The size and composition of the relic DNA pool is

determined by the input of sequences associated with the death of bacteria from the intact DNA  
82 pool and losses associated with the degradation of relic DNA sequences. With this framework  
established, we developed a sampling model to explore how diversity estimates are affected  
84 when draws come from a joint distribution of sequences derived from the intact and relic pools  
(see Methods). We found that a sufficiently large relic DNA pool is required, but not sufficient to  
86 create bias in estimates of species richness, the number of taxa in a sample (Fig. 1b). Critically,  
there must also be differences between the species abundance distributions (SAD) of the intact  
88 and relic DNA pools for bias to arise. When the relic DNA pool has a more even SAD, sampling  
from the total pool (intact + relic) leads to overestimation of true richness (Fig. 1b). Conversely,  
90 when the relic DNA pool has a less even SAD, sampling from the total pool leads to  
underestimation of true richness (Fig. 1b). Last, we developed a process-based model to identify  
92 conditions that lead to relic DNA bias. We determined that relic DNA pool size reaches a stable  
equilibrium when  $R = m I / d$ , where  $R$  is the size (number of sequences) of the relic DNA pool,  
94  $m$  is the per capita mortality rate of bacteria in the intact DNA pool,  $I$  is the size (number of  
sequences) of the intact DNA pool, and  $d$  is the per capita degradation rate of the relic DNA  
96 sequences (see Methods). From this, it can be shown that  $R$  increases with the residence time  
( $\tau = R / d$ ) of the relic DNA pool. We also used the process-based model to explore different  
98 scenarios that influence the SAD of the relic DNA pool. First, we assumed that the rate at which  
relic DNA sequences degrade is equivalent among species. Under this neutral expectation, the  
100 shapes of relic SAD and intact SAD are nearly identical. No bias arises under these conditions,  
regardless of the relic DNA pool size (Fig. 1c). Next, we simulated protection by reducing the  
102 degradation rates of relic DNA sequences belonging to randomly selected species (see Methods).  
This created a less even relic SAD, resulting in the underestimation of richness in the total DNA

104 pool (Fig. 1c). Finally, we considered the scenario where there are "hot spots" of relic DNA  
degradation. When simulating accelerated degradation rates for abundant relic DNA sequences,  
106 there was a more even relic SAD that resulted in the overestimation of richness in the total DNA  
pool (Fig. 1c).

108

**Variation in relic DNA pool size** — We collected samples from different ecosystem types (soil,  
110 sediment, water, and mammalian guts) that are known to vary in relic DNA turnover rates<sup>13,22</sup>.  
Using quantitative PCR with 16S rRNA primers, we quantified intact DNA after treating an  
112 aliquot of sample with DNase to remove relic DNA. In addition, we quantified total DNA (intact  
+ relic) pool size from a control aliquot of sample that was not treated with DNase (see Methods  
114 and Fig. S2). Consistent with previous reports<sup>20,21,23</sup>, relic DNA accounted for a substantial but  
variable portion of the total bacterial DNA pool. The proportion of relic DNA was normally  
116 distributed (mean  $\pm$  sd:  $0.33 \pm 0.218$ ) and ranged from 0 to 0.83 across 34 samples obtained from  
different ecosystem types (Fig. 2). Even though host and environmental features associated with  
118 different habitats are thought to influence relic DNA dynamics, there was only a marginally  
significant effect of ecosystem type on the proportion of relic DNA (one-way ANOVA,  $F_{3,30} =$   
120  $2.43$ ,  $P = 0.084$ )<sup>2,20</sup> which likely reflected the contrast between soil and water samples (Tukey's  
HSD, adjusted  $P = 0.058$ ). Nevertheless, the large average pool size and range in relic DNA  
122 across samples provided us with the opportunity to explore features of our model, specifically  
the magnitude and direction of bias that relic DNA should have on estimates of microbial  
124 diversity.

126 **Magnitude and direction of bias on microbial diversity** — We sequenced 16S rRNA genes  
from the intact and total DNA pools to test whether or not relic DNA biases estimates of within-  
128 sample (alpha) bacterial diversity. Despite accounting for a substantial portion of the total DNA  
(Fig. 2), relic DNA had no effect on estimates of richness, evenness, or phylogenetic diversity  
130 (PD) (Table S1). We expressed each of these alpha-diversity metrics on a per-sample basis as a  
ratio (total / intact), where values > 1 represent overestimation bias and values < 1 represent  
132 underestimation bias. Relic DNA had no effect on the diversity ratios based on the observation  
that the 95 % confidence intervals overlapped with 1.0. Furthermore, the 95 % confidence  
134 intervals of the diversity ratios overlapped across ecosystem types, indicating that the  
contribution of relic DNA to all measures of diversity was generally low irrespective of the  
136 microbial habitat. Finally, using indicator variable multiple regression, we evaluated whether the  
magnitude of bias increased with relic DNA pool size, a predicted outcome from some of our  
138 simulations (Fig. 1b, c). For all diversity ratios (richness, evenness, and PD), the slopes were not  
different from zero ( $P \geq 0.65$ ), the intercepts were not different from 1.0 ( $P \geq 0.65$ ), and the  
140 estimates were not affected by ecosystem type ( $P \geq 0.26$ ) (Fig. S5). Moreover, only a very small  
amount of variation ( $R^2 \leq 0.01$ ) was explained by relic DNA and ecosystem type in the  
142 regression models.

144 **Contribution of relic DNA to community composition** — Based on 16S rRNA gene  
sequencing of the intact and total DNA pools, relic DNA had no effect on the compositional  
146 (beta) diversity of bacterial communities within or across ecosystem types. First, the intact and  
total DNA pools were significantly and very highly correlated with one another when we  
148 performed a taxonomically based (Bray-Curtis distances) Mantel test ( $P = 0.001$ ,  $r = 0.959$ ) and

a phylogenetically based (UniFrac distances) Mantel test ( $P = 0.001$ ,  $r = 0.996$ ). Second, we  
150 tested for the effect of relic DNA on bacterial community composition for the intact and total  
DNA pools using a modified beta-dispersion metric<sup>24</sup>. Specifically, we calculated centroid  
152 distance ratios, which directly compared the dispersion between pairs of DNase-treated and  
DNase-control subsamples (see Methods and Fig. S7). With this approach, if the centroid  
154 distance ratio was  $> 1$ , we concluded that relic DNA inflated beta-diversity; if the distance ratio  
was  $< 1$ , we concluded that relic DNA homogenized beta-diversity. We found that relic DNA  
156 had no effect on the centroid distance ratios based on the observation that the 95 % confidence  
intervals overlapped with 1.0 (Figs. 4, S8). Furthermore, the 95 % confidence intervals for the  
158 centroid distance ratios overlapped across ecosystem types, indicating that the contribution of  
relic DNA to taxonomic and phylogenetic beta-diversity was low irrespective of the microbial  
160 habitat. Last, we used indicator variable multiple regression to test the prediction that bias in  
estimates of community composition would increase with increasing relic DNA pool size (see  
162 Fig. S1). The proportion of relic DNA in a sample had no effect on the slopes for centroid  
distance ratios regardless of whether they were calculated using taxonomic ( $P = 0.63$ ) or  
164 phylogenetic ( $P = 0.59$ ) distance matrices. Moreover, the intercepts for these regression  
relationships were not different from 1.0 ( $P \geq 0.74$ ) and the amount of explained variation was  
166 low ( $R^2 \leq 0.09$ ) suggesting that the overall effect of relic DNA on beta diversity was negligible  
(Fig. S8). The parameters estimates from the regression analyses were not affected by  
168 ecosystems type ( $P \geq 0.29$ ).

170

## DISCUSSION



Rarely, if ever, are biological communities completely censused. As a result, estimates of  
172 diversity are often based on incomplete sampling, which introduces uncertainty and potential  
bias<sup>25</sup>. For example, microbiologists using cultivation-independent approaches commonly  
174 estimate the diversity of a community using hundreds of thousands of sequences from samples  
that contain in excess of a billion individuals. Diversity estimation of microbial communities is  
176 further complicated by the accumulation and persistence of relic DNA, a pool of sequences that  
may not accurately reflect the composition of viable microorganisms. We developed a set of  
178 models that capture the fundamental processes regulating relic DNA dynamics along with the  
effects of sampling from a joint species abundance distribution (SAD) that contain sequences  
180 from both the relic and intact DNA pools. Our models reveal that two criteria are required in  
order for relic DNA bias to emerge. First, the relic DNA pool must be large enough so that it  
182 diminishes the probability of sampling sequences from the intact DNA pool. Second, the relic  
SAD must be distinct from the intact SAD. If these conditions are not satisfied, relic DNA has  
184 minimal effect on estimates of biodiversity. We then explored the model's expectations by  
analyzing the intact and total (relic + intact) DNA in ecosystems where relic DNA dynamics are  
186 known to vary. Despite making up a large portion of the total DNA, we found that relic DNA  
had minimal to no effect on estimates taxonomic and phylogenetic diversity, consistent with  
188 model scenarios where relic DNA degradation rates are equivalent among species.

Despite sampling a range of habitats with attributes that affect relic DNA turnover, our  
190 data suggest that estimates of taxonomic and phylogenetic diversity were unbiased by relic DNA  
(Figs. 2 and 3) even when it accounted for > 80 % of the total DNA (Figs. S5 and S8). Similar  
192 findings have been reported elsewhere. From a survey of North American soils, no bias was  
detected for taxonomic richness in 30 % of bacterial samples (7 / 31) and 55 % of fungal samples

194 (17 / 31), even when relic DNA made up a substantial portion of the total DNA pool <sup>21</sup>.

Likewise, more than 90 % of the bacterial DNA recovered from porcine lung tissues was  
196 sensitive to DNase, yet diversity in the enzymatically treated and untreated samples was the  
same indicating that large relic DNA pools do not necessarily obscure estimates of microbial  
198 diversity <sup>20</sup>. Such findings are supported by our modeling results: relic DNA has no effect on  
estimates of microbial diversity when the SADs of the relic and intact DNA pools are equivalent  
200 (Fig. 1b). When this assumption is met, the abundance distribution of relic and intact DNA pools  
should be identical. Therefore, sampling from the relic DNA pool should not bias estimates of  
202 diversity in the intact DNA pool, even if relic DNA makes up a large portion of the total DNA  
pool (Fig. 1b).

204 Our simulations identify other ecological scenarios where relic DNA bias may arise. It  
has been hypothesized that relic DNA can be protected from degradation inside of aggregates,  
206 biofilms, or other complexes that reduce contact between nucleic acids and extracellular  
enzymes. Importantly, protection can only increase bias if it alters relic DNA degradation in a  
208 species-specific manner. While most studies have emphasized the potential for protection to  
overestimate diversity, our model predicts the opposite (Fig. 1b). Protection created a less even  
210 relic SAD that resulted in the increased dominance of relic sequences associated with a few  
species. Assuming a large relic DNA pool, preferential sampling of these protected sequences  
212 will lead to the underestimation of richness, which has been observed in some instances <sup>20</sup>. In  
contrast, we identified at least one ecological scenario where relic DNA creates overestimation  
214 bias. Microscale heterogeneity in structured habitats can lead to non-uniform distributions of  
microorganisms and their metabolic activities. Our simulations indicate that the resulting "hot  
216 spots" can degrade relic DNA sequences in a density-dependent manner resulting in a more even

relic SAD. Although not significant, sediment samples trended towards positive bias for richness  
218 and PD (Fig. 3), which could potentially reflect "hot spots" of relic DNA degradation. Inflation  
bias can also arise when species in the regional pool disperse into habitats for which they are  
220 poorly adapted. When these immigrants die, they can enrich the relic DNA pool with sequences  
that are dissimilar to sequences found in the local community. A similar effect could arise when  
222 dead bacteria are transported across ecosystem boundaries, a phenomenon that occurs, for  
example, when marine snow is exported from surface waters to marine sediments<sup>26</sup>. More  
224 generally, bias may arise under non-equilibrium conditions where community turnover of the  
intact DNA pool is shorter than turnover of the relic DNA pool. Any abiotic or biotic  
226 perturbation that removes a substantial amount of living biomass could result in transient  
divergence in the composition of sequences in the intact and relic DNA pool. For example,  
228 virulent phage can dramatically reduce the abundance of bacterial prey and in the process release  
a large pulse of DNA into the environment, which could bias estimates of microbial diversity<sup>27</sup>.  
230 Last, although not fully explored here, shifts in rank abundance of taxa that are independent of  
parameters describing the SAD could also result in biased estimates of microbial diversity. In  
232 sum, there are ways to deviate from neutral expectations about the degradation of relic DNA,  
which should lead to biased estimates of microbial diversity, but sampling from a range of  
234 ecosystems suggests that such conditions are not prevalent in nature.

Our understanding of the microbial biosphere has been transformed by the development  
236 and application of molecular-based cultivation-independent techniques. The ability to rapidly  
obtain millions of gene sequences and transcripts from a range of environments has yielded  
238 valuable insight into the processes that regulate community assembly and function<sup>28,29</sup>, and has  
also paved the way for the discovery of new metabolisms<sup>30</sup>, tests for unifying patterns of

240 biodiversity<sup>31</sup>, and an updated tree of life<sup>32</sup>. There are limitations, however, associated with  
culture-independent techniques, which include inefficient nucleic-acid extraction methods<sup>32</sup> and  
242 "universal" primers that over-represent some taxonomic groups while overlooking others<sup>33</sup>.  
Sequencing of relic DNA is another important concern, which can potentially lead to the  
244 overestimation or underestimation of microbial diversity. However, this bias requires the  
decoupling of processes that regulate the compositional turnover of the relic and intact DNA  
246 pools. While some recent evidence suggests this can arise<sup>20</sup>, our findings suggest that at least in  
some ecosystems, relic DNA appears to contribute minimally to the characterization of microbial  
248 community structure.

250

## METHODS

252 **Sampling Model** — We used a set of sampling-based simulations to explore the effects of  
mixing intact and relic DNA on estimates of diversity. For each simulation, we defined a  
254 regional species pool consisting of 10,000 taxa with a lognormal abundance distribution<sup>34</sup>. The  
intact community consisted of 1,000,000 individuals sampled from this regional pool. We then  
256 combined this intact community with a relic community at proportions ranging from 0.01 to  
0.96. We altered the evenness of the regional pool from which the relic community was sampled  
258 by changing the scale parameter of the lognormal distribution. To decrease the evenness of the  
relic DNA pool, we increased the scale parameter from 0.98 to 1.8. To increase the evenness of  
260 the relic DNA pool, we decreased the parameter from 0.98 to 0.25. After mixing the intact and  
relic communities at the defined proportions, we rarified the total community to 10,000  
262 observations and calculated richness and Bray-Curtis distances to estimate compositional  
differences between the intact and total DNA pools. To estimate contribution of relic DNA to

264 diversity, we calculate ratios (total DNA / intact DNA) for richness and Bray-Curtis distances.

All simulations and estimations were performed in the R statistic environment (v 3.3.2)<sup>35</sup> using  
266 the `vegan` package as well as custom functions.

268 **Process-Based Model** — We developed a set of stochastic simulations to explore how the  
processes regulating relic DNA dynamics can give rise to bias in the estimation of microbial  
270 diversity. For each simulation, we defined a regional species pool consisting of 4,000 taxa with a  
lognormal abundance distribution<sup>34</sup>. We initiated an intact community by randomly sampling  
272 20,000 individuals from the regional species pool. Subsequent dynamics were controlled by the  
processes depicted in our conceptual model (Fig. 1a). Specifically, we simulated immigration by  
274 sampling  $j$  individuals from the regional species pool and adding them to the intact community.  
We simulated birth by randomly selecting  $I \times r$  individuals from the intact community and  
276 adding them to the intact community again, where  $I$  is the size of the intact community and  $r$  is  
the per capita birth rate. We simulated death by selecting  $(I \times m) + j$  individuals from the intact  
278 community and moving them to the relic community ( $R$ ), where  $m$  is the per capita mortality  
rate. We simulated degradation by selecting  $(R \times d) + j$  individuals from the relic community and  
280 removing them, where  $d$  is the per capita decay rate of relic DNA.

We simulated three scenarios with the process-based model. First, we considered a  
282 neutral scenario. At each time step, we simulated degradation by randomly selecting relic DNA  
sequences and removing them. Second, we considered the scenario where some relic DNA  
284 sequences have lower rates of degradation owing to protection. Each species was randomly  
assigned a degradation susceptibility probability from a beta distribution. The beta distribution is  
286 a continuous probability distribution bound by 0 and 1 with two shape parameters. We chose

shape parameters ( $\alpha = 0.7$ ,  $\beta = 0.7$ ) to provide a wide range in susceptibility probability but also  
288 have increased probabilities of low (near 0) and high (near 1) susceptibility, which created a  
wide U-shaped distribution. Susceptibility probabilities were randomly assigned to species for  
290 each model iteration and were used to weight the probability of selecting relic DNA sequences  
for degradation at each time step. Last, we considered the scenario where there are "hot spots" of  
292 relic DNA degradation, reflecting the clumped distribution of microbial taxa and their associated  
metabolic activities. At each time step, we calculated density-dependent selection probabilities  
294 by weighting the probability of selecting an individual from the relic community by its species  
abundance. Therefore, relic DNA sequencing belonging to more abundant species would be  
296 more likely to be selected for degradation. For each of the three scenarios, we ran each  
simulation for 10,000 time-steps. We used a constant immigration rate ( $j = 2,000$ ) to maintain  
298 intact community diversity, and used equal and constant birth and mortality rates ( $b$  and  $m = 0.1$ )  
to maintain intact community size. Likewise, we accounted for immigration during death and  
300 degradation to prevent uncontrolled growth. For each set of simulations, we used a range of  
decay rates ( $d$ ), which yielded relic DNA proportions between 0.05 and 0.95 according to the  
302 following equation:  $d = m/p - m$ , where  $p$  is the target proportion and  $m$  is the fixed mortality  
rate (see Supplement for derivation). At the end of each simulation, we determined the effect of  
304 relic DNA on diversity estimates by comparing the total community (intact + relic) with the  
intact community. All simulations were performed in the R statistic environment.

306

**Sample collection and DNA pools** — We collected samples from a diverse set of environmental  
308 and host-associated ecosystems. First, we sampled sediments and surface water from lakes near  
the Michigan State University, W.K. Kellogg Biological Station (KBS) in Hickory Corners,

310 Michigan, USA. Soils were sampled from the main sites and surrounding areas at the KBS Long-  
Term Ecological Research site<sup>36</sup>. We also collected fresh feces as representative gut samples  
312 from cows, dogs, horses, rabbits, and humans. In each of these ecosystem types, we obtained  
samples from 6 - 8 independent sites. In the laboratory, we applied a DNase treatment to each  
314 sample to remove relic DNA. The procedure is based on methods that have previously been used  
to quantify relic DNA in marine sediments<sup>37,38</sup>, host tissue<sup>37,38</sup>, and drinking-water biofilms  
316<sup>37,38</sup>. The DNase-based method has proven to be effective not only for removing extracellular  
DNA, but also DNA that is contained inside of dead cells<sup>37,38</sup> while not compromising the  
318 integrity of living cells<sup>37,38</sup>. We determined that the DNase method is efficient (98%) at  
removing relic DNA that was experimentally added to an environmental sample (see  
320 Supplementary Methods; Fig. S4). Thus, we refer to the DNA remaining in a subsample  
following enzymatic treatment as "intact DNA" and assume it is derived from viable cells. "Total  
322 DNA" refers to the DNA in subsample that was not treated with DNase (negative control), which  
reflects the sum of intact DNA and relic DNA.

324 For aquatic samples, we filtered 250 mL lake water through a 47 mm 0.2  $\mu$ m Supor PES  
membrane filter using 10 mm Hg vacuum. We cut the filter in half and randomly assigned one  
326 half of the filter to a DNase treatment and used the other half as the control. We then rolled up  
each piece of filter using sterile forceps and inserted it into a 2 mL centrifuge tube containing 1.5  
328 mL of pH 7.3 phosphate buffered saline (PBS). After vortexing at room temperature for 5 min,  
we removed the filter from the tube and centrifuged for 5 min at 10,000 x g. We then discarded  
330 the supernatant and resuspended the pellet in 375  $\mu$ L of PBS, which was transferred to a 2 mL  
centrifuge tube. For non-aqueous samples (soil, sediments, and feces), we directly added 0.25 g  
332 of material to a 2 mL centrifuge tube. At this stage in the procedure, aquatic and non-aquatic

samples were identically processed. For each sample in its 2 mL centrifuge tube, we added  
334 DNase digestion buffer, which consisted of 382.5  $\mu\text{L}$  of nanopure water, 5  $\mu\text{L}$  of 1 M  $\text{MgCl}_2$ , 2.5  
 $\mu\text{L}$  of bovine serum albumin (10 mg/mL), and 50  $\mu\text{L}$  of 0.5 M Tris HCl (pH 7.5). For  
336 subsamples treated with DNase, we added 40  $\mu\text{L}$  of a 10 U/ $\mu\text{L}$  stock of DNase I (Roche #  
04536282001) and 20  $\mu\text{L}$  of nanopure water resulting in a 500  $\mu\text{L}$  final working volume with an  
338 80U/ $\mu\text{L}$  DNase concentration. (See Supplementary Material and Fig. S3 for description of  
experiment to determine minimum DNase concentration needed for relic DNA removal.) For  
340 untreated subsamples, we substituted 40  $\mu\text{L}$  of nanopure water for DNase solution. For each  
sample, we measured pH using a micro pH probe (Orion 9110DJWP, Thermo Scientific) and  
342 adjusted to 7.3 - 7.7, which is in the optimum range for DNase. We then incubated the samples  
horizontally on a shaker table at 37 °C for 60 min. Following this, we transferred each sample to  
344 a 15 mL Falcon tube containing 1 mL of 1X hexadecyltrimethylammonium bromide (CTAB)  
buffer, which consisted of one part solution 1 (1 g CTAB + 0.58 g NaCl in 10 mL of nanopure  
346 water) and one part solution 2 (0.58 g NaCl + 0.82  $\text{K}_2\text{HPO}_4$  + 0.04  $\text{KH}_2\text{PO}_4$  in 10 mL of  
nanopure water). We then added 25  $\mu\text{L}$  of 0.5 M EDTA to each tube and vortexed. We stopped  
348 the DNase reaction by incubating the tubes at 75 °C for 10 min. We began DNA extraction by  
adding 1 mL of phenol:chloroform:isoamylalcohol (25:24:1) to the 2 mL tube, which was then  
350 vortexed for 10 min. This was followed by centrifuging at 7,100 x g for 10 min. We transferred  
the top aqueous layer to a 15 mL Falcon tube and combined it with an equal volume of  
352 chloroform:isoamylalcohol (24:1) and then vortexed. After centrifuging the Falcon tubes at  
7,100 x g for 5 min, we transferred 400  $\mu\text{L}$  of the top aqueous layer to a new 2 mL centrifuge  
354 tube. For clean up, we used the MO BIO PowerLyzer PowerSoil DNA Isolation Kit starting from  
Step 9. At Step 21 of the DNA Isolation Kit, we eluted in 50  $\mu\text{L}$  of Solution C6 and centrifuged



356 for 1 min at 10,000 x g. We stored extracted DNA at -20 °C for short-term storage or at -80 °C  
for long-term storage.

358 **Contribution of relic DNA to bacterial abundance** — We used 16S rRNA gene copy numbers

360 generated from quantitative PCR (qPCR) assays to estimate the proportion of relic DNA in a  
sample as  $1 - (\text{intact DNA} / \text{total DNA})$ . Briefly, the previously described qPCR assays<sup>39</sup>

362 consisted of 30  $\mu\text{L}$  reactions containing 1  $\mu\text{L}$  of DNA template, 0.5  $\mu\text{L}$  of each primer (10  
 $\mu\text{mol/L}$ ), 14.5  $\mu\text{L}$  of nuclease-free  $\text{H}_2\text{O}$ , and 13.5  $\mu\text{L}$  of 5 Prime 2.5x RealMasterMix SYBR

364 ROX. We amplified a 200 base-pair fragment of the 16S rRNA gene with Eub 338 (forward)  
and Eub518 (reverse) primers<sup>40</sup>. PCR assays were performed with an Eppendorf Mastercycler

366 Realplex<sup>2</sup> system using previously reported thermal cycle conditions<sup>40</sup>. We generated qPCR  
standards from bacterial genomic DNA (*Micrococcus* sp.) using the TOPO TA Cloning Kit

368 (Invitrogen). We extracted plasmids from transformed cells<sup>41</sup>, and used the M13 forward and  
reverse primers to generate PCR products. The PCR products were quantified and used to

370 generate a standard curve capturing a range of  $10^2 - 10^7$  gene copies/ $\mu\text{L}$ . We used a noiseband  
threshold for quantification. The coefficients of determination ( $r^2$ ) for our assays ranged from

372 0.96 and 0.99, while amplification efficiencies fell between 0.93 and 0.99. Based on melting  
curve analyses, we found no evidence for primer dimers. All unknown samples, no template

374 controls, and standards were run in triplicate on every plate. The mean coefficient of variation  
(standard deviation / mean) for our 16S rRNA qPCR assay was 0.16.

376

### **Contribution of relic DNA to bacterial diversity**

378 *Community sequencing* — We estimated the contribution of relic DNA to bacterial diversity  
using high-throughput sequencing of the 16S rRNA gene. Specifically, we amplified the V4

380 region of the 16S rRNA gene from the intact and total DNA pools of each sample using  
barcoded primers (515F and 806R) designed to work with the Illumina MiSeq platform<sup>42</sup>. We  
382 cleaned the sequence libraries using the AMPure XP purification kit, quantified the resulting  
products using the QuantIt PicoGreen kit (Invitrogen), and pooled libraries at equal molar ratios  
384 (final concentration: 20 ng per library). We then sequenced the pooled libraries with the Illumina  
MiSeq platform using paired end reads (Illumina Reagent Kit v2, 500 reaction kit) at the Indiana  
386 University Center for Genomics and Bioinformatics Sequencing Facility. Paired-end raw 16S  
rRNA sequence reads were assembled into contigs using the Needleman algorithm<sup>43</sup>. We  
388 obtained a total of 12,916,632 16S rRNA sequences. After quality trimming with a moving  
average quality score (window 50 bp, minimum quality score 35), we aligned the sequences to  
390 the Silva Database (version 123) using the Needleman algorithm. Chimeric sequences were  
removed using the UCHIME algorithm<sup>44</sup>. After this filtering, there was an average ( $\pm$  SEM) of  
392  $222,701 \pm 9,560$  sequences per site. We created operational taxonomic units (OTUs) by first  
splitting the sequences based on taxonomic class (using the RDP taxonomy) and then binning  
394 sequences into OTUs based on 97% sequence similarity. Our depth of sequencing led to a high  
degree of coverage across samples (minimum Good's Coverage = 0.98). For phylogenetic  
396 analysis, we picked representative sequences for each OTU by using the most abundant unique  
sequence. We used FastTree<sup>45</sup> to generate a phylogenetic tree from the representative sequences  
398 using the generalized time-reversible model of nucleotide evolution. We calculated phylogenetic  
distances using weighted UniFrac distances<sup>46</sup>. All initial sequence processing was completed  
400 using the software package mothur (version 1.38.1)<sup>47</sup>.

402 *Alpha Diversity* — We estimated the effects of relic DNA on richness, evenness, and  
phylogenetic diversity for the intact and total DNA pools with a sample. To estimate the number

404 of OTUs (richness), we used a resampling approach that subsampled each sample to an equal  
number of sequences per sample and summed the number of OTUs that were represented<sup>48</sup>.

406 Briefly, we subsampled to 30,000 observations, resampled 999 additional times, and then  
calculated the average richness estimates ( $\pm$  SEM) for each sample. To estimate the equitability

408 in abundance among taxa in a sample (evenness), we used the same resampling approach and  
calculated average evenness estimates ( $\pm$  SEM) using Simpson's Evenness index<sup>49</sup>. To test

410 whether relic DNA affected the phylogenetic diversity within a sample, we subsampled  
communities to 30,000 observations and then calculated Faith's *D* statistic, which sums the

412 branch lengths for each species found in a sample from the root to the tip of the phylogenetic tree<sup>50</sup>. All estimations were performed in the R statistic environment (v 3.3.2)<sup>35</sup> using the *vegan*, *ape*,  
414 *ade4*, *picante*, and *plyr* packages, along with custom functions.

416 *Beta Diversity* — We estimated the effects of relic DNA on between-sample (beta) diversity by  
comparing the taxonomic and phylogenetic diversity of bacterial communities in the intact and

418 total DNA pools. First, we conducted a Principal Coordinates Analysis (PCoA) on log<sub>10</sub>-  
transformed relative abundance data to visualize the effects of relic DNA removal (via DNase

420 treatment) on bacterial community composition within and among ecosystem types. The PCoA  
was performed with Bray-Curtis and UniFrac distances to assess taxonomic and phylogenetic

422 effects, respectively. In addition, we used PERMANOVA to test for differences in taxonomic  
and phylogenetic composition based on ecosystem type for the total DNA pool. Second, we

424 conducted a Mantel test to assess the correlation between the community resemblance matrices  
(either Bray-Curtis or UniFrac) represented by the intact and total DNA pools. Last, we tested

426 whether relic DNA altered beta-diversity within an ecosystem type by comparing centroid  
distances. To calculate this metric of sample dispersion, we determined the centroid from a

428 PCoA with either Bray-Curtis or UniFrac distances for the total DNA pool for all sites within a  
given ecosystem type. We then measured the Euclidean distances between the centroid and all  
430 samples (total and intact) to determine the centroid distances (see Fig. S7).

## 432 **ACKNOWLEDGMENTS**

We acknowledge constructive feedback from KD Webster, RZ Moger-Reischer, EK Hall, NI  
434 Wisnoski, and WR Shoemaker. We thank S. Kuenzel and B.K. Lehmkuhl for technical support.  
This work was supported by the USDA National Institute of Food and Agriculture Grant 2011-  
436 67019-3022 (JTL), the National Science Foundation Dimensions of Biodiversity Grant 1442246  
(JTL), and US Army Research Office Grant W911NF-14-1-0411 (JTL). All code and data used  
438 in this study can be found in a public GitHub repository  
(<https://www.github.com/LennonLab/relicDNA>) and the NCBI SRA.

440

442

## REFERENCES

- 1 Dell'Anno, A. & Danovaro, R. Extracellular DNA plays a key role in deep-sea ecosystem  
444 functioning. *Science* **309**, 2179-2179 (2005).
- 2 Torti, A., Lever, M. A. & Jorgensen, B. B. Origin, dynamics, and implications of  
446 extracellular DNA pools in marine sediments. *Mar. Genomics* **24**, 185-196 (2015).
- 3 Redfield, R. J. Genes for breakfast - the have-your-cake-and-eat-it-too of bacterial  
448 transformation. *J. Hered.* **84**, 400-404 (1993).
- 4 Lennon, J. T. Diversity and metabolism of marine bacteria cultivated on dissolved DNA.  
450 *Appl. Environ. Microbiol.* **73**, 2799-2805 (2007).
- 5 Klein, D. A. Microbial communities in nature: a postgenomic perspective. *Microbe* **2**,  
452 591-595 (2007).
- 6 Luna, G. M., Manini, E. & Danovaro, R. Large fraction of dead and inactive bacteria in  
454 coastal marine sediments: Comparison of protocols for determination and ecological  
significance. *Appl. Environ. Microbiol.* **68**, 3509-3513 (2002).
- 456 7 Lennon, J. T. & Jones, S. E. Microbial seed banks: the ecological and evolutionary  
implications of dormancy. *Nat. Rev. Microbiol.* **9**, 119-130 (2011).
- 458 8 Vorkapic, D., Pressler, K. & Schild, S. Multifaceted roles of extracellular DNA in  
bacterial physiology. *Curr. Genet.* **62**, 71-79 (2016).
- 460 9 Levy-Booth, D. J. *et al.* Cycling of extracellular DNA in the soil environment. *Soil Biol.*  
*Biochem.* **39**, 2977-2991 (2007).
- 462 10 Paul, J. H., Jeffrey, W. H., David, A. W., Deflaun, M. F. & Cazares, L. H. Turnover of  
464 extracellular DNA in eutrophic and oligotrophic freshwater environments of southwest  
Florida. *Appl. Environ. Microbiol.* **55**, 1823-1828 (1989).

- 466 11 Dell'Anno, A. & Corinaldesi, C. Degradation and turnover of extracellular DNA in  
marine sediments: Ecological and methodological considerations. *Appl. Environ.*  
468 *Microbiol.* **70**, 4384-4386 (2004).
- 12 Deflaun, M. F. & Paul, J. H. Detection of exogenous gene-sequences in dissolved DNA  
470 from aquatic environments. *Microb. Ecol.* **18**, 21-28 (1989).
- 13 Deflaun, M. F., Paul, J. H. & Jeffrey, W. H. Distribution and molecular-weight of  
472 dissolved DNA in subtropical estuarine and oceanic environments. *Mar. Ecol. Prog. Ser.*  
**38**, 65-73 (1987).
- 474 14 Beebee, T. J. C. Identification and analysis of nucleci acids in natural freshwaters. *Sci.*  
*Total. Env.* **135**, 123-129 (1993).
- 476 15 Romanowski, G., Lorenz, M. G. & Wackernagel, W. Adsorption of plasmid DNA to  
mineral surfaces and protection against DNase-1. *Appl. Environ. Microbiol.* **57**, 1057-  
478 1061 (1991).
- 16 Schwechheimer, C. & Kuehn, M. J. Outer-membrane vesicles from Gram-negative  
480 bacteria: biogenesis and functions. *Nat. Rev. Microbiol.* **13**, 605-619 (2015).
- 17 DeAngelis, K. M., Lindow, S. E. & Firestone, M. K. Bacterial quorum sensing and  
482 nitrogen cycling in rhizosphere soil. *FEMS Microbiol. Ecol.* **66**, 197-207 (2008).
- 18 Emerson, J. B. *et al.* Schrodinger's microbes: Tools for distinguishing the living from the  
484 dead in microbial ecosystems. *Microbiome* **5**, 86 (2017).
- 19 Sun J, Wang P. & Shi, Y. High fat diet decrease diversity of extracellular DNA of mice  
486 gut microbiota. *FASEB J* **30**, ib204 (2016).

- 20 Pezzulo, A. A. *et al.* Abundant DNase I-Sensitive Bacterial DNA in Healthy Porcine  
488 Lungs and Its Implications for the Lung Microbiome. *Appl. Environ. Microbiol.* **79**,  
5936-5941 (2013).
- 490 21 Carini, P. *et al.* Relic DNA is abundant in soil and obscures estimates of soil microbial  
diversity. *Nat. Microbiol.* **2**, 16242 (2016).
- 492 22 Morrissey, E. M. *et al.* Dynamics of extracellular DNA decomposition and bacterial  
community composition in soil. *Soil Biol. Biochem.* **86**, 42-49 (2015).
- 494 23 Corinaldesi, C., Danovaro, R. & Dell'Anno, A. Simultaneous recovery of extracellular  
and intracellular DNA suitable for molecular studies from marine sediments. *Appl.*  
496 *Environ. Microbiol.* **71**, 46-50 (2005).
- 24 Anderson, M. J., Ellingsen, K. E. & McArdle, B. H. Multivariate dispersion as a measure  
498 of beta diversity. *Ecol. Lett.* **9**, 683-693 (2006).
- 25 Magurran, A. E. *Measuring Biological Diversity.* (Wiley-Blackwell, 2004).
- 500 26 Azam, F. & Long, R. A. Oceanography - Sea snow microcosms. *Nature* **414**, 497-498  
(2001).
- 502 27 Corinaldesi, C., Tangherlini, M., Luna, G. M. & Dell'Anno, A. Extracellular DNA can  
preserve the genetic signatures of present and past viral infection events in deep  
504 hypersaline anoxic basins. *Proc. R. Soc. Lond. Biol.* **281** (2014).
- 28 Lindstrom, E. S. & Langenheder, S. Local and regional factors influencing bacterial  
506 community assembly. *Environ. Microbiol. Rep.* **4**, 1-9 (2012).
- 29 Aanderud, Z. T., Jones, S. E., Fierer, N. & Lennon, J. T. Resuscitation of the rare  
508 biosphere contributes to pulses of ecosystem activity. *Front. Microbiol.* **6** (2015).

- 30 Beja, O. *et al.* Bacterial rhodopsin: Evidence for a new type of phototrophy in the sea.  
510 *Science* **289**, 1902-1906 (2000).
- 31 Locey, K. J. & Lennon, J. T. Scaling laws predict global microbial diversity. *Proc. Natl.*  
512 *Acad. Sci. U.S.A.* **113**, 5970-5975 (2016).
- 32 Feinstein, L. M., Sul, W. J. & Blackwood, C. B. Assessment of bias associated with  
514 incomplete extraction of microbial DNA from soil. *Appl. Environ. Microbiol.* **75**, 5428-  
5433 (2009).
- 516 33 Klindworth, A. *et al.* Evaluation of general 16S ribosomal RNA gene PCR primers for  
classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res.* **41**,  
518 e1 (2013).
- 34 Shoemaker, W. R., Locey, K. J. & Lennon, J. T. A macroecological theory of microbial  
520 biodiversity. *Nat Ecol Evol* **1**, 107 (2017).
- 35 R Core Development Team R: A language and environment for statistical computing,  
522 reference index version 2.8.1. R Foundation for Statistical Computing, Vienna, Austria.  
ISBN 3-900051-07-0, URL <http://R-project.org>. (2009).
- 524 36 Hamilton, S. K., Doll, J. E. & Robertson, G. P. *The ecology of agricultural landscapes:  
Long-term research on the path to sustainability*. (Oxford University Press, 2015).
- 526 37 Dell'Anno, A., Bompadre, S. & Danovaro, R. Quantification, base composition, and fate  
of extracellular DNA in marine sediments. *Limnol. Oceanogr.* **47**, 899-905 (2002).
- 528 38 Villarreal, J. V., Jungfer, C., Obst, U. & Schwartz, T. DNase I and Proteinase K eliminate  
DNA from injured or dead bacteria but not from living bacteria in microbial reference  
530 systems and natural drinking water biofilms for subsequent molecular biology analyses.  
*J. Microbiol. Methods* **94**, 161-169 (2013).



- 532 39 Lau, J. A. & Lennon, J. T. Evolutionary ecology of plant-microbe interactions: soil  
microbial structure alters selection on plant traits. *New Phytol.* **192**, 215-224 (2011).
- 534 40 Fierer, N., Jackson, J. A., Vilgalys, R. & Jackson, R. B. Assessment of soil microbial  
community structure by use of taxon-specific quantitative PCR assays. *Appl. Environ.*  
536 *Microbiol.* **71**, 4117-4120 (2005).
- 41 Sambrook, J. & Russell, D. W. *Molecular cloning: a laboratory manual.* (Cold Spring  
538 Harbor Laboratory Press,, 2001).
- 42 Caporaso, J. G. *et al.* Ultra-high-throughput microbial community analysis on the  
540 Illumina HiSeq and MiSeq platforms. *ISME J.* **6**, 1621-1624 (2012).
- 43 Needleman S.B. & C.D., W. A general method applicable to the search for similarities in  
542 the amino acid sequence of two proteins. *J. Mol. Biol.* **48**, 443-453 (1970).
- 44 Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C. & Knight, R. UCHIME improves  
544 sensitivity and speed of chimera detection. *Bioinformatics* **27**, 2194-2200 (2011).
- 45 Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2-Approximately Maximum-  
546 Likelihood Trees for Large Alignments. *PLOS ONE* **5** (2010).
- 46 Lozupone, C. A. & Knight, R. Species divergence and the measurement of microbial  
548 diversity. *FEMS Microbiol. Rev* **32**, 557-578 (2008).
- 47 Schloss, P. D. *et al.* I Introducing mothur: open-source, platform-independent,  
550 community-supported software for describing and comparing microbial communities.  
*Appl. Environ. Microbiol.* **75**, 7537-7541 (2009).
- 552 48 Muscarella, M. E., Jones, S. E. & Lennon, J. T. Species sorting along a subsidy gradient  
alters community stability. *Ecology* (2016).

- 554 49 Smith, B. & Wilson, J. B. A consumer's guide to evenness indices. *Oikos* **76**, 70-82  
(1996).
- 556 50 Faith, D. P. onservation evaluation and phylogenetic diersity. *Biol. Conserv.* **61**, 1-10,  
(1992).
- 558

560

## FIGURE CAPTIONS

562 **Figure 1 | Modeling relic DNA dynamics.** **a** The amount of relic DNA in a microbial  
environment is determined by inputs associated with the mortality of viable individuals with  
564 intact DNA and by losses associated with the degradation of relic DNA. If the diversity of  
sequences contained in the relic DNA pool is sufficiently different from that in the intact DNA  
566 pool, then relic DNA may bias estimates of microbial biodiversity (as indicated by different  
colored boxes) when sampling from the total (intact + relic) DNA pool. **b** We developed a  
568 sampling-based simulation model to explore the effects of mixing intact and relic DNA on  
estimates of diversity. We populated intact and relic communities with individuals from a  
570 lognormal species abundance distribution (SAD). We altered the diversity of the relic  
community by changing the scale parameter of the lognormal distribution describing the SAD.  
572 We then sampled and mixed the intact and relic communities so that the relic contribution to  
total community ranged from 0.01 to 0.96. **c** To gain mechanistic insight into how bias arises, we  
574 created a stochastic process-based model that captures features that influence relic DNA  
dynamics, including: immigration, birth, death, degradation (**a**). We simulated a range of  
576 degradation rates to achieve relic DNA pool sizes with proportions ranging between 0.05 and  
0.95. To explore how degradation alters the SAD of the relic community, we explored three  
578 scenarios. First, we simulated a neutral scenario where relic DNA sequences produced by  
different species degrade at the same rate. Second, we simulated conditions where physical,  
580 chemical, or biological processes slow the degradation rate of relic DNA belonging to some  
species via protection. Third, we simulated "hot spots" where there more abundant relic DNA  
582 sequences experience faster rates of relic DNA degradation, a condition that may arise in

structured habitats where there are patchy distributions of individuals and their metabolic  
584 products (i.e., enzymes). We ran simulations for 10,000 time steps and then sampled the intact  
and relic communities. To quantify bias in diversity (**b**, **c**), we calculated "richness ratios" which  
586 reflect the number of species in the total DNA pool (intact + relic) divided by the number of  
species in the intact DNA pool in a simulation. When values richness ratios = 1, relic DNA has  
588 no effect on estimates of diversity; when richness ratios > 1, relic DNA overestimates true  
diversity; and when richness ratios < 1, relic DNA underestimates true diversity.

590

**Figure 2 | Proportion of bacterial relic DNA in different ecosystem types.** We quantified the  
592 amount of intact DNA in a sample after removing relic DNA with a DNase treatment. We then  
estimated the proportion of relic DNA as  $1 - (\text{intact DNA} / \text{total DNA})$ , where the total DNA  
594 concentration was quantified without DNase treatment. Relic DNA constituted an appreciable  
portion of the total DNA pool, but was not affected by the ecosystem type from which the  
596 sample was collected (gut, soil, sediment, and water). Grey symbols are the observed data and  
black symbols represent means  $\pm$  95% confidence intervals.

598


**Figure 3 | Bias of relic DNA on within-sample bacterial diversity in different ecosystem  
600 types. a-c** We tested for the effects of bias caused by relic DNA by calculating diversity ratios  
for (**a**) richness, (**b**) evenness, and (**c**) phylogenetic diversity. The ratios reflect the diversity of  
602 the total DNA pool (intact + relic) divided by the diversity of the intact DNA pool. Relic DNA  
did not bias any measures of diversity in any of the ecosystem types. Richness was calculated as  
604 the number of operational taxonomic units (97% sequence similarity of the 16S rRNA gene),  
evenness was calculated using Simpson's evenness index, and phylogenetic diversity was

606 calculated using Faith's *D* index. Grey symbols are the observed data and black symbols  
represent means  $\pm$  95% confidence intervals.

608


**Figure 4 | Bias of relic DNA on the among-sample bacterial diversity in different ecosystem types. a-b** We tested for the effects of bias caused by relic DNA by calculating a beta-diversity ratio based on centroid distances. Centroid distances were estimated after performing Principle Coordinates Analyses (PCoA) using taxonomic (**a**) and phylogenetic (**b**) distance metrics (Bray-Curtis and UniFrac, respectively). The centroid distance ratio was calculated on each sample within an ecosystem type and reflects the composition of the total DNA pool (intact + relic) relative to the intact DNA pool. Relic DNA had no effect on beta-diversity for any of the ecosystem types sampled. Grey symbols are the observed data and black symbols represent means  $\pm$  95% confidence intervals.

610 Fig. 1.



**Figure 1 | A model of relic DNA dynamics.** Modeling relic DNA dynamics. **a** The amount of relic DNA in a microbial environment is determined by inputs associated with the mortality of viable individuals with intact DNA and by losses associated with the degradation of relic DNA. If the diversity of sequences contained in the relic DNA pool is sufficiently different from that in the intact DNA pool, then relic DNA may bias estimates of microbial biodiversity (as indicated by different colored boxes) when sampling from the total (intact + relic) DNA pool. **b** We developed a sampling-based simulation model to explore the effects of mixing intact and relic DNA on estimates of diversity. We populated intact and relic communities with individuals from a lognormal species abundance distribution (SAD). We altered the diversity of the relic community by changing the scale parameter of the lognormal distribution describing the SAD. We then sampled and mixed the intact and relic communities so that the relic contribution to total community ranged from 0.01 to 0.96. **c** To gain mechanistic insight into how bias arises, we created a stochastic process-based model that captures features that influence relic DNA dynamics, including: immigration, birth, death, degradation (**a**). We simulated a range of degradation rates to achieve relic DNA pool sizes with proportions ranging between 0.05 and 0.95. To explore how degradation alters the SAD of the relic community, we explored three scenarios. First, we simulated a neutral scenario where relic DNA sequences produced by different species degrade at the same rate. Second, we simulated conditions where physical, chemical, or biological processes slow the degradation rate of relic DNA belonging to some species via protection. Third, we simulated "hot spots" where there more abundant relic DNA sequences experience faster rates of relic DNA degradation, a condition that may arise in structured habitats where there are patchy distributions of individuals and their metabolic products (i.e., enzymes). We ran simulations for 10,000 time steps and then sampled the intact and relic communities. To quantify bias in diversity (**b**, **c**), we calculated "richness ratios" which reflect the number of species in the total DNA pool (intact + relic) divided by the number of species in the intact DNA pool in a simulation. When values richness ratios = 1, relic DNA has no effect on estimates of diversity; when richness ratios > 1, relic DNA overestimates true diversity; and when richness ratios < 1, relic DNA underestimates true diversity.


612 Fig. 2.



**Figure 2 | Proportion of bacterial relic DNA in different ecosystem types.** We quantified the amount of intact DNA in a sample after removing relic DNA with a DNase treatment. We then estimated the proportion of relic DNA as  $1 - (\text{intact DNA} / \text{total DNA})$ , where the total DNA concentration was quantified without DNase treatment. Relic DNA constituted an appreciable portion of the total DNA pool, but was not affected by the ecosystem type from which the sample was collected (gut, soil, sediment, and water). Grey symbols are the observed data and black symbols represent means  $\pm$  95% confidence intervals.

614 **Fig. 3.**


616



**Figure 3 | Bias of relic DNA on within-sample bacterial diversity in different ecosystem types. a-c** We tested for the effects of bias caused by relic DNA by calculating diversity ratios for (a) richness, (b) evenness, and (c) phylogenetic diversity. The ratios reflect the diversity of the total DNA pool (intact + relic) divided by the diversity of the intact DNA pool. Relic DNA did not bias any measures of diversity in any of the ecosystem types. Richness was calculated as the number of operational taxonomic units (97% sequence similarity of the 16S rRNA gene), evenness was calculated using Simpson's evenness index, and phylogenetic diversity was calculated using Faith's *D* index. Grey symbols are the observed data and black symbols represent means  $\pm$  95% confidence intervals.



618 **Fig. 4.**



**Figure 4 | Bias of relic DNA on the among-sample bacterial diversity in different ecosystem types. a-b** We tested for the effects of bias caused by relic DNA by calculating a beta-diversity ratio based on centroid distances. Centroid distances were estimated after performing Principle Coordinates Analyses (PCoA) using taxonomic (**a**) and phylogenetic (**b**) distance metrics (Bray-Curtis and UniFrac, respectively). The centroid distance ratio was calculated on each sample within an ecosystem type and reflects the composition of the total DNA pool (intact + relic) relative to the intact DNA pool. Relic DNA had no effect on beta-diversity for any of the ecosystem types sampled. Grey symbols are the observed data and black symbols represent means  $\pm$  95% confidence intervals.