

Paradoxical stabilization of relative position in moving frames

Mert Özkah¹, Stuart Anstis², Bernard M. 't Hart³, Mark Wexler⁴, &
Patrick Cavanagh^{*1,3,5}

¹ Department of Psychological and Brain
Sciences, Dartmouth College, Hanover, NJ
03755, USA

² Department of Psychology, University of
California at San Diego, La Jolla, CA, USA

³ Centre for Vision Research, York University,
Toronto, ON, Canada

⁴ INCC and CNRS, Université de Paris, Paris,
France

⁵ Department of Psychology, Glendon College,
Toronto, ON M4N 3M6, Canada

*Corresponding author
Patrick Cavanagh
Department of Psychology
Glendon College
Toronto, ON M4N 3M6
Canada
patcav1@yorku.ca

ABSTRACT

To capture where things are and what they are doing, the visual system may extract the position and motion of each object relative to its surrounding frame of reference^{e.g., 1,2}. Here we report a particularly powerful example where a paradoxical stabilization is produced by a moving frame. We first take a frame that moves left and right and we flash its right edge before, and its left edge after, the frame's motion. For all frame displacements tested, the two edges are perceived as stabilized, with the left edge on the left and right edge on the right, separated by the frame's width as if the frame were not moving. This illusory stabilization holds even when the frame travels farther than its width, reversing the actual spatial order of the two flashes. Despite this stabilization, the motion of the frame is still seen, albeit much reduced, and this hides the paradoxical standstill of relative positions. In a second experiment, two probes are flashed inside the frame at the *same* physical location before and after the frame moves. Despite being physically superimposed, the probes are perceived widely separated, again as if they were seen in the frame's coordinates and the frame were stationary. This illusory separation is set by the distance of the frame's travel, independently of its speed. This paradoxical stabilization suggests a link to visual constancy across eye movements where the displacement of the entire visual scene may act as a frame to stabilize the perception of relative locations.


(Word count 249)

Key words: vision, position, motion, visual constancy

Introduction

Vision simplifies the dynamic world around us by coding motions and positions of objects relative to the frame that surrounds them¹, up to and including the frame of the whole visual field. Studies have shown that frames and backgrounds have very powerful influences on vision, changing what we judge to be “up”^{3,4} and what direction we think is straight ahead^{5,6}. When a frame is in motion, it can alter the impression of our own motion⁷ or that of an object within the frame^{1,2,8,9}.

Here we report that a moving frame also triggers a paradoxical stabilization. When a frame is in motion for a second or less, the separation between probes flashed just before and after the motion is seen as if the frame were stationary even though its edges still appear to move ([Movie 1](#), Fig. 1). These effects are the strongest illusions of position yet reported for steady gaze and we suggest that there are links between this paradoxical frame stabilization and visual constancy – our ability to see the world as stable despite the large shifts of the visual scene on our retinas every time we move our eyes.




Our first experiment examines this paradoxical stabilization of relative position and the following experiments reveal equivalent effects of this stabilization on probes flashed within the frame.

Results


Experiment 1: Flashes at the edges of a moving frame are localized as if the frame were not moving.

Participants report the perceived separation between parts of the frame flashed just before



and just after the frame moves. In the first condition, the right edge of the frame flashed when it was the left end of its path, and then the left edge when the frame was at the right end of its path (Fig. 2, left panel). In the second condition, the left edge flashed at both the left and right ends of the frame's path so that the physical separation between the flashes was equal to the frame's travel (Fig. 2, right panel). Participants adjusted a pair of markers at the upper right of the display to indicate how far apart the flashed edges appeared. The results are clear and dramatic. As the physical separation between the flashed edges changed from 12.5° to -7.5° , the perceived separation remained relatively constant (Fig. 3A) at a value only slightly less than the 12.5° physical width of the square. Importantly, the same separation was reported even when the frame was virtually static (the baseline judgment—the left most data point). The measurement technique appears to underestimate the width of the frame so that the relatively constant setting at all path lengths indicates that the perceived


separation was actually quite close to the perceived full width of the frame (average of 91.5% of baseline separation across the 4 non-baseline settings). Remarkably, this was true even



when edges had reversed their relative positions (at the two longer path lengths). A one-way ANOVA showed no significant effect of path length on the perceived spacing ($F(4,35)=1.4$, $p=0.26$). Despite this effective stabilization of relative positions, the movement of a single edge was clearly seen in condition 2 and did vary with the physical travel, being on average about 50% of the physical distance (Fig. 3B). The large but constant perceived inter-edge spacing in condition 1 is paradoxical: it would be expected only if the frame itself were perceived as stationary. But it is not – the perceived travel of the single edge showed that the motion of the frame was clearly visible, even if reduced. It is as if the locations of the flashed edges are reported in the coordinate system of the frame as if it were stationary although the frame's movement is quite apparent (but underestimated).

Experiment 2: Stabilization-induced position shifts.


Here we examine whether probes flashed inside the frame are also perceived in the coordinate system of the paradoxically stabilized frame, shifted far from their physical locations. In this experiment (Fig. 4, [Movie 2](#)) a frame moved left and right while two targets were flashed at the *same* physical location on the screen every time the motion reversed direction. The first flashed when the frame was at the left end of its travel, so that the flash was close to the right side of the frame. The second flashed when the frame was at the right



end of its travel placing the same flash location close to the left side of the frame (Fig. 1C). As before, participants adjusted a pair of markers at the upper right of the display to indicate how far apart the flashes appeared. The first condition tested the perceived separation of the flashes over a 64-fold variation of the frame's speed. Despite being physically superimposed, the two flashes appeared to be shifted away from each other by about the distance the frame had traveled, over the range of speed tested (Fig. 5A, $F(5, 42)=1.07$, $p=0.39$). Across the 6 speeds, the average shift was 96% of the path length. The flashes were therefore seen in their location in the frame as if the frame were almost stationary!

In the second condition, the speed and the frame's path length were fixed while the size of the frame varied from 10° to 25° . There was significant variation with size with the perceived shift dropping at larger frame sizes (Fig. 5B, $F(4,35)=7.30$, $p=0.0007$) perhaps because the frame was getting close to the borders of the monitor. Nevertheless, overall, the apparent offset between the flashes was close to the path length – the average across the 5 frame sizes was 109% of the path length.

In the third condition, the frame size was fixed while the path varied from 2.5° to 12.5° . On half the trials, the speed was held constant while on the other half, the duration of movement was held constant. The settings (Fig. 5C) show that the perceived offset of the flashes increased relatively linearly with the path length of the frame and did so similarly whether the speed was constant, or the duration of travel was constant (interaction $F(4,70)=1.75$, $p=0.22$). Averaged across the 5 path lengths, the illusory shift was 97% of path length.



We were able to repeat this last condition as an online experiment with York University undergraduates (see demonstration here <https://run.pavlovia.org/mthart/pubdemo/html/>), as part of a battery of cognitive tests. The monitor size and viewing distance varied from participant to participant and these values were available for about 40% of them. The results for these participants showed that the size of the frame in degrees of visual angle did not affect the outcome significantly so we averaged the results across all participants. The frame's travel and the perceived separation are reported as percent of the frame size (Fig. 5D). The effects for the 141 participants who remained after screening (see methods in Supplemental Materials) were quite similar to those for the 8 in-person participants. Perceived offset was again determined by the frame's path length and unaffected by speed and the illusory shift was 97% of the path length (averaged across path lengths).

Conclusions

Surprisingly, our data show that when a frame moves by less than about twice its width at a moderate to rapid speed, the relative distances between probes flashed before and after the motion are seen as if the frame were almost stationary. How can this be? The frame appears to be moving quite well — nobody reports that the frames are motionless. It is only our data that reveal that the frame's displacement is discounted for the judgments of the separation between the flashes. At the same time, the measurements of a single edge of the frame do show that the frame is seen to move, even though less than its physical travel. This produces the impression of robust motion, hiding the paradoxical stability of relative positions.

Why are relative positions stabilized? The effect we report here is not simply a reorganization of the visual elements in frame-specific coordinates as proposed by Johansson in 1950¹ and others^{2,8}. The critical difference is that, in Johansson's displays, the elements within a moving frame are seen to move relative to the frame, but they also are seen to share the common motion of the frame. Imagine someone waving to you from the window of a passing train, we see their hand waving up and down — its motion relative to the window — rather the sine wave it actually traces out. But we also see the person and their hand moving with the train, the motion they all have in common. In contrast, in our examples, the flashes escape the common motion — the displacement of the frame has been discounted and they have been effectively stabilized in world coordinates. We have no mechanistic explanation for this effect although it could be a consequence of the extraction of the frame-relative positions as suggested by Johansson¹. Perhaps the flashed probes do not seem to move along with the frame because they occur only before and after the motion. In contrast, probes in Johansson's displays were always presented during the frame's motion. Our frame stabilization may also be a small-scale application of a more general processes that stabilizes the visual field when the eyes move — more on this below.

What can act as a frame? Here we use an outline square, but we have also found similar results with a moving cloud of random dots where even a few dots are enough to trigger the position shift of the probes¹⁰. This suggests that anything that moves as a group probably engages motion discounting and the paradoxical stabilization it produces. Further studies

could clarify what qualifies as a group and also what transformations (e.g., translation, expansion / contraction, warping) produce stabilization.

How does this relate to other frame effects? There have been many demonstrations of the effects of frames on position, orientation, and motion decomposition, but none with effects of the magnitude we report here. The illusory displacements of almost 100% of the frame's travel here is far stronger than the original motion induction effects of Duncker² and others (review¹¹) where the stationary probe was presented continuously during the motion. Possibly Duncker's induced movement is reduced by the continuing evidence that the target is stationary during the frame's motion. In contrast, our flashes were brief and, importantly, they were not present during the motion. Moreover, Duncker's effect becomes noticeable only when the frame's motion is very slow, between motion threshold and maybe three times motion threshold¹². These speeds are much too slow to have any relevance to the abrupt displacement of the visual field that occurs with saccades. The frame effect that we see here is reliably large even for very abrupt displacements – in Fig. 3A, the right most data point is for a single jump of the frame with no intermediate positions.

Eye movement confounds. One alternative explanation of the frame stabilization is tracking eye movements. If the eyes were to smoothly pursue the frame, it would be more or less stable on the retina and flashes at the same location on the screen will fall on very different locations on the retina. This might lead to a stabilization of the frame and a perceived shift of the flashes. However, the everyday effects of pursuit are quite different – a tracked object does not appear to be stabilized, and the positions of flashes during pursuit are also compensated for the pursuit motion – although not completely¹³. Whatever the case, it is easy to verify that the effects reported here persist with fixation (see [Movie 1](#) or [2](#) and fixate a corner of the movie or another landmark on or near the display) and that they are no different with free viewing or with actual pursuit of the frame. Steady fixation on the flashed probe may reduce the effect for some viewers, and an actual fixation point near the discs will eliminate the frame's effect, probably by giving high certainty information about the flash location.

Relation to motion-induced position shifts. How are these frame effects related to motion-induced position shifts such as the flash lag^{14,15}, flash grab¹⁶ or the shortening of motion

paths¹⁷? In all these cases, motion also causes a shift in perceived position. However, there are several differences. First, the frame effects reported here are fairly constant across speeds up to quite high values (Experiments 2), whereas motion-induced position shifts for the flash lag¹⁸ and for the flash grab¹⁶ are speed dependent. Second, the perceived length of an oscillating motion path is shortened to about 70% of its true length¹⁷, perhaps due to position averaging¹⁹. We also find the frame's path shortened here (Experiment 1) – but to 50% of its actual value, almost twice the effect. More importantly, even this shortening could not explain the perceived stability of relative positions which was equivalent to a 100% path shortening – a full discounting of the frame's motion. Third, the frame stabilization had relatively global effects on position (Fig. 4B) – it was not greatly affected by the distance between the contours and the flash – in contrast, motion-induced position shifts like the flash grab decrease rapidly as the test moves away from the moving contour¹⁶. So, despite the resemblance of the motion and frame induced effects, there are qualitative differences that distinguish the two. It is likely that some stimuli²⁰ may trigger both frame stabilization and motion effects and further studies will be required to clearly understand whether the two are related and how they are different.

Visual constancy. Are the effects of a moving frame linked in any way to visual constancy, where the eyes move but the scene appears stable? Our results suggest a plausible, partial link. There have been many proposals to explain visual constancy, some that rely on extraretinal information, such as the assumption of a stable world^{21,22}, the subtraction of efference copy^{23,24}, or the remapping or updating of attended or landmark targets^{25,26,27,28, 29}. Others suggest that purely retinal information is sufficient based on the discounting of common motion or displacement³⁰. The stabilization of moving frames reported here falls in this second group of mechanisms, as the subtraction of the frame's displacement is based solely on retinal signals. However, this falls short of an explanation of visual constancy for two obvious reasons. The first is that we found that the motion of the frame itself was not discounted – it was clearly seen to move even if its amplitude was reduced (Fig. 3B). The second is that retinal motion on its own was never an adequate explanation for visual constancy. Specifically, stabilization is not seen for retinal motion in the absence of eye movement commands. When pushing your eyeball or viewing a scene in a moving mirror or an unsteady video (review³¹), the visual scene clearly moves. Moreover, if an eye movement

is attempted but produces no retinal motion due to afterimages³², paralyzing the extra-ocular muscles^{33,34}, or optical stabilization³⁵, the world nevertheless seems to move. Clearly, extraretinal signals are necessary for visual constancy; nevertheless, the robust frame stabilization reported here suggest that the frame effect would be effective contributor, along with extraretinal signals, to constancy processes.

Simulated saccades. The discounting of frame motion has been reported previously in simulated saccade studies^{36,37,38,39,40,41}. In these experiments, the display area, usually a reference ruler, is quickly shifted on a monitor or with a mirror. Probes presented at least 50 to 100 ms before the displacement are reported to be at their original location in the frame (e.g. retaining their ruler coordinates). However, because of the measurement technique, there is some doubt about whether the frame motion is discounted: the participants may have reported where they remembered the flash on the ruler at the time it flashed, well before the ruler moved. In contrast, our report technique avoids the ambiguity of the moving reference ruler and our results show that the separation between probes flashed before and after the motion is stabilized in world coordinates.

To conclude, we have demonstrated that when frames move, their displacements may be strongly suppressed, but more strikingly, probes in or on the frame that are flashed just before and after the frame's movement are seen in the frame coordinates as if the frame were stationary: the frame's motion between the first and second flash is largely discounted in perceiving the separation between the flashes. We suggest that this identifies a mechanism that contributes in part to the compensation for eye movements where the frame is the entire visual field.

(3034 words)

Acknowledgments

NSERC Canada (PC), VISTA (PC), Dartmouth PBS (PC), ANR (MW), UCSD Pathways to Retirement (SA). We would like to thank Don Macleod for helpful comments.

References

1. Johansson, G. (1950). Configurations in the perception of velocity. *Acta Psychologica*, **7**, 25-79.
2. Duncker, K. (1929). Über induzierte Bewegung (Ein Beitrag zur Theorie optisch wahrgenommener Bewegung). *Psychologische Forschung*, **12**, 180–259. Abridged and translated (1938) as “Induced Motion” in *Source Book of Gestalt Psychology* edited and translated by Ellis, W. D. London: Routledge and Kegan Paul. pp 161–172.
3. Asch, S. E., & Witkin, H. A. (1948). Studies in space orientation: I. Perception of the upright with displaced visual fields. *Journal of Experimental Psychology*, **38**, 325-337.
4. Morgan, M., Grant, S., Melmoth, D., & Solomon, J. A. (2015). Tilted frames of reference have similar effects on the perception of gravitational vertical and the planning of vertical saccadic eye movements. *Experimental Brain Research*, **233**(7), 2115-25. doi: 10.1007/s00221-015-4282-0.
5. Roelofs, C. O. (1935). Optische Lokalisation. *Archiv fur Augenheilkunde*, **109**, 395–415.
6. Matin, L., & Fox, C. R. (1989). Visually perceived eye level and perceived elevation of objects: linearly additive influences from visual field pitch and from gravity. *Vision Research*, **29**(3), 315-24. doi: 10.1016/0042-6989(89)90080-1.
7. Warren, H. C. (1895). Sensations of rotation. *Psychological Review*, **2**(3), 273–276. doi: 10.1037/h0074437
8. Wallach, H. (1959). The perception of motion. *Scientific American*, **201**, 56-60.
9. Agaoglu, M. N., Herzog, M. H., & Ögmen, H. (2015). Field-like interactions between motion-based reference frames. *Attention, Perception, & Psychophysics*, **77**, 2082-2097. doi: 10.3758/s13414-015-0890-9
10. Cavanagh, P., Wexler, M., & Anstis, S. (2020). Frame-induced position shifts. *Journal of Vision*, **20**(11):607. doi: 10.1167/jov.20.11.607.
11. Reinhardt-Rutland, A. H. (1988). Induced movement in the visual modality: an overview. *Psychological Bulletin*, **103**, 57–71.
12. Nakayama, K., & Tyler, W. (1978). Relative motion induced between stationary lines. *Vision Research*, **18**, 1663–1668.
13. Wertheimer, A. H. (1987). Retinal and extraretinal information in movement perception: how to invert the Fihlene illusion. *Perception*, **16**, 299-308.

14. Hogendoorn, H. (2020). Motion extrapolation in visual processing: lessons from 25 years of flash-lag debate. *Journal of Neuroscience*, **40**(30), 5698-5705. doi: 10.1523/JNEUROSCI.0275-20.2020
15. Cavanagh, P. & Anstis, S. (2013). The flash grab effect. *Vision Research*, **91**, 8-20. doi: 10.1016/j.visres.2013.07.007
16. Sinico, M., Parovel, G., Casco, C., & Anstis, S. (2009). Perceived shrinkage of motion path. *Journal of Experimental Psychology: Human Perception and Performance*, **35**(4), 948-957. doi: 10.1037/a0014257
17. Wojtach, W. T., Sung, K., Truong, S., & Purves, D. (2008). An empirical explanation of the flash-lag effect. *Proceedings of the National Academy of Sciences*, **105**(42), 15338-16343. doi:10.1073/pnas.0808916105
18. Krekelberg, B., & Lappe, M. (2000). A model of the perceived relative positions of moving objects based upon a slow averaging process. *Vision Research*, **40**, 201-215.
19. Anstis, S., & Cavanagh, P. (2017). Moving backgrounds massively change the apparent size, shape, and orientation of flashed test squares. *I-Perception*, **8**(6), 1-4. doi: 10.1177/2041669517737561
20. MacKay, D.M. (1973) Visual stability and voluntary eye movement. In R. Jung (ed.), *Handbook of sensory physiology, Vol. VII/3*. Heidelberg/New York: Springer Verlag, pp. 307-331.
21. O'Regan, K. (1992). Solving the "real" mysteries of visual perception: the world as an outside memory. *Canadian Journal of Psychology*, **46**(3), 461-488.
22. Helmholtz, H. V. (1867). *Handbuch der Physiologischen Optik*, 1st Edn. Leipzig: Voss. Translated as (1967). *Treatise on physiological optics, Volumes 1 and 2* (J. P. C. Southall, Trans.). New York: Dover.
23. von Holst E, Mittelstaedt H (1950) Das Reafferenzprinzip (Wechselwirkung zwischen Zentralnervensystem und Peripherie). *Naturwissenschaften*, **37**, 464–476.
24. Wurtz, R. H. (2008). Neuronal mechanisms of visual stability. *Vision Research*, **48**, 2070–2089.
25. Wurtz, R. H. (2018). Corollary Discharge Contributions to Perceptual Continuity Across Saccades. *Annual Review Vision Science*, **4**, 215-237. doi: 10.1146/annurev-vision-102016-061207

26. Collins, T., Rolfs, M., Deubel, H., & Cavanagh, P. (2009). Post-saccadic location judgments reveal remapping of saccade targets to foveal locations. *Journal of Vision*, **9**(5):29, 1-9.
27. Deubel, H., Koch, C., & Bridgeman, B. (2010). Landmarks facilitate visual space constancy across saccades and during fixation. *Vision Research*, **50**, 249–59
28. Cavanagh, P., Hunt, A., Afraz, A., & Rolfs, M. (2010). Visual stability based on remapping of attention pointers. *Trends in Cognitive Sciences*, **14**, 147-153.
29. Murakami, I., & Cavanagh, P. (1998). A jitter aftereffect reveals motion-based stabilization of vision. *Nature*, **395**, 798-801.
30. Bridgeman, B. (2010). Space constancy: the rise and fall of perceptual compensation. In R. Nijhawan and B. Khurana (eds.) *Space and Time in Perception and Action*. Cambridge: Cambridge University Press, pp. 94–108.
31. Mack, A., Bachant, J. (1969). Perceived movement of the afterimage during eye movements. *Perception & Psychophysics*, **6**, 379–384. doi: 10.3758/BF03212795
32. Mack, A., Bachant, J. (1969). Perceived movement of the afterimage during eye movements. *Perception & Psychophysics*, **6**, 379–384. doi: 10.3758/BF03212795
33. Mach, E. (1886). *Beitrage zur Analyse der Empfindungen*. 1st ed. Jena:Verlag von Gustav Fischer. Translated by Fischer J. 1959. *The analysis of sensations*. New York:Dover.
34. Stevens, J. K., Emerson, R. C., Gerstein, R. L., Kailos, T., Neufeld, G. R., Nichols, C. W., & Rosenquist, A. C. (1976). Paralysis of the awake human: Visual perceptions. *Vision Research*, **16**, 93–98.
35. Ditchburn, R. W., & Ginsborg, B. L. (1952). Vision with a stabilized retinal image. *Nature*, **170**, 36-37. doi: 10.1038/170036a0.
36. Sperling, G. & Speelman, R. (1965). Visual spatial localization during object motion, apparent object motion, and image motion produced by eye movements. *Journal of the Optical Society of America*, **55**, 1576.
37. MacKay, D. M. (1970). Mislocation of test flashes during saccadic image displacements. *Nature*, **227**(5259), 731-3. doi: 10.1038/227731a0
38. O'Regan, J. K. (1984). Retinal versus extraretinal influences in flash localization during saccadic eye movements in the presence of a visible background. *Perception & Psychophysics*, **36**(1), 1-14.

39. Ostendorf, F., Fischer, C., Gaymard, B., & Ploner, C. J. (2006). Perisaccadic mislocalization without saccadic eye movements. *Neuroscience*, **137**(3), 737-45. doi: 10.1016/j.neuroscience.2005.09.032
40. Honda, H. (1995). Visual mislocalization produced by a rapid image displacement on the retina: examination by means of dichotic presentation of a target and its background scene. *Vision Research*, **35**, 3021–3028.
41. Morrone, M. C., Ross, J., & Burr, D. (1997). Apparent position of visual targets during real and simulated saccadic eye movements. *Journal of Neuroscience*, **17**, 7941–7953.

Methods

In-person Experiments

Participants

Eight individuals, including two of the authors, participated in the in-person experiments of this study (1 female; age range: 22-75, mean age = 46 ± 7.4). All participants other than the two authors were naive to the purpose of this study and had normal or corrected-to-normal vision. Written, informed consent, approved by the York University HPRC and the Committee for the Protection of Human Subjects at Dartmouth College, was obtained from each participant prior to their experimental sessions.

Apparatus

All stimuli were generated on an Apple Macintosh G4 computer with custom software written in C using the Vision Shell Graphics Libraries (Comtois, 2003). The display was presented on an LCD monitor with 60 Hz refresh rate and resolution of 800 x 600 pixels. The size of the display area was $40^\circ \times 30^\circ$ (degrees of visual angle). Response adjustments were made with a track pad or mouse. Head movements were restrained with a chin rest and the viewing distance was 57 cm.

Stimuli

The screen was filled with a uniform mid-grey background while the lighter square frame had 50% contrast (Michelson) with the background. The square's size varied depending on the experiment, but its contour always subtended 0.6° . The frame's motion path was centered horizontally on the display and the vertical center of the frame was 3.75° below the display's vertical midpoint to provide space for the measurement markers at the top right. Probes were alternately red and blue. In the first experiment, the probes were superimposed on the left or right contour of the frame and had the same width and height as the contour. In the second and third experiments, the two flashed probes were discs of 1.5° in diameter. In the continuous motion condition of experiment 3, the single probe was red and was also 1.5° in diameter. In all conditions, two adjustment markers were present in the upper right of the display, 10° horizontally from the screen center and 16.5° above the midpoint. In Experiment


1, the markers were red and blue vertical bars 0.6° wide and 1.5° tall; in Experiment 2, they were discs with the same size and color as the probes.

Procedure

Common to both experiments, each trial began with a beep following which the frame was present continuously and moved repeatedly back and forth horizontally. With one exception, the red and blue probes flashed alternately each time the motion reversed direction. They were presented for 33 ms, centered in the pause of the frame's motion at the end of each transit. The pause at the end of the motion was adjusted to avoid the deterioration of the frame's effect seen for short intervals between the repeating flashes (Cavanagh, Wexler, & Anstis, 2019). In any condition where the motion duration was less than 333 ms, the pause was increased from 33 ms (default value) to keep the inter-flash interval at 366 ms. Participants were instructed to look around the display wherever they wanted as they made their setting but to avoid fixating directly on a probe. Using a mouse or a trackpad, they adjusted the two markers at the top right until their separation matched that of the flashed discs (or the angle of the continuous path). When they were satisfied with their match, they pressed the space bar, and the next trial began. There were 4 repetitions of each trial. The responses were self-paced, participants could take a break at any time. The two experiments were run separately, and the sessions lasted about 10 and 20 minutes, respectively.

Experiment 1: Frame edges. Left vs right edge condition. The frame was 12.5° across, the motion duration was 166 ms and the pause was 200 ms. The path length took 1 of 5 values: 0.25° , 5° , 10° , 15° , and 20° . The shortest value was given a small offset so that it had some reversal transient. When the frame was at the left end of the trajectory, its right side briefly flashed red; when the frame reached the right end of the trajectory, its left side briefly flashed blue. Participants adjusted the two markers to indicate the separation they saw between the red and blue edges. Note that there was no overlap between frame at its leftmost and rightmost positions for the path lengths of 15° and 20° so the red right edge and blue left edges of the frame reversed locations in screen coordinates. *Left edge condition.* Same as the first condition except that it was the left edge that flashed at both ends of the frame's travel. It flashed red at the leftmost end and red at the rightmost end. All the trials for the 2 conditions were randomly intermixed for a total of (10×4) 40 trials.

Experiment 2: Flashed discs. Speed condition. The frame was 15° across and the path length was 10°. The motion took 1 of 6 durations: 16 ms, 66 ms, 133 ms, 266 ms, 533 ms, and 1066 ms. The motion paused at each end of the travel for at least 33 ms and the flash of 33 ms was centered in that pause. The pause durations for the 6 motion durations were 350 ms, 300 ms, 233 ms, 100 ms, 33 ms, and 33 ms, respectively. *Size condition.* The path length was 5° and the motion duration was 166 ms with a pause of 200 ms at each reversal. The frame size took one of 5 values: 10°, 13.75°, 17.5°, 21.25°, and 25°. *Path length condition.* The frame was 15° in size. Path length took 1 of 5 values: 2.5°, 5°, 7.5°, 10°, and 12.5°. On half the trials, the motion duration was fixed at 500 ms with a 33 ms pause so that the speed increased as the path length increased. On the other half of the trials, the motion duration was adjusted to keep the speed of the moving frame constant at 30°/s, so 83 ms, 166 ms, 250 ms, 333 ms, 416 ms, with pauses of 283 ms, 200 ms, 33 ms, 33 ms, and 33 ms, respectively. All the trials for the 3 conditions were randomly intermixed for a total of (21 x 4) 84 trials.



Analysis

The 4 settings for each condition were averaged and then the mean and standard error of the mean across participants were calculated and plotted in Figs. 3 and 5. When standard error for a Participant was high, a second session was run. The participant means for each condition were entered in one-way and two-way ANOVAs reported in the text.

Online Experiment

Participants

Participants were recruited from the York University population, and provided prior, written, informed consent. We selected only participants with self-reported normal or corrected-to-normal vision. All procedures were approved by the York University Human Participant Review Committee. We applied several inclusion criteria (see below). There were 274 participants with a complete data set and 141 remained following screening. The mean age was 22.1 ± 6.2 years (range: 17-50) including 106 females. 121 were right-handed, 15 left-handed, and 5 other.

Apparatus

The task was created in PsychoPy Builder (Pierce et al., 2019) and ran on the Pavlovia platform (<https://pavlovia.org/>). It was accessed from a Qualtrics questionnaire, embedded with other tasks as well as questions about the participant, the device used, and the presence of any surrounding distractions. We selected only those who indicated using a laptop or a PC (i.e. no phone or tablet), and who reported using a mouse or trackpad for the task (no touch screen), and who did not report any major disruptions during the task. Since each person used a different device, the screen size and viewing distance was variable. We asked people to measure the distance from the bridge of their nose to the center of the screen and to adjust an on-screen rectangle to a credit card pressed to the screen. Not all of the participants were able to do this, but for those who did (N=59), we calculated the size of the stimuli in degrees visual angle. The analysis reported in the main text showed that the effects were not influenced by the visual angle of the stimulus allowing us to average results across all participants.

Stimuli

On the bottom of the participant's browser window were instructions that remained throughout the session and the number of completed trials over the total number of trials was shown at the top left of the screen. On the left were a pair of discs, one red, one blue, for adjustment that were also presented continuously. On the right side, the frame moved left and right and red and blue probes were flashed at each reversal of direction. See demonstration version here <https://run.pavlovia.org/mthart/pubdemo/html/>. The frame was a square whose height and width were 24.5% of the smaller of the height or width of the browser window, running in full screen mode. All further spatial values are given in percent of the frame width. The width of the frame's contour was 8% of the frame size. The diameter

of the adjustment and the flashed discs was 10.2% of the frame's width. The separation of the two discs on the left could be controlled by moving their mouse or track pad left and right to indicate the perceived distance between the two flashed discs on the right. All 214 participants' operating systems reported a refresh rate of 60 Hz and the dots were flashed for a duration that should ensure 2 frames (67 ms) with a 30 Hz refresh rate, so 4 frames (67 ms) for the 60 Hz refresh. Four catch trials were included with probes alone and no frame.

Procedure

There were exactly 22 trials in the experiment. In 12 of the trials, the probes were flashed at the same location. There were again 5 different path lengths: 41%, 49%, 57%, 65%, and 73% of the frame's width combined with the two conditions: constant speed and constant duration. In the constant speed case, the motion duration increased in step with the path length (500 ms, 600 ms, 700 ms, 800 ms, and 900 ms, respectively) making the frame's speed a constant 82% of the frame's width per second. In the constant duration case, the duration was 700 ms throughout, resulting in 5 different speeds. There was one trial for each combination, except for 57% and 700 ms, present in both conditions, which had two trials in each set. There were an additional 6 stimuli with a vertical offset of 61% of the frame's width between the two flashed probes. Three path lengths for the frame's movement—41%, 57% and 73% of the frame's width—were combined in both constant speed and constant duration trials, one of each. For constant speed, the motion duration increasing with the horizontal offset to give a constant horizontal velocity of 82% of the frame's width per second. For the 3 constant duration trials, there was a fixed 700 ms motion duration. These data are not reported but are available with the rest of the data on the OSF site (see below). Finally, there were 4 catch trials where the probes were flashed with no frame present. Two of these had no vertical offset but a horizontal offset of 61% of the absent frame's width, one with the red probe on the left, one with the red probe on the right. Two had only a vertical offset of 61% of the absent frame's width and no horizontal offset one with red on top and one with blue on top. These data were intended to screen participants for accuracy but all responses were reasonable.

The 22 stimuli were ordered randomly for each participant, with a 1 s inter-trial interval. Participant were asked to change the horizontal position of the two continuously present dots on the left of the screen so that their horizontal offset (as well as angle in some stimuli)

matched that of the two flashed dots on the right. When they were satisfied with their setting, they pressed the space bar, ending the trial and starting the next. It took participants on average about 6.5 minutes to complete the task.

Given the nature of online experiments, participants may not have been fully focused on the task, or may have misunderstood the short, written instructions. Based on the self-report criteria described above, 33 participants were removed for reports such as “just clicked as quickly as possible,” “didn’t understand the instructions,” “interrupted during test.” Of the remaining 241, 100 participants were removed for not meeting performance criteria. To eliminate outliers who may have been pressing the space bar without making adjustments we required that settings on the 12 main trials had to all be less than 3 times the maximum frame movement, and that a maximum of two settings could be 0. Of the original 274 participants, 141 remained. These trials were kept in the data.

Analysis

We used descriptive statistics (means and standard errors) to plot the effects of the frame’s five different movement amplitudes and the two main conditions (constant velocity and constant speed) on perceived distance between the red and blue dots. As with the main data, we also fit a Linear Mixed Effects model to the responses, using an REML criterion with a Nelder-Mead optimizer, participant as a random effect, condition as a categorical fixed effect and frame movement amplitude as a numeric fixed effect with a standard linear link function. These analyses do not add anything substantive to the results as plotted and so are not included in the main text. They are available with the raw data on the OSF site.

Data and Code Availability

The raw data, summary statistics, ANOVAs and Linear Mixed Effects Analyses, as well as the code and the 2 movies are (or will soon be) available at OSF (<https://osf.io/y5xrn/wiki/home/>).

Supplemental Materials

Movies

Movie 1: Edge spacing

- The movie starts slowly to make the positions of the flashes clear: red on the left, blue on the right
- However, when the frame reaches full speed, you may see the blue flash to the left of the red.
- The markers above are adjusted to match the perceived separation.
- The frame will fade out briefly to reveal again that red is physically left of blue.

Movie 1: Paradoxical stabilization of a moving frame. The right edge flashes red at the left end of the frame's travel and the left edge flashes blue at the right end. Participants matched the perceived separation between the flashes with the markers on the upper right. In this example, the frame travel is longer than the frame size so red flash is physically to the left of the blue flash. Nevertheless, blue is seen left of red. Double click on the movie to start or go here <https://vimeo.com/492667100> or here <https://cavlab.net/Demos/Frame>

Movie 2: Superimposed flashes

- The red and blue flashes are *always* at the same location.
- When the frame is present and moving, the blue flash may appear to be to the left of the red.
- The markers above are adjusted to match the perceived separation.
- The frame will fade out briefly to reveal that red is physically superimposed on blue.

Movie 2: Superimposed flashed probes. The two discs, one red and one blue, always flash at the same location but when the moving frame is visible, they are pulled to the left and right by almost as much as the frame's displacement. This is not affected much by fixation (try fixating the corners of the movie). Double click on the movie to start or go here <https://vimeo.com/492667545> or here <https://cavlab.net/Demos/Frame>.