

Skeletal muscle mitochondrial respiration in a model of age-related osteoarthritis is impaired after dietary rapamycin

Christian J. Elliehausen^{*1,2,3}, Dennis M. Minton^{*1,2,3}, Alexander D. Nichol³, Adam R. Konopka^{1,2,3}

¹Department of Medicine, Division of Geriatrics and Gerontology, University of Wisconsin-Madison, ²Geriatric Research, Education, and Clinical Center, William S. Middleton Memorial Veterans Hospital, ³Department of Kinesiology and Community Health, University of Illinois at Urbana-Champaign

*indicates equal contribution

Keywords: Aging, reactive oxygen species, mTOR, metformin, mitochondria

Corresponding Author:

Adam R. Konopka, PhD

Assistant Professor

Department of Medicine

Division of Geriatrics and Gerontology

University of Wisconsin-Madison

akonopka@medicine.wisc.edu

Abstract

A decline in skeletal muscle mitochondrial function is associated with the loss of skeletal muscle size and function during knee osteoarthritis (OA). We have recently reported that the 12-weeks of dietary rapamycin (Rap, 14ppm), with or without metformin (Met, 1000ppm), increased plasma glucose and OA severity in male Dunkin Hartley (DH) guinea pigs, a model of naturally occurring, age-related OA. The purpose of the current study was to determine if increased OA severity after dietary Rap and Rap+Met was accompanied by impaired skeletal muscle mitochondrial function. Mitochondrial respiration and hydrogen peroxide (H_2O_2) emissions were evaluated in permeabilized muscle fibers via high-resolution respirometry and fluorometry using either a saturating bolus or titration of ADP. Rap and Rap+Met decreased complex I (CI)-linked respiration and increased ADP sensitivity, consistent with previous findings in patients with end-stage OA. Rap also tended to decrease mitochondrial H_2O_2 emissions, however, this was no longer apparent after normalizing to respiration. The decrease in CI-linked respiration was accompanied with lower CI protein abundance. This is the first inquiry into how lifespan extending treatments Rap and Rap+Met can influence skeletal muscle mitochondria in a model of age-related OA. Collectively, our data suggest that Rap with or without Met inhibits CI-linked capacity and increases ADP sensitivity in DH guinea pigs that have greater OA severity.

INTRODUCTION

Osteoarthritis (OA) is amongst the top 10 diseases that limit human healthspan and lifespan (Murray et al., 2013). However, there are currently no disease modifying therapies for OA. Age is the greatest risk factor for naturally occurring, primary OA. OA is the leading cause of disability in older adults affecting 250 million people worldwide and creating a significant individual and socioeconomic burden (Hunter et al., 2014). OA is characterized as a disease of the whole-joint, including articular cartilage, subchondral bone and periarticular skeletal muscle (Loeser et al., 2012). The loss of skeletal muscle size (Fink et al., 2007; Serrão et al., 2015), function (Øiestad et al., 2015; Thorstensson et al., 2004), and quality (Kumar et al., 2014; Noehren et al., 2018) are features of OA that contributes to joint pain, physical inactivity, and comorbidities. Although it is difficult to dissociate the loss of muscle size and function with OA as a result of muscle disuse, there is evidence to suggest that loss of quadriceps muscle strength may occur prior to and be predictive of incident radiographic and symptomatic knee OA (Segal & Glass, 2011; Slemenda et al., 1997).

The age-related decline of skeletal muscle mitochondrial function is associated with impaired skeletal muscle contractile function (Gonzalez-Freire et al., 2018), walking speed (Coen et al., 2013), and fatigability (Santanasto et al., 2015). Older adults with high fatigability also had lower mitochondrial respiration compared to older adults with low fatigability. Nearly 50% of older individuals with high fatigability were diagnosed with OA (Santanasto et al., 2015). Consistent with these findings, patients with end-stage OA about to receive knee arthroplasty had lower skeletal muscle mitochondrial content and maximal complex IV activity compared to young and age-matched non-OA controls (Safdar et al., 2010). Additionally, OA patients have impaired redox homeostasis as evident by decreased activity of the antioxidant MnSOD and increased oxidative damage (Safdar et al., 2010).

The onset of naturally occurring (primary) OA in humans is insidious and unpredictable which limits the study of OA to end-stage disease and makes it difficult to study the cellular processes that are involved in the onset, early progression, and treatment of OA. To circumvent this limitation, we use the Dunkin Hartley (DH) guinea pig model of primary OA. The DH guinea pig predictably develops naturally occurring OA by 5 months of age and demonstrates an age-associated progression to moderate OA by 8-9 months of age (Bendele & Hulman, 1988; Radakovich et al., 2018). The DH guinea pig highly resembles the OA pathology observed in humans as evident by cartilage and bone lesions that typically originate in the medial tibia and spread laterally with age (Huebner & Kraus, 2006; Kapadia et al., 2000). In addition to OA, the DH guinea pig displays a number of recognized age-related changes in human skeletal muscle

including a decline in skeletal muscle density, muscle fiber size and mitochondrial protein synthesis rates (Musci et al., 2020). These characteristics make the DH guinea pig a desirable model to study the onset, progression and treatment of primary, age-related OA.

The FDA approved drug rapamycin (Rap) can extend lifespan in mice (Harrison et al., 2009) and when metformin is added to Rap (Rap+Met) there may be an even greater effect on lifespan extension (Strong et al., 2016). Treatments that extend lifespan are also intended to extend healthspan, the time spent without chronic age-related disease and disability. However, the therapeutic potential of Rap and Met for specific age-related diseases is incompletely understood. Despite previous work showing protective effects of Rap or Met on injury induced OA in young mice, we have recently shown that dietary Rap and Rap+Met treatment increased plasma glucose and worsened age-related OA severity in DH guinea pigs (Minton et al., 2021). Based on these findings, we sought to determine if the detrimental effects of Rap and Rap+Met on OA pathology and glucose homeostasis would be accompanied by impaired skeletal muscle mitochondrial bioenergetics.

METHODS

Animal Use

All animal procedures were approved by the Institutional Animal Care and Use Committee at the University of Illinois Urbana-Champaign. Data collection and analysis were completed at University of Wisconsin-Madison and William S. Middleton Memorial Veterans Hospital. The details of the experimental design and animal husbandry have been previously described in (Minton et al., 2021). In brief, Dunkin Hartley Guinea Pigs were obtained from Charles River and during acclimation provided a standard chow diet (Envigo 2040) fortified with vitamin C (1050 ppm) and Vitamin D (1.5 IU/kg) *ad libitum*. At 5 months of age, DH guinea pigs were then randomized to continue the control diet (n=8), or standard chow enriched with encapsulated Rapamycin (14 ppm, n=8) or the combination of encapsulated Rapamycin+Metformin (14 ppm, 1000 ppm, n=7) for 12-weeks. Microencapsulated Rapamycin (Rapamycin holdings) and Metformin HCl (AK Scientific I506) were compounded into the diet by Envigo. Guinea pigs provided experimental diets enriched with Rap or the combination of Rap+Met were allowed continued *ad libitum* access to food and water. Food provided to the control group was calculated to match the food consumption of the experimental diet to minimize the influence of food intake on dependent variables.

Tissue Collection

Following 3 months of treatment, two animals were sacrificed daily. Food was removed from cages 2-3 hours prior to euthanasia. Animals were anesthetized in a chamber containing 5% isoflurane and maintained using a mask with 1.5-3% isoflurane. Once reflexively unresponsive, the thoracic cavity was opened and whole blood was collected via cardiac venipuncture in K₂EDTA coated tubes. The heart was then excised. The right soleus was removed from the hind limb and immediately placed in ice-cold BIOPS solution (2.77 mM CaK₂-EGTA, 7.23 mM K₂-EGTA, 20 mM imidazole, 20 mM taurine, 50 mM K-MES, 0.5 mM dithiothreitol, 6.56 mM MgCl₂, 5.77 mM ATP, and 15 mM phosphocreatine, pH 7.1). The left soleus was frozen in liquid nitrogen and stored at -80C until processing.

High-resolution respirometry and fluorometry

High-resolution respirometry and fluorometry measurements were performed in duplicate using two chambers of the Oxygraph-2k (O2K; OROBOROS INSTRUMENTS, Innsbruck, Austria). Air calibration and instrument background oxygen flux were performed before each experiment. After the addition of the permeabilized fibers, the O2K was calibrated to measure to

measure H₂O₂ emissions with three injections of H₂O₂ (40μM). Temperature within the chambers was maintained at 37° C and oxygen concentrations were kept within the range of 350 - 250 μmol/mL. Reoxygenation was performed when oxygen concentrations approached 250 μmol/mL to ensure oxygen availability was not a limiting factor. Data are presented as the average between duplicates and respiration rates (O₂ flux) and H₂O₂ emissions were expressed relative to tissue wet weight (pmol/s/mg).

Tissue Preparation

Consistent with muscle from patients with knee OA, muscle from the Hartley guinea pig contained a visible prevalence of non-contractile tissue (Noehren et al., 2018). During mechanical permeabilization, connective and adipose tissue were removed. Fibers were then chemically permeabilized in BLOPS with saponin (50μg/mL) for 30 minutes. Permeabilized muscle fibers were rinsed with MiR05 (0.5 mM EGTA, 3 mM MgCl₂, 60 mM K-lactobionate, 20 mM taurine, 10 mM KH₂PO₄, 20 mM HEPES, 110 mM sucrose, 1 g/l BSA essentially fatty acid free, pH 7.1) plus 25μM blebbistatin, a myosin II-specific inhibitor to prevent muscle fiber contraction (Perry et al., 2011). Fibers were then blotted on filter paper to remove excess buffer and weighed on a microscale (Mettler Toledo X105). Fibers bundles (1.5-2.5 mg) were then placed into the chambers containing MiR05 plus 12.5μM blebbistatin.

Mitochondrial Bioenergetics

Two different protocols were used to assess mitochondrial function. The first protocol was supported by complex-I linked substrates, pyruvate (5mM), glutamate (10mM) and malate (0.5mM) to evaluate complex I (CI)-linked LEAK (CI_L) respiration followed by a bolus of ADP (5mM) to stimulate maximal complex I-linked OXPHOS (CI_P). Subsequent additions included cytochrome c (10mM) to test mitochondrial membrane integrity, succinate (10mM) for complex I plus II supported OXPHOS (CI+II_P), and Carbonyl cyanide m-chlorophenyl hydrazone (CCCP) (0.25-0.75 μM) to stimulate maximal uncoupled electron transport system (ETS; CI+II_E) capacity. Next, the complex-I inhibitor rotenone (0.5μM) was added so that the remaining respiration was reflective of CII_E. Mitochondrial respiration was stopped by the complex III inhibitor antimycin A (2.5μM) to measure residual oxygen consumption (ROX). SUIT 1 was used to measure respiration in 4 animals per group.

We next utilized an ADP titration protocol to evaluate mitochondrial respiration and H₂O₂ emissions in a different set of 4 animals per group. This protocol was also supported by pyruvate (5mM), glutamate (10mM) and malate (0.5mM) to stimulate LEAK respiration and evaluate CI-

linked H₂O₂ emissions. Subsequently, ADP was serially injected to reach concentrations of 0.125, 0.25, 0.56, 1.2, 2.4, 5.6, and 11.8 mM followed by sequential addition of cytochrome c, succinate, and CCCP to determine maximal CI+II_P and CI+II_E as previously performed (Konopka et al., 2017, 2019). Using Michaelis-Menten kinetics, V_{max} and the apparent Km for ADP were calculated. We also determined the area under the curve (AUC) for respiration and H₂O₂ emissions during titration of ADP. For both protocols, the ratio of maximal OXPHOS to maximal uncoupled ETS capacity (P/E) was used to gain insight into intrinsic mitochondrial function independent of changes in mitochondrial abundance. We also expressed H₂O₂ emissions relative to respiration as an indicator of electron leak.

Western Blotting

Soleus and liver tissue were homogenized in liquid nitrogen cooled mortar and pestle followed by bead homogenization in RIPA lysis buffer (150mM NaCl, 0.1mM EDTA, 50mM Tris, 0.1% wt/vol deoxycholate, 0.1% wt/vol SDS, 1% vol/vol Triton X-100) with 1X Halt Protease and Phosphatase Inhibitor Cocktail (ThermoFisher Scientific: 78442) as previously described (Minton et al., 2021) 10 µg of protein was loaded for OXPHOS proteins using a cocktail of subunits from the 5 electron transport system complexes, CI-NDUFB8, CII-SDH8, CIII-UQCRC2, CIV-MTCO1, CV-ATP5A (Abcam, MS604). 20 µg of soleus and liver protein loaded for all other proteins. Samples prepared in reducing conditions with β-mercaptoethanol in 2x Laemmli Sample Buffer (BioRad 1610737) and heated at 95C. The samples for OXPHOS blots were heated at 50C to preserve Complex I, II, IV proteins which are sensitive to high heat. Proteins were separated on 4-15% TGX Precast Gels (BioRad, 4561083) and transferred to PVDF membranes (Invitrogen). Membranes were blocked in TBS-Tween 20 (0.1%) (TBST) with 5% BSA and then incubated overnight with primary antibodies for OXPHOS (Abcam MS604), P-RPS6 Ser235/236 (Cell Signaling, 4858), RPS6 (Cell Signaling, 2217), P-AKT S473 (Cell Signaling, 4060), AKT (Cell Signaling, 4685) and B-Actin (Abcam Ab6276) at 1:500, 1:2000, 1:1000, and 1:5000 respectively. Despite reactivity with other guinea pig tissues, we were not able to detect AMPK (Cell Signaling, 2532) and p-AMPK (Cell Signaling, 50081) in DH guinea pig skeletal muscle therefore we probed liver tissue. OXPHOS was diluted in 1% Nonfat milk/PBS following manufacturer recommendation and all others in 5% BSA/TBST. Membranes were washed with TBST between incubations. Membranes were then incubated with IgG HRP conjugated Anti-Mouse (Abcam, 6728) and Anti-Rabbit (Cell Signaling, 7074) secondary antibodies at 1:10,000. Membranes exposed to SuperSignal Pico Substrate (Fisher, PI34095) and imaged with BioRad ChemiDox

XRS+ or UVP Biospectrum 500. Membranes were stripped with Restore Stripping Buffer following manufacturer recommendations (ThermoFisher Scientific 21059). Densitometric calculations determined using Image Lab (BioRad) or VisionWorks (Analytikjena).

Statistics

Data are presented as mean \pm standard error of the mean (SEM). Significance was set at $P < 0.05$. Michaelis-Menten kinetics were used to determine the estimated apparent K_m and maximal respiration (V_{max}) in Prism 9.0 (GraphPad Software, Inc., La Jolla, CA, USA). One-way ANOVA was used. When significant main effects were present a two-stage linear step-up procedure of Benjamini, Krieger and Yekutieli post-hoc test was chosen to account for multiple comparisons. A priori, our primary aim was to test the effects of each treatment compared to control and not to determine differences between treatments.

RESULTS

DH Guinea Pig Characteristics

We have previously published the characteristics and OA score from the same cohort of DH guinea pigs (Minton et al., 2021). We summarize those data in Table 1 for ease of readership. Briefly, Rap and Rap+Met animals weighed 15 and 22% ($P < 0.05$) less than control animals. Rap and Rap+Met treated animals had increased fasting plasma glucose by 64 and 39% ($P < 0.05$). OA severity measured by medial tibia OARSI score was nearly 2-fold greater in Rap and Rap+Met compared to age matched controls ($P < 0.05$).

Mitochondrial Respiration

ADP bolus: When provided a saturating bolus of ADP there was no statistical differences in mitochondrial respiration between DH guinea pigs treated with Rap and Rap+Met compared to control (Figure 2A). There were also no differences between treatments compared to control in the P/E ratio (Figure 2B).

ADP Titration: Using the ADP titration protocol revealed statistically significant differences in DH guinea pigs treated with Rap and Rap+Met compared to control. Treatments did not influence CI_L respiration. Compared to control, Rap and Rap+Met treated DH guinea pigs had lower ($P < 0.05$) submaximal and maximal CI -linked respiration starting at a sub-saturating dose of 1 mM ADP and continuing until a maximal, saturating dose of 12 mM ADP (Figure 3A). The inhibitory effects of Rap and Rap+Met were further supported by a lower V_{max} ($P < 0.001$) and AUC ($P < 0.01$) versus control (Figure 3B, C). Rapamycin also induced a leftward shift in the respiration curve (% of

maximum) consistent with a lower apparent K_m of ADP in the Rap and Rap+Met ($P<0.05$) treated DH guinea pigs compared to control (Figure 3D, E). A lower apparent K_m of ADP represents a greater ADP sensitivity with Rap treatment. The differences in mitochondrial respiration between Rap and Rap+Met versus control were no longer apparent during $CI+II_P$ ($P=0.1$), $CI+II_E$, or CII_E (Figure 3F). Furthermore, the ratio of P/E was not significantly different after Rap or Rap+Met treatment (Figure 3G).

Mitochondrial H_2O_2 emissions

Mitochondrial H_2O_2 emissions during CI-supported LEAK respiration tended to be lower in the Rap ($P=0.11$) and Rap+Met ($P=0.12$) treatments versus control (Figure 4A). This trend continued with titration of ADP as evident by a lower Mitochondrial H_2O_2 emissions AUC in guinea pigs treated with Rap ($P=0.08$) and Rap+Met ($P=0.10$) (Figure 4B, C). However, when mitochondrial H_2O_2 emissions were expressed relative to mitochondrial O_2 flux, there were no longer trends for differences between groups (Figure 4D, E).

Mitochondrial Content

To estimate mitochondrial protein abundance, we evaluated subunit content of complexes I through V of the ETS. CI protein content was lower ($P<0.05$) in the Rap and tended to be lower ($P=0.09$) in the Rap+Met treatment groups (Figure 4A). There were no other changes in protein abundance for any other complexes between DH guinea pigs receiving treatment compared to control.

Nutrient Signaling

To evaluate whether dietary Rap and Rap+Met modified mTOR signaling pathways in skeletal muscle of the DH guinea pig, we investigated phosphorylation of RpS6 S235/236 and AKT S473 as downstream targets of mTOR complex 1 (mTORC1) and mTOR complex II (mTORC2). Rap and Rap+Met diminished mTORC1 signaling as evident by an 83 and 87% ($P<0.01$) decrease in phosphorylated RpS6 (Figure 4B) respectively. Rap and Rap+Met did not inhibit mTORC2 signaling as evident by a non-significant 20 and 40% decrease in AKT S473 phosphorylation (Figure 4C). For unknown reasons, we were unable to detect total or phosphorylated AMPK in skeletal muscle of DH guinea pigs. We therefore explored the impact of Rap and Rap+Met on AMPK signaling in the liver. While we were able to detect AMPK in the liver, no treatment altered AMPK phosphorylation (Data not shown).

DISCUSSION

The main findings of this study are that at doses previously shown to slow aging in mice, dietary Rap and Rap+Met modified skeletal muscle mitochondrial bioenergetics in the DH guinea pig model of age-related primary OA. DH guinea pigs treated with Rap with or without Met had 1) lower submaximal and maximal CI-linked respiration, 2) greater ADP sensitivity, indicated by a lower apparent K_m of ADP, and 3) less CI protein content. The rapamycin-induced changes in mitochondrial function recapitulate similar findings in human patients with severe, end-stage OA. Consistent with this notion, the impaired mitochondrial function in Rap treated DH guinea pigs were accompanied by elevated hyperglycemia and increased OA pathology. Collectively our data indicate that dietary Rap with or without Met appears to worsen skeletal muscle mitochondrial function, glycemic control, and OA pathology in DH guinea pigs.

Rapamycin impairs mitochondrial function in a model of age-related OA

Interventions targeted to improve mitochondrial biogenesis and function have been proposed to increase longevity and delay musculoskeletal disease progression. Rap has demonstrated the ability to improve lifespan and extend healthspan in aging models (Harrison et al., 2009; Miller et al., 2011) but the effects on the mitochondria and in models of specific age-related disease are poorly understood. In the current study, we observed divergent findings depending on the protocol used to evaluate mitochondrial respiration in permeabilized muscle fibers. When using a bolus of ADP, there were no differences in maximal mitochondrial respiration between any treatment and control. These findings are not surprising and are in agreement with our previous study that showed by using a titration and not a bolus of ADP, we were able to detect subtle differences in skeletal muscle mitochondrial respiration after healthspan extending treatments in older adults (Konopka et al., 2019). Titrating ADP revealed that dietary Rap seemed to impair mitochondrial function in DH guinea pigs as evident by decreased submaximal and maximal CI-linked respiration and increased ADP sensitivity. These findings are in line with previous work to suggest that Rap decreased respiration in cultured muscle cells (Ye et al., 2012, 2013) and mice (Cunningham et al., 2007). In the context of aging, we initially viewed greater ADP sensitivity after Rap and Rap+Met treatment as beneficial since older adults have lower ADP sensitivity (Holloway et al., 2018). However, in patients with severe, end-stage OA about to undergo joint replacement, skeletal muscle mitochondrial ADP sensitivity was elevated and mitochondrial complex IV activity was lower compared to healthy, age-matched controls (Eimre et al., 2006; Safdar et al., 2010). We have previously found that the off-target side effects of dietary Rap and Rap+Met were associated with increased OA severity in DH guinea pigs. Therefore, our data in DH guinea pigs

treated with Rap and Rap+Met are consistent with end-stage human OA and indicate a potential connection between greater OA severity and impaired skeletal muscle mitochondrial bioenergetics.

Alterations to adenylate transport proteins through either change in protein content or post-translational modifications contribute to the regulation of ADP sensitivity (Holloway et al., 2018) (Miotto et al., 2018). Transport of adenylates (ADP, ATP, etc.) into the mitochondria is mediated by voltage-dependent anion channels (VDAC), mitochondrial creatine kinase and adenine nucleotide transport proteins. Previous work indicates that mTORC1 phosphorylates Bcl-IX at serine 62 which complexes with VDAC1 to increase substrate and adenylate transport across the outer mitochondrial membrane (Ramanathan & Schreiber, 2009). Attenuation of mTORC1 signaling by Rap, as observed in the current study, can disassociate Bcl-IX from VDAC limiting ADP/ATP transport and could be one mechanism involved in lowering the apparent K_m of ADP and limiting mitochondrial respiratory capacity.

Rapamycin reduces CI abundance

The inhibition of CI-supported respiration after Rap treatment was accompanied by a decrease in CI protein abundance. The P/E ratio is considered a reflection of intrinsic mitochondrial function independent from changes in protein content. Therefore, no change in P/E ratio after Rap supports the notion that a decrease in CI content contributes to lower CI-linked respiration. Previous work has demonstrated that rapamycin decreases transcription factors involved in mitochondrial biogenesis (Cunningham et al., 2007; Ye et al., 2012, 2013) with no change in OXPHOS proteins. Rap preserves bulk mitochondrial protein synthesis rates (Drake et al., 2013) while proteomic approaches reveal that turnover of proteins within the ETS are subunit specific (Karunadharm et al., 2015; Wolff et al., 2020). Rap increased the half-life of the CI subunit NDUFB8 in skeletal and cardiac muscle (Dai et al., 2014; Karunadharm et al., 2015) which is the specific subunit probed in our western blot analysis for CI. Therefore, changes in NDUFB8 protein turnover could be one factor contributing to lower abundance and changes to mitochondrial respiration. Although it remains unclear how Rap alters mitochondrial content and function, it may be linked to changes in protein turnover and assembly of ETS subunits. Future work is needed to connect ETS proteostasis to functional outcomes to further understand the mechanisms promoting healthspan extension.

Rapamycin decreases mitochondrial H₂O₂ emissions in a model of age-related OA

Mitochondrial H₂O₂ emissions in permeabilized muscle fibers are the result of H₂O₂ production and antioxidant scavenging capacity. Rap treatment tended to lower mitochondrial H₂O₂ emissions in the DH guinea pig. However, it should be noted that decreased mitochondrial H₂O₂ emissions with Rap treatment were driven primarily in the absence of ADP during CI-linked LEAK respiration. Future studies should consider also using substrates with convergent electron supply to the Q-junction through CII to induce maximal mitochondrial H₂O₂ emissions to clarify if Rap also improves the sensitivity of ADP to suppress H₂O₂ emissions. When mitochondrial H₂O₂ emissions were expressed relative to respiration, there were no longer differences between Rap and control. The ratio of mitochondrial H₂O₂ emissions to respiration is often an index to reflect electron leak. Since Rap similarly decreases both respiration and H₂O₂ emissions, these data would suggest that Rap may influence flux through OXPHOS.

Elevated plasma glucose after 12 weeks of rapamycin treatment

A common side effect of chronic rapamycin treatment is impaired glucose metabolism and insulin resistance driven by off target inhibition of mTORC2 (Arriola Apelo, Pumper, et al., 2016; Lamming et al., 2012; Ye et al., 2012). Consistent with these findings, we show that 12-weeks of dietary Rap and Rap+Met increased plasma glucose and non-significantly attenuated skeletal muscle mTORC2 signaling as evident by a 20% and 40% decrease in AKT S473 phosphorylation. We have also found that increased plasma glucose was correlated to increased OA severity in DH guinea pigs. Intermittent rapamycin dosing schedules or alternative rapamycin analogs that selectively inhibit mTORC1 minimize off-target side metabolic effects mediated by inhibition of mTORC2 (Arriola Apelo, Neuman, et al., 2016). Therefore, future work is needed to understand if different rapamycin or rapalog dosing regimens may prevent detrimental side effects such as impaired mitochondrial function and hyperglycemia and be able to delay or prevent age-related OA pathology.

Conclusion

Within the current study we observed that DH guinea pigs treated with Rap either alone or in combination with Met, had lower mitochondrial CI-linked respiration and greater ADP sensitivity. These findings are consistent with previous work in patients with end-stage OA that have lower mitochondrial respiration and elevated ADP sensitivity compared to age-matched non-OA controls. Taken together with our previous findings that Rap-induced hyperglycemia was

associated with greater OA severity, our data suggests long-term treatment with dietary rapamycin (14ppm) may not be suitable for the prevention or treatment of age-related OA in DH guinea pigs. Additional work is needed to determine if alternative rapamycin dosing regimens such as lower concentrations, intermittent administration, or rapamycin analogs may be more effective to safely maximize healthspan extending effects on OA pathology by minimizing off-target side effects associated with chronic use of dietary rapamycin.

References

- Arriola Apelo, S. I., Neuman, J. C., Baar, E. L., Syed, F. A., Cummings, N. E., Brar, H. K., Pumper, C. P., Kimple, M. E., & Lamming, D. W. (2016). Alternative rapamycin treatment regimens mitigate the impact of rapamycin on glucose homeostasis and the immune system. *Aging Cell*, *15*(1), 28–38. <https://doi.org/10.1111/ace1.12405>
- Arriola Apelo, S. I., Pumper, C. P., Baar, E. L., Cummings, N. E., & Lamming, D. W. (2016). Intermittent administration of rapamycin extends the life span of female C57BL/6J Mice. *Journals of Gerontology - Series A Biological Sciences and Medical Sciences*, *71*(7), 876–881. <https://doi.org/10.1093/gerona/glw064>
- Bendele, A. M., & Hulman, J. F. (1988). Spontaneous cartilage degeneration in guinea pigs. *Arthritis & Rheumatism*, *31*(4), 561–565. <https://doi.org/10.1002/art.1780310416>
- Coen, P. M., Jubrias, S. A., Distefano, G., Amati, F., Mackey, D. C., Glynn, N. W., Manini, T. M., Wohlgemuth, S. E., Leeuwenburgh, C., Cummings, S. R., Newman, A. B., Ferrucci, L., Toledo, F. G. S., Shankland, E., Conley, K. E., & Goodpaster, B. H. (2013). Skeletal muscle mitochondrial energetics are associated with maximal aerobic capacity and walking speed in older adults. *Journals of Gerontology - Series A Biological Sciences and Medical Sciences*, *68*(4), 447–455. <https://doi.org/10.1093/gerona/gls196>
- Cunningham, J. T., Rodgers, J. T., Arlow, D. H., Vazquez, F., Mootha, V. K., & Puigserver, P. (2007). mTOR controls mitochondrial oxidative function through a YY1-PGC-1 α transcriptional complex. *Nature*, *450*(7170), 736–740. <https://doi.org/10.1038/nature06322>
- Dai, D. F., Karunadharma, P. P., Chiao, Y. A., Basisty, N., Crispin, D., Hsieh, E. J., Chen, T., Gu, H., Djukovic, D., Raftery, D., Beyer, R. P., Maccoss, M. J., & Rabinovitch, P. S. (2014). Altered proteome turnover and remodeling by short-term caloric restriction or rapamycin rejuvenate the aging heart. *Aging Cell*, *13*(3), 529–539. <https://doi.org/10.1111/ace1.12203>
- Drake, J. C., Peelor, F. F., Biela, L. M., Watkins, M. K., Miller, R. A., Hamilton, K. L., & Miller, B. F. (2013). Assessment of mitochondrial biogenesis and mtorc1 signaling during chronic rapamycin feeding in male and female mice. *Journals of Gerontology - Series A Biological*

Sciences and Medical Sciences, 68(12 A), 1493–1501.

<https://doi.org/10.1093/gerona/glt047>

- Eimre, M., Puhke, R., Alev, K., Seppet, E., Sikkut, A., Peet, N., Kadaja, L., Lenzner, A., Haviko, T., Seene, T., Saks, V. A., & Seppet, E. K. (2006). Altered mitochondrial apparent affinity for ADP and impaired function of mitochondrial creatine kinase in gluteus medius of patients with hip osteoarthritis. *American Journal of Physiology - Regulatory Integrative and Comparative Physiology*, 290(5), 1271–1275. <https://doi.org/10.1152/ajpregu.00651.2005>
- Fink, B., Egl, M., Singer, J., Fuerst, M., Bubenheim, M., & Neuen-Jacob, E. (2007). Morphologic changes in the vastus medialis muscle in patients with osteoarthritis of the knee. *Arthritis and Rheumatism*, 56(11), 3626–3633. <https://doi.org/10.1002/art.22960>
- Gonzalez-Freire, M., Scalzo, P., D'Agostino, J., Moore, Z. A., Diaz-Ruiz, A., Fabbri, E., Zane, A., Chen, B., Becker, K. G., Lehmann, E., Zukley, L., Chia, C. W., Tanaka, T., Coen, P. M., Bernier, M., de Cabo, R., & Ferrucci, L. (2018). Skeletal muscle ex vivo mitochondrial respiration parallels decline in vivo oxidative capacity, cardiorespiratory fitness, and muscle strength: The Baltimore Longitudinal Study of Aging. *Aging Cell*, 17(2), 1–11. <https://doi.org/10.1111/acel.12725>
- Harrison, D. E., Strong, R., Sharp, Z. D., Nelson, J. F., Astle, C. M., Flurkey, K., Nadon, N. L., Wilkinson, J. E., Frenkel, K., Carter, C. S., Pahor, M., Javors, M. A., Fernandez, E., & Miller, R. A. (2009). Rapamycin fed late in life extends lifespan in genetically heterogeneous mice. *Nature*, 460(7253), 392–395. <https://doi.org/10.1038/nature08221>
- Holloway, G. P., Holwerda, A. M., Miotto, P. M., Dirks, M. L., Verdijk, L. B., & van Loon, L. J. C. (2018). Age-Associated Impairments in Mitochondrial ADP Sensitivity Contribute to Redox Stress in Senescent Human Skeletal Muscle. *Cell Reports*, 22(11), 2837–2848. <https://doi.org/10.1016/j.celrep.2018.02.069>
- Huebner, J. L., & Kraus, V. B. (2006). Assessment of the utility of biomarkers of osteoarthritis in the guinea pig. *Osteoarthritis and Cartilage*, 14(9), 923–930. <https://doi.org/10.1016/j.joca.2006.03.007>
- Hunter, D. J., Schofield, D., & Callander, E. (2014). The individual and socioeconomic impact of osteoarthritis. *Nature Reviews Rheumatology*, 10(7), 437–441. <https://doi.org/10.1038/nrrheum.2014.44>
- Kapadia, R. D., Badger, A. M., Levin, J. M., Swift, B., Bhattacharyya, A., Dodds, R. A., Coatney, R. W., & Lark, M. W. (2000). Meniscal ossification in spontaneous osteoarthritis in the guinea-pig. *Osteoarthritis and Cartilage*, 8(5), 374–377. <https://doi.org/10.1053/joca.1999.0312>

- Karunadharma, P. P., Basisty, N., Chiao, Y. A., Dai, D. F., Drake, R., Levy, N., Koh, W. J., Emond, M. J., Kruse, S., Marcinek, D., Maccoss, M. J., & Rabinovitch, P. S. (2015). Respiratory chain protein turnover rates in mice are highly heterogeneous but strikingly conserved across tissues, ages, and treatments. *FASEB Journal*, *29*(8), 3582–3592. <https://doi.org/10.1096/fj.15-272666>
- Konopka, A. R., Castor, W. M., Wolff, C. A., Musci, R. V., Reid, J. J., Laurin, J. L., Valenti, Z. J., Hamilton, K. L., & Miller, B. F. (2017). Skeletal muscle mitochondrial protein synthesis and respiration in response to the energetic stress of an ultra-endurance race. *Journal of Applied Physiology*, *123*, 1516–1524.
- Konopka, A. R., Laurin, J. L., Schoenberg, H. M., Reid, J. J., Castor, W. M., Wolff, C. A., Musci, R. V., Safairad, O. D., Linden, M. A., Biela, L. M., Bailey, S. M., Hamilton, K. L., & Miller, B. F. (2019). Metformin inhibits mitochondrial adaptations to aerobic exercise training in older adults. *Aging Cell*, *18*(1). <https://doi.org/10.1111/ace1.12880>
- Kumar, D., Karampinos, D. C., MacLeod, T. D., Lin, W., Nardo, L., Li, X., Link, T. M., Majumdar, S., & Souza, R. B. (2014). Quadriceps intramuscular fat fraction rather than muscle size is associated with knee osteoarthritis. *Osteoarthritis and Cartilage*, *22*(2), 226–234. <https://doi.org/10.1016/j.joca.2013.12.005>
- Lamming, D. W., Ye, L., Katajisto, P., Goncalves, M. D., Saitoh, M., Stevens, D. M., Davis, J. G., Salmon, A. B., Sabatini, D. M., & Baur, J. A. (2012). Rapamycin-Induced Insulin Resistance Is Mediated by mTORC2 Loss and Uncoupled from Longevity. *Science*, *335*(March).
- Loeser, R. F., Goldring, S. R., Scanzello, C. R., & Goldring, M. B. (2012). Osteoarthritis: A disease of the joint as an organ. *Arthritis and Rheumatism*, *64*(6), 1697–1707. <https://doi.org/10.1002/art.34453>
- Miller, R. A., Harrison, D. E., Astle, C. M., Baur, J. A., Boyd, A. R., De Cabo, R., Fernandez, E., Flurkey, K., Javors, M. A., Nelson, J. F., Orihuela, C. J., Pletcher, S., Sharp, Z. D., Sinclair, D., Starnes, J. W., Wilkinson, J. E., Nadon, N. L., & Strong, R. (2011). Rapamycin, but not resveratrol or simvastatin, extends life span of genetically heterogeneous mice. *Journals of Gerontology - Series A Biological Sciences and Medical Sciences*, *66 A*(2), 191–201. <https://doi.org/10.1093/gerona/glq178>
- Minton, D. M., Elliehausen, C. J., Javors, M. A., Santangelo, K. S., & Konopka, A. R. (2021). Rapamycin induced hyperglycemia is associated with exacerbated age-related osteoarthritis. *BioRxiv*. <https://doi.org/https://doi.org/10.1101/2021.05.21.445179> ; this
- Miotto, P. M., LeBlanc, P. J., & Holloway, G. P. (2018). High-fat diet causes mitochondrial

dysfunction as a result of impaired ADP sensitivity. *Diabetes*, 67(11), 2199–2205.

<https://doi.org/10.2337/db18-0417>

Murray, C. J. L., Abraham, J., Ali, M. K., Alvarado, M., Atkinson, C., Baddour, L. M., Bartels, D. H., Benjamin, E. J., Bhalla, K., Birbeck, G., Bolliger, I., Burstein, R., Carnahan, E., Chen, H., Chou, D., Chugh, S. S., Cohen, A., Colson, K. E., Cooper, L. T., ... Zabetian, A. (2013). The State of US health, 1990-2010: Burden of diseases, injuries, and risk factors. *JAMA - Journal of the American Medical Association*, 310(6), 591–608.

<https://doi.org/10.1001/jama.2013.13805>

Musci, R. V., Walsh, M. A., Konopka, A. R., Wolff, C. A., Peelor, F. F., Reiser, R. F., Santangelo, K. S., & Hamilton, K. L. (2020). The Dunkin Hartley Guinea Pig Is a Model of Primary Osteoarthritis That Also Exhibits Early Onset Myofiber Remodeling That Resembles Human Musculoskeletal Aging. *Frontiers in Physiology*, 11(October), 1–18.

<https://doi.org/10.3389/fphys.2020.571372>

Noehren, B., Kosmac, K., Walton, R. G., Murach, K. A., Lyles, M. F., Loeser, R. F., Peterson, C. A., & Messier, S. P. (2018). Alterations in quadriceps muscle cellular and molecular properties in adults with moderate knee osteoarthritis. *Osteoarthritis and Cartilage*, 26(10), 1359–1368. <https://doi.org/10.1016/j.joca.2018.05.011>

Øiestad, B. E., Juhl, C. B., Eitzen, I., & Thorlund, J. B. (2015). Knee extensor muscle weakness is a risk factor for development of knee osteoarthritis. A systematic review and meta-analysis. *Osteoarthritis and Cartilage*, 23(2), 171–177.

<https://doi.org/10.1016/j.joca.2014.10.008>

Perry, C. G. R., Kane, D. A., Lin, C. Te, Kozy, R., Cathey, B. L., Lark, D. S., Kane, C. L., Brophy, P. M., Gavin, T. P., Anderson, E. J., & Neuffer, P. D. (2011). Inhibiting myosin-ATPase reveals a dynamic range of mitochondrial respiratory control in skeletal muscle. *Biochemical Journal*, 437(2), 215–222. <https://doi.org/10.1042/BJ20110366>

Radakovich, L. B., Marolf, A. J., Shannon, J. P., Pannone, S. C., Sherk, V. D., & Santangelo, K. S. (2018). Development of a microcomputed tomography scoring system to characterize disease progression in the Hartley guinea pig model of spontaneous osteoarthritis.

Connective Tissue Research, 59(6), 523–533.

<https://doi.org/10.1080/03008207.2017.1409218>

Ramanathan, A., & Schreiber, S. L. (2009). Direct control of mitochondrial function by mTOR. *Proceedings of the National Academy of Sciences of the United States of America*, 106(52), 22229–22232. <https://doi.org/10.1073/pnas.0912074106>

Safdar, A., Hamadeh, M. J., Kaczor, J. J., Raha, S., deBeer, J., & Tarnopolsky, M. A. (2010).

- Aberrant mitochondrial homeostasis in the skeletal muscle of sedentary older adults. *PLoS ONE*, 5(5), 31–33. <https://doi.org/10.1371/journal.pone.0010778>
- Santanasto, A. J., Glynn, N. W., Jubrias, S. A., Conley, K. E., Boudreau, R. M., Amati, F., Mackey, D. C., Simonsick, E. M., Strotmeyer, E. S., Coen, P. M., Goodpaster, B. H., & Newman, A. B. (2015). Skeletal Muscle Mitochondrial Function and Fatigability in Older Adults. *Journals of Gerontology - Series A Biological Sciences and Medical Sciences*, 70(11), 1379–1385. <https://doi.org/10.1093/gerona/glu134>
- Segal, N. A., & Glass, N. A. (2011). Is quadriceps muscle weakness a risk factor for incident or progressive knee osteoarthritis? *Physician and Sportsmedicine*, 39(4), 44–50. <https://doi.org/10.3810/psm.2011.11.1938>
- Serrão, P. R. M. S., Vasilceac, F. A., Gramani-Say, K., Lessi, G. C., Oliveira, A. B., Reiff, R. B. M., Mattiello-Sverzut, A. C., & Mattiello, S. M. (2015). Men with early degrees of knee osteoarthritis present functional and morphological impairments of the quadriceps femoris muscle. *American Journal of Physical Medicine and Rehabilitation*, 94(1), 70–81. <https://doi.org/10.1097/PHM.0000000000000143>
- Slemenda, C., Brandt, K. D., Heilman, D. K., Mazzuca, S., Braunstein, E. M., Katz, B. P., & Wolinsky, F. D. (1997). Quadriceps weakness and osteoarthritis of the knee. *Annals of Internal Medicine*, 127(2), 97–104. <https://doi.org/10.7326/0003-4819-127-2-199707150-00001>
- Strong, R., Miller, R. A., Antebi, A., Astle, C. M., Bogue, M., Denzel, M. S., Fernandez, E., Flurkey, K., Hamilton, K. L., Lamming, D. W., Javors, M. A., de Magalhães, J. P., Martinez, P. A., McCord, J. M., Miller, B. F., Müller, M., Nelson, J. F., Ndukum, J., Rainger, G. E., ... Harrison, D. E. (2016). Longer lifespan in male mice treated with a weakly estrogenic agonist, an antioxidant, an α -glucosidase inhibitor or a Nrf2-inducer. *Aging Cell*, 15(5), 872–884. <https://doi.org/10.1111/accel.12496>
- Thorstensson, C. A., Petersson, I. F., Jacobsson, L. T. H., Boegård, T. L., & Roos, E. M. (2004). Reduced functional performance in the lower extremity predicted radiographic knee osteoarthritis five years later. *Annals of the Rheumatic Diseases*, 63(4), 402–407. <https://doi.org/10.1136/ard.2003.007583>
- Wolff, C. A., Lawrence, M. M., Porter, H., Zhang, Q., Reid, J. J., Laurin, J. L., Musci, R. V., Linden, M. A., Peelor, F. F., Wren, J. D., Creery, J. S., Cutler, K. J., Carson, R. H., Price, J. C., Hamilton, K. L., & Miller, B. F. (2020). Sex differences in changes of protein synthesis with rapamycin treatment are minimized when metformin is added to rapamycin. *GeroScience*. <https://doi.org/10.1007/s11357-020-00243-8>

Ye, L., Varamini, B., Lamming, D. W., Sabatini, D. M., & Baur, J. A. (2012). Rapamycin has a biphasic effect on insulin sensitivity in C2C12 myotubes due to sequential disruption of mTORC1 and mTORC2. *Frontiers in Genetics*, 3(SEP), 1–10.

<https://doi.org/10.3389/fgene.2012.00177>

Ye, L., Widlund, A. L., Sims, C. A., Lamming, D. W., Guan, Y., Davis, J. G., Sabatini, D. M., Harrison, D. E., Vang, O., & Baur, J. A. (2013). Rapamycin doses sufficient to extend lifespan do not compromise muscle mitochondrial content or endurance. *Aging*, 12(7), 6486–6487. <https://doi.org/10.18632/aging.102976>

Figure Legends

Figure 1. **A)** Skeletal muscle mitochondrial respiration during a SUIT protocol with an ADP bolus. **B)** The ratio of coupled OXPHOS to uncoupled ETS capacity (P/E). Con (n=4), Rap (n=4), and Rap+Met (n=3). Data are presented as mean \pm SEM.

Figure 2. **A)** Complex I (CI)-linked respiration, **B)** Vmax and **C)** area under the curve (AUC) during an ADP titration protocol. **D)** CI-linked respiration expressed as a percentage of maximal respiration and used to determine **E)** the apparent Km of ADP. **F)** CI+II OXPHOS, ETS and ROX after the ADP titration protocol and **G)** the P/E ratio. Con (n=4), Rap (n=4), and Rap+Met (n=4). *P<0.05 vs. Con, **P<0.01 vs Con, ***P<0.001 vs Con. Data are presented as mean \pm SEM

Figure 3. **A)** Skeletal muscle mitochondrial hydrogen peroxide (H₂O₂) emissions during CI-linked LEAK respiration and **B)** ADP titration. **C)** H₂O₂ emissions area under the ADP titration curve (AUC). **D)** Skeletal muscle H₂O₂ emissions expressed relative to respiration during CI-linked LEAK respiration and **E)** during titration of ADP. Con (n=4), Rap (n=4), and Rap+Met (n=4). Data are presented as mean \pm SEM.

Figure 4. Skeletal muscle OXPHOS and nutrient-sensing signaling proteins **A)** Protein content of the ETS complex I-V. Control is indicated by the dotted line and all data is expressed relative to the mean of the control. **B)** Protein content of P-RPS6/Total and **C)** P-AKT/Total. **D)** Protein content determined by western blot. Con (n=8), Rap (n=8), and Rap+Met (n=6). *P<0.05 vs. Con. **P<0.01 vs Con. Data are presented as mean \pm SEM.

Table 1 Physical and Metabolic Characteristics of Dunkin Hartley Guinea Pigs


	CONTROL	RAP	RAP+MET
Body Weight (g)	992 ± 90	846 ± 49*	775 ± 100*
Medial Tibia OARSI Score	4.6 ± 1.4	8.9 ± 3.0*	8.7 ± 3.6*
Fasted Blood Glucose (mg/dL)	242 ± 55	396 ± 61*	335 ± 53*
Blood Rapamycin (ng/mL)	0.4 ± 0	72 ± 8	78 ± 10
Blood Metformin (ng/mL)	2 ± 0	N/A	282 ± 54

Note: Control values for Rap and Met in blood circulation are set to lowest readable value.

***P<0.05 vs Con.**

Figure 1.

A.



B.




Figure 2.




Figure 3.






Figure 4.


A.



B.



C.



D.

