

Intellectual phenotypes in autism strongly correlate with gene dosage changes and exon locations of truncating mutations

Andrew H. Chiang^{1,2,3}, Jonathan Chang^{1,2,3}, and Dennis Vitkup^{1,2*}

¹ Department of Biomedical Informatics, Columbia University, New York, New York, USA

² Department of Systems Biology, Center for Computational Biology and Bioinformatics, Columbia University, New York, New York, USA

³ These authors contributed equally to this work

Correspondence to DV at dv2121@columbia.edu

Abstract

Autism spectrum disorders (ASD) are a group of related neurodevelopmental diseases displaying significant genetic and phenotypic heterogeneity¹⁻⁴. Despite recent progress in understanding ASD genetics, the nature of phenotypic heterogeneity across probands is currently unclear^{5,6}. It is also unknown how autism-associated mutations affect specific disorder phenotypes or whether similar genetic insults lead to similar phenotypic consequences. Likely gene-disrupting (LGD) *de novo* mutations affecting the same gene often result in substantially different intellectual quotient (IQ) phenotypes. Nevertheless, we find that truncating mutations affecting the same exon frequently lead to strikingly similar intellectual phenotypes in unrelated simplex ASD probands. Analogous patterns are observed for several other important ASD phenotypes. These findings suggest that exons, rather than genes, often represent a unit of effective phenotypic impact for truncating mutations. Our analysis shows that similar phenotypic effects are likely mediated by similar perturbations to the expression of splicing isoforms and corresponding gene dosage changes. For genes with recurrent truncating mutations, predicted changes in expression dosage strongly correlate with relative phenotypic consequences. Further analysis demonstrates that LGD mutations in the same exon often lead to similar perturbations of gene and isoform expression across human tissues. Therefore, analogous phenotypic patterns may be also observed in other developmental disorders triggered by highly penetrant genetic insults.

In this study, we focused on severely damaging, so-called likely gene-disrupting (LGD) mutations, which include nonsense, splice site, and frameshift variants. We used genetic and phenotypic data, including exome *de novo* mutations and corresponding phenotypes of ASD probands⁷, for more than 2,500 families from the Simons Simplex Collection (SSC). *De novo* LGD mutations are observed at significantly higher rates in SSC probands compared to unaffected siblings^{8,9}. This demonstrates a substantial contribution of these mutations to disease etiology in simplex ASD families, i.e. families with only a single affected child among siblings. We primarily considered in the paper the impact of *de novo* LGD mutations on several well-studied intellectual phenotypes: full-scale (FSIQ), nonverbal (NVIQ), and verbal (VIQ) intelligence quotients^{8,10,11}. Notably, these scores are standardized by age and normalized across a broad range of phenotypes⁷.




Figure 1: IQ differences between pairs of probands with *de novo* LGD mutations in the same gene. Each point in the figures corresponds to a pair of probands with *de novo* LGD mutations in the same gene. The x-axis represents the nucleotide distance between LGD mutations. The y-axis represents the absolute difference in IQs (full-scale, nonverbal, or verbal IQ) between affected probands. Moving averages are plotted in red.

We first investigated the variability of intellectual phenotypes associated with *de novo* LGD mutations in the same gene. The IQ differences between probands with mutations in the same gene were slightly smaller than differences between all pairs of probands. Specifically, the mean pairwise differences for probands with mutations in the same gene were: 28.3 for FSIQ (~11% smaller compared to all pairs of ASD probands, Mann-Whitney *U* one-tail test $P = 0.2$), 25.7 NVIQ (~12% smaller, $P = 0.14$), and 34.9 VIQ points (~1.1% smaller, $P = 0.5$). We next explored whether probands with LGD mutations at similar locations within the same gene resulted, on average, in more similar phenotypes (Fig. 1). Indeed, IQ differences between probands with LGD mutations ≤ 1000 base pairs apart, for example, were significantly smaller than differences between probands with more distant mutations; ≤ 1 kbp FSIQ/NVIQ/VIQ average difference 11.5, 10.4, 20.6 points; > 1 kbp average difference 31.4, 28.6, 37.5 points (MWU one-tail test $P = 0.002, 0.005, 0.01$). However, across the entire range of nucleotide

distances between LGD mutations, we did not observe significant correlations between IQ differences and mutation proximity (FSIQ/NVIQ/VIQ Spearman's $\rho = 0.09, 0.1, 0.03, P = 0.5, 0.4, 0.8$).




Figure 2: Average difference in IQ between SSC probands. Each bar shows the average IQ difference between pairs of probands from different groups. From right to left, the bars represent the average IQ difference between pairs of probands in the entire SSC population (light green), between probands with *de novo* LGD mutations in the same gene (dark green), between probands with *de novo* LGD mutations in the same exon (red), between probands of the same gender and with *de novo* LGD mutations in the same exon (orange). Error bars represent the SEM.

To explain the observed patterns of phenotypic similarity, we next considered the exon-intron structure of target genes. Specifically, we investigated truncating mutations affecting the same exon in unrelated ASD probands; we took into account LGD mutations in the exon's coding sequence as well as disruptions of the exon's flanking canonical splice sites, since such splice site mutations should affect the same transcript isoforms (Supplementary Fig. 1). Interestingly, the analysis of 16 unrelated ASD probands (8 pairs) with such mutations showed that they have strikingly more similar phenotypes (Fig. 2, red bars) compared to probands with LGD mutations in the same gene (Fig. 2, dark green bars); same exon FSIQ/NVIQ/VIQ average IQ difference 8.9, 8.3, 17.3 points, same gene average difference 28.3, 25.7, 34.9 points (Mann-Whitney U one-tail test $P = 0.003, 0.005, 0.016$). Notably, the phenotypic similarity only extended to truncating mutations in the same exon. The average IQ differences between probands with mutations in neighboring exons were not significantly different compared to mutations in non-neighboring exons (MWU one-tail test $P = 0.6, 0.18, 0.8$; Supplementary Fig. 2). Because of well-known

gender differences in autism susceptibility¹¹⁻¹³, we also compared IQ differences between probands of the same gender harboring truncating mutations in the same exon (Fig. 2, orange bars) to IQ differences between probands of different genders; same gender FSIQ/NVIQ/VIQ average difference 5.4, 7.2, 12.2, different gender average difference 14.7, 10, 25.7 (MWU one-tail test $P = 0.04, 0.29, 0.07$). Thus, stratification by gender further decreases the phenotypic differences between probands with LGD mutations in the same exon.

We next explored the relationship between phenotypic similarity and the proximity of truncating mutations in the corresponding protein sequences. This analysis revealed that probands with LGD mutations in the same exon often had similar IQs, despite being affected by truncating mutations separated by scores to hundreds of amino acids in protein sequence (Fig. 3a; Supplementary Fig. 3). Notably, probands with LGD mutations in the same exon were more phenotypically similar than probands with LGD mutations separated by comparable amino acid distances but in different exons (FSIQ/NVIQ/VIQ distance-matched permutation test $P = 0.010, 0.002, 0.018$; Supplementary Fig. 4). We also investigated whether *de novo* mutations truncating a larger fraction of protein sequences resulted, on average, in more severe intellectual phenotypes. The analysis showed no significant correlations between the fraction of truncated protein and the severity of intellectual phenotypes (Fig. 3b); FSIQ/NVIQ/VIQ Pearson's $R = 0.05, 0.05, 0.06$ ($P = 0.35, 0.35, 0.28$; Supplementary Fig. 5). We also did not find any significant biases in the distribution of truncating *de novo* mutations across protein sequences compared with the distribution of synonymous *de novo* mutations (Kolmogorov-Smirnov two-tail test $P = 0.9$; Supplementary Fig. 6).




Figure 3: Amino acid position of *de novo* LGD mutations in protein sequence and probands' IQs. **(a)** Amino acid distance between LGD mutations in protein sequence versus differences in nonverbal IQ. Each point corresponds to a pair of probands with LGD mutations in the same gene. The x-axis represents the amino acid distance between LGD mutations, and the y-axis represents the difference between the corresponding probands' nonverbal IQs (NVIQ). Red points represent pairs of probands with LGD mutations in the same exon, and white points represent pairs of probands with mutation in the same gene but different exons. **(b)** Relative fraction of protein sequence truncated by LGD mutations versus

probands' NVIQs. Each point corresponds to a single individual affected by an LGD mutation. The x-axis represents the fraction of protein sequence (i.e. fraction from the first amino acid) truncated by the mutation. The y-axis represents the corresponding NVIQ. Red line represents a moving average of the data.

The results presented above suggest that it is the occurrence of *de novo* LGD mutations in the same exon, rather than simply the proximity of mutation sites in nucleotide or amino acid sequence, that leads to similar phenotypic consequences. To explain this observation, we hypothesized that truncating mutations in the same exon usually affect, due to nonsense-mediated decay (NMD)¹⁴, the expression of the same splicing isoforms. Therefore, such mutations should lead to similar functional impacts through similar effects on overall gene dosage and the expression levels of affected transcriptional isoforms. To explore this mechanistic model, we used data from the Genotype and Tissue Expression (GTEx) Consortium^{15,16}, which collected exome sequencing and human tissue-specific gene expression data from hundreds of individuals and across multiple tissues. Using ~4,400 LGD variants in coding regions and corresponding RNA-seq data, we compared the expression changes resulting from LGD variants in the same and different exons of the same gene (Fig. 4). For each truncating variant, we analyzed allele-specific read counts¹⁷ and then used an empirical Bayes approach to infer the effects of NMD on gene expression (see Methods). This analysis demonstrated that the average gene dosage changes were more than 7 times more similar for individuals with LGD variants in the same exon compared to individuals with LGD variants in different exons of the same gene (Fig 4a); 2.2% versus 17.3% average difference in overall gene dosage decrease (Mann-Whitney U one-tail test $P < 2 \times 10^{-16}$). Moreover, by analyzing GTEx data for each tissue separately, we consistently found drastically more similar dosage changes resulting from LGD variants in the same exons (Fig. 4a).

Distinct splicing isoforms often have different functional properties^{18,19}. Consequently, LGD variants may affect phenotypes not only through NMD-induced changes in overall gene dosage, but also by altering the expression levels of different splicing isoforms. To analyze changes in the relative expression of specific isoforms, we used GTEx variants and calculated the angular distance metric between vectors describing isoform-specific expression changes (see Methods). This analysis confirmed that changes in relative isoform expression are significantly (~5 fold) more similar for LGD variants in the same exon compared to variants in different exons (Fig. 4b); 0.1 versus 0.46 average angular distance (Mann-Whitney U one-tail test $P < 2 \times 10^{-16}$). The results were also consistent across tissues (Fig. 4b). Overall, the analyses of GTEx data demonstrate that the changes in expression due to truncating variants in the same exon are indeed substantially more similar than the changes due to variants in different exons of the same gene.




Figure 4: Gene expression changes due to LGD variants in the same exon and in the same gene but different exons. **(a)** Expression changes for genes harboring LGD variants were calculated based on data from the Genotype and Tissue Expression (GTEx) Consortium¹⁵. The average difference in gene expression change between GTEx LGD variants in the same exon (red bars) and in the same gene but different exons (blue bars) across human tissues. Error bars represent the SEM. **(b)** The average difference in isoform-specific expression change due to GTEx LGD variants. Differences in expression change across transcriptional isoforms were quantified using the angular distance metric between vectors representing isoform-specific expression changes. The average differences in isoform expression were calculated for LGD variants in the same exon (red bars) and in the same gene but different exons (blue bars) across human tissues. The height of each bar represents the average angular distance between isoform-specific expression changes across pairs of variants. Error bars represent the SEM.

Truncating variants in highly expressed exons should lead, through NMD, to relatively larger decreases in overall gene dosage. To confirm this hypothesis, we used RNA-seq data from GTEx to quantify the relative exon expression for each exon harboring a truncating variant. To calculate relative exon

expression, we normalized GTEx expression values of each exon by GTEx expression values of the corresponding gene. Indeed, we observed a strong correlation between the relative expression levels of exons harboring LGD variants and the corresponding changes in overall gene dosage (Fig. 5; Pearson's $R = 0.69$, $P < 2 \times 10^{-16}$; Spearman's $\rho = 0.81$, $P < 2 \times 10^{-16}$; see Methods). Notably, the resulting dosage changes may mediate the relationship between the expression levels of target exons and the corresponding phenotypic effects of truncating mutations.




Figure 5: Relationship between the relative expression of exons harboring an LGD variant and the variant-induced decrease in overall gene expression. Each point corresponds to an LGD variant from the Genotype and Tissue Consortium (GTEx) dataset, with gene and exon expression measured in one of ten human tissues. The x-axis represents the relative expression of the exon harboring the LGD variant in a tissue; the relative exon expression was calculated as the ratio between exon expression and total gene expression (see Methods). The y-axis represents the decrease in overall gene expression due to nonsense-mediated decay (see Methods). Red line represents a moving average of the data.

To investigate how NMD-induced changes in gene dosage affect intellectual phenotypes in ASD, we used the BrainSpan dataset²⁰, which contains exon-specific expression from human brain tissues. The BrainSpan data allowed us to estimate expression dosage changes resulting from LGD mutations in different exons of ASD-associated genes (see Methods). Notably, it is likely that there is substantial variability in the sensitivity of intellectual phenotypes to dosage changes across human genes. Therefore,

to quantify the IQ sensitivities for genes with recurrent truncating mutations in SSC, we considered a simple linear dosage model. Specifically, we assumed that changes in probands' IQs are linearly proportional to decreases in gene dosage; we further assumed the average neurotypical IQ (100) for wild type gene dosage. Using this model, we estimated the sensitivity of IQs to dosage changes for each gene with recurrent truncating ASD mutations (Supplementary Fig. 7; see Methods). Calculated in this way, the IQ sensitivity for a gene is equal to the estimated phenotypic effect of a truncating mutation in an exon with average expression. We then normalized the observed phenotypic effects of each mutation by the target gene's IQ sensitivity. This analysis revealed that mutation-induced dosage changes are indeed strongly correlated with the normalized phenotypic effects; FSIQ/NVIQ/VIQ Pearson's $R = 0.56, 0.65, 0.51$, permutation test $P = 0.03, 0.017, 0.02$; (Fig. 6a; Supplementary Fig. 8). Reassuringly, no or very weak correlations were obtained using randomly permuted data, i.e. when truncating mutations were randomly re-assigned to different exons in the same gene (average FSIQ/NVIQ/VIQ Pearson's $R = 0.10, 0.16, 0.00$; $SD = 0.22, 0.20, 0.21$; see Methods). Since the heritability of intelligence is known to significantly increase with age²¹, we also investigated how the results depend on the age of probands. When we restricted our analysis to the older half of probands in SSC (median age 8.35 years), the strength of the correlations between the predicted dosage changes and normalized phenotypic consequences increased further; FSIQ/NVIQ/VIQ Pearson's $R = 0.71, 0.78, 0.63$; permutation test $P = 0.03, 0.016, 0.04$ (Fig. 6b; Supplementary Fig. 9). The strong correlations between target exon expression and intellectual ASD phenotypes suggest that, when gene-specific effects are taken into account, a significant fraction (30%-40%) of the relative phenotypic effects of *de novo* LGD mutations can be explained by the resulting dosage changes in target genes.

Next, we evaluated the ability of our linear dosage model to predict the effects of LGD mutations on non-normalized IQs. For each gene with multiple truncating mutations, we used our regression model to perform leave-one-out predictions of each mutation's effect on proband IQ scores (see Methods). Notably, the prediction errors of the dosage model were significantly smaller than the differences in IQ scores between probands with LGD mutations in the same gene; FSIQ/NVIQ/VIQ median prediction error 13.7, 10.8, 22.0 points; same gene median IQ difference 24.0, 22.0, 30.5 points; MWU one-tail test $P = 0.017, 0.009, 0.03$ (Fig. 6c; Supplementary Fig. 10). The predictions based on probands of the same gender had significantly smaller errors compared to predictions based on probands of the opposite gender; same gender FSIQ/NVIQ/VIQ median error 11.4, 9.9, 20.7 points; different gender median error 19, 17, 31.3 points (MWU one-tail test $P = 0.08, 0.015, 0.05$). Moreover, the prediction errors strikingly decreased for older probands; for example, for probands older than 12 years, median FSIQ/NVIQ/VIQ error 7.3, 7.6, 10.0 points (Fig. 6c, Supplementary Fig. 10, Supplementary Fig. 11).




Figure 6: Relationship between the relative expression of exons harboring LGD mutations and the corresponding decrease in probands' intellectual phenotypes. **(a)** Each point corresponds to a proband with an LGD mutation in a gene; only genes with multiple LGD mutations in SSC were considered. The x-axis represents the relative expression (exon expression normalized by the total gene expression) of the exon harboring the LGD mutation. The y-axis represents the normalized effect of each mutation on the affected proband's nonverbal IQ (see Methods). The regression line across all points is shown in red. *P*-values were calculated based on randomly shuffled data (see Methods). **(b)** Same as (a), but with the analysis restricted to the older half of probands in SSC (median age 8.35 years). **(c)** Boxplots represent the distribution of errors in predicting the effects of LGD mutations on nonverbal IQ (see Methods) compared to the differences in IQ scores between probands with LGD mutations in the same gene (blue); prediction errors are shown for all probands (green) and for probands older than 12 years (purple). Only genes with multiple LGD mutations in SSC were considered. The ends of each solid box represent the upper and lower quartiles; the horizontal lines inside each box represent the medians; and the whiskers represent the 5th and 95th percentiles.

Although we primarily analyzed the impact of ASD mutations on intellectual phenotypes, similar dosage and isoform expression changes in affected genes may also lead to analogous patterns for other phenotypes^{22,23}. Indeed, we observed similar results for several other key ASD phenotypes (Fig. 7). For example, probands with truncating mutations in the same exon exhibited more similar adaptive behavior abilities compared to probands with mutations in the same gene (Fig. 7a, Supplementary Fig. 12); Vineland Adaptive Behavior Scales, composite standard score of 6.9 versus 12 points average differences (Mann-Whitney U one-tail test $P = 0.04$). Likewise, motor skills were more similar for probands with truncating mutations in the same exon (Fig. 7b); Purdue Pegboard Test, 1.2 versus 3.0 for the average difference in normalized tasks completed with both hands (MWU one-tail test $P = 0.02$, Supplementary Fig. 13, see Methods). The age at which probands first walked was also more similar; median difference of 1 versus 3 months (MWU one-tail test $P = 0.1$). Finally, coordination scores in the Social Responsiveness Scale questionnaire were more closely related (Fig. 7c); 0.6 versus 1.1 for the average difference in normalized response (MWU one-tail test $P = 0.05$; Supplementary Fig. 14).




Figure 7: Differences in non-IQ phenotypes between probands with *de novo* LGD mutations affecting the same exon or the same gene. Each barplot shows the average difference in phenotypic score between pairs of ASD probands with mutations in the same gene (green) and in the same exon (red). **(a)** Differences in VABS-II (Vineland Adaptive Behavior Scales, 2nd ed.; composite standard score) scores. **(b)** Differences in Purdue Pegboard scores. **(c)** Differences in normalized SRS coordination score. Purdue Pegboard test scores and SRS scores were adjusted to account for probands' age and gender (see Methods). Error bars represent the SEM.

Discussion

Previous studies explored phenotypic similarity in syndromic forms of ASD due to mutations in specific genes²⁴⁻²⁸. Nevertheless, across a large collection of contributing genes, the nature of the substantial phenotypic heterogeneity in ASD remains unclear. Our study reveals several main sources of the observed heterogeneity in simplex ASD cases triggered by highly penetrant truncating mutations. There is a substantial variability in the IQ sensitivity to dosage and isoform expression changes across human genes (Supplementary Fig. 7). We also estimate that, due to the imperfect efficiency of NMD, truncating mutations usually result in relatively mild changes in gene dosage, with the average decrease in overall expression ~15-30% (Supplementary Fig. 15; see Methods). Nevertheless, when gene-specific sensitivities are taken into account, the relative phenotypic effects are significantly correlated with expression dosage changes, which depend on the target exon expression (Fig. 6). Furthermore, even perturbations leading to similar dosage changes in the same gene may result in diverse phenotypes, if different functional isoforms are affected. When the same isoforms are perturbed, as is the case for LGD mutations in the same exon, the phenotypic diversity in unrelated probands decreases even further (Fig. 2). Overall, these results demonstrate that for truncating *de novo* mutations, exons, rather than genes, represent a unit of effective phenotypic impact. It is also likely that differences in genetic background and environment represent other important sources of phenotypic variability²⁹⁻³¹. As the heritability of IQ phenotypes usually increases with age, it is reassuring that we observe a substantially higher correlation between phenotypes and gene dosage changes for older probands (Fig. 6b).

In the present study, we focused specifically on simplex cases of ASD, in which *de novo* LGD mutations are highly penetrant. In more diverse cohorts, individuals with LGD mutations in the same exon will likely display substantially greater phenotypic heterogeneity. For example, the Simons Variation in Individuals Project identified broad spectra of phenotypes associated with specific variants in the general population³²⁻³⁵. We also observed significantly larger phenotypic variability for probands from sequenced family trios, i.e. families without unaffected siblings (Supplementary Fig. 16). For these probands, the enrichment of *de novo* LGD mutations is likely to be substantially lower and the contribution from genetic background larger³⁶, thus resulting in more pronounced phenotypic variability.

Our study may have important implications for the future of precision medicine^{29,37,38}. The presented results suggest that by sequencing and phenotyping sufficiently large patient cohorts harboring truncating mutations in different exons of contributing ASD genes, it may be possible to understand likely phenotypic consequences, at least for cases resulting from highly penetrant *de novo* LGD mutations in simplex families. Furthermore, because we observe similar patterns of expression changes across multiple human tissues, medically relevant phenotypic analyses may be also extended to other developmental disorders caused by highly penetrant truncating mutations.

References

- 1 American Psychiatric Association (DSM-5 Task Force). *Diagnostic and Statistical Manual of Mental Disorders: DSM-5*. 5th edn, (American Psychiatric Association, 2013).
- 2 Krumm, N., O'Roak, B. J., Shendure, J. & Eichler, E. E. A de novo convergence of autism genetics and molecular neuroscience. *Trends in neurosciences* **37**, 95-105, doi:10.1016/j.tins.2013.11.005 (2014).
- 3 Ronemus, M., Iossifov, I., Levy, D. & Wigler, M. The role of de novo mutations in the genetics of autism spectrum disorders. *Nature reviews. Genetics* **15**, 133-141, doi:10.1038/nrg3585 (2014).
- 4 de la Torre-Ubieta, L., Won, H., Stein, J. L. & Geschwind, D. H. Advancing the understanding of autism disease mechanisms through genetics. *Nature medicine* **22**, 345-361, doi:10.1038/nm.4071 (2016).
- 5 Jeste, S. S. & Geschwind, D. H. Disentangling the heterogeneity of autism spectrum disorder through genetic findings. *Nature Reviews Neurology* **10**, 74, doi:10.1038/nrneurol.2013.278 (2014).
- 6 Talkowski, M. E., Minikel, E. V. & Gusella, J. F. Autism Spectrum Disorder Genetics: Diverse Genes with Diverse Clinical Outcomes. *Harvard Review of Psychiatry* **22** (2014).
- 7 Fischbach, G. D. & Lord, C. The Simons Simplex Collection: a resource for identification of autism genetic risk factors. *Neuron* **68**, 192-195, doi:10.1016/j.neuron.2010.10.006 (2010).
- 8 Iossifov, I. *et al.* The contribution of de novo coding mutations to autism spectrum disorder. *Nature* **515**, 216-221, doi:10.1038/nature13908 (2014).
- 9 Sanders, S. J. *et al.* De novo mutations revealed by whole-exome sequencing are strongly associated with autism. *Nature* **485**, 237-241, doi:10.1038/nature10945 (2012).
- 10 O'Roak, B. J. *et al.* Sporadic autism exomes reveal a highly interconnected protein network of de novo mutations. *Nature* **485**, 246-250, doi:10.1038/nature10989 (2012).
- 11 Chang, J., Gilman, S. R., Chiang, A. H., Sanders, S. J. & Vitkup, D. Genotype to phenotype relationships in autism spectrum disorders. *Nat Neurosci* **18**, 191-198, doi:10.1038/nn.3907 (2015).
- 12 Fombonne, E. Epidemiology of Pervasive Developmental Disorders. *Pediatric Research* **65**, 591, doi:10.1203/PDR.0b013e31819e7203 (2009).
- 13 Robinson, E. B., Lichtenstein, P., Anckarsäter, H., Happé, F. & Ronald, A. Examining and interpreting the female protective effect against autistic behavior. *Proceedings of the National Academy of Sciences* **110**, 5258-5262, doi:10.1073/pnas.1211070110 (2013).
- 14 Chang, Y. F., Imam, J. S. & Wilkinson, M. F. The nonsense-mediated decay RNA surveillance pathway. *Annual review of biochemistry* **76**, 51-74, doi:10.1146/annurev.biochem.76.050106.093909 (2007).
- 15 GTEx Consortium. Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science (New York, N.Y.)* **348**, 648-660, doi:10.1126/science.1262110 (2015).
- 16 Mele, M. *et al.* Human genomics. The human transcriptome across tissues and individuals. *Science (New York, N.Y.)* **348**, 660-665, doi:10.1126/science.aaa0355 (2015).
- 17 Rivas, M. A. *et al.* Human genomics. Effect of predicted protein-truncating genetic variants on the human transcriptome. *Science (New York, N.Y.)* **348**, 666-669, doi:10.1126/science.1261877 (2015).
- 18 Keren, H., Lev-Maor, G. & Ast, G. Alternative splicing and evolution: diversification, exon definition and function. *Nature reviews. Genetics* **11**, 345-355, doi:10.1038/nrg2776 (2010).
- 19 Yang, X. *et al.* Widespread expansion of protein interaction capabilities by alternative splicing. *Cell* **164**, 805-817, doi:10.1016/j.cell.2016.01.029 (2016).

- 20 Kang, H. J. *et al.* Spatio-temporal transcriptome of the human brain. *Nature* **478**, 483-489, doi:doi:10.1038/nature10523 (2011).
- 21 Haworth, C. M. A. *et al.* The heritability of general cognitive ability increases linearly from childhood to young adulthood. *Molecular Psychiatry* **15**, 1112, doi:10.1038/mp.2009.55 (2009).
- 22 Buja, A. *et al.* Damaging de novo mutations diminish motor skills in children on the autism spectrum. *Proceedings of the National Academy of Sciences* (2018).
- 23 Bishop, S. L. *et al.* Identification of Developmental and Behavioral Markers Associated with Genetic Abnormalities in Autism Spectrum Disorder. *The American journal of psychiatry* **174**, 576-585, doi:10.1176/appi.ajp.2017.16101115 (2017).
- 24 Sztainberg, Y. & Zoghbi, H. Y. Lessons learned from studying syndromic autism spectrum disorders. *Nat Neurosci* **19**, 1408-1417, doi:10.1038/nn.4420 (2016).
- 25 Bernier, R. *et al.* Disruptive CHD8 mutations define a subtype of autism early in development. *Cell* **158**, 263-276 (2014).
- 26 Helsmoortel, C. *et al.* A SWI/SNF-related autism syndrome caused by de novo mutations in ADNP. *Nature genetics* **46**, 380-384 (2014).
- 27 Van Bon, B. *et al.* Disruptive de novo mutations of DYRK1A lead to a syndromic form of autism and ID. *Molecular psychiatry* (2015).
- 28 Ben-Shalom, R. *et al.* Opposing Effects on NaV1.2 Function Underlie Differences Between SCN2A Variants Observed in Individuals With Autism Spectrum Disorder or Infantile Seizures. *Biological Psychiatry* **82**, 224-232 (2017).
- 29 Gandal, M. J., Leppa, V., Won, H., Parikshak, N. N. & Geschwind, D. H. The road to precision psychiatry: translating genetics into disease mechanisms. *Nature Neuroscience* **19**, 1397, doi:10.1038/nn.4409 (2016).
- 30 Robinson, E. B. *et al.* Autism spectrum disorder severity reflects the average contribution of de novo and familial influences. *Proceedings of the National Academy of Sciences* **111**, 15161 (2014).
- 31 Robinson, E. B. *et al.* Genetic risk for autism spectrum disorders and neuropsychiatric variation in the general population. *Nature Genetics* **48**, 552, doi:10.1038/ng.3529 (2016).
- 32 Simons VIP Consortium. Simons Variation in Individuals Project (Simons VIP): a genetics-first approach to studying autism spectrum and related neurodevelopmental disorders. *Neuron* **73**, 1063-1067, doi:10.1016/j.neuron.2012.02.014 (2012).
- 33 Qureshi, A. Y. *et al.* Opposing brain differences in 16p11.2 deletion and duplication carriers. *The Journal of Neuroscience* **34**, 11199-11211 (2014).
- 34 Hanson, E. *et al.* The cognitive and behavioral phenotype of the 16p11.2 deletion in a clinically ascertained population. *Biol Psychiatry* **77**, 785-793, doi:10.1016/j.biopsych.2014.04.021 (2015).
- 35 D'Angelo, D. *et al.* Defining the Effect of the 16p11.2 Duplication on Cognition, Behavior, and Medical Comorbidities. *JAMA psychiatry* **73**, 20-30, doi:10.1001/jamapsychiatry.2015.2123 (2016).
- 36 Zhao, X. *et al.* A unified genetic theory for sporadic and inherited autism. *Proc Natl Acad Sci U S A* **104**, 12831-12836, doi:10.1073/pnas.0705803104 (2007).
- 37 Collins, F. S. & Varmus, H. A new initiative on precision medicine. *N Engl J Med* **372**, 793-795, doi:10.1056/NEJMp1500523 (2015).
- 38 Geschwind, D. H. & State, M. W. Gene hunting in autism spectrum disorder: on the path to precision medicine. *The Lancet Neurology* **14**, 1109-1120 (2015).