

Directed evolution of *Escherichia coli* with lower-than-natural plasmid mutation rates

Daniel E. Deatherage, Dacia Leon, Álvaro E. Rodriguez, Salma K. Omar,
Jeffrey E. Barrick*

Department of Molecular Biosciences, Center for Systems and Synthetic Biology
The University of Texas at Austin, Austin, Texas 78712, U.S.A.

*Correspondence: jbarrick@cm.utexas.edu

Unwanted evolution of designed DNA sequences limits metabolic and genome engineering efforts. Engineered functions that are burdensome to a host cell and slow its replication are rapidly inactivated by mutations, and unplanned mutations with unpredictable effects often accumulate alongside designed changes in large-scale genome editing projects. We developed a directed evolution strategy, Periodic Reselection for Evolutionarily Reliable Variants (PResERV), to discover mutations that prolong the function of a DNA sequence in an engineered organism. Here, we used PResERV to isolate *E. coli* cells that replicate ColE1 plasmids with higher fidelity. We found mutations in DNA polymerases I and IV and in RNase E that reduce plasmid mutation rates by 6- to 30-fold. Using these antimutator *E. coli* and applying PResERV to other chassis organisms in the future promises to prevent evolutionary failures and unpredictability to provide a more stable genetic foundation for synthetic biology.

Populations of cells engineered to function as factories or biosensors experience a failure mode that is peculiar to living systems: they evolve. Unwanted evolution is a foundational problem for bioengineering that limits the efficiency and predictability of metabolic and genome engineering efforts¹⁻³. Often an engineered function diverts resources from cellular replication or otherwise interferes with growth or homeostasis. In these cases, ‘broken’ cells with mutations that inactivate the engineered function can rapidly outcompete the original design⁴⁻⁶. The rate at which an engineered function decays within a cell population in this manner can be summarized as an evolutionary lifetime or half-life⁴, which depends on the rates at which various mutational failure modes occur and their fitness benefits^{5,6}. It is sometimes possible to edit a genome to eliminate or reduce certain types of mutations^{7,8} or to devise a way of reducing the burden of an engineered function^{1,9}. However, given the complexity of DNA replication and repair processes

and the multifarious ways that an engineered function can burden a host cell, it can be difficult to improve the reliability of a cell.

Directed evolution is an effective strategy for optimizing the performance of complex systems with many interacting components, even when they include unknown factors. For example, it has been used to engineer novel enzymes that outperform their natural counterparts¹⁰ and to tune artificial gene circuits to effectively perform logic operations¹¹. Given the similarly complex constraints underlying cellular mutagenesis and the fitness burdens of engineered functions, we reasoned that a directed evolution procedure, PResERV (**Fig. 1a**), could be an effective strategy for improving the evolutionary reliability of an engineered cell. In PResERV, one artificially selects for cells that exhibit improved maintenance of a burdensome engineered function over tens to hundreds of cell divisions. We expected that PResERV might isolate cells with mutations that either reduced the rate at which failure mutations occurred or the fitness burden of the engineered function, or both, possibly in ways that would generalize to stabilizing other engineered functions in the evolved cells.

Results

We applied PResERV to *E. coli* K-12 strain BW25113¹² transformed with pSKO4, a high-copy-number pBR322 plasmid (ColE1-type origin) that encodes GFP under control of an inducible promoter¹³. GFP expression is a generic proxy for a costly engineered function in this scenario. A UV-mutagenized library of cells containing pSKO4 was propagated through daily 1:1000 serial transfers in the presence of antibiotic selection for plasmid retention. Under these conditions, cells with mutations in pSKO4 that inactivate or reduce costly GFP expression evolve, outcompete fully fluorescent cells, and constitute a majority of the population within a

few days¹³. The fluorescence of cells in the PResERV population was periodically monitored by flow cytometry. When ~75% or more of the cells exhibited reduced GFP signal, cell sorting was used to isolate ~10⁵ cells that remained as fluorescent as the ancestor to continue the population. We subjected this population to a total of 8 sorts spread throughout 30 regrowth cycles (**Fig. 1b**).

Six *E. coli* clones were isolated from the final population for further characterization. Five of these maintained more fully-fluorescent cells for more cell doublings than the unevolved wild-type strain with the original pSKO4 plasmid (**Fig. 1c**). Mutations that evolved in the plasmid, the *E. coli* chromosome, or both could have been responsible for these improvements. To determine which was the case, we cured these cells of their plasmids and retransformed them with the original pSKO4 plasmid, and we also isolated plasmids from each of the evolved strains and transformed them into unevolved wild-type host cells. For four of these strains the improvement in the evolutionary lifetime of GFP expression appeared to be mainly associated with mutations located on the *E. coli* chromosome (**Supplementary Fig. 1**).

We sequenced the genomes of these four evolved clones to understand the genetic basis of their improved reliability (**Supplementary Table 1**). In agreement with the re-transformation tests, no mutations were found in the pSKO4 plasmid in any of these strains. Each contained from four to ten mutations in the *E. coli* chromosome. These mutations could lead to improved maintenance of GFP expression by reducing the burden of GFP expression from the plasmid or by reducing the rate at which mutations that inactivate GFP arise.

Two of these strains (AER7 and AER8) had eight identical mutations while a third strain (AER9) had these same eight mutations plus two additional ones. All three shared mutations in *polB* and *rne* that were candidates for affecting evolutionary stability. PolB (Pol IV) is a stress-induced error-prone DNA polymerase that lacks proofreading activity. The *polB* gene sustained

two mutations in these PResERV strains: a missense mutation (H597Y) and a nonsense mutation earlier in the reading frame (S558*). Deletion of *polB* in the clean-genome *E. coli* strain MDS42 has been shown to lead to ~30% lower chromosomal mutation rates⁷. The *rne* gene (encoding RNase E) contains a missense mutation (L222S) in all three strains. RNase E regulates the copy number of ColE1 origin plasmids in *E. coli* by processing the RNAI antisense regulator of the RNAII replication primer¹⁴. Mutants defective in *rne* accumulate higher levels of RNA I and have reduced plasmid copy number¹⁵.

The fourth sequenced strain (AER12) had a completely different set of four mutations, which included a missense mutation in *polA* (H734Y). PolA (Pol I) is the DNA polymerase that is utilized primarily for filling gaps during lagging strand synthesis and in DNA repair in *E. coli*. Antimutator variants of PolA that lower the frequencies of mutations observed on a ColE1 reporter plasmid have been identified previously by screening a sequence library created by mutagenizing a PolA variant lacking 3'→5' exonuclease proofreading activity¹⁶. An H734Y substitution was also observed among the 592 polymerase variants tested in that study.

To test whether these three mutations contributed to the increased evolutionary reliability of the PResERV strains, we tested strains in which we reverted the evolved alleles back to their wild-type sequences. We then propagated multiple populations of the ancestor, two evolved clones (the clone with the *polA* mutation and one of the three clones containing mutations in *polB* and *rne*), and four revertant strains (one for each mutation and also a strain in which *polB* and *rne* were both reverted) under the same conditions as the initial evolution experiment and monitored the loss of fluorescence over the course of ~100 cell doublings (**Supplementary Fig. 2**). Mutations in *polA* and *rne* appeared to be responsible for most or all of the improved stability, as reverting these mutations reduced the evolutionary lifetime of GFP expression back

to a level similar to that observed in the ancestral strain. In contrast, reverting the *polB* mutation alone or in combination with reverting *rne* did not appreciably affect GFP stability.

We next used Luria-Delbrück fluctuation tests¹⁷ to determine if the increase in evolutionary reliability in these strains was associated with a decrease in mutation rates. We first measured the rates of point mutations that reverted a stop codon in a β -lactamase gene cloned into another pBR322-based plasmid¹⁸. Mutation rates to carbenicillin resistance, which requires mutating this stop codon to a sense codon, were significantly lower in each of the two focal evolved clones compared to the ancestor (**Fig. 2a**). In agreement with the GFP stability measurements, reversion of either the *polA* or *rne* mutation raised the mutation rate to that of the ancestral strain, and reversion of the *polB* mutation had no detectable effect on the mutation rate. We also measured chromosomal mutation rates in two further sets of fluctuation tests, selecting either for mutants resistant to rifampicin or to D-cycloserine. We did not find any significant differences versus the wild-type strain in these assays (**Supplementary Fig. 3**). Thus, it appears that PResERV discovered *E. coli* mutants that primarily display a lower plasmid mutation rate with much less of an effect, if any, on mutation rates in the chromosome.

Given previous reports on the effects of *rne* loss-of-function mutations¹⁵, we were concerned that a reduction in ColE1 plasmid copy number in the PResERV evolved cells could give a false signal of improvement in our two assays. First, it would lower the GFP expression burden on host cells and thereby increase the number of cell doublings it would take for mutants to outcompete cells with the wild-type plasmid (i.e., it would increase the apparent evolutionary stability). Second, with fewer plasmids per cell there would be a smaller chance that any given cell would experience a mutation in one of its plasmids would lead to resistance in the stop codon reversion assay (i.e., it would reduce the apparent mutation rate per cell). Therefore, we

measured the copy number of the pBR322 plasmid used in the mutation rate assay in the two focal evolved clones and four reconstructed strains using qPCR (**Fig. 2b**), and we also examined the per-cell GFP fluorescence from the pSKO4 plasmid in each strain (**Fig. 2c**).

The *polA* mutation did seem to reduce plasmid copy number somewhat. The evolved *polA* strain (AER12) had marginally fewer plasmids per chromosomal DNA copy when compared to wild-type ($p = 0.0666$, one-tailed t -test) and also exhibited reduced GFP fluorescent intensity ($p = 0.0142$, one-tailed t -test on log-transformed GFP+ subpopulation medians). Interestingly, other mutations in the evolved strain appeared to counteract the effects of the *polA* mutation, as reverting just this mutation to the wild-type allele increased both copy number ($p = 0.0011$) and GFP intensity ($p = 0.0003$). GFP signal ($p = 0.0108$) and perhaps copy number ($p = 0.0893$) were even greater in this *polA* revertant that still contained all other evolved mutations than they were in the original wild-type *E. coli* strain. Overall, these results suggest that there is a trade-off in the evolved *polA* mutant between plasmid copy number and mutation rate, but the reduction in plasmid copy number of ~2- to 4-fold is much less than the ~30-fold reduction in mutation rates.

In contrast, plasmid copy number was unchanged from wild-type in the evolved *rne polB* strain (AER8) and all three reconstructed strains reverting those mutations singly and in combination (one-way ANOVA, $F_{4,10} = 0.4393$, $p = 0.7777$). Here, too, there was evidence that other mutations in these strains may have increased GFP intensity, as all four of the AER8-derived strains considered together had indistinguishable fluorescence intensities (one-way ANOVA, $F_{4,10} = 0.4393$, $p = 0.7777$) that were, as a group, significantly greater than that of the wild-type ($p = 0.0112$, one-tailed t -test). Therefore, the *polB* and *rne* mutations reduced plasmid mutation rates by ~6-fold with no detectable trade-off in terms of plasmid copy number.

Discussion

Natural selection has optimized the mutation rates of organisms to balance the deleterious effects of most mutations with the need for at least some supply of new mutations to fuel adaptive evolution¹⁹. Evolution is unable to further reduce mutation rates in natural populations once the strength of selection against the deleterious genetic load from mutations is too weak to overcome genetic drift²⁰. We have shown that by artificially imposing strong selection for cells that do not give rise to mutations in a burdensome plasmid that we can overcome this ‘drift barrier’ and isolate *E. coli* that have lower-than-natural plasmid mutation rates.

Despite years of studying the mechanisms of DNA repair and replication, we do not know the fundamental physiological limits on mutation rates that could potentially be achieved by tuning or augmenting these processes. Some studies have identified antimutator polymerase variants¹⁶ or housecleaning proteins that can be overexpressed to reduce mutation rates²¹. Often, however, these studies are conducted in cells that are defective in DNA repair, so the mutations that they find may only suppress a deficit, rather than improving upon already low mutation rates in a wild-type cell. Future applications of PResERV with the burdensome reporter gene encoded in the chromosome, rather than on a plasmid, might enrich for host cell variants that have a higher fidelity for replicating all components of the bacterial genome. This approach could also potentially be generalized to other cell types used for industrial bioproduction, such as yeast or Chinese hamster ovary cells. Ultimately, lowering mutation rates can improve the foundations of synthetic biology so that engineered cells will function more predictably and reliably.

Methods

Culture conditions. *E. coli* was grown as 10 mL cultures in 50 mL Erlenmeyer flasks with incubation at 37°C and 120 r.p.m. orbital shaking over a diameter of 1 inch unless otherwise noted. The Miller formulation of Lysogeny Broth (LB) was used (10 g/L tryptone, 5 g/L yeast extract, and 10 g/L NaCl). Media were supplemented with 100 µg/mL carbenicillin (Crb), 20 µg/mL chloramphenicol (Cam), 50 µg/mL kanamycin (Kan), 100 µg/mL rifampicin (Rif), and 1 mM Isopropyl β-D-1-thiogalactopyranoside (IPTG), as indicated. Bacterial cultures were frozen at –80°C after adding glycerol as a cryoprotectant to a final concentration of 13.3% (v/v).

Strains and plasmids. The progenitor strain (BW25113) of the Keio knockout collection¹² was transformed with pSKO4. This plasmid encodes the redesigned I7101 (R0010+E0240) circuit that was edited to remove unstable repeat sequences in a prior study by Sleight et al. on the BioBrick cloning vector pSB1A2 backbone¹³. It is a ColE1 group plasmid with a pBR322 origin of replication. Plasmid pTEM-1.D254tag encodes β-lactamase with the codon for an amino acid at a surface-exposed position in its structure at which multiple amino acid substitutions are compatible with enzyme function replaced with a TAG stop codon¹⁸. It has a pBR322 origin of replication and also encodes the *rop* protein.

UV mutagenesis. BW25113 cells containing pSKO4 were cultured overnight to stationary phase in LB-Crb. Then, these cultures were pelleted by centrifugation and resuspended in an equal volume of sterile saline. Eleven 120 µl droplets of these cell suspensions were spotted on petri dishes and subjected to 27,500 µJ of UV radiation in a UVP CL-1000 crosslinker. After UV exposure, 100 µl from each droplet was combined and pelleted by centrifugation to collect

$\sim 2.5 \times 10^6$ surviving cells. These cells were inoculated into 10 mL of LB-Crb and grown to a final density of $\sim 2 \times 10^9$ cells/mL. This mutagenized library was archived as a frozen stock.

PRESERV directed evolution procedure. All growth steps were conducted in 10 mL of LB-IPTG-Crb. We used 0.1 mL of the mutagenized library to found the experimental population. After overnight growth to saturation, we propagated the population through daily 1:1000 dilutions of saturated cultures into fresh media followed by regrowth. GFP expression was monitored using a BD Fortessa flow cytometer. Periodically, overnight cultures were diluted to $\sim 2.5 \times 10^6$ cells/mL in HPLC grade water and stained with 150 nM of the nucleic acid dye SYTO 17 (Life Technologies). The GFP+ portion of the population was calculated as the percentage of SYTO 17 positive cells with at least the ancestral level of GFP fluorescence by flow cytometry. When only ~ 15 - 25% or fewer of cells in the population were GFP+, instead of a normal transfer, the population was diluted to $\sim 2.5 \times 10^6$ cells/mL in HPLC grade water and between $\sim 4 \times 10^4$ and $\sim 5 \times 10^5$ GFP+ cells were sorted into 10 mL of fresh LB-Crb using a BD FACS Aria IIIu. The SYTO 17 dye was found to decrease cell viability, so we did not use gating on this counterstain in these sorting steps, at the cost of less efficient enrichment of GFP+ over GFP- cells.

Isolation of evolved cells and plasmids. The evolved population was plated on LB-IPTG-Crb and six visibly GFP+ colonies were selected at random for further study. Each of these clonal isolates was grown overnight in LB-IPTG-Crb before isolating its plasmid and creating a frozen stock. To select for plasmid loss, we also diluted these cultures 1:1000 into media lacking Crb (LB-IPTG). After overnight growth, dilutions of the resulting cultures were then plated on LB-IPTG agar. GFP- colonies were patched onto LB-Crb agar to ensure the lack of fluorescence was

caused by loss of plasmid rather than a GFP mutation. One colony which had been cured of its plasmid was selected from each of the original six evolved clones and re-transformed with the ancestral plasmid. The ancestral BW25113 strain was separately transformed with the plasmids isolated from each of the six evolved clones. GFP decay experiments were carried out on the resulting twelve strains to examine them for evidence of increased evolutionary stability.

GFP decay curves. For each of the twelve strains tested, individual colonies were used to inoculate nine replicate *E. coli* populations. In order to more accurately estimate the number of cell doublings elapsed since the single-cell bottleneck, care was taken to transfer all cells in each colony to the first liquid culture by excising and transferring the piece of agar underneath and around each colony. Each population was then subjected to daily transfers under the same conditions as the PResERV experiment, while monitoring GFP fluorescence using SYTO17 staining and the BD Fortessa flow cytometer as describe above. For creating graphs of the percentage of cells remaining GFP⁺ over time, flow cytometry data were analyzed in R using the flowCore Bioconductor package (v1.42.3)²². Among the events successfully exhibiting a SYTO17 signal, cells were classified as GFP⁺ if they were above a uniform signal intensity across all samples in an experiment that was determined by examining the wild-type distribution. For graphing and comparing the initial GFP fluorescence in each strain, median intensity values for the GFP⁺ subpopulation were log₂ transformed before performing statistical analyses.

Genome sequencing. DNA was extracted from stationary phase *E. coli* cultures using the PureLink Genomic DNA Mini kit from Life Technologies. Purified DNA was fragmented using the Covaris AFA system and Illumina sequencing libraries were prepared using the NEBNext

DNA Library Prep Reagent Set for Illumina kit from New England Biolabs. An Illumina HiSeq 2500 was used to generate 2×125 paired-end reads from each sample at The University of Texas at Austin Genome Sequencing Analysis Facility (GSAF). FASTQ files have been deposited in the NCBI sequence read archive (SRP090775). Mutations in each of the evolved strains were predicted by using *breseq* (version 0.32.0a)²³ to compare the Illumina reads to the *E. coli* BW25113 reference genome (GenBank:CP009273.1). Genetic differences between this reference sequence and all four sequenced samples were assumed to have existed in the ancestral *E. coli* strain used to initiate the PResERV experiment and are not reported.

Strain reconstruction. Donor strains from the i-Deconvoluter library²⁴ were used to revert the three candidate evolved alleles we tested in two evolved clones (AER8 and AER12) back to wild type sequences using P1 transduction as previously described²⁵, except that only 2 μ l of lysate was used in each transduction. Lysate from SMR20954, SMR20794, and SMR20838 was combined with the appropriate evolved strain with a mutation in *polA*, *polB*, or *rne*, respectively, and plated on LB agar plates containing Kan. Resultant colonies were screened for correct replacement via Sanger sequencing before FLP recombinase was used to remove the linked Kan^R cassette used for selection of transductants as previously described²⁶. The strain with *rne* reverted to the wild-type allele was subjected to a second round of P1 transduction to also revert the *polB* mutation, and the Kan^R cassette was again removed via FLP recombination.

Mutation rate measurements. Luria-Delbrück fluctuation tests were carried out to measure mutation rates¹⁷. For plasmid mutation rates, strains cured of plasmid pSKO4 and modified as noted were transformed with plasmid pTEM-1.D254tag. Cultures were grown in LB-Cam to select for retention of this plasmid and plated on LB-Cam agar additionally supplemented with

500 µg/ml Crb to select for mutants. Thus, this assay measures the aggregate rate of all mutations that revert this stop codon to a permitted sense codon. For chromosomal mutation rates, LB agar containing 100 µg/ml rifampicin (Rif) or 60 µg/ml D-cycloserine (DCS) was used for the selective conditions. Rif resistance requires specific point mutations in the *rpoB* gene²⁷. In minimal media DCS resistance requires a loss-of function mutation specifically in the *cycA* gene²⁸, but mutations in additional targets may also be possible in the rich LB media used here.

Crb^R plasmid and Rif^R chromosomal mutation rates were determined by taking an overnight culture of a strain and transferring ~1,000 cells from a dilution in sterile saline to multiple 0.2 mL test tubes containing non-selective media. After growth of the replicate test tube cultures overnight to saturation, the entire volume of each one was either plated on a selective LB agar plate or a dilution was plated on a non-selective LB agar plate. A total of 6 non-selective and 48 or 12 selective plates were used for plasmid and chromosomal mutations respectively. Mutant colonies on the selective plates were counted after 48 h of incubation. For DCS chromosomal mutation rates, 1.0 mL LB cultures and 12 selective media plates were used. Additionally, only a portion (25 µl) of these cultures were plated on the selection LB agar. Colony counts on selective and non-selective plates were used to estimate mutation rates using the rSalvador R package (v1.7)²⁹. We used its likelihood ratio methods for calculating confidence intervals and the statistical significance of differences between mutation rate estimates.

Plasmid copy number determination. Cells containing the pTEM-1.D254tag plasmid were revived in LB-Cam with overnight growth from frozen stocks and then diluted 1000-fold into fresh media and harvested at exponential phase (OD₆₀₀ ~0.5). Three cell pellets from different biological replicates were collected for each strain. Primer pairs were designed to amplify products from either the antibiotic resistance gene (*cat*) in the pTEM-1.D254tag plasmid or the

ftsZ gene in the *E. coli* chromosome (**Supplementary Table 2**). Two amplification reactions for each of the two primer pairs were conducted using Power SYBR Green PCR Master Mix reagents and a ViiA7 Real-Time PCR System (ThermoFisher). Plasmid copy number per chromosome was estimated using the $\Delta\Delta C_t$ method after averaging technical replicates for each biological replicate³⁰. Graphing and statistical analyses were performed using log2 transformed values of copy number estimates for each biological replicate. The mean copy number estimated for the pTEM-1.D254tag plasmid in the wild-type BW25113 strain was 410.

References

1. Sandoval, C. M. *et al.* Use of pantothenate as a metabolic switch increases the genetic stability of farnesene producing *Saccharomyces cerevisiae*. *Metab. Eng.* **25**, 1–12 (2014).
2. Teng, X. *et al.* Genome-wide consequences of deleting any single gene. *Mol. Cell* **52**, 485–494 (2013).
3. Gibson, D. G. *et al.* Creation of a bacterial cell controlled by a chemically synthesized genome. *Science* **329**, 52–56 (2010).
4. Canton, B., Labno, A. & Endy, D. Refinement and standardization of synthetic biological parts and devices. *Nat. Biotechnol.* **26**, 787–793 (2008).
5. Renda, B. A., Hammerling, M. J. & Barrick, J. E. Engineering reduced evolutionary potential for synthetic biology. *Mol. Biosyst.* **10**, 1668–1678 (2014).
6. Arkin, A. P. & Fletcher, D. A. Fast, cheap and somewhat in control. *Genome Biol.* **7**, 114 (2006).
7. Csörgo, B., Fehér, T., Tímár, E., Blattner, F. R. & Pósfai, G. Low-mutation-rate, reduced-genome *Escherichia coli*: an improved host for faithful maintenance of engineered genetic constructs. *Microb. Cell Fact.* **11**, 11 (2012).
8. Jack, B. R. *et al.* Predicting the genetic stability of engineered DNA sequences with the EFM calculator. *ACS Synth. Biol.* **4**, 939–943 (2014).
9. Ceroni, F., Algar, R., Stan, G.-B. & Ellis, T. Quantifying cellular capacity identifies gene expression designs with reduced burden. *Nat. Methods* **12**, 415–418 (2015).
10. Romero, P. A. & Arnold, F. H. Exploring protein fitness landscapes by directed evolution. *Nat. Rev. Mol. Cell Biol.* **10**, 866–876 (2009).

11. Haseltine, E. L. & Arnold, F. H. Synthetic gene circuits: design with directed evolution. *Annu. Rev. Biophys. Biomol. Struct.* **36**, 1–19 (2007).
12. Baba, T. *et al.* Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol. Syst. Biol.* **2**, 2006.0008 (2006).
13. Sleight, S. C., Bartley, B. A., Lieviant, J. A. & Sauro, H. M. Designing and engineering evolutionary robust genetic circuits. *J. Biol. Eng.* **4**, 12 (2010).
14. Lin-Chao, S. & Cohen, S. N. The rate of processing and degradation of antisense RNAI regulates the replication of ColE1-type plasmids in vivo. *Cell* **65**, 1233–1242 (1991).
15. Tomcsányi, T. & Apirion, D. Processing enzyme ribonuclease E specifically cleaves RNA I. An inhibitor of primer formation in plasmid DNA synthesis. *J. Mol. Biol.* **185**, 713–720 (1985).
16. Loh, E., Choe, J. & Loeb, L. A. Highly tolerated amino acid substitutions increase the fidelity of *Escherichia coli* DNA polymerase I. *J. Biol. Chem.* **282**, 12201–9 (2007).
17. Rosche, W. A. & Foster, P. L. Determining mutation rates in bacterial populations. *Methods* **20**, 4–17 (2000).
18. Tack, D. S. *et al.* Addicting diverse bacteria to a noncanonical amino acid. *Nat. Chem. Biol.* **12**, 138–140 (2016).
19. Sniegowski, P. D., Gerrish, P. J., Johnson, T. & Shaver, A. The evolution of mutation rates: separating causes from consequences. *BioEssays* **22**, 1057–1066 (2000).
20. Lynch, M. *et al.* Genetic drift, selection and the evolution of the mutation rate. *Nat. Rev. Genet.* **17**, 704–714 (2016).
21. Kamiya, H. *et al.* Suppression of spontaneous and hydrogen peroxide-induced mutations by a MutT-type nucleotide pool sanitization enzyme, the *Escherichia coli* Orf135 protein.

- Genes to Cells* **8**, 941–950 (2003).
22. Ellis, B. *et al.* flowCore: Basic structures for flow cytometry data. R package version 1.42.3. (2018).
 23. Deatherage, D. E. & Barrick, J. E. Identification of mutations in laboratory-evolved microbes from next-generation sequencing data using *breseq*. *Methods Mol. Biol.* **1151**, 165–188 (2014).
 24. Nehring, R. B. *et al.* An ultra-dense library resource for rapid deconvolution of mutations that cause phenotypes in *Escherichia coli*. *Nucleic Acids Res.* **44**, 1–11 (2015).
 25. Thomason, L. C., Costantino, N. & Court, D. L. *E. coli* genome manipulation by P1 transduction. *Curr. Protoc. Mol. Biol.* **Chapter 1**, Unit 1.17 (2007).
 26. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. USA* **97**, 6640–6645 (2000).
 27. Jin, D. J. & Zhou, Y. N. Mutational analysis of structure-function relationship of RNA polymerase in *Escherichia coli*. *Methods Enzymol.* **273**, 300–19 (1996).
 28. Fehér, T., Cseh, B., Umenhoffer, K., Karcagi, I. & Pósfai, G. Characterization of *cycA* mutants of *Escherichia coli*. An assay for measuring in vivo mutation rates. *Mutat. Res.* **595**, 184–190 (2006).
 29. Zheng, Q. rSalvador 1.6: an R tool for the Luria-Delbrück fluctuation assay. Available at: <http://eeeeeric.github.io/rSalvador>.
 30. Lee, C., Kim, J., Shin, S. G. & Hwang, S. Absolute and relative QPCR quantification of plasmid copy number in *Escherichia coli*. *J. Biotechnol.* **123**, 273–280 (2006).

Acknowledgments

We thank Drew Tack and Andy Ellington for providing pTEM-1.D254tag. This research was funded by the DARPA BRICS program (HR0011-15-C0095), an NSF CAREER Award (CBET-1554179), and the NSF BEACON Center for the Study of Evolution in Action (DBI-0939454). D.L. was supported by an ASM Watkins graduate research fellowship, and Á.E.R. was supported by a CONACyT doctoral scholarship (476261).

Author Contributions

D.E.D. and S.K.O. performed PResERV experiments. D.E.D. and Á.E.R. characterized GFP decay. D.E.D. and D.L. reconstructed strains to test the effects of evolved alleles and measured mutation rates. D.L. measured plasmid copy number. D.E.D. and J.E.B. conceived PResERV and wrote the manuscript with contributions from D.L., Á.E.R., and S.K.O.

Competing Financial Interests

The authors declare no competing financial interests.




Figure 1

Figure 1 | Periodic Reselection for Evolutionarily Reliable Variants (PResERV) method

applied to an *E. coli* plasmid. (a) PResERV begins with a cell expressing GFP to such a high level that it imposes a significant fitness burden. After mutagenesis, the cell population is cultured until cells with mutations that reduce GFP expression take over the majority of the population. Periodically, the population is sorted to retain only those cells that remain as fluorescent as the original strain, enriching for mutant host cells that have reduced mutation rates or a lower fitness cost for GFP expression. Once the evolutionary stability of GFP expression improves, fluorescent cells are isolated and their genomes are sequenced to identify the causal genetic variants. **(b)** Propagation and sorting regimen used to perform PResERV on an *E. coli* population in which GFP was expressed from plasmid pSKO4, a high-copy plasmid with a pBR322 origin of replication. The red diamond denotes the ancestral strain that was UV-mutagenized prior to PResERV. Dashed grey and solid green bifurcating lines show when the population was sorted to retain fully fluorescent cells. The blue circle indicates when fluorescent clones were isolated and sequenced. **(c)** Populations initiated from six clones isolated from the various populations at the end of PResERV (blue and purple solid lines) were allowed to evolve alongside six replicates of the non-mutagenized, wild-type *E. coli* strain-plasmid combination (red dashed lines). Cells were considered GFP⁺ if they maintained a fluorescent intensity above a threshold level that was constant across all tested strains.




Figure 2

Figure 2 | Plasmid mutation rates, plasmid copy number, and GFP fluorescence in evolved

and reconstructed *E. coli* strains. (a) Plasmid mutation rates of wild-type and evolved strains with mutant *polA*, *polB* and *rne* alleles (mut) reverted to wild-type alleles (wt), individually or in combination. Strains related to the evolved clone with a *polA* mutation (AER12) were tested in one experiment (left panel), and strains related to the evolved clone with *polB* and *rne* mutations (AER8) were tested in another experiment (right panel). Each experiment included the wild-type *E. coli* ancestor for comparison (dashed lines). Error bars are 95% confidence intervals estimated by the likelihood ratio methods from Luria-Delbrück fluctuation tests (see **Methods**). (b) Plasmid copy number for wild-type, evolved, and reconstructed strains determined by qPCR. The horizontal dashed line indicates the estimated copy number in the wild-type strain. Error bars show the standard error of the mean on log-transformed values from three biological replicates. (c) Initial GFP fluorescence of wild-type, evolved, and reconstructed strains as measured by flow cytometry. The median per-cell fluorescence intensity of the GFP+ subpopulation of cells was determined for each of nine replicate cultures immediately after outgrowth in liquid culture (after ~35 cell doublings). The graphed values are the log-averaged values of these medians. Error bars are 95% confidence intervals calculated assuming the logarithms of the medians are normally distributed. The horizontal dashed line shows the value for the wild-type strain.