

Motor nuclei innervating eye muscles spared in mouse model of SOD1-linked ALS

Eleanor V. Thomas¹, Maria Nagy^{2,3}, Wayne A. Fenton², and

Arthur L. Horwich^{2,3}

¹Interdepartmental Program in Neuroscience, Yale University School of Medicine,

New Haven, CT 06510

²Dept Genetics, Yale University School of Medicine, New Haven, CT 06510

³Howard Hughes Medical Institute

Correspondence: arthur.horwich@yale.edu

Summary.

The eye muscles of humans with either inherited or sporadic forms of ALS are relatively unaffected during disease progression, a function of sparing of the cranial nerve motor neurons supplying them. Here we observe that cranial nerve nuclei are also spared in a mouse model of inherited SOD1-linked ALS. We examined the cranial nerve motor nuclei in a mouse strain, G85R SOD1YFP, which carries a high copy transgene encoding a mutant human SOD1-YFP fusion protein, that exhibits florid YFP-fluorescent aggregates in spinal cord motor neurons and paralyzes by 6 months of age. We observed in the cranial nerve nuclei that innervate the eye, 3N (oculomotor), 4N (trochlear), and 6N (abducens), that there was little (4N, 6N) or no (3N) aggregation, in comparison with other motor nuclei, 5N (trigeminal), 7N (facial), and 12N (hypoglossal), in the latter two of which florid aggregation was observed. Correspondingly, the number of ChAT positive motor neurons in 3N of G85R SOD1YFP relative to that in 3N of ChAT-EGFP mice showed that there was no loss of motor neurons over time, whereas in 12N there was progressive loss of motor neurons, amounting to a loss of ~30% from G85R SOD1YFP by end-stage. Thus, the sparing of extraocular motor neurons as occurs in humans with ALS appears to be replicated in our SOD1-linked ALS mouse strain, supporting the validity of the mouse model for studying this aspect of selective motor system loss in ALS. Comparisons of extraocular motor neurons (e.g., from 3N), resistant to ALS pathology, with other cranial motor neurons (e.g., from 12N), sensitive to such pathology, may thus be of value in understanding mechanisms of protection vs. susceptibility of motor neurons.

The sparing of eye movements in ALS patients is well-known and has been related to lack of involvement of the motor nuclei innervating the extraocular muscles, namely, cranial nerve nuclei 3N (oculomotor), 4N (trochlear), and 6N (abducens) (ref.1 and references therein). We asked whether our mouse model of ALS, bearing a high copy number transgene of a human genomic clone of G85R mutant SOD1 (including the human promoter) fused in its last exon with YFP (2,3), might likewise exhibit sparing of the same cranial nerve motor nuclei as compared with the other cranial motor nuclei. Notably, spinal cord motor neurons in our mouse strain exhibit large YFP fluorescent aggregates by 1 month of age, which increase in number subsequently and are associated with motor neuron cell death, amounting to 50% loss by 3 months of age and 75% loss by end-stage (3). The mice exhibit lower extremity symptoms by 3 months of age - twitching or pulling in when picked up by the tail – and uniformly paralyze by 6 months of age (4). We thus examined the cranial nerve nuclei at these two times, in each case examining 3 mice. At both 3 months and 6 months (end-stage), we observed no aggregation in ChAT positive motor neurons in 3N (see representative images in Fig.1A and 1B, and quantitation in bar graphs below and Table 1). In particular, 3N exhibited no aggregates in greater than 500 ChAT-positive motor neurons examined, ~200 examined from three mice at 3 months and ~300 examined from three mice at end-stage. The two other extraocular nuclei, 4N and 6N, exhibited rare aggregates at 3 months of age and essentially none at end-stage (see Fig.1 bar graphs and Table 1). By contrast, 7N and 12N exhibited florid aggregation at both times (Fig.1A, 1B, and Table 1), affecting ~20-30% of neurons, whereas 5N was less affected, exhibiting a level of aggregation resembling that of 4N and 6N at 3 months, but greater than 3N, 4N, and 6N at end-stage.

To inspect for whether ChAT-positive motor neuron survival was related to aggregation, we compared 3N and 12N from ChAT promoter-EGFP mice (using GFP fluorescence) with G85R SOD1YFP mice (using immunostaining with anti-ChAT) at 3 months and end-stage, counting motor neurons in 20 micron coronal sections taken at 100 micron intervals down the length of these nuclei, totaling 5 sections for 3N and 10 sections for 12N (see images in 2A and 2B for representative sections). The total of ChAT-positive motor neurons was then determined for each of 3 animals from the three groups. As shown in the bar graphs of Fig.2A, comparing ChAT-EGFP with G85R SOD1YFP 3 mo and endstage, the number of motor neurons in 3N was indistinguishable (the means were 297, 290, 327, non-significant p values with t-test). The lack of motor neuron loss in 3N thus mirrors the lack of aggregation. By contrast, the mean motor neuron count for 12N was reduced significantly in end-stage G85R SOD1YFP relative to ChAT-EGFP (507 vs. 708, $p = 0.01$). Thus for 12N, the occurrence of substantial aggregation was associated with significant cell loss, amounting to ~30%.

In sum, the foregoing data indicate that the cranial nerves supplying extraocular muscles in our SOD1-linked ALS mice are spared of aggregation relative to the other cranial motor nuclei, and where motor neuron survival was scored, there was no cell loss in 3N (oculomotor) vs. significant cell loss in 12N (hypoglossal) by end-stage. Thus, there appears to be a relationship between aggregation and cell loss, albeit it remains to be seen whether the former can drive the latter. It may rather be that the same cellular pathophysiology that drives aggregation can, at the same time, direct cell death. Regardless, the difference in behavior between 3N and 12N can serve as a basis for trying to understand mechanism(s) of protection/sensitivity. We presume that ultimately other mouse models of ALS will likely

exhibit the same extraocular motor sparing and that, more generally, at least some of the features of motor collapse in SOD1-linked ALS mice can serve to provide understanding of “sporadic” versions of ALS where, potentially, at least some similar steps of motor neuron loss are occurring.

Acknowledgments

We thank HHMI for supporting this work. E.V.T. was supported by the Nelson Fund.

We are especially grateful to Elizabeth Engle, Children’s Hospital/Harvard Medical School, for suggesting inspection of cranial nuclei in G85R SOD1YFP mice for aggregates.

References

1. Cwik, V.A., ALS clinical motor signs and symptoms, *in* Amyotrophic Lateral Sclerosis, Mitsumoto, Przedborski, Gordon (eds), 2006, Taylor and Francis, chapter 5, p.111.
2. Wang, J., Farr, G.W., Zeiss, C.J., Rodriguez-Gil, D.J., Wilson, J.H., et al. (2009). Progressive aggregation despite chaperone associations of a mutant SOD1-YFP in transgenic mice with ALS. *Proc. Natl. Acad. Sci. USA* 106, 1392-1397.
3. Hadzipasic, M., Tahvildari, B., Nagy, M., Bian, M., Horwich, A.L., and McCormick, D.A. (2014). Selective degeneration of a physiological subtype of spinal motor neuron in mice with SOD1-linked ALS. *Proc. Natl. Acad. Sci. USA* 111, 16883-16888.



4. Nagy, M., Fenton, W.A., Li, D., Furtak, K., and Horwich, A.L. (2016). Extended survival of misfolded G85R SOD1-linked ALS mice by transgenic expression of chaperone Hsp110.

Proc. Natl. Acad. Sci. USA 113, 5424-5428.

5. Thomas, E.V., Fenton, W.A., McGrath, J., and Horwich, A.L. (2017) Transfer of pathogenic and non-pathogenic cytosolic proteins between spinal cord motor neurons in vivo in chimeric mice.

Proc. Natl. Acad. Sci. USA 114(15):E3139-E3148.

A. 3 mo G85R SOD1YFP



B. End-stage G85R SOD1YFP





Fig.1. Aggregation of SOD1YFP in cranial nerve motor nuclei.

Panels A and B show aggregation in ChAT positive motor neurons from cranial nerve nuclei of G85R SOD1YFP mice at 3 months (panel A) and end-stage (panel B), with representative images shown (top) from 3N (oculomotor), 7N (facial), and 12N (hypoglossal), and bar graphs for 3N, 4N, 6N (extraocular innervation) and 5N, 7N, 12N (other motor nuclei, excluding the motor division of 10N, which did exhibit aggregates). Scale bars are 20 microns and apply to all images. Aggregates are more abundant in the latter group at both 3 months and end-stage, but 5N is less affected and exceeds the extraocular nuclei only at end-stage.

Sample preparation, ChAT immunostaining, and microscopy were carried out as in ref.5.

Aggregates were counted in 20 micron coronal sections from 3 mice at 3 months and 3 mice at end-stage. The neuron counts used to generate the bar graphs are shown in Table 1.





Fig. 2

Fig.2. Survival of motor neurons in cranial nerve nuclei 3 and 12 (oculomotor and hypoglossal) of G85R SOD1YFP ALS mice.

Panels A and B show ChAT positive motor neurons of 3N and 12N, respectively, determined either from direct GFP fluorescence of ChAT-EGFP mice or anti-ChAT antibody staining of sections from G85R SOD1YFP mice of 3 months or end-stage. At left are low magnification images (scale bar 200 microns) and at right are high resolution images of the indicated areas (scale bar 20 microns). At right are box plots of the motor neuron counts corresponding to the respective mice (n = 3 for each designation). The upper and lower extents of the boxes indicate the 75th and 25th percentiles of the data, respectively. The lines inside the boxes indicate the means, and the whiskers ± 1.5 S.D. *p* values determined by paired t-tests (Origin). n.s., non-significant.

ChAT positive motor neurons were counted in 20 micron (coronal) sections taken every 100 microns along the length of the respective nuclei (5 such sections from 3N and 10 from 12N). The sections were stained with anti-ChAT antibody (5), positive cells counted, and counts summed across the sections for the cranial nucleus of a given animal. The numbers were compared with those in the corresponding cranial nuclei of a ChAT promoter-EGFP mouse. 3 mice each of ChAT-EGFP (3 mo), 3 mo G85R SOD1YFP, and end-stage G85R SOD1YFP were analyzed.

Table 1. Percentage of motor neurons with aggregates in cranial nerve nuclei of G85R SOD1YFP mice at 3 months of age and at end-stage.

3 mo G85R SOD1YFP	3N	4N	6N	5N	7N	12N
Mouse 1 (cn 298)	0% (0/79)	8% (5/59)	3% (1/36)	10% (6/60)	41% (48/118)	42% (47/112)
Mouse 2 (cn 247)	0% (0/76)	2% (1/43)	9% (1/11)	5% (4/78)	14% (32/226)	40% (44/109)
Mouse 3 (cn 290)	0% (0/70)	0% (0/44)	0% (0/25)	0% (0/124)	10% (24/251)	17% (17/98)

End stage G85R SOD1YFP	3N	4N	6N	5N	7N	12N
Mouse 1 (cn 262)	0% (0/92)	0% (0/54)	0% (0/75)	9% (6/69)	30% (41/138)	34% (34/99)
Mouse 2 (cn 269)	0% (0/93)	4% (2/54)	2% (1/67)	7% (6/84)	19% (27/143)	37% (37/101)
Mouse 3 (cn 268)	0% (0/90)	0% (0/22)	0% (0/62)	5% (1/22)	22% (38/175)	32% (32/99)

cn = copy number of G85R SOD1YFP transgene.