

1 E6 proteins from high-risk HPV, low-risk HPV, and animal papillomaviruses activate the
2 Wnt/ β -catenin pathway through E6AP-dependent degradation of NHERF1

3

4 Camille M. Drews¹, Samuel Case¹, Scott B. Vande Pol^{1*}

5

6 ¹Department of Pathology, University of Virginia, Charlottesville, VA, USA

7

8 Short title: High and low-risk E6 degrade NHERF1 to activate Wnt signaling

9

10 * Corresponding author

11 Email: vandepol@virginia.edu (SVP)

12

13

14

15

16

17

18

19

20

21

22

23

24 **Abstract**

25 High-risk human papillomavirus (HPV) E6 proteins associate with the cellular
26 ubiquitin ligase E6-Associated Protein (E6AP), and then recruit both p53 and certain
27 cellular PDZ proteins for ubiquitination and degradation by the proteasome. Low-risk
28 HPV E6 proteins also associate with E6AP, yet fail to recruit p53 or PDZ proteins; their
29 E6AP-dependent targets have so far been uncharacterized. We found a cellular PDZ
30 protein called Na⁺/H⁺ Exchanger Regulatory Factor 1 (NHERF1) is targeted for
31 degradation by both high and low-risk HPV E6 proteins as well as E6 proteins from
32 diverse non-primate mammalian species. NHERF1 was degraded by E6 in a manner
33 dependent upon E6AP ubiquitin ligase activity but independent of PDZ interactions. A
34 novel structural domain of E6, independent of the p53 recognition domain, was
35 necessary to associate with and degrade NHERF1, and the NHERF1 EB domain was
36 required for E6-mediated degradation. Degradation of NHERF1 by E6 activated
37 canonical Wnt/ β -catenin signaling, a key pathway that regulates cell growth and
38 proliferation. Expression levels of NHERF1 increased with increasing cell confluency.
39 This is the first study in which a cellular protein has been identified that is targeted for
40 degradation by both high and low-risk HPV E6 as well as E6 proteins from diverse
41 animal papillomaviruses. This suggests that NHERF1 plays a role in regulating
42 squamous epithelial growth and further suggests that the interaction of E6 proteins with
43 NHERF1 could be a common therapeutic target for multiple papillomavirus types.

44

45

46

47 **Author summary**

48 Papillomaviruses cause benign squamous epithelial tumors through the action of virally
49 encoded oncoproteins termed E6 and E7, which are classified as either high or low-risk
50 based upon the propensity of the tumor to evolve into cancer. E6 proteins from both
51 high and low-risk HPVs interact with a cellular ubiquitin ligase called E6AP. High-risk E6
52 proteins hijack E6AP ubiquitin ligase activity to target p53 for degradation. Degradation
53 targets of the low-risk E6 proteins in complex with E6AP have not been described.
54 Here, we describe a protein called NHERF1 that is targeted for degradation by both
55 high and low-risk E6 proteins, as well as E6 proteins from diverse animal species.
56 Degradation of NHERF1 resulted in activation of an oncogenic cellular signaling
57 pathway called Wnt. Identification of NHERF1 as a highly conserved E6 degradation
58 target could inform therapies directed against both low-risk HPVs and cancer-inducing
59 high-risk HPVs.

60

61

62

63

64

65

66

67

68

69

70 **Introduction**

71 Human papillomaviruses (HPVs) are small DNA tumor viruses that cause
72 squamous epithelial papillomas in which the virus replicates. The papillomas are initially
73 benign and the host is usually able to clear the underlying HPV infection over time.
74 However, a subset of HPV infections may result in lesions that persist and grow to
75 harmful size or that have a propensity to evolve into carcinomas [1]. The cancer-causing
76 HPV types are called high-risk and the most commonly occurring high-risk types are
77 HPV16 and HPV18. Worldwide, high-risk HPVs are responsible for 5% of cancers, with
78 cervical cancer being the most common [2]. HPV types that are not associated with
79 malignancies are termed low-risk HPV; although low-risk for malignancies, the size and
80 location of the benign papillomas can render these lesions medically serious [3].

81 Beyond HPVs, papillomaviruses have been isolated from mammalian species
82 including rodents, primates, bats, cetaceans, and ungulates [4], and are clustered into
83 related genera based upon the divergence of the L1 capsid protein nucleotide sequence
84 (both high and low-risk mucosal HPV types discussed in this study belong to the
85 primate Alpha genera) [5]. Most non-human papillomaviruses encode E6 proteins that
86 are similar in predicted fold to high-risk HPV16 E6 [6]. When diverse mammalian
87 papillomaviruses are clustered based on their E6 sequence similarity, two main groups
88 of papillomaviruses emerge: those that encode E6 proteins that bind to the Notch co-
89 activator Mastermind Like 1 (MAML1), and those that bind to a cellular E3 ubiquitin
90 ligase called E6-Associated Protein (E6AP) [7]. An E6 protein that preferentially binds
91 MAML1 suppresses MAML1 transcriptional activation, while an E6 protein that
92 preferentially binds E6AP stimulates E6AP E3 ubiquitin ligase activity to then target

93 additional cellular proteins recruited by E6 to E6AP for ubiquitination and degradation by
94 the proteasome [7].

95 The difference between the propensity of high and low-risk HPVs to cause
96 cancer is secondary to differences between their respective E6 and E7 oncoproteins [8].
97 E6 and E7 from both high and low-risk HPVs bind cellular E3 ubiquitin ligases and
98 hijack their ubiquitin ligase activity to perturb certain cellular proteins that are recruited
99 by E6 or E7 [9]. Both high-risk and low-risk E7 proteins interact with ubiquitin ligases of
100 the cullin and N-end rule families and target the degradation of additional cellular
101 proteins recruited to E7 such as pocket-family proteins (RB, RBL1, and RBL2) and
102 PTPN14 [10, 11]. High and low-risk E7 proteins target certain cellular proteins in
103 common (such as the RBL2 p130 pocket protein) [12]. However only high-risk HPV E7
104 types interact with and target the degradation of the retinoblastoma (RB) pocket protein
105 [13, 14], which has implications for the carcinogenic properties of high-risk E7. High and
106 low-risk Alpha genera HPV E6 proteins interact with the cellular E3 ubiquitin ligase
107 E6AP [15-17], but only cellular proteins targeted for degradation by the high-risk E6
108 protein (such as p53) are well established [18, 19].

109 Another striking difference between high and low-risk E6 is the presence of a
110 PDZ binding motif (PBM) at the extreme carboxy terminus of high-risk E6 proteins [20-
111 22]]. The high-risk PBM enables E6 to interact with a group of cellular proteins termed
112 PDZ proteins, all of which contain PDZ (PSD-90/Dlg1/ZO-1) homology domains [23].
113 The targeted degradation of cellular proteins that are recruited through interaction with
114 the high-risk E6 PBM has been controversial, but the E6 PBM functionally promotes
115 retention of the viral DNA plasmid within infected cells [24]; the E6 PBM function can be

116 rescued by disruption of p53 function [25]. Although low-risk E6 proteins bind E6AP,
117 they do not have a PBM at the carboxy-terminus [20], do not interact with p53 [17], and
118 no cellular targets of the low-risk E6+E6AP complex have been described. Such cellular
119 targets would be presumed to be of exceptional interest since they would be common to
120 both high and low-risk E6 proteins, just as RBL2 is a common target of the high and
121 low-risk HPV E7 protein.

122 In this study, we identify the PDZ-adaptor protein NHERF1 as degraded by both
123 high and low-risk E6 proteins, in a manner dependent upon the ubiquitin ligase activity
124 of E6AP and the proteasome. Other E6 proteins from diverse species where E6 could
125 bind E6AP were also able to initiate NHERF1 degradation, indicating the conservation
126 of this function. Interaction of NHERF1 with E6 required prior association of E6 with
127 E6AP, and we identified a novel interaction domain within 16E6 that is required. Finally,
128 the targeted degradation of NHERF1 by both low and high-risk E6 proteins resulted in
129 the activation of canonical Wnt signaling, connecting the degradation of NHERF1 by E6
130 to the activation of an oncogenic signaling pathway.

131

132 **Results**

133 **NHERF1 protein levels are reduced by high and low-risk E6 proteins in the** 134 **presence of E6AP_WT.**

135 NHERF1 was previously shown to be degraded by HPV16 E6 (16E6) (but not by
136 HPV18 E6 (18E6) or HPV11 E6 (11E6)) through an interaction requiring the PBM of
137 16E6 [26]. In our proteomic studies of cellular proteins that associate with the 16E6 and
138 18E6 PBMs [27], we did not identify NHERF1, but in other experiments observed a

139 reduction of NHERF1 protein levels by 16E6, 18E6, and 11E6. In order to characterize
140 the reduction of NHERF1 by these E6 proteins, we performed transient transfections
141 into E6AP-null 8B9 cells reconstituted with either WT E6AP (E6AP_WT) or a mutant
142 E6AP defective in ubiquitin ligase activity (E6AP_Ub⁻). E6APs were co-transfected with
143 plasmids encoding p53, NHERF1, and 16E6, 16E6 deleted of the PBM (16E6 Δ PBM),
144 11E6, or 18E6. Consistent with published literature, p53 was degraded by high-risk E6
145 proteins (16E6 and 18E6) independently of a PBM [25] and dependent upon E6AP
146 ubiquitin ligase activity [28] (Fig 1). Expression of 11E6 together with E6AP_WT
147 resulted in a lack of p53 degradation by low-risk E6 (11E6), corroborating published
148 findings [17, 29]. However, NHERF1 protein levels were reduced by each observed E6
149 protein (Fig 1A), in contrast to what has been previously published [26].

150 To ensure the reduction of NHERF1 by either high or low-risk E6 proteins was
151 not due to an overexpression artifact, we performed an E6 titration experiment (Fig 1B).
152 Representative western blots from which the quantification in Fig 1B was derived are
153 shown in S1 Fig. Three different E6 proteins were used: 16E6_WT, 16E6 Δ PBM, and
154 11E6_WT. We used p53 as a control for 16E6-mediated degradation, as multiple
155 studies have shown low-risk E6 proteins (11E6) do not degrade p53 [17, 29] while high-
156 risk 16E6 is able to degrade p53 independent of the presence of a PBM [25]. Observing
157 the degradation of p53 in cells expressing variable amounts of E6 provided a guide for
158 physiologically relevant E6 expression levels. NHERF1 and p53 protein levels were
159 similarly reduced by both 16E6_WT and 16E6 Δ PBM at the various E6 titrations (Fig
160 1B). 11E6_WT was unable to initiate the degradation of p53 but targeted NHERF1 at

161 levels similar to those required by 16E6. Deletion of the 16E6 PBM did not impact the
162 degradation of p53 or reduction of NHERF1 protein levels by 16E6.

163 **NHERF1 protein levels are sensitive to cell confluency.**

164 We established that NHERF1 protein levels are reduced by E6 in a transient
165 transfection system. To determine whether low levels of stable 16E6-expression could
166 initiate the reduction of NHERF1 protein levels, we retrovirally transduced normal
167 immortalized keratinocytes with either empty vector or 16E6_WT and observed
168 NHERF1 protein levels. Initially, our results were variable. We hypothesized that
169 keratinocyte confluency may affect NHERF1 protein levels. To test this possibility, we
170 seeded vector-transduced and 16E6_WT-transduced keratinocytes at three different
171 cell densities: 5×10^3 cells/cm² (very sub-confluent), 1.3×10^4 cells/cm² (sub-confluent),
172 and 2.6×10^4 cells/cm² (mid-confluent). NHERF1 protein levels increased with an
173 increase in cell density and 16E6_WT consistently reduced NHERF1 (Fig 2A). In order
174 to determine if changes in NHERF1 levels with confluency were secondary to changes
175 in NHERF1 RNA levels, we performed qPCR on RNA extracted from keratinocytes
176 retrovirally transduced with vector or 16E6_WT and plated as in Fig 2A. Interestingly,
177 NHERF1 RNA levels did not differ between keratinocytes seeded at different densities
178 expressing either empty vector or 16E6_WT (Fig 2B).

179 **E6-mediated degradation of NHERF1 occurs via the proteasome.**

180 Because the ability of each E6 protein to reduce NHERF1 protein levels was also
181 dependent upon the ubiquitin ligase activity of E6AP (Fig 1A), we hypothesized that E6
182 reduction of NHERF1 levels would be secondary to proteasome activity. We seeded
183 retrovirally transduced keratinocytes expressing either empty vector or 16E6_WT at

184 similar confluency and treated with either DMSO, mitomycin C (MMC) to induce p53
185 [30], or the proteasome inhibitor MG132 at differing concentrations for 8 hours. As
186 expected, p53 levels increased in vector keratinocytes treated with MMC compared to
187 untreated cells as well as in 16E6_WT cells exposed to increasing concentrations of
188 MG132 [28] (Fig 3). NHERF1 protein levels increased significantly in a dose dependent
189 manner upon treatment with MG132 in parallel to that seen with p53. (Fig 3, lanes 3-8).
190 This indicated that NHERF1 is degraded through the proteasome by E6 in a manner
191 dependent upon WT E6AP.

192 **E6-mediated degradation of NHERF1 is conserved across papillomaviruses from**
193 **diverse hosts.**

194 The observation that NHERF1 was targeted by both high and low-risk HPV E6
195 proteins suggested that NHERF1 may also be a target of diverse non-primate E6
196 proteins. We examined the ability of E6 proteins from multiple different genera and
197 different mammalian species to target NHERF1 for degradation (Fig 4). E6 proteins that
198 preferentially bind MAML1 were unable to degrade NHERF1 (Fig 4A and 4B). All of the
199 tested Alpha (primate), Dyodelta (boar), and Dyopi (porpoise) genera E6 proteins that
200 bind E6AP targeted NHERF1. While E6AP-binding was necessary it was not sufficient,
201 as E6 proteins from Omega (polar bear, UmPV1) and Omikron (cetaceans, PphPV1
202 and TtPV5) did not degrade NHERF1 (Fig 4A). Interestingly, E6 proteins that bind E6AP
203 but did not target NHERF1 degradation sequence-clustered separately from E6 proteins
204 that did target NHERF1 degradation, suggesting evolutionary divergence of this function
205 (Fig 4B).

206 **A novel 16E6 substrate interaction domain is required for 16E6 degradation of**
207 **NHERF1.**

208 Because the ability of E6 to degrade NHERF1 was not dependent upon the
209 presence of a PBM (Figs 1 and 4), we attempted to identify which residue(s) of 16E6
210 were required to mediate degradation of NHERF1. The crystal structure of 16E6
211 complexed with the E6-binding peptide from E6AP [31] (Fig 6A) was examined to
212 identify amino acids that were at least 20% exposed, resulting in over eighty candidate
213 residues (S2 Fig). Candidate residues were individually mutated in the context of the
214 16E6 gene and the resulting point mutants were screened for their ability to degrade
215 NHERF1 in the presence of E6AP_WT in transiently transfected C33 cells. To ensure
216 our point mutants were not functionally defective (i.e. could not fold properly or could not
217 interact with E6AP), we also screened the mutants for ability to degrade p53. A
218 selection of mutants and the results of the screen are shown in Fig 5. Four mutants
219 stood out as selectively defective in their ability to degrade NHERF1 (Fig 5B) while still
220 being able to degrade p53 (Fig 5C): F69A, K72A, F69R and a double mutant:
221 F69A/K72A. As evidenced in the crystal structure of 16E6, the side chains of F69 and
222 K72 (Fig 6B) are located along the connecting alpha-helix that links the amino-terminal
223 and carboxy-terminal zinc-structured domains of 16E6. The F69 and K72 side chains
224 are aligned and adjacent on the connecting helix, which is on the opposite side of 16E6
225 from the p53 interaction surface [32] (Fig 6C).

226 We had identified the 16E6_F69A/K72A mutant in a transient transfection
227 screen. To ensure the identified 16E6_F69A/K72A double mutant was selectively
228 defective for degrading NHERF1 in the context of a stable cell line, keratinocytes

229 retrovirally transduced with empty vector, 16E6_WT, 16E6 Δ PBM, 16E6_F69A/K72A, or
230 11E6_WT were seeded at equal confluency and lysates prepared. Keratinocytes
231 expressing 16E6_WT, 16E6 Δ PBM, and 11E6_WT degraded NHERF1 (Fig 7, lanes 2, 3,
232 5). However, keratinocytes expressing 16E6_F69A/K72A were unable to stimulate the
233 degradation of NHERF1 (Fig 7, lane 4), indicating a novel substrate interaction domain
234 important for 16E6-mediated degradation of NHERF1.

235 **Degradation of NHERF1 by 16E6 requires the NHERF1 EB domain.**

236 Because the PBM of E6 proteins is not required to initiate the degradation of
237 NHERF1 (Figs 1, 4, and 7), we hypothesized that neither of the PDZ domains of
238 NHERF1 would be required for 16E6 to initiate NHERF1 degradation. We truncated
239 NHERF1 and deleted several characterized domains within the protein [33, 34] (Fig 8A).
240 E6AP-null 8B9 cells were co-transfected with 16E6_WT, NHERF1 truncations,
241 HA_GFP, and either E6AP_Ub⁻ or E6AP_WT. NHERF1 protein levels were quantified,
242 and then normalized to the internal transfection control (HA_GFP). The various
243 NHERF1 truncations displayed different expression levels. To account for these
244 variations, levels of NHERF1 truncations in the presence of E6AP_Ub⁻ were set to
245 100% and the expression level of the corresponding NHERF1 truncation in the
246 presence of E6AP_WT was normalized accordingly (Fig 8B and 8C, bar graphs). All
247 NHERF1 truncations containing the EB domain were targeted for degradation by 16E6
248 in the presence of E6AP_WT (highlighted in green in Fig 8A). Truncations of NHERF1
249 that lacked the EB domain were not targeted for degradation by 16E6 (highlighted in red
250 in Fig 8A). In addition, the NHERF1 PBM was not required for 16E6 mediated
251 degradation (Fig 8C, lanes 5 vs. 6 and 9 vs. 10).

252 We identified the NHERF1 EB domain as a requirement for 16E6 mediated
253 degradation and the importance of 16E6 residues F69 and K72. In order to examine the
254 interactions between the 16E6+E6AP+NHERF1 complex, all three proteins were
255 expressed in a yeast three-hybrid system so as to detect the heterotrimeric complex.
256 We fused 16E6_WT and ubiquitin ligase dead E6AP (E6AP_Ub⁻) to the LexA DNA
257 binding domain and co-expressed this fusion with either vector, 16E6_WT, or
258 16E6_F69A/K72A in yeast containing a LexA responsive LacZ reporter. These yeast
259 were then mated to yeast expressing native p53 or Gal4 (G4) transactivator fusions to
260 NHERF1 121-358 (containing the EB domain), NHERF1 121-297 (deleted of the EB
261 domain), 16E6_WT, or the tyrosine phosphatase PTPN3 (a PDZ protein) (Fig 9). The
262 LexA_16E6 fusion co-expressed with p53 (in the absence of E6AP) resulted in very
263 weak activation of the LacZ reporter (spot 4B) while co-expression with G4_PTPN3
264 resulted in strong transactivation (spot 6B), but no interaction with NHERF1 (spots 2B
265 and 3B). We then co-expressed 16E6 and E6AP by using a LexA_E6AP_Ub⁻ fusion
266 together with native 16E6. When LexA_E6AP_Ub⁻, untagged 16E6_WT, and p53 were
267 co-expressed, a strong activation of the LacZ reporter was observed (Fig 9, spot 4D),
268 illustrating that while p53 has a weak direct interaction with 16E6, it interacts strongly
269 with 16E6 bound to E6AP. This activation was also seen with 16E6_F69A/K72A in the
270 presence of LexA_E6AP_Ub⁻ and p53 (Fig 9, spot 4E), indicating the preserved ability
271 of the 16E6_F69A/K72A double mutant to bind E6AP and recruit p53. When
272 LexA_E6AP_Ub⁻, 16E6_WT, and G4_NHERF1 121-358 (contains the EB domain) were
273 co-expressed, we observed activation of the LacZ reporter, indicating the recruitment of
274 NHERF1 to E6AP by 16E6_WT (Fig 9, spot 2D). Truncating the EB domain from the

275 G4_NHERF1 (G4_NHERF1 121-297, spot 3D) or the use of the 16E6_F69A/K72A
276 double mutant (spot 2E) ablated the reporter transactivation, indicating the requirement
277 of the EB domain and the importance of 16E6 residues F69 and K72 in the interaction
278 of the 16E6+E6AP+NHERF1 complex.

279 **E6AP-dependent NHERF1 degradation by E6 activates the canonical Wnt/ β -**
280 **catenin pathway.**

281 It has been shown that high-risk HPV E6 proteins augment the canonical Wnt/ β -
282 catenin signaling pathway [35-39]. Additionally, it has been shown that NHERF1 inhibits
283 the canonical Wnt/ β -catenin signaling pathway through multiple mechanisms. NHERF1
284 forms a complex with β -catenin [40] and can also bind to the intracellular PBM of certain
285 isoforms of Frizzled [41], a G-protein coupled receptor important in the activation of the
286 canonical Wnt signaling pathway. Therefore, we hypothesized E6 degradation of
287 NHERF1 would activate the Wnt/ β -catenin signaling pathway in cells expressing E6. To
288 test this possibility, we utilized the TOP/FOP luciferase reporter assay. 16E6,
289 16E6 Δ PBM, 11E6, and 18E6 all stimulated the activity of the Wnt/ β -catenin pathway
290 over vector-transfected cells (Fig 10). However, cells transfected with 16E6_F69A/K72A
291 were unable to augment the canonical Wnt pathway over vector levels, indicating that
292 the ability of E6 to degrade NHERF1 is required for E6 activation of the canonical
293 Wnt/ β -catenin signaling pathway.

294 The earlier Accardi et al. study proposed that expression of 16E7 sensitized
295 NHERF1 for degradation by the induction of NHERF1 phosphorylation [26]. Our
296 experiments did not show either E7 induction of slow-migrating NHERF1

297 phosphorylated isoforms or an enhancement of E6-NHERF1 degradation upon co-
298 expression of E7 (S3 and S4 Figs).

299

300 **Discussion**

301 E6 proteins from papillomaviruses can be separated into distinct groups: those
302 that bind MAML1 and repress Notch signaling, and those that bind E6AP and hijack its
303 ubiquitin ligase activity [7, 42-44]. E6 proteins from papillomaviruses in the Alpha,
304 DyoDelta, Dyopi, Omega, and Omikron genera all behave similarly in that they bind
305 E6AP and activate its ubiquitin ligase activity [7]. Here, we describe the degradation of
306 NHERF1 by E6 proteins from both high and low-risk HPVs, as well as from
307 papillomaviruses from multiple divergent mammalian species. The ability of these E6s
308 to degrade NHERF1 is dependent upon E6AP (Fig 1) and the proteasome (Fig 3). In
309 addition, we identify two amino acids in 16E6 (F69 and K72) that are necessary for E6-
310 mediated degradation of NHERF1. These two residues are aligned, and adjacent in the
311 outwardly oriented face of the E6 connecting alpha helix, suggesting a novel interaction
312 domain (Fig 6B and 6C). NHERF1 degradation by E6 requires the NHERF1 EB domain,
313 but does not require the PBM at the extreme carboxy terminus of NHERF1 (Fig 8B and
314 8C). The ability of E6 proteins to degrade NHERF1 augments the canonical Wnt/ β -
315 catenin signaling (Fig 10), an oncogenic pathway frequently active in cancer.

316 NHERF1 is the product of the SLC9A3R1 gene. SLC9A3R1 mRNA is broadly
317 expressed in epithelia, with the highest mRNA expression in kidney, gut, and
318 esophagus. NHERF1 is not developmentally essential, although mice have considerably
319 reduced lifespans [45]. NHERF1-null mice are prone to phosphate wasting, brittle bone

320 structure, and hydrocephaly [45] due to the mislocalization of proteins with which
321 NHERF1 normally associates [45-47]. NHERF1 contains two PDZ domains and an EB
322 domain at the carboxy terminus through which it interacts with ezrin, radixin, and
323 moesin to link itself, and proteins to which it is bound, to the actin cytoskeleton network
324 [48]. While the functions of NHERF1 are varied due to its role as a scaffold, multiple
325 studies indicate it regulates cell growth and differentiation, two key cellular functions that
326 papillomaviruses disrupt in the process of viral infection.

327 Whether NHERF1 is a tumor suppressor or an oncogene has been debated in
328 the literature. There are numerous papers regarding NHERF1 human cancer
329 phenotypes, but they are collectively inconsistent [49, 50]. NHERF1-null mice do not
330 have a direct cancer phenotype, but have lengthened intestines [51], indicating a growth
331 regulatory function of NHERF1. The diminished life span of NHERF1 null mice could
332 limit observation of cancer traits. However, a recent in vivo study provided strong
333 genetic support for NHERF1 as a tumor suppressor. APC^{Min/+} mice bred as either
334 heterozygote or knockout for NHERF1 experience considerably shorter survival than
335 their NHERF1-expressing counterparts due to increased tumor burden, demonstrating a
336 tumor suppressor phenotype for NHERF1 [51]. Additionally, these APC^{Min/+} mice lacking
337 NHERF1 have greater activation of Wnt/ β -catenin signaling, suggesting NHERF1 acts
338 as a negative regulator of this oncogenic pathway. NHERF1-associated proteins that
339 plausibly could regulate cell proliferation are numerous and include β -catenin [40],
340 Frizzled [41], G-protein coupled receptors (β -adrenergic type 2, [52]), receptor tyrosine
341 kinases (PDGFR, [53]), phosphatases (PTEN, [54]), transcriptional coactivators (YAP1,

342 [55]), ion channels (Kir1.1 and CFTR, [56]), phospholipase-C [57], and actin anchoring
343 proteins (ezrin, radixin, and moesin, [48]).

344 Several studies have indicated that HPV E6 proteins can activate canonical
345 Wnt/ β -catenin signaling [35-39]. Our work expands and builds upon the scope of
346 these studies. The ability of E6 to degrade NHERF1 and activate Wnt signaling may
347 aid in propagation of papillomaviruses by enhancing the stimulation of cellular
348 proliferation and promoting cell survival. There are numerous cell growth regulatory
349 avenues that E6 could manipulate by degrading NHERF1 and within this study we
350 have explored one possibility: the canonical Wnt/ β -catenin signaling pathway (Fig
351 10); other possibilities will be the subject of future studies.

352 The EB domain of NHERF1 is required for E6-mediated degradation in the
353 presence of E6AP (Fig 8B and 8C). This domain is responsible for linking NHERF1 to
354 the actin cytoskeleton network via interaction with ERM proteins [48]. NHERF1 has a
355 PBM at its extreme carboxy terminus and when the EB domain is not bound to ERM
356 proteins, the NHERF1 PBM can self-associate with the NHERF1 PDZ2 domain,
357 resulting in a closed NHERF1 conformation [33]. The head-to-tail closed NHERF1
358 confirmation is not required for E6-mediated degradation, as an NHERF1 Δ PBM mutant
359 was still targeted for degradation by E6 in the presence of E6AP_WT (Fig 8C). Nor was
360 the 16E6 PBM required for degradation of NHERF1 (Figs 1, 4, and 7), contrary to a
361 prior report [26].

362 In addition to the requirement of the NHERF1 EB domain, two E6 residues, F69
363 and K72, are necessary for E6-mediated NHERF1 degradation (Figs 5 and 7). Crucially,
364 the 16E6_F69A/K72A double mutant can still initiate the degradation of p53, indicating it

365 is still able to bind E6AP, recruit p53 to the complex, and trigger ubiquitination. The F69
366 and K72 residues are also required to form a tri-molecular complex between E6AP, E6,
367 and NHERF1 in yeast (Fig 9, spot 2E vs. 2D). Like the association of E6 with p53,
368 NHERF1 does not interact directly with E6, but requires prior association of E6 with
369 E6AP, indicating that NHERF1 requires an altered conformation of E6 that is secondary
370 to E6 binding to E6AP [58].

371 As we were testing the ability of E6 proteins to degrade NHERF1 in stable
372 keratinocyte cell lines, we discovered that NHERF1 protein levels are sensitive to cell
373 confluency (Fig 2A). The relationship between NHERF1 and cell confluency may
374 contribute to the lack of identification of NHERF1 as a degradation target of low-risk E6
375 proteins in the past, as well as differences between our studies and a prior publication
376 [26]. It is likely that this observation underlies disparate findings between different
377 laboratories regarding NHERF1 cancer associated traits [49, 50]. Future studies of
378 NHERF1 must take into account and carefully control cell densities when performing
379 experiments.

380 Binding to E6AP is necessary for E6-induced degradation of NHERF1, but it is
381 not sufficient, as three tested E6 proteins that bind E6AP do not target NHERF1 for
382 degradation: UmPV1 E6 (polar bear), PphPV1 E6 (porpoise), and TtPV5 E6 (bottlenose
383 dolphin) (Fig 4A). Interestingly, the three E6 proteins that do not degrade NHERF1
384 cluster together in phylogenetic relatedness (Fig 4B). We utilized transfected human
385 NHERF1 throughout our study, so it is possible that the inability of these three E6
386 proteins to target NHERF1 for degradation may be due to evolutionary divergence in the
387 NHERF1 homologs. Future studies will explore if the lack of degradation of human

388 NHERF1 by UmPV1, PphPV1, and TtPV5 is due to evolutionary divergence of the
389 respective NHERF1 proteins compared to human NHERF1. It would be of interest to
390 determine if NHERF1 is a “universal” target of E6 proteins that act through association
391 with E6AP.

392 Discovery of NHERF1 as a novel target for not only high and low-risk mucosal
393 and cutaneous HPV E6 as well as a wide range of E6 proteins across divergent host
394 species indicates a significant and previously undescribed role for NHERF1 in
395 papillomavirus biology. That NHERF1 is a conserved target of papillomavirus E6
396 proteins further elevates the importance of NHERF1 as a cell growth regulator. Finally,
397 the identification of this highly conserved E6 degradation target may represent a novel
398 avenue for therapeutic intervention against both low and high-risk HPV.

399

400 **Materials and methods**

401 **Cells and cell culture**

402 E6AP-null 8B9 cells (a gift of Dr. Lawrence banks, ICGEB, Italy) [59] and HPV-negative
403 C33A cervical cancer cells (ATCC) were maintained and transfected using
404 polyethylenimine (PEI) as previously described [58]. Normal immortalized keratinocytes
405 (NIKS, obtained from ATCC) are spontaneously immortalized foreskin keratinocytes [60]
406 that were cocultured with mitomycin C-treated 3T3 feeder cells in F medium as
407 described previously [61]. NIKS were retrovirally transduced with replication-defective
408 murine retroviruses based on pLXSN [62] as previously described [25]. Retrovirally
409 transduced NIKS cells were counted and seeded at equal confluency in each
410 experiment.

411 **Plasmids**

412 Epitope tagged E6AP, GFP, E6, and NHERF1 were all transiently expressed from the
413 pcDNA3 plasmid. HA-tagged NHERF1 originated from Vijaya Ramesh's laboratory
414 (from Addgene, plasmid 11635). 16E6 point mutants were created using QuikChange
415 primer design (Agilent Technologies). NHERF1 truncations were PCR generated and
416 sequenced.

417 **Antibodies and Western blots**

418 12 well plates of transfected mammalian cells were lysed in 0.5X IPEGAL as described
419 previously [7]. Transduced NIKS were lysed in 1% SDS, 5mM EDTA, and 1 mM sodium
420 vanadate and equilibrated for protein content (Biorad assay kit). All lysates were
421 resolved by SDS-PAGE electrophoresis and transferred to PVDF membranes.

422 Antibodies: anti-HA (Bethyl Laboratories, Inc.), anti-FLAG M2 (Sigma), anti-p53 Ab-8
423 (ThermoFisher Scientific), anti-16E6 6G6 (a generous gift from Arbor Vita Corporation),
424 anti-SLC9A3R1 (Sigma), anti-GAPDH (Cell Signaling Technology), and anti-MYC 9B11
425 (Cell Signaling Technology).

426 **RT-PCR**

427 Retrovirally transduced NIKS were plated at different cell densities and harvested
428 following a TRizol RNA harvest protocol (Invitrogen). cDNA was generated using
429 random hexamers. Quantitative real-time PCR was performed on the cDNA using iQ™
430 SYBR® Green Supermix (BioRad #1708880). The primers targeted the SLC9A3R1
431 gene (BioRad Assay ID: qHsaCEP0050521) and the GAPDH gene (BioRad Assay ID:
432 qHsaCEP0041396). Relative values were analyzed using the $\Delta\Delta C_T$ method (where C_T
433 is the threshold cycle) and GAPDH as a control.

434 **Wnt/ β -catenin luciferase reporter assay**

435 C33A cells plated at 70% confluency were transiently transfected with DNA of the
436 TOPFLASH or control FOPFLASH (containing mutated TCF/ β -catenin binding sites; 1
437 ug) plasmid, Renilla luciferase (0.005 ug) plasmid (used to evaluate transfection
438 efficiency), FLAG_E6AP_WT (0.35 ug) plasmid, and the indicated E6 plasmids (0.3 ug).
439 18 hrs post-transfection, media was removed and Wnt3A conditioned media was added
440 for 8.5 hours to stimulate the Wnt pathway. Luciferase levels were measured using the
441 Dual-Luciferase® Reporter Assay System (Promega) and a Cytation1 Plate Reader
442 (software version 3.04.17). FOPFLASH luciferase readings were low, and were
443 subtracted from the paired TOPFLASH readouts. 10% fetal bovine serum Wnt3A
444 conditioned media was generated using L Wnt-3A murine fibroblasts (ATCC, CRL-
445 2647) as previously described [63].

446 **Phylogenetic analysis**

447 Multiple protein sequence files were downloaded from the Papillomavirus Episteme [64]
448 and aligned using the EMBL-EBI MUSCLE (Multiple Sequence Comparison by Log-
449 Expectation) program [65]. The phylogenetic tree was generated as a neighbour-joining
450 tree without distance corrections within the MUSCLE program [65].

451 **Yeast expression**

452 Modified LexA-based yeast three-hybrid assays were performed as previously
453 described [58].

454

455 **Acknowledgements**

456 We thank Nicole Brimer for assistance with technical challenges that were encountered
457 throughout this study and Kelly Drews for extensive discussions and for critical reading
458 of the manuscript.

459

460 **References**

- 461 1. zur Hausen H. Papillomaviruses in the causation of human cancers - a brief
462 historical account. *Virology*. 2009;384(2):260-5.
- 463 2. de Martel C, Plummer M, Vignat J, Franceschi S. Worldwide burden of cancer
464 attributable to HPV by site, country and HPV type. *Int J Cancer*. 2017;141(4):664.
- 465 3. Egawa N, Doorbar J. The low-risk papillomaviruses. *Virus Res*. 2017;231:119-27.
- 466 4. Rector A, Van Ranst M. Animal papillomaviruses. *J Virol*. 2013;445(1):213--23.
- 467 5. de Villiers EM, Fauquet C, Broker TR, Bernard HU, zur Hausen H. Classification
468 of papillomaviruses. *Virology*. 2004;324(1):17-27.
- 469 6. Vande Pol SB. Papillomavirus E6 Oncoproteins Take Common Structural
470 Approaches to Solve Different Biological Problems. *PLoS Pathog*.
471 2015;11(10):e1005138.
- 472 7. Brimer N, Drews CM, Vande Pol SB. Association of papillomavirus E6 proteins
473 with either MAML1 or E6AP clusters E6 proteins by structure, function, and
474 evolutionary relatedness. *PLoS Pathog*. 2017;13(12):e1006781.
- 475 8. Klingelhutz AJ, Roman A. Cellular transformation by human papillomaviruses:
476 lessons learned by comparing high- and low-risk viruses. *Virology*.
477 2012;424(2):77-98.

- 478 9. Davey NE, Trave G, Gibson TJ. How viruses hijack cell regulation. Trends
479 Biochem Sci. 2011;36(3):159-69.
- 480 10. Roman A, Munger K. The papillomavirus E7 proteins. Virology. 2013;445(1-
481 2):138-68.
- 482 11. White EA, Münger K, Howley PM. High-Risk Human Papillomavirus E7 Proteins
483 Target PTPN14 for Degradation. MBio. 2016;7(5):01530--16.
- 484 12. Zhang B, Chen W, Roman A. The E7 proteins of low- and high-risk human
485 papillomaviruses share the ability to target the pRB family member p130 for
486 degradation. Proc Natl Acad Sci U S A. 2006;103(2):437.
- 487 13. Münger K, Werness BA, Dyson N, Phelps WC, Harlow E, Howley PM. Complex
488 formation of human papillomavirus E7 proteins with the retinoblastoma tumor
489 suppressor gene product. EMBO J. 1989;8(13):4099--105.
- 490 14. Boyer SN, Wazer DE, Band V. E7 Protein of Human Papilloma Virus-16 Induces
491 Degradation of Retinoblastoma Protein through the Ubiquitin-Proteasome
492 Pathway. Cancer Res. 1996;56(20):4620--4.
- 493 15. Huibregtse JM, Scheffner M, Howley PM. A cellular protein mediates association
494 of p53 with the E6 oncoprotein of human papillomavirus types 16 or 18. EMBO J.
495 1991;10(13):4129-35.
- 496 16. Huibregtse JM, Scheffner M, Howley PM. Cloning and expression of the cDNA
497 for E6-AP, a protein that mediates the interaction of the human papillomavirus E6
498 oncoprotein with p53. Mol Cell Biol. 1993;13(2):775-84.
- 499 17. Brimer N, Lyons C, Vande Pol SB. Association of E6AP (UBE3A) with human
500 papillomavirus type 11 E6 protein. Virology. 2007;358(2):303-10.

- 501 18. Cooper B, Schneider S, Bohl J, Jiang Y, Beaudet A, Vande Pol S. Requirement
502 of E6AP and the features of human papillomavirus E6 necessary to support
503 degradation of p53. *Virology*. 2003;306(1):87-99.
- 504 19. Vande Pol SB, Klingelutz AJ. Papillomavirus E6 oncoproteins. *Virology*.
505 2013;445(1-2):115-37.
- 506 20. Lee SS, Weiss RS, Javier RT. Binding of human virus oncoproteins to
507 hDlg/SAP97, a mammalian homolog of the *Drosophila* discs large tumor
508 suppressor protein. *Proc Natl Acad Sci U S A*. 1997;94(13):6670.
- 509 21. Songyang Z, Fanning AS, Fu C, Xu J, Marfatia SM, Chishti AH, et al. Recognition
510 of Unique Carboxyl-Terminal Motifs by Distinct PDZ Domains. *Science*.
511 1997;275(5296):73--7.
- 512 22. Kiyono T, Hiraiwa A, Fujita M, Hayashi Y, Akiyama T, Ishibashi M. Binding of
513 high-risk human papillomavirus E6 oncoproteins to the human homologue of the
514 *Drosophila* discs large tumor suppressor protein. *Proc Natl Acad Sci U S A*.
515 1997;94(21):11612--6.
- 516 23. Ganti K, Broniarczyk J, Manoubi W, Massimi P, Mittal S, Pim D, et al. The
517 Human Papillomavirus E6 PDZ Binding Motif: From Life Cycle to Malignancy.
518 *Viruses*. 2015;7(7):3530.
- 519 24. Lee C, Laimins LA. Role of the PDZ domain-binding motif of the oncoprotein E6
520 in the pathogenesis of human papillomavirus type 31. *J Virol*.
521 2004;78(22):12366--77.

- 522 25. Brimer N, Vande Pol SB. Papillomavirus E6 PDZ interactions can be replaced by
523 repression of p53 to promote episomal human papillomavirus genome
524 maintenance. *J Virol.* 2014;88(5):3027-30.
- 525 26. Accardi R, Rubino R, Scalise M, Gheit T, Shahzad N, Thomas M, et al. E6 and
526 E7 from human papillomavirus type 16 cooperate to target the PDZ protein Na/H
527 exchange regulatory factor 1. *J Virol.* 2011;85(16):8208-16.
- 528 27. Webb Strickland S, Brimer N, Lyons C, Vande Pol SB. Human Papillomavirus E6
529 interaction with cellular PDZ domain proteins modulates YAP nuclear
530 localization. *Virology.* 2018;516:127-38.
- 531 28. Scheffner M, Huibregtse JM, Vierstra RD, Howley PM. The HPV-16 E6 and E6-
532 AP complex functions as a ubiquitin-protein ligase in the ubiquitination of p53.
533 *Cell.* 1993;75(3):495-505.
- 534 29. Scheffner M, Werness BA, Huibregtse JM, Levine AJ, Howley PM. The E6
535 oncoprotein encoded by human papillomavirus types 16 and 18 promotes the
536 degradation of p53. *Cell.* 1990;63(6):1129-36.
- 537 30. Abbas T, Olivier M, Lopez J, Houser S, Xiao G, Kumar GS, et al. Differential
538 Activation of p53 by the Various Adducts of Mitomycin C. *J Biol Chem.*
539 2002;277(43):40513--9.
- 540 31. Zanier K, Charbonnier S, Sidi AO, McEwen AG, Ferrario MG, Poussin-
541 Courmontagne P, et al. Structural basis for hijacking of cellular LxxLL motifs by
542 papillomavirus E6 oncoproteins. *Science.* 2013;339(6120):694-8.

- 543 32. Martinez-Zapien D, Ruiz FX, Poirson J, Mitschler A, Ramirez J, Forster A, et al.
544 Structure of the E6/E6AP/p53 complex required for HPV-mediated degradation of
545 p53. *Nature*. 2016;529(7587):541-5.
- 546 33. Morales FC, Takahashi Y, Momin S, Adams H, Chen X, Georgescu MM.
547 NHERF1/EBP50 head-to-tail intramolecular interaction masks association with
548 PDZ domain ligands. *Mol Cell Biol*. 2007;27(7):2527-37.
- 549 34. Bhattacharya S, Dai Z, Li J, Baxter S, Callaway DJE, Cowburn D, et al. A
550 Conformational Switch in the Scaffolding Protein NHERF1 Controls Autoinhibition
551 and Complex Formation. *J Biol Chem*. 2010;285(13):9981--94.
- 552 35. Bello JO, Nieva LO, Paredes AC, Gonzalez AM, Zavaleta LR, Lizano M.
553 Regulation of the Wnt/beta-Catenin Signaling Pathway by Human Papillomavirus
554 E6 and E7 Oncoproteins. *Viruses*. 2015;7(8):4734-55.
- 555 36. Bonilla-Delgado J, Bulut G, Liu X, Cortes-Malagon EM, Schlegel R, Flores-
556 Maldonado C, et al. The E6 oncoprotein from HPV16 enhances the canonical
557 Wnt/beta-catenin pathway in skin epidermis in vivo. *Mol Cancer Res*.
558 2012;10(2):250-8.
- 559 37. Lichtig H, Gilboa DA, Jackman A, Gonen P, Levav-Cohen Y, Haupt Y, et al.
560 HPV16 E6 augments Wnt signaling in an E6AP-dependent manner. *Virology*.
561 2010;396(1):47-58.
- 562 38. Sominsky S, Kuslansky Y, Shapiro B, Jackman A, Haupt Y, Rosin-Arbesfeld R,
563 et al. HPV16 E6 and E6AP differentially cooperate to stimulate or augment Wnt
564 signaling. *Virology*. 2014;468-470:510-23.

- 565 39. Sominsky S, Shterzer N, Jackman A, Shapiro B, Yaniv A, Sherman L. E6
566 proteins of alpha and beta cutaneous HPV types differ in their ability to potentiate
567 Wnt signaling. *Virology*. 2017;509:11-22.
- 568 40. Kreimann EL, Morales FC, de Orbeta-Cruz J, Takahashi Y, Adams H, Liu TJ, et
569 al. Cortical stabilization of beta-catenin contributes to NHERF1/EBP50 tumor
570 suppressor function. *Oncogene*. 2007;26(36):5290-9.
- 571 41. Wheeler DS, Barrick SR, Grubisha MJ, Brufsky AM, Friedman PA, Romero G.
572 Direct interaction between NHERF1 and Frizzled regulates beta-catenin
573 signaling. *Oncogene*. 2011;30(1):32-42.
- 574 42. Brimer N, Lyons C, Wallberg AE, Vande Pol SB. Cutaneous papillomavirus E6
575 oncoproteins associate with MAML1 to repress transactivation and NOTCH
576 signaling. *Oncogene*. 2012;31(43):4639-46.
- 577 43. Meyers JM, Spangle JM, Munger K. The HPV8 E6 protein interferes with NOTCH
578 activation during keratinocyte differentiation. *J Virol*. 2013.
- 579 44. Tan MJ, White EA, Sowa ME, Harper JW, Aster JC, Howley PM. Cutaneous
580 beta-human papillomavirus E6 proteins bind Mastermind-like coactivators and
581 repress Notch signaling. *Proc Natl Acad Sci U S A*. 2012;109(23):E1473-80.
- 582 45. Shenolikar S, Voltz JW, Minkoff CM, Wade JB, Weinman EJ. Targeted disruption
583 of the mouse NHERF-1 gene promotes internalization of proximal tubule sodium-
584 phosphate cotransporter type IIa and renal phosphate wasting. *Proc Natl Acad
585 Sci U S A*. 2002;99(17):11470-5.
- 586 46. Cunningham R, Brazie M, Kanumuru S, E X, Biswas R, Wang F, et al. Sodium-
587 hydrogen exchanger regulatory factor-1 interacts with mouse urate transporter 1

- 588 to regulate renal proximal tubule uric acid transport. *J Am Soc Nephrol.*
589 2007;18(5):1419--25.
- 590 47. Weinman EJ, Wang Y, Wang F, Greer C, Steplock D, Shenolikar S. A C-terminal
591 PDZ motif in NHE3 binds NHERF-1 and enhances cAMP inhibition of sodium-
592 hydrogen exchange. *Biochemistry.* 2003;42(43):12662--8.
- 593 48. Reczek D, Berryman M, Bretscher A. Identification of EBP50: A PDZ-containing
594 phosphoprotein that associates with members of the ezrin-radixin-moesin family.
595 *J Cell Biol.* 1997;139(1):169-79.
- 596 49. Centonze M, Saponaro C, Mangia A. NHERF1 Between Promises and Hopes:
597 Overview on Cancer and Prospective Openings. *Transl Oncol.* 2018;11(2):374-
598 90.
- 599 50. Vaquero J, Nguyen Ho-Boulidoires TH, Claperon A, Fouassier L. Role of the
600 PDZ-scaffold protein NHERF1/EBP50 in cancer biology: from signaling
601 regulation to clinical relevance. *Oncogene.* 2017;36(22):3067-79.
- 602 51. Georgescu MM, Gagea M, Cote G. NHERF1/EBP50 Suppresses Wnt-beta-
603 Catenin Pathway-Driven Intestinal Neoplasia. *Neoplasia.* 2016;18(8):512-23.
- 604 52. Hall RA, Premont RT, Chow CW, Blitzer JT, Pitcher JA, Claing A, et al. The
605 beta2-adrenergic receptor interacts with the Na⁺/H⁺-exchanger regulatory factor
606 to control Na⁺/H⁺ exchange. *Nature.* 1998;392(6676):626-30.
- 607 53. Maudsley S, Zamah AM, Rahman N, Blitzer JT, Luttrell LM, Lefkowitz RJ, et al.
608 Platelet-derived growth factor receptor association with Na⁽⁺⁾/H⁽⁺⁾ exchanger
609 regulatory factor potentiates receptor activity. *Mol Cell Biol.* 2000;20(22):8352-63.

- 610 54. Takahashi Y, Morales FC, Kreimann EL, Georgescu MM. PTEN tumor
611 suppressor associates with NHERF proteins to attenuate PDGF receptor
612 signaling. *EMBO J.* 2006;25(4):910-20.
- 613 55. Mohler PJ, Kreda SM, Boucher RC, Sudol M, Stutts MJ, Milgram SL. Yes-
614 associated protein 65 localizes p62(c-Yes) to the apical compartment of airway
615 epithelia by association with EBP50. *J Cell Biol.* 1999;147(4):879-90.
- 616 56. Yoo D, Flagg TP, Olsen O, Raghuram V, Foskett JK, Welling PA. Assembly and
617 trafficking of a multiprotein ROMK (Kir 1.1) channel complex by PDZ interactions.
618 *J Biol Chem.* 2004;279(8):6863-73.
- 619 57. Tang Y, Tang J, Chen Z, Trost C, Flockerzi V, Li M, et al. Association of
620 mammalian trp4 and phospholipase C isozymes with a PDZ domain-containing
621 protein, NHERF. *J Biol Chem.* 2000;275(48):37559-64.
- 622 58. Ansari T, Brimer N, Vande Pol SB. Peptide interactions stabilize and restructure
623 human papillomavirus type 16 E6 to interact with p53. *J Virol.*
624 2012;86(20):11386-91.
- 625 59. Massimi P, Shai A, Lambert P, Banks L. HPV E6 degradation of p53 and PDZ
626 containing substrates in an E6AP null background. *Oncogene.*
627 2008;27(12):1800-4.
- 628 60. Allen-Hoffmann BL, Schlosser SJ, Ivarie CA, Sattler CA, Meisner LF, O'Connor
629 SL. Normal growth and differentiation in a spontaneously immortalized near-
630 diploid human keratinocyte cell line, NIKS. *J Invest Dermatol.* 2000;114(3):444-
631 55.

- 632 61. Lambert PF, Ozbun MA, Collins A, Holmgren S, Lee D, Nakahara T. Using an
633 immortalized cell line to study the HPV life cycle in organotypic "raft" cultures.
634 *Methods Mol Med.* 2005;119:141-55.
- 635 62. Miller AD, Rosman GJ. Improved retroviral vectors for gene transfer and
636 expression. *Biotechniques.* 1989;7(9):980-2, 4-6, 9-90.
- 637 63. Wickramasinghe SN. Culture of animal cells. A manual of basic technique. 3rd
638 ed. New York: R. Ian Freshney, Wiley-Liss, Inc.; 1996. 75-6 p.
- 639 64. Van Doorslaer K, Li Z, Xirasagar S, Maes P, Kaminsky D, Liou D, et al. The
640 Papillomavirus Episteme: a major update to the papillomavirus sequence
641 database. *Nucleic Acids Res.* 2017;45(D1):499--506.
- 642 65. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high
643 throughput. *Nucleic Acids Res.* 2004;32(5):1792--7.

644

645 **Figure legends**

646 **Fig 1. NHERF1 protein levels are reduced by both high and low-risk E6 proteins.**

647 (A) NHERF1 protein levels are reduced in an E6 and E6AP dependent manner.

648 Plasmids encoding the indicated FLAG_E6AP (1 ug), HA_NHERF1 (0.5 ug), human

649 p53 (0.5 ug), HA_GFP (0.08 ug), and the listed E6 proteins (1 ug) were transiently

650 transfected into E6AP-null 8B9 cells and HA-NHERF1 expression was analyzed by

651 western blot. Reduction of NHERF1 protein levels by high or low-risk E6 requires ligase

652 active E6AP (E6AP_WT) but does not require the E6 PDZ binding motif (PBM).

653 FLAG_18E6* is a truncated splice isoform of 18E6. E6AP_Ub⁻ denotes an E6AP

654 mutant defective for ubiquitin ligase activity. Quantitation is the result of three

655 independent experiments (N=3) where NHERF1 levels are normalized to co-transfected
656 HA_GFP. Shown is a single representative blot. Vertical black line in blots represents
657 removal of an irrelevant sample. The means of triplicate independent experiments \pm
658 standard error are shown. N=3. * <0.05 , ** <0.01 by Student's t-test. (B) Reduction of
659 NHERF1 protein is not an overexpression artifact. Titrations of the indicated E6 proteins
660 were co-transfected with FLAG_E6AP_WT (1 ug), HA_GFP (0.02 ug), and either
661 HA_NHERF1 (0.5 ug) or p53 (0.5 ug) in murine 8B9 cells. With increased E6
662 expression, NHERF1 decreased for each E6 protein parallel with p53. As expected, p53
663 degradation was observed for the high-risk 16E6 proteins (both WT and Δ PBM) but not
664 by low-risk 11E6 protein despite reduction of NHERF1 protein levels by 11E6. The
665 means of triplicate independent experiments \pm standard error are shown.

666

667 **Fig 2. NHERF1 protein levels increase with increased cell density.** (A) Protein
668 levels of endogenous NHERF1 increase with cell confluency. Keratinocytes retrovirally
669 transduced with either vector or 16E6_WT were counted and plated at the indicated cell
670 densities. As confluency increased, NHERF1 protein levels also increased, though still
671 reduced in the presence of 16E6_WT. The means of triplicate independent experiments
672 \pm standard error are shown. (B) NHERF1 RNA levels are not changed by cell
673 confluency or by the presence of 16E6_WT. Total RNA was extracted from
674 keratinocytes retrovirally transduced with either vector or 16E6_WT and plated at the
675 indicated cell densities. cDNA was reverse transcribed and NHERF1 RNA levels
676 determined by qPCR. The means of triplicate independent experiments \pm standard error
677 are shown.

678

679 **Fig 3. Degradation of NHERF1 by 16E6 requires proteasome function.**

680 Keratinocytes retrovirally transduced with either vector or 16E6_WT were seeded at
681 equal confluency. Cells were treated with DMSO, mitomycin C (MMC), or the
682 proteasome inhibitor MG132 at varying concentrations for 8 hours as indicated. MG132
683 significantly rescued NHERF1 protein levels in a dose dependent manner. MMC
684 treatment was used to induce p53 levels, which were observed as a positive control.
685 Quantification was normalized to vector-transduced cells treated with DMSO. The
686 means of triplicate independent experiments \pm standard error are shown. N=3, * <0.05 ,
687 ** <0.01 , *** <0.001 , n.s. = no significance by Student's t-test for samples compared to
688 untreated 16E6 keratinocytes (lane 3).

689

690 **Fig 4. E6 proteins from evolutionarily diverse species target NHERF1. (A) E6**

691 proteins from divergent animal species degrade NHERF1 via E6AP. HA_NHERF1 (0.4
692 ug), HA_GFP (0.1 ug), FLAG_E6AP_WT (0.35 ug), and the indicated FLAG_E6 (0.3 ug)
693 plasmids were co-transfected into C33 cells. E6 proteins are classified based on their
694 known preference for binding E6AP or MAML as indicated. NHERF1 was degraded by
695 E6 proteins isolated from numerous different mammalian species. Many, but not all, of
696 the E6 proteins that bind E6AP targeted NHERF1 for degradation, while E6 proteins
697 that bind MAML1 did not. HA_NHERF1 protein levels in the presence of the indicated
698 E6 proteins were normalized to co-transfected HA_GFP as an internal transfection
699 control. A single representative blot and the means of five independent experiments \pm
700 standard error are shown. N=5, ** <0.01 , *** <0.001 by Student's t-test. (B) E6 proteins

701 that degrade NHERF1 cluster phylogenetically. The E6 proteins from the listed
702 papillomaviruses were subjected to a multiple sequence alignment and then clustered
703 phylogenetically using the program MUSCLE [65]. For E6 physical association, blue
704 denotes MAML1 and light purple denotes E6AP. The preferential association of three
705 E6 proteins is unknown. Ability to degrade NHERF1 is denoted in green and lack of
706 ability to degrade NHERF1 is indicated by red. Interestingly, E6 proteins that can bind
707 E6AP but not degrade NHERF1 cluster differently from other E6 proteins that cannot
708 degrade NHERF1. The genera of each papillomavirus is listed. Western blot indicating
709 NHERF1 expression in the presence of HPV1 E6, HPV8 E6, and SfPV1 E6 is shown in
710 S3 Fig. H = *Homo sapiens* (human), Mm = *Macaca mulata* (rhesus monkey), Ss = *Sus*
711 *scrofa* (wild boar), Pph = *Phocoena phocoena* (harbor porpoise), Um = *Ursus maritimus*
712 (polar bear), Tt = *Tursiops truncatus* (bottlenose dolphin), Oc = *Oryctolagus cuniculus*
713 (rabbit), Mc = *Mastomys coucha* (mouse), Ma = *Mesocricetus auratus* (golden hamster),
714 Sf = *Sylvilagus floridanus* (Cottontail rabbit; CRPV1). Caption credit: Brimer N, Drews
715 CM, Vande Pol SB. Association of papillomavirus E6 proteins with either MAML1 or
716 E6AP clusters E6 proteins by structure, function, and evolutionary relatedness. PLoS
717 Pathog. 2017;13(12):e1006781.

718

719 **Fig 5. 16E6 mutagenesis screen identified mutants selectively defective in their**
720 **ability to degrade NHERF1. (A)** Amino acids F69 and K72 are important for
721 degradation of NHERF1 by 16E6. Plasmids encoding untagged 16E6_WT or 16E6
722 mutants (0.3 ug) were co-transfected with FLAG_E6AP (0.35 ug), HA_NHERF1 (0.4
723 ug), MYC_p53 (0.25 ug), and HA_GFP (0.08 ug) into C33 cells and HA_NHERF1 levels

724 determined by western blot. Multiple 16E6 proteins were identified that were unable to
725 degrade NHERF1 but were still capable of degrading p53. **(B)** HA_NHERF1 and **(C)**
726 p53 protein levels were quantified and normalized to co-transfected HA_GFP as an
727 internal transfection control.

728

729 **Fig 6. Amino acid side chains F69 and K72 define a novel substrate interaction**
730 **domain on 16E6.** **(A)** HPV16 E6 structure (PDB file 4GIZ) showing the amino-terminal
731 zinc-structured domain in green, connecting alpha helix in yellow, and the carboxy-
732 terminal zinc-structured domain in blue. The E6 protein is complexed with the LXXLL
733 peptide of E6AP (pictured in light pink). **(B)** The E6 protein depicted in A is rotated 45°
734 clockwise (C.W.) and the F69 and K72 residues and their side chains are highlighted in
735 red. **(C)** A similar view as part B is shown complexed with the core p53 DNA binding
736 domain (grey). The E6 interaction face with p53 is opposite the F69 and K72 residues.
737

738 **Fig 7. NHERF1 degradation by E6 proteins from both high and low-risk**
739 **papillomaviruses in stable keratinocytes.** Keratinocytes retrovirally transduced with
740 the indicated E6 proteins were seeded at equal confluency and endogenous NHERF1
741 protein levels were normalized to GAPDH. 16E6_WT, 16E6 deleted of its PBM (Δ PBM),
742 and 11E6_WT all degraded NHERF1. The 16E6_F69A/K72A double mutant did not
743 target NHERF1 for degradation. The means of triplicate independent experiments \pm
744 standard error and one representative blot are shown. N=3, *<0.05, **<0.01, n.s. = no
745 significance by Student's t-test.

746

747 **Fig 8. NHERF1 truncations identify the EB domain as necessary for NHERF1**
748 **degradation by 16E6. (A)** Schematic of NHERF1 truncations. NHERF1 proteins that
749 were successfully degraded by 16E6_WT are depicted in green while truncations that
750 were not degraded are depicted in red. **(B and C)** NHERF1 truncations containing the
751 EB domain were degraded, while those lacking the EB domain were not. The listed
752 HA_NHERF1 truncations (shown in A in the order loaded in B and C, 0.8 ug), untagged
753 16E6_WT (1 ug), FLAG_GFP (0.08 ug), and either FLAG_E6AP_WT (1.2 ug) or
754 FLAG_E6AP_Ub⁻ (1.2 ug, defective for ubiquitin ligase activity) were co-transfected in
755 E6AP-null 8B9 cells. HA_NHERF1 levels were quantified and normalized to FLAG_GFP
756 as an internal transfection control. The bar graph below the blot represents
757 quantification of each listed HA_NHERF1 truncation. In panel C, the WT NHERF1 in
758 lanes 2-4 contains an amino terminal 1X HA tag while the WT NHERF1 in lanes 17 and
759 18 contains an amino terminal 2X HA tag. All of the NHERF1 truncations contain amino
760 terminal 2X HA tags. Levels of HA_NHERF1 truncations in the presence of
761 FLAG_E6AP_WT were normalized to their corresponding expression in the presence of
762 FLAG_E6AP_Ub⁻ to account for the differing expression levels. UT = untransfected.
763

764 **Fig 9. The E6-E6AP-NHERF1 complex can be modeled in yeast.** Yeast three-hybrid
765 plasmids expressing the LexA DNA binding domain fused to either 16E6_WT or
766 E6AP_Ub⁻ were co-expressed in yeast (bait) together with either vector, 16E6_WT, or
767 16E6_F69A/K72A as indicated. The bait yeast were mated to prey yeast expressing
768 Gal4 activation domain (G4), or G4 fused to 16E6_WT, PTPN3, truncations of NHERF1,
769 or native p53 and diploids selected. Positive controls for 16E6 expression included the

770 established interaction of the 16E6 PBM with the PDZ domain of tyrosine phosphatase
771 PTPN3 and 16E6-E6AP complex interaction with p53. 16E6_WT recruited NHERF1,
772 p53, and PTPN3 to LexA_E6AP_Ub⁻. The recruitment of NHERF1 to LexA_E6AP_Ub⁻
773 by 16E6 was specifically lost upon mutation of residues F69 and K72, however, p53 and
774 PTPN3 recruitment were maintained. 16E6_WT recruitment of NHERF1 was not seen
775 with an NHERF1 truncation lacking the EB domain (G4_NHERF1 121-297). Caption
776 credit: Ansari T, Brimer N, Vande Pol SB. Peptide interactions stabilize and restructure
777 human papillomavirus type 16 E6 to interact with p53. J Virol. 2012;86(20):11386-91.
778

779 **Fig 10. Activation of the canonical Wnt/ β -catenin pathway is augmented by E6**
780 **proteins that can degrade NHERF1.** The listed E6 proteins were co-expressed with
781 FLAG_E6AP_WT, the TOPFLASH or FOPFLASH luciferase reporter, and a renilla
782 luciferase internal transfection control plasmid in C33A cells. Transfected cells were
783 treated with Wnt3A conditioned media for 8.5 hours, lysed in 1X passive lysis buffer
784 (Promega), and measured for luciferase and renilla luminescence. Fold activation was
785 determined by normalizing the TOPFLASH luminescence by the FOPFLASH
786 luminescence. Each E6 protein that could degrade NHERF1 (16E6_WT, 16E6_ Δ PBM,
787 11E6_WT, and 18E6_WT) augmented the canonical Wnt pathway. 16E6_F69A/K72A,
788 which cannot degrade NHERF1, failed to increase Wnt pathway activation over vector
789 levels. Statistical significance was determined from three independent experiments by
790 Student's t-test (**0.001, n.s. = no significance).

791

792

793 **Supporting information legends**

794 **S1 Fig. Reduction of NHERF1 protein levels is not an overexpression artifact.**

795 Titrations of the indicated three different E6 proteins (16E6_WT, 16E6_ΔPBM, and
796 11E6_WT) were co-transfected with FLAG_E6AP_WT (1 ug), HA_GFP (0.02 ug), and
797 either HA_NHERF1 (0.5 ug) or p53 (0.5 ug) in E6AP-null 8B9 cells. A representative
798 blot of the triplicate experiments for each E6 protein is shown. Increased E6 expression
799 for 16E6_WT, 16E6ΔPBM, and 11E6_WT resulted in decreased NHERF1 protein
800 levels. Both 16E6_WT and 16E6ΔPBM degrade p53 with increasing E6 expression.
801 Overexpression of 11E6_WT (>0.1 ug E6) resulted in degradation of co-expressed
802 E6AP_WT.

803

804 **S2 Fig. 16E6 point mutants screened to determine amino acid(s) necessary for** 805 **NHERF1 degradation.** The 16E6 crystal structure (PDB file 4GIZ) was examined for

806 residues that were at least 20% exposed as determined by the Swiss PDB Viewer.
807 Point mutants of these identified amino acids were then screened to identify which
808 residue(s) resulted in an E6 protein that was selectively defective for degrading
809 NHERF1 but retained degradation of p53. Residues of interest are indicated in red.

810

811 **S3 Fig. The E7 papillomavirus protein does not induce phospho-specific isoforms**

812 **of NHERF1.** Keratinocytes retrovirally transduced with vector or the indicated E7 and/or
813 E6 proteins were seeded at equal confluency. Levels of endogenous NHERF1 were
814 determined by western blot. Levels of phosphorylated NHERF1 (pNHERF1) were
815 unchanged in keratinocytes expressing empty vector compared to the various E7

816 proteins. Keratinocytes expressing the E6 protein from high-risk (HPV16) and low-risk
817 (HPV11) degraded NHERF1. H = *Homo sapiens* (human), Sf = *Sylvilagus floridanus*
818 (Cottontail rabbit; CRPV1). Caption credit: Accardi R, Rubino R, Scalise M, Gheit T,
819 Shahzad N, Thomas M, et al. E6 and E7 from human papillomavirus type 16 cooperate
820 to target the PDZ protein Na/H exchange regulatory factor 1. J Virol. 2011;85(16):8208-
821 16.

822

823 **S4 Fig. The presence of the E7 oncoprotein does not enhance NHERF1**

824 **degradation by E6 proteins.** C33A cells were co-transfected with the following
825 plasmids: HA_NHERF1 (0.4 ug), FLAG_E6AP_WT (0.35 ug), HA_GFP (0.08 ug), the
826 indicated E6 protein (0.3 ug), and the indicated E7 protein (0.3 ug). HA_NHERF1 levels
827 were determined by western blot. FLAG_18E6* is a truncated splice isoform of 18E6.
828 Quantitation is derived from three experimental replicates. A representative blot and
829 means of triplicate independent experiments \pm standard error are shown. N=3. **<0.01,
830 ***<0.001, ****0.0001, n.s. = no significance by Student's t-test.





Fig1






Fig2









Fig3

A.**B.****Fig4**

A.**B.****C.**




Fig6




Fig7




Fig8




Fig9

Fold Activation (TOP/FOP)




Fig10