

Forest yeast sampling biases

1 Quantifying the efficiency and biases of forest *Saccharomyces* sampling strategies

2

3 Primrose J. Boynton*¹, Vienna Kowallik², Doreen Landermann¹, Eva H.

4 Stukenbrock^{1,3}

5

6 1) Max-Planck Institute for Evolutionary Biology, Environmental Genomics Research

7 Group, Plön, Germany

8 2) Okinawa Institute of Science and Technology, Okinawa, Japan

9 3) Christian-Albrechts Universität, Botanisches Institut, Kiel, Germany

10

11 * To whom correspondence should be addressed:

12 pboynton@evolbio.mpg.de

13 August-Thienemann-Str. 2, 24306 Plön, Germany

14

15 Keywords: *Saccharomyces paradoxus*, *Saccharomyces cerevisiae*, forest, enrichment

16 culture, soil, environmental isolates

17 **Abstract**

18 *Saccharomyces* yeasts are emerging as model organisms for ecology and
19 evolution, and researchers need environmental *Saccharomyces* isolates to test
20 ecological and evolutionary hypotheses. However, methods for isolating
21 *Saccharomyces* from nature have not been standardized and isolation methods can
22 influence the genotypes and phenotypes of studied strains. We developed a direct
23 isolation method for forest floor *Saccharomyces* and compared its success and
24 phenotypic biases to a previously published enrichment-based isolation method. In a
25 European forest, direct isolation was more successful at isolating *S. paradoxus*, but
26 also more labor intensive, than enrichment culturing. Average growth rates of *S.*
27 *paradoxus* isolates collected using the two methods did not differ at the enrichment
28 isolation temperature, but variances in growth rates did: direct isolation produced a
29 collection of *S. paradoxus* isolates with less variation in growth rates than enrichment
30 culturing. In other words, enrichment culturing sampled more phenotypic diversity
31 than direct isolation. Enrichment culturing also sampled more *Saccharomyces* species
32 diversity than direct isolation, including our only isolations of rare *S. cerevisiae*.
33 Enrichment culturing may sample higher *Saccharomyces* phenotypic and species
34 diversity because of variations in interactions between yeasts and the other microbes
35 that were present in the soil and leaf litter samples. We recommend direct culturing
36 for researchers interested in randomly sampling their study habitats and enrichment
37 culturing for researchers interested in discovering new *Saccharomyces* phenotypes or
38 rare *Saccharomyces* species from natural environments. We include step-by-step
39 sampling protocols in the supplemental materials.

40


41 **Introduction**

42 Naturally-occurring *Saccharomyces* populations are models for ecology and
43 evolution (Boynton & Greig, 2014). Use of these models has led to exciting
44 discoveries about the ecology and evolution of microbial phenotypes; for example,
45 adaptation to climate can lead to speciation (Leducq et al., 2014), domesticated *S.*
46 *cerevisiae* is more phenotypically diverse than wild *S. paradoxus* (Warringer et al.,
47 2011), and interspecific hybrids can have high fitnesses in stressful environments
48 (Bernardes, Stelkens, & Greig, 2017; Stelkens, Brockhurst, Hurst, Miller, & Greig,
49 2014). These studies made inferences based on the phenotypes and genotypes of
50 isolates collected from wild and domesticated substrates. And *Saccharomyces*
51 substrates are diverse: wild substrates include tree bark, insect guts, leaf litter, soil,
52 fruits, and parasitic *Cyttaria* galls, (Kowallik & Greig, 2016; Libkind et al., 2011;
53 Mortimer & Polsinelli, 1999; Sampaio & Goncalves, 2008; Stefanini et al., 2012), and
54 domesticated substrates include wine, beer, bread, kimchi, kombucha, palm wine, and
55 pulque, among many other substrates (Boynton & Greig, 2016; Carbonetto,
56 Ramsayer, Nidelet, Legrand, & Sicard, 2018; Estrada-Godina et al., 2001; Ezeronye
57 & Okerentugba, 2001; Gallone et al., 2016; Greenwalt, Steinkraus, & Ledford, 2000;
58 Jeong, Jung, Lee, Jin, & Jeon, 2013). One challenge of environmental yeast sampling
59 is to minimize sampling biases so researchers can assure that differences among
60 source environments, not sampling techniques, are responsible for observed
61 phenotypic patterns.

62 Enrichment culturing is a reliable and frequently-used method for isolating
63 difficult-to-culture microbes, including *Saccharomyces*, from natural environments
64 (Schlegel & Jannasch, 1967; Sniegowski, Dombrowski, & Fingerman, 2002) (Figure
65 1A). Microbiologists have been relying on enrichment cultures for over a century

Forest yeast sampling biases

A) Enrichment cultures



B) Direct plating




Figure 1: Schematic illustration of sampling strategies used to isolate *Saccharomyces* for this project. A) Enrichment culturing B) Direct plating. Photo: Doreen Landermann

66 (Beijernick, 1961), and have used enrichment culturing to isolate many of the model
67 *Saccharomyces* strains commonly used in laboratory studies (Johnson et al., 2004;
68 Liti et al., 2009; Sniegowski et al., 2002). To isolate a microbe using enrichment
69 culturing, a researcher adds a small amount of natural material to a growth medium
70 designed to be hospitable to the target microbe and inhospitable to other microbes
71 (Liti, Warringer, & Blomberg, 2017; Schlegel & Jannasch, 1967). If the enrichment
72 medium is well-designed, the target microbe is expected to grow in abundance, and
73 after some incubation time, this enrichment culture can be streaked to solid media and
74 colonies of the target microbe can be easily isolated. An alternative to enrichment
75 culturing is to spread a microbial substrate directly onto a selective solid medium,
76 with or without dilution, and to pick colonies which morphologically resemble the
77 target microbe (Stefanini et al., 2012) (Figure 1B).

78 Because it can be difficult to isolate *Saccharomyces* from natural substrates,
79 many investigations of wild *Saccharomyces* rely on enrichment culturing, usually in
80 high-sugar, acidic media (Charron, Leducq, Bertin, Dube, & Landry, 2014; Robinson,
81 Pinharanda, & Bensasson, 2016; Sniegowski et al., 2002; Sweeney, Kuehne, &

Forest yeast sampling biases

82 Sniegowski, 2004). Comparative studies of *Saccharomyces* genomes have been
83 carried out using *Saccharomyces* strains isolated using disparate strategies, including
84 both enrichment and direct culturing (Liti et al., 2009; Peter et al., 2018). However,
85 isolation strategy can influence the genotypes and phenotypes of isolated microbes
86 (Stefani et al., 2015), and we were concerned about the biases that might be
87 introduced during enrichment culturing of *Saccharomyces* yeasts. Specifically, we
88 were concerned that isolation method might bias results in studies of *Saccharomyces*
89 phenotypes. For example, enrichment culturing might select for individuals with high
90 relative fitness in the enrichment medium. Such potential biases in sampled yeast
91 phenotypes are likely to lead to biases in sampled genotypes because genetic
92 information is responsible for expressed phenotypes. Isolation biases have also been
93 suggested as potential explanations for differences between results of culture-
94 dependent and culture-independent studies of environmental *Saccharomyces*
95 (Alsammar et al., 2018).

96 This study's goals were to compare isolation success between enrichment
97 culturing and a direct culturing strategy, and to quantify biases in *Saccharomyces*
98 phenotypes (and therefore genotypes) that might be introduced when sampling a
99 forest environment. We tested the assumption that it is easier to sample
100 *Saccharomyces* from forest substrates using enrichment cultures than direct plating.
101 We also compared growth rates between *S. paradoxus* isolated using enrichment and
102 direct strategies. Enrichment culturing might decrease or increase sampled *S.*
103 *paradoxus* phenotypic diversity compared to direct plating, thereby decreasing or
104 increasing variance among *S. paradoxus* growth rates. For example, variance among
105 growth rates would be low (and average growth rates high) among *S. paradoxus*
106 isolated using enrichment cultures if the enrichment temperature and media select for

Forest yeast sampling biases

107 the fastest growing *S. paradoxus* present in every sample. Conversely, variance
108 among growth rates would be high for *S. paradoxus* isolated using enrichment
109 cultures if diversity in the non-*Saccharomyces* microbial communities present on
110 sampled substrates select for diverse *S. paradoxus* among samples.

111 To test these predictions, we compared *Saccharomyces* sampling success and
112 isolates' growth rates among soil and leaf litter samples from a well-studied northern
113 German forest (Kowallik & Greig, 2016; Kowallik, Miller, & Greig, 2015). A
114 previous study showed that *S. paradoxus*, the wild sister species of the lab model *S.*
115 *cerevisiae*, is readily isolated using enrichment cultures from oak leaf litter in this
116 forest (Kowallik & Greig, 2016). We were also previously able to isolate *S.*
117 *paradoxus* directly from these forest substrates without enrichment (Kowallik, 2015).

118

119 **Methods**

120 Field sampling and yeast isolation

121 All isolates were sampled from a mixed hardwood and conifer forest in
122 Nehnten, Schleswig-Holstein, northern Germany (Nehmtener Forst). We sampled
123 leaf litter and soil material from close to the bases of ten oak trees at four sampling
124 dates (Table 1), although not all trees were sampled at every date. Trees were between
125 12 and 744 m from one another. At each date, samples were collected from leaf litter
126 and the top organic layer of soil within one meter of the base of each tree. Paired leaf
127 litter and soil samples were collected on the north, south, east, and west side of each
128 tree at all collection days except 7 April, when samples were collected at an arbitrary
129 two of the four cardinal directions.

130 Material was collected simultaneously for the **direct plating** and **enrichment**
131 collections at each sampling point (Figure 1). First, leaf litter was collected by

Forest yeast sampling biases

132 Table 1: Number of sampling points from each tree at each timepoint (each sampling
133 point includes one enrichment and one direct plating sample)

	21 March 2017		7 April 2017		12 June 2017		10 July 2017	
	soil	litter	soil	litter	soil	litter	soil	litter
Tree 1	4	4	2	2	4	4	4	4
Tree 2	4	4	2	2	4	4	4	4
Tree 3	4	4	2	2	4	4	4	4
Tree 4	4	4	2	2	4	4	4	4
Tree 5	4	4					4	4
Tree 6			2	2	4	4	4	4
Tree 7			2	2	4	4	4	4
Tree 8			2	2	4	4	4	4
Tree 9					4	4	4	4
Tree 10							4	4

134

135 aseptically transferring litter into sterile collection tubes: approximately 5 ml of
136 compressed leaf litter was collected for the direct plating method and approximately
137 2 ml for the enrichment method. Then, the remaining leaf litter was removed from the
138 soil surface and the top approximate 2 cm of soil (mostly composed of soil organic
139 layer) were aseptically transferred into sterile collection tubes. As for leaf litter,
140 approximately 5 ml of compressed soil was collected for the direct plating method
141 and approximately 2 ml for the enrichment method. Instruments were sterilized
142 between samples using 70% ethanol. Samples were transported between the field and
143 lab at ambient temperature and processed within four hours of collection.

144 For **direct plating** (Figure 1A), material was mixed with 20 ml sterile water in
145 a sterile 50 ml tube, the mixture was vigorously mixed for at least 10 seconds with a
146 vortex mixer on its highest setting, and 0.2 ml of the resulting dirty liquid was
147 pipetted on each of two plates containing solid modified selective media PIM1 (3 g
148 yeast extract, 5 g peptone, 10 g sucrose, 3 g malt extract, 1 mg chloramphenicol, 80
149 ml ethanol, 5.2 ml 1 M HCl, and 20 g agar per liter) (Kowallik & Greig, 2016;
150 Sniegowski et al., 2002). Liquid was spread on plates using sterile glass beads, and

Forest yeast sampling biases

151 plates were left open in a laminar flow hood until dry. Plates were incubated for three
152 days at 30 °C before colonies were picked.

153 For **enrichments** (Figure 1B), material was mixed with 10 ml liquid selective
154 media PIM1 (composition as for solid PIM1 but without agar) in a 15-ml sterile tube,
155 mixtures were inverted, and tubes were incubated, slightly open and without shaking,
156 at 30 °C. After 10 days, a sterile wooden stick was inserted into each enrichment tube
157 and a small amount of liquid was streaked onto a single plate with solid selective
158 media PIM2 (20 g Methyl-(alpha)-D-glucopyranoside, 1 ml 5% Antifoam Y-30
159 emulsion, 6.7 g Yeast Nitrogen Base without amino acids, 4 ml 1M HCl, and 20 g
160 agar per liter) (Kowallik & Greig, 2016; Sniegowski et al., 2002), and plates were
161 incubated 4 days at 30°C before colonies were picked.

162 We include these procedures as step-by-step protocols for the convenience of
163 future researchers in the supplementary materials (Supplemental File 1).

164

165 Yeast identification

166 After incubation, we streaked colonies with yeast-like morphology to fresh
167 YPD media (10 g yeast extract, 20 g peptone, 20 g dextrose, and 25 g agar per liter).
168 For each method, up to 6 (March and April sampling days) or 12 (June and July
169 sampling days) colonies per sample were selected. After one day of growth on YPD at
170 30 °C, cultures were frozen at -80 °C in 20% glycerol and a small amount of each
171 culture was transferred to sporulation media (20 g potassium acetate, 2.2g yeast
172 extract, 0.5 g dextrose, 870 mg complete amino acid mixture, and 25 g agar per liter).
173 Any cultures with bacteria-like morphology on YPD media (slimy culture and/or cells
174 smaller than 1 micron across) were not frozen and were discarded. Sporulation

Forest yeast sampling biases

175 cultures were incubated for at least three days at room temperature before being
176 screened under a compound microscope for *Saccharomyces*-like asci (tetrads).

177 All cultures producing tetrads were identified using sequencing of the internal
178 transcribed sequence (ITS), a region neighboring rRNA-coding DNA (Schoch et al.,
179 2012). We sequenced every strain using the ITS1/ITS4 primer pair (White, Bruns,
180 Lee, & Taylor, 1990). PCR mixes were 7-15 μ l in volume and contained one yeast
181 colony, 0.5 μ M each primer, and either 50% Phusion® High-Fidelity PCR master mix
182 with HF buffer or 1x HF-buffer, 100 μ M dNTP mix, 3% DMSO, and 1 U/50 μ l
183 Phusion DNA polymerase. PCR reactions were cycled at 98 °C for 30 s, then 35
184 cycles of 98 °C for 5 s, 62 °C for 20 s, and 72 °C for 30 s, plus a 10 min terminal
185 extension at 72 °C. PCR products were cleaned using illustra™ ExoProStar™
186 according to the manufacturer's instructions, and sequenced on an ABI 3130xl
187 sequencer.

188 ITS sequences were compared to sequences from the type or neotype strains of
189 *S. paradoxus*, *S. cerevisiae*, *S. kudriavzevii*, and *S. mikatae* (Genbank accession
190 numbers NR_138272.1, NR_111007.1, KY105195.1, and KY105198.1). If a
191 sequence did not align with *Saccharomyces* sequences, we compared the sequence
192 with all sequences in the NCBI database from type strains using BLAST (Zhang,
193 Schwartz, Wagner, & Miller, 2000). If the sequence aligned with *Saccharomyces*
194 sequences but had more than one base pair different from its closest match, we
195 supplemented ITS sequences with sequences from the gene for translation elongation
196 factor 1 using primers EF1-983F and EF1-2212R (Rehner & Buckley, 2005) using the
197 protocols above, but with a PCR annealing temperature of 57 °C. In some cases,
198 cultures originating from apparent single colonies were in fact mixtures of two yeast

199 species. We counted these colonies as *Saccharomyces* if sequences from one of the
200 species was *Saccharomyces*.

201

202 Growth rates

203 We compared the distributions of maximum growth rates between two groups
204 of *S. paradoxus* strains: strains collected using enrichment culturing and strains
205 collected using direct plating. To avoid confounding effects of environmental source
206 (*i.e.*, combination of substrate, date collected, and tree), we compared growth rates for
207 pairs of *S. paradoxus* strains originating from the same environmental source. In other
208 words, we collected a dataset of *S. paradoxus* growth rates from two groups of strains
209 with equal representations of combinations of substrate, date collected, and tree, and
210 differing only in the method used to isolate the strains. To ensure that all isolates were
211 pure *S. paradoxus* cultures, we streaked all isolates used for growth rate
212 measurements to single-colony cultures a second time and reidentified these cultures
213 by mating them with a *S. paradoxus* tester strain (NCYC 3708, α , *ura3::KANMX*,
214 *ho::HYGMX*). In total, 110 isolates (55 from each sampling method) were measured.

215 Growth rates were measured using an Epoch 2 microplate reader (Biotek
216 Instrument, Inc., Winooski, VT, USA) and calculated using the included Gen5
217 software version 3.03.14 (Biotek Instrument, Inc., Winooski, VT, USA). We first
218 inoculated strains in 0.2 ml liquid YPD media (composition as for solid YPD, but
219 without agar) in a 96-well microplate and incubated cultures without shaking or
220 measurement in the microplate reader at 30 °C for 24 hours to condition strains to
221 microplate reader conditions. We then transferred 2 μ l from each culture to 198 μ l
222 fresh liquid YEPD in a new microplate and incubated the new microplate under the
223 same conditions for 20-24 hours. OD₆₆₀ was measured during the second incubation

224 every ten minutes, and maximum growth rate ($\text{mOD}_{660}/\text{min}$) was calculated from the
225 maximum slope of each growth curve over four points (30 min total) using Gen5
226 software. Reported growth rates for each isolate are means of three replicates.

227

228 Statistical analyses

229 We compared sampling success across substrates (leaf litter or soil) and
230 methods (direct plating or enrichment) using a generalized linear mixed-effects model
231 with probability of isolating *Saccharomyces* (including *S. paradoxus* and *S.*
232 *cerevisiae*) as the response variable, substrate and method as fixed effects, and tree
233 and date as random effects. We selected the best model using a top-down strategy,
234 comparing Akaike's Information Criteria (AIC) after removing predictors from a full
235 model one by one.

236 We compared growth rate distributions by first comparing variances using
237 Levene's test (Levene, 1960) for homogeneity of variance, and then comparing
238 medians using a paired Wilcoxon signed rank test. Statistics were computed using R
239 version 3.3.1 (R Development Core Team, 2016) and the car and lme4 packages
240 (Bates, Machler, Bolker, & Walker, 2015; R Development Core Team, 2016).
241 Graphics were produced using the ggplot2 package (Wickham, 2016).

242

243 **Results**

244 Influence of sampling method on success isolating *Saccharomyces*

245 Direct plating was more successful than enrichment culturing for isolating
246 *Saccharomyces* from natural substrates ($z = 6.1$, $p < .001$) (Tables 2, 3, Figure 2). We
247 found *Saccharomyces* isolates in 45% of direct plating cultures and 19% of
248 enrichment cultures. However, enrichment culturing produced the only *S. cerevisiae*

Forest yeast sampling biases

249 found in this study: we found six *S. cerevisiae* isolates from a single enrichment
 250 culture from tree 3 in March of 2017. All other *Saccharomyces* isolates found in this
 251 study were *S. paradoxus*. Other detected yeast species included *Saccharomycodes*
 252 *ludwigii*, *Torulaspora delbrueckii*, *Pichia membranifaciens*, and *Hanseniaspora*
 253 *osmophila*, all of which have previously been found alongside *Saccharomyces* yeasts
 254 in beverage fermentations (Domizio et al., 2011; Gschaedler, 2017).

255 Table 2: Model selection (Mixed-effects generalized linear model)

model	fixed effects	random effects	AIC	compared to	better model
1	method + substrate + method:substrate	1 tree + 1 month	447.20		
2	method + substrate + method:substrate	1 month	470.94	model 1	1
3	method + substrate + method:substrate	1 tree	451.99	model 1	1
4	method + substrate	1 tree + 1 month	446.76	model 1	4
5	method	1 tree + 1 month	480.34	model 4	4
6	substrate	1 tree + 1 month	485.71	model 4	4


256

257 Table 3: Model summary table (model 4)

	Estimate	Std. Error	z	p
intercept	-2.5324	0.4599	-5.506	< .001
method (plating)	1.5494	0.2559	6.054	< .001
substrate (soil)	1.4445	0.2542	5.683	< .001

258

A) Enrichment cultures



B) Direct plating




Figure 2: Percentages of samples in which *Saccharomyces* could be detected using A) enrichment cultures or B) direct plating. Bars represent all samples for each category of sampling, and colors represent *Saccharomyces* species. *S. paradoxus* and *S. cerevisiae* were the only detected *Saccharomyces* species.

259 While the direct plating method was more successful than the enrichment
260 method, it was also more labor-intensive (Table 4). We screened 3.4 times as many
261 colonies for tetrads when using the direct plating method than we did using the
262 enrichment method. Only 32% of the processed direct plating colonies were *S.*
263 *paradoxus*, compared to 74% of enrichment colonies.

264

265 Table 4: Colonies processed and sampling success for each method

method	total colonies picked	sequenced colonies with	
		<i>Saccharomyces</i> -like ascus morphology	<i>Saccharomyces</i> isolates
enrichment	284	246 (87%)	211 (74%)
plating	969	344 (35%)	307 (32%)

266

267 Both methods isolated *Saccharomyces* colonies from both substrates, most
268 trees, and all timepoints (Figure 2). We had significantly more sampling success on
269 soil than leaf litter substrates ($z = 5.7$, $p < .001$, Table 3), but other relationships
270 among sampling success, sampling method, and sampling environments were
271 idiosyncratic. For example, direct plating did not produce any *Saccharomyces*
272 isolates from tree 6, while three enrichment samples from this tree isolated *S.*
273 *paradoxus*, and enrichments produced more *Saccharomyces* isolates in March than
274 direct plating did (Figure 2). Because our sampling effort was not the same for all
275 trees at all months, we did not model tree habitat or sampling month as fixed effects;
276 instead, we modeled these parameters as random effects, and found that models
277 including tree and month fit the data better than models without tree and month (Table
278 2). A list of sequenced yeasts from each sample is included in the supplemental
279 materials (Supplemental File 2).

280

281 Phenotypes of sampled *S. paradoxus*

Forest yeast sampling biases

282 Variances in growth rate, but not median growth rates, were different between
283 strains isolated using the two methods (Figure 3). Growth rates of *S. paradoxus*
284 isolated using enrichment culturing had a larger variance than growth rates of *S.*
285 *paradoxus* isolated using direct plating (Levine's test $F_{1,108} = 5.42$, $p = .02$). Median
286 growth rates for the two groups of *S. paradoxus* strains did not differ (Wilcoxon
287 signed rank test $V = 6320$, $p = .39$). Measured growth rates of each strain are included
288 in the supplemental materials (Supplemental File 2).




Figure 3: Histograms representing distributions of growth rates for *S. paradoxus* clones isolated using A) enrichment culturing or B) direct plating.

289

290 Discussion

291 Direct plating detects *S. paradoxus* more frequently than enrichment culturing

Forest yeast sampling biases

292 Enrichment culturing did not increase *Saccharomyces* sampling success from
293 forest leaf litter and soil over direct plating, in spite of researchers' long history of
294 using enrichment culturing to isolate *Saccharomyces* from forest environments
295 (Kowallik & Greig, 2016; Naumov, Naumova, & Sniegowski, 1998; Sniegowski et
296 al., 2002). We expect reliable *Saccharomyces* isolation from this forest using direct
297 plating to be a result of high *S. paradoxus* abundance on forest floor substrates.
298 Indeed, it is common to find hundreds to tens of thousands of *S. paradoxus* cells per
299 gram of leaf litter near the bases of oak trees in this forest (Kowallik & Greig, 2016).
300 We expect direct plating to be less successful in environments in which
301 *Saccharomyces* are rarer, and note that enrichment culturing is frequently used to
302 isolate *Saccharomyces* from tree bark, which may be a habitat with lower
303 *Saccharomyces* density than the forest floor habitats we sampled (Kowallik et al.,
304 2015; Sniegowski et al., 2002).

305

306 Isolation using enrichment culturing samples more phenotypic diversity than direct
307 plating

308 Conditions in enrichment cultures resulted in isolating a different (albeit
309 higher) phenotypic diversity than random colony selection on plate cultures did
310 (Figure 3). There are several potential methodological and ecological explanations for
311 the high phenotypic diversity in enrichment isolates. But we expect that interactions
312 with microbes *during the enrichment culturing* are most likely to be responsible for
313 the increased phenotypic diversity among enrichment-sourced *S. paradoxus* isolates.

314 Microbes that potentially interact with *S. paradoxus* were doubtless present in
315 our enrichment cultures, and it is realistic to expect these microbes to influence the
316 phenotypes of the *S. paradoxus* that were ultimately recovered. Our enrichment

Forest yeast sampling biases

317 cultures contained all of the microbes that were present in the soil and leaf litter
318 samples, and soil and leaf litter include a wide diversity of bacterial and fungal taxa
319 that can interact with fungi (Curd, Martiny, Li, & Smith, 2018; Glassman et al., 2018;
320 Santonja et al., 2018). For example, some agricultural soils contain *Bacillus* and
321 *Pseudomonas* species that secrete compounds toxic to phytopathogenic fungi (Islam,
322 Jeong, Lee, & Song, 2012; Petatán-Sagahón et al., 2011). Conversely, some soil
323 bacteria promote mycorrhizal fungal growth and host plant colonization (Xie et al.,
324 2018).

325 We expect a variety of similar inhibition and facilitation interactions to
326 determine the identities of the *Saccharomyces* strains that ultimately reached high
327 frequencies in enrichment cultures. Bacteria and fungi co-occurring with *S. paradoxus*
328 on oak bark can both inhibit and facilitate *S. paradoxus* growth (Kowallik et al.,
329 2015). These effects are dependent on temperature and interacting microbe identities,
330 and likely also depend on other environmental conditions and *S. paradoxus* genotype.
331 The *Saccharomyces* strains recovered at the end of our enrichment cultures were
332 probably the fittest *Saccharomyces* strains present in the cultures, but this high
333 relative fitness was as likely to be a result of interactions with co-occurring microbes
334 that occur by chance in the same enrichment cultures as it was to be the result of
335 intrinsic growth rate. Microbial diversity during enrichment may similarly explain our
336 idiosyncratic sampling success across months and trees (Figure 2). For example, it is
337 possible that a bacterium that facilitates rare *S. paradoxus* growth in the enrichment
338 medium was more common in spring than summer months, resulting in higher
339 enrichment sampling success in spring than summer. Unfortunately, we did not
340 measure the microbial diversity of the enrichment cultures and do not know which
341 microbes may have interacted with *Saccharomyces*.

Forest yeast sampling biases

342 It is also possible, although in our opinion less likely, that methodological
343 biases resulted in relatively low phenotypic diversity among directly plated *S.*
344 *paradoxus* compared to enrichment *S. paradoxus*. While we aimed to randomly select
345 colonies with *Saccharomyces*-like morphologies from direct culture plates, biases in
346 colony picking could have selected for low variance in *S. paradoxus* phenotypes. For
347 example, if a *S. paradoxus* colony had an unusual morphology, it might have been
348 mistaken for a bacterial colony and not isolated. However, direct culturing isolated
349 more non-*Saccharomyces* yeast isolates than enrichment culturing did, and we expect
350 interspecific morphological variation to be higher than intraspecific morphological
351 variation. We therefore consider it unlikely that biases in colony picking decreased
352 phenotypic variance among *S. paradoxus* isolates.

353 We continue to expect plate culturing to be a more random sampling of
354 *Saccharomyces* diversity than enrichment culturing. Colonies are physically isolated
355 on a plate and unlikely to influence each other's growth, and plated colonies come
356 from a well-mixed mixture of environmental substrate and sterile water. We therefore
357 expect *Saccharomyces* colonies on plates to accurately reflect the diversity of
358 *Saccharomyces* strains present in nature, as long as all strains are able to grow on the
359 selective media chosen. Biases in sample *success* can still be introduced to direct
360 plating samples by the presence of morphologically similar yeast species. On petri
361 dishes, a high density of non-*Saccharomyces* yeasts with morphologies similar to
362 *Saccharomyces* could have prevented us from detecting *Saccharomyces*. This issue
363 can be mitigated by designing a more selective isolation medium, picking more
364 colonies per plate (we picked up to 6 or 12 colonies across two plates), or replica
365 plating selective plates to media containing a color indicator for the target yeast (*e.g.*,
366 Wallerstein media) (Hall, 1971).

367

368 Recommendations for future yeast sampling

369 Researchers should consider both resources available for sampling and study
370 goals when choosing a *Saccharomyces* field sampling strategy. Our results identified
371 a tradeoff between resources spent on sampling and resources spent on sequencing:
372 enrichment culturing was less successful than direct plating at finding *Saccharomyces*
373 per sample collected, but more successful per ITS region sequenced (Figure 2, Table
374 4). Researchers with a few precious samples are therefore better off isolating
375 *Saccharomyces* using direct plating than enrichments, especially if *Saccharomyces* is
376 common on their substrates. Conversely, if samples are easy to get but funds available
377 for sequencing are limited, researchers may prefer to use enrichment culturing.
378 Researchers with limited time or freezer space who would like assurances that most
379 picked colonies are *Saccharomyces* may also prefer enrichment culturing.

380 However, the scientific question to be answered by environmental
381 *Saccharomyces* samples may be more important than sampling and sequencing
382 resource limitations, especially as sequencing becomes cheaper. Direct plating
383 samples a more random collection of *Saccharomyces* cells from environmental
384 samples than enrichment culturing, and we recommend that researchers who need
385 random environmental samples to answer ecological and evolutionary questions rely
386 on direct plating. Researchers targeting phenotypic diversity, especially for applied
387 yeast biology (*e.g.*, food microbiology, drug discovery) may uncover more diversity
388 by isolating environmental *Saccharomyces* using enrichment cultures. Researchers
389 interested in detecting rare *Saccharomyces* species in an environment (*e.g.*, *S.*
390 *cerevisiae* from our study forest, *S. mikatae* and *S. eubayanus* from European forests)

391 (Alsammar et al., 2018) may also have more success using enrichment culturing or a
392 combination of enrichment and direct plating strategies.

393

394 **Conclusions**

395 Isolation protocols do indeed influence characteristics of isolated microbes. As
396 researchers continue to develop *Saccharomyces* yeasts as model organisms for
397 ecology and evolution, they must also consider how isolation history of environmental
398 strains can influence the ecological stories the strains can tell. The results of this study
399 highlight the need for consistent sampling within studies, and are a warning to
400 researchers comparing phenotypes (and perhaps also genotypes) among
401 *Saccharomyces* strains from different sources. Much of what we know about
402 *Saccharomyces* evolution has come from culture collections (Strope et al., 2015;
403 Warringer et al., 2011)—these collections are invaluable, as are the data and
404 conclusions they enable, but sampling information about deposited strains is not
405 always available. As technology improves and both genome sequencing and
406 phenotype assays become cheaper and more accessible (Porter & Hajibabaei, 2018;
407 Stewart et al., 2018; van Dijk, Auger, Jaszczyszyn, & Thermes, 2014; Zackrisson et
408 al., 2016), environmental sampling may emerge as a limiting step to studying
409 *Saccharomyces* ecology and evolution. Researchers should carefully consider the
410 consequences of their chosen sampling strategies as they conduct this exciting
411 microbiological research.

412

413 **Acknowledgements**

414 We would like to thank Danielle Stevens and Tjorben Nawroth for help in the field,
415 Jenna Gallie and the Gallie lab for help with growth curves, and Amine Hassani for

416 helpful conversations on microbial diversity in enrichment cultures. Thank you to
417 Christoph Freiherr von Fürstenberg-Plessen for permission to work in the Nehnten
418 forest. This work was supported through a Max Planck Fellowship to Eva H.
419 Stukenbrock.

420

421 **Data Accessibility**

422 All data for this project are included in Supplementary File 2.

423

424 **Figure Legends**

425 Figure 1: Schematic illustration of sampling strategies used to isolate *Saccharomyces*
426 for this project. A) Enrichment culturing B) Direct plating. Photo: Doreen
427 Landermann.

428

429 Figure 2: Percentages of samples in which *Saccharomyces* could be detected using A)
430 enrichment cultures or B) direct plating. Bars represent all samples for each category
431 of sampling, and colors represent *Saccharomyces* species. *S. paradoxus* and *S.*
432 *cerevisiae* were the only detected *Saccharomyces* species.

433

434 Figure 3: Histograms representing distributions of growth rates for *S. paradoxus*
435 clones isolated using A) enrichment culturing or B) direct plating.

436

437 **Citations**

438 Alsammar, H. F., Naseeb, S., Brancia, L. B., Gilman, R. T., Wang, P., & Delneri, D.
439 (2018). Targeted metagenomics approach to capture the biodiversity of
440 *Saccharomyces* genus in wild environments. *Environmental Microbiology*
441 *Reports*. doi:10.1111/1758-2229.12724
442 Bates, D., Machler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting Linear Mixed-
443 Effects Models Using lme4. *Journal of Statistical Software*, 67(1), 1-48.

- 444 Beijernick, M. W. (1961). Enrichment culture studies with urea bacteria (T. D. Brock,
445 Trans.). In T. D. Brock (Ed.), *Milestones in Microbiology*. London, United
446 Kingdom: Prentice-Hall International.
- 447 Bernardes, J. P., Stelkens, R. B., & Greig, D. (2017). Heterosis in hybrids within and
448 between yeast species. *Journal of Evolutionary Biology*, 30(3), 538-548.
449 doi:10.1111/jeb.13023
- 450 Boynton, P. J., & Greig, D. (2014). The ecology and evolution of non-domesticated
451 *Saccharomyces* species. *Yeast*, 31(12), 449-462. doi:10.1002/yea.3040
- 452 Boynton, P. J., & Greig, D. (2016). Species richness influences wine ecosystem
453 function through a dominant species. *Fungal Ecology*, 22, 61-72.
454 doi:10.1016/j.funeco.2016.04.008
- 455 Carbonetto, B., Ramsayer, J., Nidelet, T., Legrand, J., & Sicard, D. (2018). Bakery
456 yeasts, a new model for studies in ecology and evolution. *Yeast*, 35(11), 591-
457 603. doi:10.1002/yea.3350
- 458 Charron, G., Leducq, J. B., Bertin, C., Dube, A. K., & Landry, C. R. (2014).
459 Exploring the northern limit of the distribution of *Saccharomyces cerevisiae*
460 and *Saccharomyces paradoxus* in North America. *FEMS Yeast Research*,
461 14(2), 281-288. doi:10.1111/1567-1364.12100
- 462 Curd, E. E., Martiny, J. B. H., Li, H. Y., & Smith, T. B. (2018). Bacterial diversity is
463 positively correlated with soil heterogeneity. *Ecosphere*, 9(1), 16.
464 doi:10.1002/ecs2.2079
- 465 Domizio, P., Romani, C., Comitini, F., Gobbi, M., Lencioni, L., Mannazzu, I., &
466 Ciani, M. (2011). Potential spoilage non-Saccharomyces yeasts in mixed
467 cultures with *Saccharomyces cerevisiae*. *Annals of Microbiology*, 61(1), 137-
468 144. doi:10.1007/s13213-010-0125-1
- 469 Estrada-Godina, A. R., Cruz-Guerrero, A. E., Lappe, P., Ulloa, M., Garcia-Garibay,
470 M., & Gomez-Ruiz, L. (2001). Isolation and identification of killer yeasts
471 from Agave sap (aguamiel) and pulque. *World Journal of Microbiology &*
472 *Biotechnology*, 17(6), 557-560. doi:10.1023/a:1012210106203
- 473 Ezeronye, O. U., & Okerentugba, P. O. (2001). Genetic and physiological variants of
474 yeast selected from palm wine. *Mycopathologia*, 152(2), 85-89.
475 doi:10.1023/a:1012323721012
- 476 Gallone, B., Steensels, J., Prahl, T., Soriaga, L., Saels, V., Herrera-Malaver, B., . . .
477 Verstrepen, K. J. (2016). Domestication and Divergence of *Saccharomyces*
478 *cerevisiae* Beer Yeasts. *Cell*, 166(6), 1397+. doi:10.1016/j.cell.2016.08.020
- 479 Glassman, S. I., Weihe, C., Li, J. H., Albright, M. B. N., Looby, C. I., Martiny, A. C.,
480 . . . Martiny, J. B. H. (2018). Decomposition responses to climate depend on
481 microbial community composition. *Proceedings of the National Academy of*
482 *Sciences of the United States of America*, 115(47), 11994-11999.
483 doi:10.1073/pnas.1811269115
- 484 Greenwalt, C. J., Steinkraus, K. H., & Ledford, R. A. (2000). Kombucha, the
485 fermented tea: Microbiology, composition, and claimed health effects. *Journal*
486 *of Food Protection*, 63(7), 976-981. doi:10.4315/0362-028x-63.7.976
- 487 Gschaedler, A. (2017). Contribution of non-conventional yeasts in alcoholic
488 beverages. *Current Opinion in Food Science*, 13, 73-77.
489 doi:10.1016/j.cofs.2017.02.004
- 490 Hall, J. F. (1971). Detection of wild yeasts in the brewery. *Journal of the Institute of*
491 *Brewing*, 77(6), 513-516. doi:10.1002/j.2050-0416.1971.tb03413.x
- 492 Islam, M. R., Jeong, Y. T., Lee, Y. S., & Song, C. H. (2012). Isolation and
493 Identification of Antifungal Compounds from *Bacillus subtilis* C9 Inhibiting

- 494 the Growth of Plant Pathogenic Fungi. *Mycobiology*, 40(1), 59-66.
495 doi:10.5941/myco.2012.40.1.059
- 496 Jeong, S. H., Jung, J. Y., Lee, S. H., Jin, H. M., & Jeon, C. O. (2013). Microbial
497 succession and metabolite changes during fermentation of dongchimi,
498 traditional Korean watery kimchi. *International Journal of Food*
499 *Microbiology*, 164(1), 46-53. doi:10.1016/j.ijfoodmicro.2013.03.016
- 500 Johnson, L. J., Koufopanou, V., Goddard, M. R., Hetherington, R., Schafer, S. M., &
501 Burt, A. (2004). Population genetics of the wild yeast *Saccharomyces*
502 *paradoxus*. *Genetics*, 166(1), 43-52. doi:10.1534/genetics.166.1.43
- 503 Kowallik, V. (2015). *The natural ecology of Saccharomyces yeasts*. (Dr. rer. nat.),
504 Christian Albrechts University of Kiel,
- 505 Kowallik, V., & Greig, D. (2016). A systematic forest survey showing an association
506 of *Saccharomyces paradoxus* with oak leaf litter. *Environmental Microbiology*
507 *Reports*, 8(5), 833-841. doi:10.1111/1758-2229.12446
- 508 Kowallik, V., Miller, E., & Greig, D. (2015). The interaction of *Saccharomyces*
509 *paradoxus* with its natural competitors on oak bark. *Molecular Ecology*, 24(7),
510 1596-1610. doi:10.1111/mec.13120
- 511 Leducq, J. B., Charron, G., Samani, P., Dube, A. K., Sylvester, K., James, B., . . .
512 Landry, C. R. (2014). Local climatic adaptation in a widespread
513 microorganism. *Proceedings of the Royal Society B-Biological Sciences*,
514 281(1777), 9. doi:10.1098/rspb.2013.2472
- 515 Levene, H. (1960). Robust tests for equality of variances. In I. Olkin & H. Hotelling
516 (Eds.), *Contributions to Probability and Statistics: Essays in Honor of Harold*
517 *Hotelling* (pp. 278-292): Stanford University Press.
- 518 Libkind, D., Hittinger, C. T., Valerio, E., Goncalves, C., Dover, J., Johnston, M., . . .
519 Sampaio, J. P. (2011). Microbe domestication and the identification of the
520 wild genetic stock of lager-brewing yeast. *Proceedings of the National*
521 *Academy of Sciences of the United States of America*, 108(35), 14539-14544.
522 doi:10.1073/pnas.1105430108
- 523 Liti, G., Carter, D. M., Moses, A. M., Warringer, J., Parts, L., James, S. A., . . . Louis,
524 E. J. (2009). Population genomics of domestic and wild yeasts. *Nature*,
525 458(7236), 337-341. doi:10.1038/nature07743
- 526 Liti, G., Warringer, J., & Blomberg, A. (2017). Isolation and Laboratory
527 Domestication of Natural Yeast Strains. *Cold Spring Harb Protoc*, 2017(8),
528 pdb.prot089052. doi:10.1101/pdb.prot089052
- 529 Mortimer, R., & Polsinelli, M. (1999). On the origins of wine yeast. *Research in*
530 *Microbiology*, 150(3), 199-204. doi:10.1016/s0923-2508(99)80036-9
- 531 Naumov, G. I., Naumova, E. S., & Sniegowski, P. D. (1998). *Saccharomyces*
532 *paradoxus* and *Saccharomyces cerevisiae* are associated with exudates of
533 North American oaks. *Canadian Journal of Microbiology*, 44(11), 1045-1050.
534 doi:10.1139/cjm-44-11-1045
- 535 Petatán-Sagahón, I., Anducho-Reyes, M. A., Silva-Rojas, H. V., Arana-Cuenca, A.,
536 Tellez-Jurado, A., Cárdenas-Álvarez, I. O., & Mercado-Flores, Y. (2011).
537 Isolation of Bacteria with Antifungal Activity against the Phytopathogenic
538 Fungi *Stenocarpella maydis* and *Stenocarpella macrospora*. *International*
539 *Journal of Molecular Sciences*, 12(9), 5522-5537. doi:10.3390/ijms12095522
- 540 Peter, J., De Chiara, M., Friedrich, A., Yue, J. X., Pflieger, D., Bergstrom, A., . . .
541 Schacherer, J. (2018). Genome evolution across 1,011 *Saccharomyces*
542 *cerevisiae* isolates. *Nature*, 556(7701), 339-+. doi:10.1038/s41586-018-0030-
543 5

- 544 Porter, T. M., & Hajibabaei, M. (2018). Scaling up: A guide to high-throughput
545 genomic approaches for biodiversity analysis. *Molecular Ecology*, 27(2), 313-
546 338. doi:10.1111/mec.14478
- 547 R Development Core Team. (2016). R: A language and Environment for Statistical
548 Computing. Vienna, Austria: R Foundation for Statistical Computing.
- 549 Rehner, S. A., & Buckley, E. (2005). A *Beauveria* phylogeny inferred from nuclear
550 ITS and EF1-alpha sequences: evidence for cryptic diversification and links to
551 *Cordyceps teleomorphs*. *Mycologia*, 97(1), 84-98.
552 doi:10.3852/mycologia.97.1.84
- 553 Robinson, H. A., Pinharanda, A., & Bensasson, D. (2016). Summer temperature can
554 predict the distribution of wild yeast populations. *Ecology and Evolution*, 6(4),
555 1236-1250. doi:10.1002/ece3.1919
- 556 Sampaio, J. P., & Goncalves, P. (2008). Natural populations of *Saccharomyces*
557 *kudriavzevii* in Portugal are associated with oak bark and are sympatric with *S.*
558 *cerevisiae* and *S. paradoxus*. *Applied and Environmental Microbiology*, 74(7),
559 2144-2152. doi:10.1128/aem.02396-07
- 560 Santonja, M., Foucault, Q., Rancon, A., Gauquelin, T., Fernandez, C., Baldy, V., &
561 Mirleau, P. (2018). Contrasting responses of bacterial and fungal communities
562 to plant litter diversity in a Mediterranean oak forest. *Soil Biology &*
563 *Biochemistry*, 125, 27-36. doi:10.1016/j.soilbio.2018.06.020
- 564 Schlegel, H. G., & Jannasch, H. W. (1967). Enrichment cultures. *Annual Reviews*
565 *Microbiology*, 21, 49-70. doi:10.1146/annurev.mi.21.100167.000405
- 566 Schoch, C. L., Seifert, K. A., Huhndorf, S., Robert, V., Spouge, J. L., Levesque, C.
567 A., . . . Fungal Barcoding, C. (2012). Nuclear ribosomal internal transcribed
568 spacer (ITS) region as a universal DNA barcode marker for Fungi.
569 *Proceedings of the National Academy of Sciences of the United States of*
570 *America*, 109(16), 6241-6246. doi:10.1073/pnas.1117018109
- 571 Sniegowski, P. D., Dombrowski, P. G., & Fingerman, E. (2002). *Saccharomyces*
572 *cerevisiae* and *Saccharomyces paradoxus* coexist in a natural woodland site in
573 North America and display different levels of reproductive isolation from
574 European conspecifics. *FEMS Yeast Research*, 1(4), 299-306.
575 doi:10.1111/j.1567-1364.2002.tb00048.x
- 576 Stefani, F. O. P., Bell, T. H., Marchand, C., de la Providencia, I. E., El Yassimi, A.,
577 St-Arnaud, M., & Hijri, M. (2015). Culture-Dependent and -Independent
578 Methods Capture Different Microbial Community Fractions in Hydrocarbon-
579 Contaminated Soils. *PLoS One*, 10(6), 16. doi:10.1371/journal.pone.0128272
- 580 Stefanini, I., Dapporto, L., Legras, J. L., Calabretta, A., Di Paola, M., De Filippo, C., .
581 . . Cavalieri, D. (2012). Role of social wasps in *Saccharomyces cerevisiae*
582 ecology and evolution. *Proceedings of the National Academy of Sciences of*
583 *the United States of America*, 109(33), 13398-13403.
584 doi:10.1073/pnas.1208362109
- 585 Stelkens, R. B., Brockhurst, M. A., Hurst, G. D. D., Miller, E. L., & Greig, D. (2014).
586 The effect of hybrid transgression on environmental tolerance in experimental
587 yeast crosses. *Journal of Evolutionary Biology*, 27(11), 2507-2519.
588 doi:10.1111/jeb.12494
- 589 Stewart, E. L., Croll, D., Lendenmann, M. H., Sanchez-Vallet, A., Hartmann, F. E.,
590 Palma-Guerrero, J., . . . McDonald, B. A. (2018). Quantitative trait locus
591 mapping reveals complex genetic architecture of quantitative virulence in the
592 wheat pathogen *Zymoseptoria tritici*. *Molecular Plant Pathology*, 19(1), 201-
593 216. doi:10.1111/mpp.12515

Forest yeast sampling biases

- 594 Strobe, P. K., Skelly, D. A., Kozmin, S. G., Mahadevan, G., Stone, E. A., Magwene,
595 P. M., . . . McCusker, J. H. (2015). The 100-genomes strains, an *S. cerevisiae*
596 resource that illuminates its natural phenotypic and genotypic variation and
597 emergence as an opportunistic pathogen. *Genome Research*, 25(5), 762-774.
598 doi:10.1101/gr.185538.114
- 599 Sweeney, J. Y., Kuehne, H. A., & Sniegowski, P. D. (2004). Sympatric natural
600 *Saccharomyces cerevisiae* and *S. paradoxus* populations have different
601 thermal growth profiles. *FEMS Yeast Research*, 4(4-5), 521-525.
602 doi:10.1016/s1567-1356(03)00171-5
- 603 van Dijk, E. L., Auger, H., Jaszczyszyn, Y., & Thermes, C. (2014). Ten years of next-
604 generation sequencing technology. *Trends in Genetics*, 30(9), 418-426.
605 doi:10.1016/j.tig.2014.07.001
- 606 Warringer, J., Zorgo, E., Cubillos, F. A., Zia, A., Gjuvsland, A., Simpson, J. T., . . .
607 Blomberg, A. (2011). Trait Variation in Yeast Is Defined by Population
608 History. *Plos Genetics*, 7(6), 30. doi:10.1371/journal.pgen.1002111
- 609 White, T. J., Bruns, T., Lee, S., & Taylor, J. (1990). Amplification and Direct
610 Sequencing fo Fungal Ribosomal RNA Genes for Phylogenetics. In M. A.
611 Innis, D. H. Gelfand, J. J. Sninsky, & T. J. White (Eds.), *PCR Protocols: a*
612 *Guide to Methods and Applications* (pp. 315-322): Academic Press.
- 613 Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*: Springer.
- 614 Xie, L., Lehvavirta, S., Timonen, S., Kasurinen, J., Niemikapee, J., & Valkonen, J. P.
615 T. (2018). Species-specific synergistic effects of two plant growth-Promoting
616 microbes on green roof plant biomass and photosynthetic efficiency. *PLoS*
617 *One*, 13(12), 16. doi:10.1371/journal.pone.0209432
- 618 Zackrisson, M., Hallin, J., Ottosson, L. G., Dahl, P., Fernandez-Parada, E.,
619 Landstrom, E., . . . Blomberg, A. (2016). Scan-o-matic: High-Resolution
620 Microbial Phenomics at a Massive Scale. *G3-Genes Genomes Genetics*, 6(9),
621 3003-3014. doi:10.1534/g3.116.032342
- 622 Zhang, Z., Schwartz, S., Wagner, L., & Miller, W. (2000). A greedy algorithm for
623 aligning DNA sequences. *Journal of Computational Biology*, 7(1-2), 203-214.
624 doi:10.1089/10665270050081478
625