

1 Corresponding Author:
2 Dr. Megan L. Fritz
3 112 Derieux Place
4 1549 Thomas Hall
5 Department of Entomology
6 North Carolina State University
7 Raleigh, NC 27695
8 mlfritz@ncsu.edu
9 (919) 515-1651
10

11 **Application of a dense genetic map for assessment of genomic responses to selection and**
12 **inbreeding in *Heliothis virescens*.**

13

14 Megan L. Fritz^{1,2,§}, Sandra Paa¹, Jennifer Baltzegar², and Fred Gould^{1,2}

15 ¹Department of Entomology, North Carolina State University, Raleigh, NC, USA

16 ²Department of Biological Sciences, Program in Genetics, North Carolina State University, Raleigh,
17 NC, USA

18 [§]Corresponding Author

19 **Abstract**

20 Adaptation of pest species to laboratory conditions and selection for resistance to toxins in the
21 laboratory are expected to cause inbreeding and genetic bottlenecks that reduce genetic variation.
22 *Heliothis virescens*, a major cotton pest, has been colonized in the laboratory many times, and a few
23 laboratory colonies have been selected for Bt resistance. We developed 350 bp Double-Digest
24 Restriction-site Associated DNA-sequencing (ddRAD-seq) molecular markers to examine and compare
25 changes in genetic variation associated with laboratory adaptation, artificial selection, and inbreeding in
26 this non-model insect species. We found that allelic and nucleotide diversity declined dramatically in
27 laboratory-reared *H. virescens* as compared with field-collected populations. The declines were
28 primarily due to the loss of low frequency alleles present in field-collected *H. virescens*. A further,
29 albeit modest decline in genetic diversity was observed in a Bt-selected population. The greatest
30 decline was seen in *H. virescens* that were sib-mated for 10 generations, where more than 80% of loci
31 were fixed for a single allele. To determine which regions of the genome were resistant to fixation in
32 our sib-mated and Bt-selected lines, we generated a dense intraspecific linkage map containing 3 PCR-
33 based, and 659 ddRAD-seq markers. Markers that retained polymorphism were observed in small
34 clusters spread over multiple linkage groups. These markers are likely associated with genomic
35 regions under balancing selection, thus preventing fixation of deleterious alleles.

36 **Keywords:** *Heliothis virescens*, genetic variation, colonization, inbreeding, ddRAD-seq, linkage map

37 **Introduction**

38 Laboratory-reared insect colonies are important resources for many types of entomological
39 experiments. They are used to quantify physiological or behavioral differences between insect
40 populations or species (Dekker *et al.*, 2006; Dobzhansky & Spassky, 1954; Fritz *et al.*, 2015; Groot *et*
41 *al.*, 2005; Shaw *et al.*, 2000; Sokolowski, 1980; Tomaru *et al.*, 2000), identify the genetic architecture
42 of insect traits (Gahan *et al.*, 2010, Mackay *et al.*, 2012, Oppenheim *et al.*, 2012), develop insect
43 populations that express desirable traits (Collins, 1984; Goldman *et al.*, 1986; Gould *et al.*, 1995; Hoy,
44 1989; Pradeep *et al.*, 2005), and generate genetically modified species as a means of pest control (de
45 Valdez *et al.*, 2011). A major concern for researchers maintaining insect colonies is the degree to which
46 adaptation to the laboratory environment affects insect genotypic, and thereby phenotypic diversity
47 (Boller, 1972; Huettel, 1976).

48 The phenotypic consequences of adaptation to the laboratory depend upon the trait of interest,
49 and range from undetectable to severe (Baeshen *et al.*, 2014; Fox *et al.*, 2007; Gerloff *et al.*, 2003;
50 Raulston, 1975; Roush, 1986). Observed phenotypic changes can be attributed to inadvertent selection
51 for traits that are favorable in the laboratory environment (Roush, 1986), inbreeding depression (*i.e.*
52 reduction in fitness caused by matings between related individuals; reviewed in Charlesworth & Willis,
53 2009; Mackauer, 1976), or the interaction of the two. Indeed, the selection that occurs during colony
54 establishment creates conditions conducive to inbreeding (Roush, 1986). Families with higher fitness
55 under laboratory conditions contribute disproportionately to the reproductive pool, thereby increasing
56 the probability of matings between related individuals. Where selection is very strong, as in the
57 production of an insecticide resistant colony, measures must often be taken to minimize the effects of
58 inbreeding and thereby inbreeding depression (Gould *et al.*, 1995). Overall, the expectation is that the
59 selection and inbreeding that takes place during insect colonization results in an overall loss of genetic

60 diversity (Munstermann, 1994), and concomitant genome-wide increase in homozygosity (reviewed in
61 Etzel & Legner, 1999).

62 Previous studies that have examined genetic differences between field-collected, laboratory-
63 adapted, and inbred populations of non-model insects have primarily focused on Dipteran species and
64 were limited to small numbers of molecular markers (Mukhopadhyay *et al.*, 1997; Munstermann, 1994;
65 Norris *et al.*, 2001). Such small numbers of markers allow for estimation of the genome-wide average
66 change in genetic variability across populations, but cannot be used to examine fine-scale patterns of
67 genomic change. Examination of these patterns allows for identification of where and how genetic
68 variation, the raw material necessary for environmental adaptation, is maintained (Dobzhansky &
69 Spassky, 1954). The relatively recent development of high-throughput sequencing combined with
70 reduced-representation DNA library preparation techniques allows for the discovery of hundreds to
71 thousands of new molecular markers, even in species for which genomic data are absent (Davey *et al.*,
72 2011). Here we used Double-Digest Restriction-Site Associated DNA Sequencing (ddRAD-seq;
73 Peterson *et al.*, 2012), one type of reduced representation library, for *de novo* construction of molecular
74 markers in the non-model species, *Heliothis virescens*.

75 The tobacco budworm, *H. virescens*, is an historically important pest of cotton throughout much
76 of the Southeastern United States (Blanco, 2012). This non-model Lepidopteran species has been
77 colonized a number of times for investigations of mating and host-selection behaviors (Sheck & Gould,
78 1995; Sheck & Gould, 1996; Sheck *et al.*, 2006), as well as detecting the underlying genetic basis for
79 insecticide resistance (Gahan *et al.*, 2001; Gahan *et al.*, 2010; Taylor *et al.*, 1993). We used our newly
80 developed ddRAD-seq markers to examine and compare the effects of colonization, selection, and sib-
81 mating on *H. virescens* genome-wide measures of genetic diversity. To examine fine-scale patterns of
82 change in genetic diversity, we also used our ddRAD-seq markers to generate a dense intraspecific

83 genetic map for *H. virescens*. This map consists of 659 high quality 350-bp markers which will serve
84 as an important genomic resource to the entomological community.

85 Overall, our research aims to:

- 86 1. Quantify overall patterns of change in genomic diversity across field-collected, laboratory-
87 reared (non-selected), Bt-selected, and sib-mated *H. virescens*.
- 88 2. Determine whether the observed degree of inbreeding in our sib-mated *H. virescens* calculated
89 from our ddRAD-seq genotypic data matched theoretical expectations (Falconer & Mackay
90 1996).
- 91 3. Examine fine-scale patterns of genomic change by identifying which genomic regions were
92 resistant to fixation in Bt-selected and sib-mated lines using ddRAD-enabled linkage mapping.

93

94 **Results**

95 We sequenced 204 *H. virescens* individuals from a total of 6 populations that were used in a
96 population-level analysis of genomic change associated with laboratory colonization, artificial selection
97 and inbreeding. These populations were comprised of 2 field-collected, 2 laboratory-reared, and 1 Bt-
98 selected population, as well as specimens from a single inbred family following 10 generations of full-
99 sibling mating (Table 1). Three of these populations, YDK, YHD2, and the inbred line were founded
100 from a collection in Yadkin County, NC, in 1988 (Gould *et al.*, 1995), but were thereafter subjected to
101 different rearing conditions, allowing us to make comparisons of population-genomic change within
102 the same genetic background. In addition, 99 individuals (2 parents and 97 progeny) were sequenced
103 for linkage analysis. This produced a total of 105,487,499 Illumina MiSeq reads (38,221,995 and
104 67,265,504 for linkage- and population-level analyses, respectively) that passed quality filters (data
105 available upon request).

106 *Genomic diversity among H. virescens populations*

107 On average, 338,892 sequencing reads (*s.d.* = 113397) were produced per individual, and the
108 variation in average read counts was spread uniformly across populations (Supplementary Figure 1).
109 Ninety-four percent of individuals had read counts between 90,000 and 688,000, and were fed into the
110 Stacks pipeline (Catchen *et al.*, 2011; 2013) for *de novo* locus construction (Supplementary Figure 1).
111 Loci constructed by Stacks were sequenced to a 6× average depth of coverage per individual. In total,
112 4,281 polymorphic 350-bp ddRAD-seq markers (hereafter loci) were detected in at least one individual
113 per population across all populations. Two well-documented challenges commonly encountered when
114 working with moderate coverage reduced representation library data like ours are: 1) uneven
115 distribution of missing data across sets of loci (Davey *et al.*, 2012; Xu *et al.*, 2014), and 2) under-
116 sampling of heterozygotes (Li *et al.*, 2009; Nielsen *et al.*, 2011). Both reduce confidence in final
117 genotypes called by genotyping-by-sequencing SNP calling algorithms, including the algorithm used in
118 Stacks. To overcome these challenges, we examined several subsets of these 4,281 polymorphic loci
119 for our downstream population genomic analyses. These subsets contained between 125-1231 loci, and
120 were chosen based upon the overall proportion missing genotype calls in the subset. The smallest
121 subset consisted of loci for which over 75% of individuals per population had genotypic data present
122 and were therefore likely sequenced to greatest depth of coverage. Each larger subset allowed
123 additional loci at the expense of coverage (*i.e.* more missing genotypic data were allowed; Table 2). By
124 using multiple datasets, we were able to examine whether the presence of missing genotype calls
125 influenced overall estimates of genomic diversity across populations.

126 We first examined all subsets of loci and determined the mean and maximum numbers of alleles
127 per locus (Table 2). Across all sequenced populations (*n* = 192 total *H. virescens*), the mean numbers
128 of unique alleles detected per locus ranged from 29 to 34 depending upon the number of loci included

129 in the analysis. As more loci were included, the average number of unique alleles detected per locus
130 decreased. However, the maximum number of unique alleles detected increased from 86 in the
131 smallest subset of loci to 94 in the 3 larger subsets. We also examined the proportion of loci that were
132 fixed (*i.e.* only a single allele present) across populations. Across subsets, few loci were fixed for a
133 single allele in laboratory-reared (5.6-10.9%), Bt-selected (5.3-7.3%) and field-collected (0-2.4%)
134 populations (Table 2). Yet over 80% of loci were fixed in the inbred line following 10 generations of
135 sib-mating (Table 2). Of the 125 loci with the fewest missing genotype calls, 86% were fixed in the
136 inbred line. Expanding the number of loci to include those with more missing genotypes ($n= 378, 573,$
137 1231) reduced the percentage of fixed loci in the inbred line by up to 5% (Table 2).

138 We then determined the mean number of unique alleles present per locus for each subset of loci
139 (Supplementary Figure 2). In general, we found no within population differences in the mean numbers
140 of unique alleles detected among subsets of loci, and therefore we used a single, conservatively chosen
141 subset of loci ($n = 378$) where at least 10 individuals were genotyped per population per locus for
142 further analysis. The mean numbers of unique alleles per locus were 2.1 for the inbred line, 5.3 for the
143 Bt-selected population, 5.4 and 4.4 for the non-selected, laboratory-reared populations (YDK and
144 BENZ, respectively), and 18.4 and 17 for the field-collected populations (LA and TX, respectively).
145 However, our sample sizes (*i.e.* numbers of individuals sequenced; see Table 1) differed for each
146 population, and it was unclear whether differences between the aforementioned means were caused by
147 sample size or population-level differences. To determine whether the mean number of unique alleles
148 per locus differed according to population, where sample sizes were equal across populations, we
149 randomly sampled 6, 12, 18, and 24 alleles per population per locus without replacement. We found
150 that increasing the total number of alleles sampled led to an increase in the mean numbers of unique
151 alleles per locus for all but the inbred line (Figure 1). Regardless of the number of alleles sampled,

152 field populations always exhibited the greatest allelic diversity, followed by selected and non-selected
153 colony populations. The lowest allelic diversity was observed in the inbred line.

154 When 18 alleles were randomly sampled per population per locus, we detected an average of
155 just over 1 unique allele per locus in the inbred line, indicating that most loci were fixed for a single
156 allele. For the inbred line, 52 loci did not reach fixation. Of these, forty-seven had 2 unique alleles,
157 four had 3 unique alleles, and one had 4 unique alleles out of the 18 that were randomly sampled. On
158 average, Bt-selected and non-selected colony strains each had *ca.* 3 unique alleles per locus, and field-
159 collected populations had *ca.* 9 unique alleles per locus (Figure 1). The majority of unique alleles
160 present in the field-collected populations (70.3% and 68.7% for LA and TX populations, respectively)
161 were observed only once (of 18 alleles; Figure 2). Low frequency alleles were less common in the Bt-
162 selected and non-selected laboratory populations (35.1% and 21.5%, respectively) and rare (0.7%) in
163 the sib-mated inbred line (Figure 2).

164 To further quantify and compare genetic diversity by population, we used the same random
165 sampling regime ($n = 18$ alleles without replacement), paired with the same conservative subset of loci
166 ($n = 378$) to calculate sample-size corrected S_K (Charlesworth & Charlesworth, 2010), and π (Nei 1978;
167 Nei & Li, 1979). These two measures are complementary: π is calculated as the proportion of
168 nucleotides that differ per two randomly chosen DNA sequences, averaged across all pairwise
169 comparisons per marker per population, and S_K is calculated as the number of unique single nucleotide
170 variants in a population at a single locus. When averaged across all markers ($n = 378$), the number of
171 variant sites (S_K) per 350 bp marker was 0.15 for the inbred line, and the maximum S_K was 2.95. Bt-
172 selected and non-selected laboratory populations had, on average, just over 1 nucleotide variant per 350
173 bp locus, with a maximum of *ca.* 6. Field-collected populations had the greatest number of variant sites
174 per 350 bp locus, where the genome-wide average was just over 5 nucleotide variants per locus, with a

175 maximum of *ca.* 15. Similar trends were observed for genome-wide and maximum nucleotide diversity
176 (π) values. Relative to the laboratory-reared populations, genome-wide estimates of π were nearly an
177 order of magnitude lower for the inbred line. The genome-wide π estimate for laboratory-reared
178 populations ranged from 4.0×10^{-3} (Bt-selected population) to 6.7×10^{-3} (non-selected, YDK population),
179 and 6.2×10^{-4} for the inbred line. Field-collected populations exhibited genome-wide π estimates of
180 9.3×10^{-3} and 9.2×10^{-3} for the LA and TX populations, respectively. All genome-wide and maximum π
181 and S_K estimates, along with their corresponding 95% non-parametric bootstrapped confidence intervals
182 ($N = 5000$) are reported in Table 3.

183

184 *Inbreeding among laboratory-reared H. virescens populations*

185 To determine whether the observed degree of homozygosity in our inbred line was consistent
186 with that which would be expected following 10 generations of sib-mating, we compared the
187 inbreeding coefficient F , as calculated according to pedigree- (Falconer & Mackay, 1996) and DNA
188 marker-based information (Keller & Waller, 2002; Kim *et al.*, 2007). The expected inbreeding
189 coefficient (F_t), following 10 generations of sib-mating was 0.89. This expected value fell within the
190 bootstrapped 95% confidence intervals for marker-based inbreeding coefficients (F_{IT}) calculated from
191 all subsets of markers. This indicated that there was no significant difference between the expected
192 inbreeding coefficient and the observed inbreeding coefficient calculated using ddRAD-seq marker
193 data. The genome-wide F_{IT} values (95% CIs) were 0.92 (0.88, 0.96), 0.89 (0.86, 0.92), 0.89 (0.87,
194 0.92), and 0.88 (0.86, 0.89), for the inbred line as calculated from 125, 378, 573, and 1231 ddRAD-seq
195 loci, respectively.

196

197

198 *Linkage mapping*

199 Few genomic resources are available for *H. virescens*. Therefore, we determined the genomic
200 location of loci which were resistant to fixation by generating a dense ddRAD-seq linkage map. The
201 map was produced using a male informative cross (reviewed in Baxter *et al.*, 2009), and was comprised
202 of 659 informative polymorphic loci generated by ddRAD-seq, plus 3 partial gene sequences of
203 *ABCC2*, *HevCaLP*, and *Desat1*. Adding these partial gene sequences to our linkage map, all with
204 known locations in the *B. mori* genome allowed us to validate marker groupings for our linkage map.
205 All informative ddRAD-seq loci were grouped into 33 linkage groups, two more than the expected 30
206 *H. virescens* autosomes, and one segregating Z chromosome from the hybrid male parent used in our
207 cross. Linkage groups ranged in size from 7cM to 110cM (Figure 3), and yielded a total map length of
208 1919.5 cM. On average, there were 20 ddRAD-seq loci per linkage group, and the average spacing
209 was one locus per 3.5 cM. The smallest and largest linkage groups contained 3 and 53 loci,
210 respectively. The *HevCaLP*, *Desat1*, and *ABCC2* genes were grouped with linkage groups 15, 16, and
211 22, respectively. These linkage groups corresponded to *B. mori* chromosomes 6, 23, and 15 (Table 4),
212 where these candidate genes are known to reside (Gahan *et al.*, 2001; Gahan *et al.*, 2010; Mita *et al.*,
213 2004).

214 In total, 99 of the 659 mapped ddRAD-seq loci could be aligned uniquely to a single locus in
215 the *B. mori* genome. Twenty-two linkage groups contained ddRAD-seq loci that could be aligned to a
216 single *B. mori* chromosome (Table 4), while 8 linkage groups did not contain any that could be aligned.
217 Linkage groups 19, 25, and 33 contained ddRAD-seq loci that aligned uniquely to more than one *B.*
218 *mori* chromosome. This was unlikely caused by spurious associations between ddRAD-seq loci;
219 increasing the LOD score to 8 failed to break up associations for those three linkage groups.

220

221 *Identification of genomic regions resistant to fixation*

222 To determine where heterozygosity was being maintained in the genome, we examined
223 nucleotide diversity at the 659 mapped ddRAD-seq loci for Bt-selected (YHD2), non-selected (YDK),
224 and inbred line. Of these 659 mapped loci, 302 (46%) were previously included in our population-level
225 analyses, and 357 were newly discovered. Nucleotide diversity (π) was only calculated for a
226 population if at least 3 individuals were genotyped at a locus. Therefore only 441 loci were examined
227 for the inbred line, and 658 loci were examined for both Bt-selected and non-selected populations. In
228 total, 60 (13%) of the mapped loci retained polymorphism in the inbred line, whereas 645 (98%) and
229 583 (86%) of mapped loci retained polymorphism in the Bt-selected and non-selected (YDK)
230 populations, respectively.

231 DdRAD-seq loci that retained polymorphism in the inbred line were spread over 16 linkage
232 groups (Figure 4), and often clustered together within a linkage group (Supplementary Figure 3). We
233 used Tajima's D, a statistical test which identifies departures from the neutral model of molecular
234 evolution, to determine whether any of these polymorphic loci were under balancing selection in the
235 inbred line. Of these 60 polymorphic sites, 20 had statistically significant Tajima's D values ($q > 0.05$)
236 indicative of balancing selection (Table 5). Linkage groups 3, 11, 14, and 15 each contained multiple
237 polymorphic sites that appeared to be under balancing selection. Whereas 5 polymorphic loci on
238 linkage group 11 were spread across the length of the linkage group, linkage groups 3, 14, and 15
239 contained multiple polymorphic loci clustered together within 5cM of one another.

240 We examined which, if any, of these polymorphic loci with strongly positive Tajima's D values
241 were shared between the inbred line, the Bt-selected population, and their ancestral laboratory-reared
242 *H. virescens* population (YDK). Following a bayesian adjustment for multiple comparisons (Efron
243 2001), no loci showed significantly positive Tajima's D values for the Bt-selected population, but 35

244 loci had positive Tajima's D values in the non-selected YDK population. While only 3 loci under
245 balancing selection were shared between the inbred line and the non-selected (YDK) populations (n =
246 3), many of the same genomic regions were shared between populations. For example, both
247 populations shared signatures of balancing selection on chromosome 15, for locus Hv20150 at 61 cM.
248 A complete list of the shared genomic regions showing signatures of balancing selection can be found
249 in Table 6.

250

251 **Discussion**

252 Here, we examined the degree to which colonization, artificial selection, and intense inbreeding
253 influence genome-wide and fine-scale patterns of diversity. In the absence of a publicly available *H.*
254 *virescens* reference genome, we used ddRAD-seq *de novo* locus construction to identify multiple
255 subsets of polymorphic loci ranging in size from 125 (where at least 75% of individuals per population
256 contained genotypic data), to 1231 (where at least 25% of individuals had genotypes present per
257 population). Genome-wide measures of allelic diversity, F_{IT} values, and the degree of homozygosity
258 were either unaffected (Supplementary Figure 2), or minimally affected (Table 2) when markers with
259 high levels of missing genotypic data were present. Therefore, any biased genotype calls made by the
260 Stacks SNP calling algorithm due to our moderate depth of sequencing coverage had little impact on
261 our overall genome-wide estimates of diversity. Our results demonstrate that moderate coverage
262 ddRAD-seq data can be used with confidence when conducting population genomic comparisons of
263 genome-wide means.

264 We observed a precipitous decline in nucleotide and allelic diversity following long-term
265 laboratory colonization, selection, and inbreeding for *H. virescens*. Despite the decline in genomic
266 diversity for non-selected and Bt-selected laboratory-reared populations, fewer than 10% of loci were

267 fixed. While our Bt-selected population did not retain the level of genetic diversity that their ancestral
268 (YDK) laboratory-reared population did, they consistently had higher measures of genomic diversity
269 than did the non-selected (BENZ) population. Retention of higher levels of polymorphism in our Bt-
270 selected line was likely due to the measures taken during its generation to ensure genomic diversity was
271 maintained in the face of strong selection (Gould, 1995). Alternatively, strong genetic bottlenecks in
272 the non-selected (BENZ) population prior to their use in our study could explain why our Bt-selected
273 line was more genetically diverse than the non-selected line. Overall, differences among Bt-selected
274 and non-selected laboratory-reared populations were modest; when 18 total alleles were sampled,
275 laboratory-reared populations retained *ca.* 3 alleles per 350-bp locus relative to 9 alleles per 350-bp
276 locus present in field-collected *H. virescens*. However, few low frequency alleles remained in the
277 laboratory-reared populations relative to the field-collected populations (Figure 2), which has been
278 observed elsewhere (Munstermann, 1994). For this reason, laboratory-reared populations are generally
279 considered inbred (Roush, 1986). In the case of *H. virescens*, our results clearly show that a great deal
280 of genomic diversity is retained, even following decades in colony. Few genome assembly algorithms
281 accommodate polymorphism well (Kajitani *et al.*, 2014), and it is clear that the reductions in
282 heterozygosity in our inbred line will be useful for production of a high quality *H. virescens* reference
283 genome assembly.

284 Following 10 generations of inbreeding, > 80% of markers went to fixation in our sib-mated *H.*
285 *virescens* population. Indeed, our inbreeding coefficient F_{IT} , as observed from our ddRAD-seq data,
286 met theoretical expectations for all subsets of loci. Our *H. virescens* laboratory population was more
287 amenable to inbreeding than other insect species (Munstermann, 1994; Rumball *et al.*, 1994; Turissini
288 *et al.*, 2014; You *et al.*, 2013), despite their relatively high levels of genomic diversity (Figure 1) and
289 genetic load (Supplementary Figure 4). Higher than expected allelic diversity has been observed in

290 several other insect species following experimental inbreeding attempts (Munstermann, 1994; Rumball
291 *et al.*, 1994; Turissini *et al.*, 2014; You *et al.*, 2013). As one example, only 57% of the *An. gambiae*
292 genome went to fixation, as observed according to SNP markers, following 10 generations of
293 inbreeding (Turissini *et al.*, 2014). Observed differences between our *H. virescens* population and
294 other insects could be species specific, but is more likely related to the proportion of the genome
295 containing balanced lethal systems (Falconer & Mackay, 1996).

296 To determine where heterozygosity was being maintained in our inbred line, we developed a
297 high density genetic linkage map for *H. virescens*. Our map contained 659 newly developed 350 bp
298 ddRAD-seq markers that are long enough for future primer design and direct sequencing. This map
299 represents a new tool for an historically important pest species that lacks genomic resources. No
300 ddRAD-seq markers remained unlinked following mapping, which indirectly speaks to the quality of
301 our linkage map. However, the number of groups in our linkage map was 2 more than expected
302 ($n = 31$). This is likely due to the relatively small mapping population size used in this work (Silva *et*
303 *al.*, 2007). Other mapping studies that analyzed segregating populations of a similar size have also
304 reported genetic maps with excess numbers of linkage groups (Pootakham *et al.*, 2015; Singh *et al.*,
305 2009). Another possible explanation for the disparity between our observed and expected number of
306 linkage groups is that polymorphic markers may not be evenly distributed over the chromosomes
307 (Paterson, 1996). Perhaps the markers in several of our smaller linkage groups are found on opposite
308 ends of *H. virescens* chromosomes, and thus are unlikely to show strong associations.

309 When we applied our linkage map to examine fine-scale patterns of genomic change following
310 10 generations of sib-mating, we found several linkage groups contained clusters of loci that did not go
311 to fixation (Figure 4). Twenty out of the 60 ddRAD-seq loci not fixed for a single allele showed
312 signatures of balancing selection (Table 5). These loci were likely genetically linked to balanced lethal

313 systems, or perhaps chromosomal inversions. Twelve loci spanning the entire length of linkage group
314 11 (syntenic to *B. mori* chromosome 9) did not go to fixation in the inbred line, and 5 appeared to be
315 under balancing selection. Furthermore, several loci on this linkage group were under balancing
316 selection in our non-selected (YDK) population, suggesting that a large chromosomal inversion may be
317 present on *H. virescens* linkage group 11.

318 Several smaller genomic regions with signatures of balancing selection were also shared among
319 our non-selected population (YDK) and inbred line (Table 6). These regions are likely associated with
320 deleterious alleles that were sampled from the original field population in Yadkin County, NC from
321 which our YDK population derived. It is possible that those deleterious alleles were never purged from
322 YDK, but instead increased in frequency creating several balanced lethal systems. YDK is ancestral to
323 the inbred line, so these balanced lethals were likely transmitted the inbred line as it was generated.

324

325 **Conclusions**

326 This work serves as one of the most thorough attempts to quantify the effects of genomic
327 responses to selection and inbreeding in a non-model insect species. We demonstrated that laboratory-
328 reared *H. virescens* have reduced allelic and nucleotide diversity relative to field-collected populations,
329 and that inbreeding further diminishes genetic diversity. Although we identified several loci that did
330 not go to fixation in *H. virescens* following 10 generations of inbreeding, our ddRAD-seq marker-based
331 F_{IT} values met theoretical expectations. The genomic regions that retained polymorphism were likely
332 due to deleterious alleles that were present in the founding individuals, and would unlikely go to
333 fixation. This work demonstrates the difficulty involved in producing fully homozygous insect strains,
334 which are currently critical to producing high-quality, complete reference genomes.

335

336 **Methods**

337 *Field-collected H. virescens*

338 Adult male moths were collected from Bossier Parish, Louisiana, and College Station, Texas
339 using pheromone-baited live traps. Collections took place in LA from May through September, 2012,
340 and in TX from May through October, 2012. Moths from each collection date were immediately
341 placed in bottles of 95% ethanol for long-term storage. All bottles were held at -20 °C until DNA
342 isolations took place.

343

344 *H.virescens colonies*

345 *H.virescens* were collected from Yadkin County, NC in 1988 (Gould *et al.*, 1995). This original
346 population founded two of the colonies used in this study. YHD2 was selected for high levels of Bt
347 resistance for 4 years on MVP- treated (0.864 mg/mL diet; Mycogen, San Diego, CA) corn-soy diet
348 (Gould *et al.*, 1995), whereas a non-selected population (YDK) was reared on corn-soy diet alone. A
349 third population (BENZ) originating from Stoneville, MS, was acquired from Benzon Research
350 Incorporated (Carlisle, PA). BENZ *H. virescens* were acquired in their pupal stage, and newly eclosed
351 adults were used for population-level comparisons. To produce an inbred population, single pair
352 matings (SPMs) were set up between YDK siblings for 10 generations. An initial 37 SPMs were used
353 to establish 29 lineages in filial generation one (8 single pair matings did not produce progeny). When
354 SPMs failed to produce offspring, likely due to inbreeding depression, surviving lineages were
355 expanded (Supplementary Figure 4). This was done to extend inbreeding for as many generations as
356 possible, thus promoting as complete a reduction in heterozygosity as possible. Adult males from each
357 laboratory-reared population were killed by freezing (-20 °C), and stored at -80 °C until DNA
358 isolation.

359

360 *Mapping cross*

361 A non-selected female from the BENZ population was crossed to a Bt-selected (YHD2) male in
362 a single pair mating. One hybrid male offspring was then back-crossed to a Bt-selected (YHD2)
363 female, and their progeny were reared to adulthood on untreated corn-soy diet according to Joyner and
364 Gould (1985). Of the 120 progeny, 97 reached adulthood. Parents and their 97 adult progeny were
365 killed by freezing and stored until DNA isolation as described above.

366

367 *Genomic DNA library preparation*

368 All DNA was isolated from the adult thorax using a Qiagen Dneasy Blood and Tissue Kit
369 (Qiagen, Inc., Valencia, CA, U.S.A.). Genomic DNA samples were prepared for Illumina sequencing
370 according to the Poland *et al.*, (2012) protocol with minor modifications. Two-hundred ng of DNA per
371 individual were digested with EcoRI and MspI. For each individual, the overhang sites were ligated to
372 standard Truseq Universal adapters (Illumina, Inc. San Diego, CA). These adapters, which were
373 ligated to EcoRI overhang sites, contained one of 48 unique barcodes (Elshire *et al.*, 2011;
374 Supplementary Table 1). DNA fragments from each individual were assigned a unique barcode, and
375 individuals were combined into pools of no more than 48 individuals. A Pippin Prep (Sage Science,
376 Inc., Beverly, MA) was used to select adapter-ligated DNA fragments ranging from 450-650 bp from
377 each pool. Size-selected DNA fragments were amplified in a Peltier PTC200 thermalcycler (here and
378 throughout) using Illumina primers (Supplementary Table 2) under the following conditions: 72 °C for
379 5min, 18 cycles of 98 °C for 30 sec, 65 °C for 20 sec, 72 °C for 30 sec, followed by 72 °C for 5 min.
380 For each pool, 1 of 4 Illumina indices was added via PCR to the MspI adapter. Therefore, sequences
381 from each individual could be identified by the unique combination of barcode and index. A complete

382 list of barcodes and indices used in this study can be found in the Supplementary Tables 1 and 2,
383 respectively. Amplified libraries were pooled, cleaned with a Qiaquick PCR Purification Kit (Qiagen,
384 Inc., Valencia, CA, U.S.A.), and diluted to 4nM prior to sequencing. Prepared genomic DNA libraries
385 constructed from 303 *H. virescens* individuals were spread across 9 full and partial Illumina MiSeq
386 runs.

387

388 *De novo marker formation*

389 Overlapping paired-end reads were merged with FLASH (Magoc & Salzberg, 2011), and Stacks
390 v. 1.09 (Catchen *et al.*, 2011; 2013) was used for demultiplexing and *de novo* formation of loci.
391 Merged paired-end reads were filtered for quality using the process_radtags script. Further quality
392 filtering entailed removal of reads when: 1) they did not have an intact EcoRI cut site, 2) had a quality
393 score < 30, or 3) were smaller than 350 bp. We did not allow process_radtags to rescue reads where
394 barcode sequences contained an error. All remaining merged reads were truncated at a length of 350
395 bp, and fed into the Stacks pipeline.

396

397 *Stacks parameter settings*

398 Reads from all individuals were run through ustacks with the following parameter settings: -m
399 3, -M 14 (allowing for 5% nucleotide mismatch rate between alleles per individual), -max_locus_stacks
400 2, --alpha 0.05. A consensus catalog of loci was first formed using the parents of the mapping cross
401 with cstacks, where the -n 14 parameter allowed for a 5% between individual nucleotide mismatch rate.
402 For the mapping family, genotype calls were made using sstacks prior to field- and colony-strain alleles
403 being added to the catalog. Progeny genotypes were automatically corrected using the Stacks
404 genotypes script. Twenty-four individuals of each colony and field-collected strain collected in 2012

405 were later added to the catalog, and all field-collected and laboratory-reared populations were also
406 genotyped using sstacks.

407

408 *Data analyses*

409 All population genomic and linkage analysis were conducted in R version 3.1.2 (R core team,
410 2014).

411

412 *Genomic diversity among H. virescens populations*

413 In total, we sequenced the 13 surviving males from an inbred line subjected to 10 generations of
414 sib-mating, 42-46 males per colony strain, and 30 males per field-collected population (Table 1). Prior
415 to running sequence data through the Stacks pipeline, we checked individual read counts across
416 populations to ensure uniformity (Supplementary Figure 1). Twelve of the 204 individuals sequenced
417 had too few (< 90,000) or too many reads (> 688,000) and were removed from the dataset prior to
418 analysis, following Bi *et al.* (2013). From our Stacks output, we constructed 4 different sets of
419 consensus loci present across populations. These subsets, containing a core overlapping set of 125 loci,
420 and increasing in size from 125 to 1231, consisted of marker sets with varying percentages of missing
421 genotype calls (range = 11.2-29.5%) (Table 2). We used these 4 different subsets to examine and
422 compare changes in genomic diversity across populations.

423 We estimated the mean number of unique alleles present per locus, and corresponding 95% non-
424 parametric bootstrapped confidence intervals (N = 5000) across populations using a custom-written R
425 script. Within a population, each locus was randomly sampled without replacement to a depth of 6, 12,
426 18, and 24 haplotypes, and then unique alleles per locus were counted. Due to their small sample size
427 resulting from intensive inbreeding, we only sampled 6, 12, and 18 haplotypes per locus for the inbred

428 line. We focused our analysis on the subset of loci containing 378 consensus loci because genotype
429 calls were present for at least 10 individuals per population. Increasing the number of loci analyzed did
430 not significantly change the mean number of unique alleles per population (Supplementary Figure 2).
431 We also calculated two measures of nucleotide diversity per 350 bp locus using the R package, pegas
432 (v. 0.6; Paradis, 2010): π (Nei, 1987) and S_K corrected for sample size (Charlesworth & Charlesworth,
433 2010; Watterson, 1975). We then generated population-level genome-wide means and 95% non-
434 parametric bootstrapped ($N = 5000$) confidence intervals for each metric (Table 3).

435

436 *Estimating the inbreeding coefficient*

437 To estimate our marker-based inbreeding coefficient, we examined multiple sets of loci (Table
438 2) and found that trends across all datasets were similar (data not shown). However, we reported F_{IT}
439 values from a set of 378 loci because the dataset contains few missing genotypes per population, while
440 still making inferences from several hundred loci. We calculated F_{IT} for the inbred line relative to the
441 non-selected (YDK) population after Keller and Waller (2002), where $(1-F_{IS})(1-F_{ST}) = 1-F_{IT}$. F_{IS} was
442 the level of inbreeding within the inbred line, calculated as $F_{IS} = 1-(H_O/H_E)$, where H_O and H_E were
443 calculated for each locus using the R package adegenet (v. 1.4-2; Jombart, 2008). F_{ST} was the
444 accumulated effect of inbreeding over time, calculated as $1-(H_E(\text{inbred line})/H_E(\text{YDK}))$.

445

446 *Amplification and genotyping of PCR-based markers for the mapping family*

447 Progeny from the mapping family were genotyped at three additional loci, and these loci were
448 mapped alongside our ddRAD-seq markers to validate our linkage groupings. We targeted the
449 previously described *H. virescens* genes *DesatI*, *ABCC2*, and *HevCaLP* via PCR followed by gel
450 electrophoresis, or direct sequencing. Amplification and genotyping protocols were as follows.

451 A 468 bp fragment from *DesatI* was amplified in a 30 μ l reaction with forward and reverse
452 primers [5'-TGAGGGACCATCGTCTCCAT-3'] and [5'-CACTGCTACATTTTGGGCAG-3'],
453 respectively (Ward, 2009). Each reaction contained 6 μ L of 5 \times GoTaq buffer (ProMega), 29 μ M per
454 dNTP, 92 ng per primer, 0.75 U GoTaq polymerase, and *ca.* 1 μ g genomic DNA. Sample DNA was
455 amplified alongside a negative control (here and throughout), where pcr-grade H₂O was substituted for
456 genomic DNA. Reactions were incubated at 95°C for 1 min followed by 35 cycles of 95°C for 1min,
457 52°C for 1min, and 72°C for 2 min. PCR products were purified using a standard ethanol precipitation,
458 and directly sequenced on an ABI3730xl (Applied Biosystems, San Francisco, CA). A single
459 nucleotide polymorphism (cytosine to tyrosine substitution) at bp 36 was found in the YHD2 parent of
460 the mapping cross. Offspring were genotyped at this locus using PolyPhred (Nickerson *et al.*, 1997),
461 and genotype calls were visually confirmed using consed (Gordon *et al.*, 1998).

462 An intronic region of the *ABCC2* gene previously described by Gahan *et al.* (2010) was
463 amplified using primers Hs-ABC2dU02-F1 [5' – TGGTTACAAGAAATAGAAAATGCAAC-3'] and
464 Hs-ABC2eU03-R2 [5' – CTTTCAAACCTGAACCGCATCAC – '3]. Each 30 μ l reaction volume
465 consisted of 6 μ L of 5 \times GoTaq buffer, 29 μ M per dNTP, 73 ng per primer, 0.75 U GoTaq polymerase,
466 and 1 μ g genomic DNA. Reactions were held at 95°C for 2 min followed by 30 cycles of 95°C for 30
467 sec, 58°C for 30 sec, and 72°C for 40 sec, and the resulting products were cleaned via ethanol
468 precipitation. Following sequencing on an ABI3730xl, chromatogram files were visualized using
469 FinchTV (version 1.3.1, PerkinElmer, Inc., Seattle, WA). As described by Gahan *et al.* (2010), the
470 YHD2 parent was homozygous for a 22 bp deletion, whereas the F₁ parent was heterozygous for this
471 deletion. Therefore, we examined the segregation of this deletion in the mapping family offspring,
472 which was detectable by the presence of a TAT sequence near amplicon bp 40.

473 Finally, the *HevCaLP* locus described by Gahan *et al.* (2007) was amplified in a multiplexed

474 reaction using three primers: the universal reverse primer [5'-
475 ATACGAGCTGACGACACGCTGGGAGA-3'], one forward primer that targets a retrotransposon
476 insertion conferring resistance to *Bacillus thuringiensis* [5' –
477 CGCAACGCGGATCTACTCTTGTCACC – 3'], and another forward primer that targets wild-type
478 sequence [5' – AAGTGTCCCAGTCGATGCTGAA – 3']. An initial 20- μ l reaction contained 4 μ l 5 \times
479 GoTaq buffer, 29 μ M per dNTP, 56 ng per primer, 0.5 U GoTaq polymerase, and 1 μ g genomic DNA.
480 Reactions were incubated at 95°C for 2 min, followed by 30 cycles of 95°C for 30 sec, 58°C for 20 sec,
481 and 72°C for 40 sec. A reconditioning reaction, aimed at reducing heteroduplex formation, was set up
482 as above, but incubated for 3 rather than 30 cycles. These reactions were capable of producing two
483 amplicons, which differed in length by 76 bp. The YHD2 parent was homozygous for the long
484 amplicon (*ca.* 800 bp) containing the insertion that confers resistance to *Bacillus thuringiensis*, whereas
485 the F₁ parent was heterozygous for a long and short amplicon. PCR products from mapping family
486 offspring were run on a 2% agarose gel alongside Hyperladder I (Bioline, Taunton, MA) for
487 visualization and genotype scoring.

488

489 *Linkage mapping*

490 Double-digest RAD-seq markers present in fewer than 75% of the mapping family offspring
491 were filtered out, and the remainder were checked using a chi-square test for mendelian segregation (α
492 = 0.01). PCR-based markers, as well as those ddRAD-seq markers that segregated in a mendelian
493 fashion were assigned to linkage groups (LOD = 5, maximum recombination fraction = 0.3) using the
494 onemap package (Margarido *et al.*, 2007) in R. We validated groupings by aligning all markers to the
495 *Bombyx mori* genome using Blastn in Kaikobase version 3.2.2 (<http://sgp.dna.affrc.go.jp/KAIKObase/>).
496 Furthermore, we confirmed that the locations of the *Desat1*, *ABCC2*, and *HevCaLP* pcr-based markers,

497 as well as ddRAD-seq markers found in their respective linkage groups aligned to the same *B. mori*
498 chromosomes (Table 4). Markers on each linkage group were ordered using the recombination
499 counting and ordering algorithm (RECORD; Van Os *et al.*, 2005). RECORD was chosen based upon
500 previous studies demonstrating the reliability of its performance (Collard *et al.*, 2009, Mollinari *et al.*,
501 2009).

502

503 *Assessment of fine-scale differences in nucleotide diversity across laboratory-reared populations*

504 Markers that retained polymorphism in the inbred line were examined for signatures of
505 balancing selection in the following populations: non-selected (YDK), Bt-selected (YHD2), and the
506 inbred line. We used a Tajima's D test as calculated by the R package, pegas (v. 0.6; Paradis, 2010). A
507 bayesian false discovery rate methodology (Efron *et al.*, 2001) implemented with the R package fdrtool
508 (v.1.2.15; Strimmer 2008) was applied to Tajima's D p-values to account for multiple hypothesis tests.

509

510 **Acknowledgements**

511 Thanks to Dr. R. Whetten of North Carolina State University, and Dr. J. Schaff of the NCSU
512 Genomic Sciences Lab for their insightful suggestions on ways to improve our methods. Dr. S.
513 Micinski, Dr. J. Lopez, and Dr. J. Westbrook collected the moths used in this project. R. Waples
514 provided one of the custom scripts used in our data pipeline. This project is supported by the
515 Biotechnology Risk Assessment Program competitive grant number 2012-33522-19793 from the
516 USDA - National Institute of Food and Agriculture.

517

518

519

520 **References**

521

522 Baeshen, R., Ekechukwu, N., Toure, M., Paton, D., Coulibaly, M., Traoré, S. and Tripet, F. (2014).
523 Differential effects of inbreeding and selection on male reproductive phenotype associated with the
524 colonization and laboratory maintenance of *Anopheles gambiae*. *Malar J*, 13(1), p.19.

525 Baxter, S.W., McMillan, O., Chamberlain, N., French-Constant, R.H., Jiggins, C.D. (2009). Prospects
526 for Locating Adaptive Genes in Lepidopteran Genomes. From Molecular Biology and Genetics of the
527 Lepidoptera. CRC Press, Boca Raton, FL, USA. ed. by Goldsmith MR, Marec F. pp. 105-118.

528 Bi, K., Linderoth, T., Vanderpool, D., Good, J., Nielsen, R. and Moritz, C. (2013). Unlocking the vault:
529 next-generation museum population genomics. *Molecular Ecology*, 22(24), pp.6018-6032.

530 Blanco, C. (2012). *Heliothis virescens* and Bt cotton in the United States. *GM Crops & Food*, 3(3),
531 pp.201-212.

532 Boller, E. (1972). Behavioral aspects of mass-rearing of insects. *Entomophaga*, 17(1), pp.9-25.

533 Catchen, J., Amores, A., Hohenlohe, P., Cresko, W., Postlethwait, J. and De Koning, D. (2011). Stacks:
534 Building and Genotyping Loci De Novo From Short-Read Sequences. *G3: Genes Genomes Genetics*,
535 1(3), pp.171-182.

536 Catchen, J., Hohenlohe, P., Bassham, S., Amores, A. and Cresko, W. (2013). Stacks: an analysis tool set
537 for population genomics. *Molecular Ecology*, 22(11), pp.3124-3140.

538 Charlesworth, B. and Charlesworth, D. (1999). The genetic basis of inbreeding depression. *Genetical*
539 *Research*, 74(3), pp.329-340.

540 Charlesworth, B., Charlesworth, D. (2010). Elements of Evolutionary Genetics. Roberts and Company
541 Publishers, Greenwood, CO, USA. pp. 29.

542 Charlesworth, D., and Willis, J.H. (2009). The genetics of inbreeding depression. *Nature Reviews*
543 *Genetics*, 10, pp.783-796.

544 Collard, B., Mace, E., McPhail, M., Wenzl, P., Cakir, M., Fox, G., Poulsen, D., Jordan, D. (2009). How
545 accurate are the marker orders in crop linkage maps generated from large marker datasets? *Crop and*
546 *Pasture Science*, 60(4), pp.362-372.

547 Collins A.M. Artificial selection of desired characteristics in insects. Advances and Challenges in
548 Insect Rearing, Eds. EG King and NC Leppla. USDA Technical Bulletin. Jan. 1984 pp. 9-19.

549 Davey, J.W., Cezard, T., Fuentes-Utrilla, P., Eland, C., Gharbi, K., and Blaxter, M.L. (2012). Special
550 features of RAD sequencing data: implications for genotyping. *Molecular Ecology*, 22, pp.3151-3164.

551 Dekker, T., Ibba, I., Siju, K.P., Stensmyr, M.C., Hansson, B.S. (2006). Olfactory shifts parallel
552 superspecialism for toxic fruit in *Drosophila melanogaster* sibling, *D. sechellia*. *Current Biology*,
553 16(1), pp.101-109.

554 de Valdez, M.F.W., Nimmo, D., Betz, J., Gong, H.F., James, A.A., Alphey, L. and Black, W.C. (2011).
555 Genetic elimination of dengue vector mosquitoes. *Proceedings of the National Academy of Sciences*,
556 *USA*, 108(12), pp.4772-4775.


- 557 Dobzhansky, T., and Spassky, N. (1954). Environmental modification of heterosis in *Drosophila*
558 *pseudoobscura*. *Proceedings of the National Academy of Sciences, USA*, 40(6), pp.407-415.
- 559 Efron, B. (2001). Empirical bayes analysis of a microarray experiment. *Journal of the American*
560 *Statistical Association*, 102, pp.93-103.
- 561 Elshire, R.J., Glaubitz, J.C., Sun, Q., Poland, J.A., Kawamoto, K., Buckler, E.S., Mitchell, S.E. (2011).
562 A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. *PloS ONE*,
563 6(5), pp. e19379.
- 564 Etzel, L.K., and Legner, E.F. (1999). Culture and Colonization, Chapter 7 (pp. 125-176) in Handbook
565 of Biological Control, Academic Press, San Diego, CA, USA. ed. Bellows TS and Fisher TW.
- 566 Falconer, D.S., and Mackay TFC (1996) Introduction to Quantitative Genetics. Fourth Edition,
567 Longman Group Ltd. Essex, England. pp. 89.
- 568 Fox, C.W., Scheibly, K.L., Smith, B.P., Wallin, W.G. (2007). Inbreeding depression in two seed-feeding
569 beetles *Callosobruchus maculatus* and *Stator limbatus* (Coleoptera: Chrysomelidae). *Bulletin of*
570 *Entomological Research*, 97(1), pp.49-54.
- 571 Gahan, L.J., Gould, F., Heckel, D.G. (2001). Identification of a gene associated with Bt resistance in
572 *Heliothis virescens*. *Science*, 293, pp.857-860.
- 573 Gahan, L.J., Pauchet, Y., Vogel, H., Heckel, D.G. (2010). An ABC transporter mutation is correlated
574 with insect resistance to *Bacillus thuringiensis* Cry1Ac toxin. *PloS Genetics*, 6(12), pp. e1001248.
- 575 Gerloff, C.U., Ottmer, B.K., Schmid-Hempel, B. (2003). Effects of inbreeding on immune response and
576 body size in a social insect, *Bombus terrestris*. *Functional Ecology*, 17(5), pp. 582-589.
- 577 Goldman, I. F., Arnold, J., and Carlton, B. C. (1986). Selection for resistance to *Bacillus thuringiensis*
578 subspecies israelensis in field and laboratory populations of the mosquito *Aedes aegypti*. *Journal of*
579 *Invertebrate Pathology*, 47(3), pp. 317-324.
- 580 Goldsmith, M.F., and Marec, F. (2010). Molecular Biology and Genetics of the Lepidoptera. CRC
581 Press, Boca Raton, FL. pp.28-29.
- 582 Gordon, D., Abajian, C., Green, P. (1998). Consed: a graphical tool for sequence finishing. *Genome*
583 *Research*, 8, pp. 195-202.
- 584 Gould, F., Anderson, A., Reynolds, A., Bumgarner, L., Moar, W. (1995). Selection and genetic analysis
585 of a *Heliothis virescens* (Lepidoptera: Noctuidae) strain with high levels of resistance to *Bacillus*
586 *thuringiensis* toxins. *Journal of Economic Entomology*, 88(6), pp.1545-1559.
- 587 Groot, A., Gemeno, C., Brownie, C., Gould, F., and Schal, C. (2005). Male and female antennal
588 responses in *Heliothis virescens* and *H. subflexa* to conspecific and heterospecific sex pheromone
589 compounds. *Environmental Entomology*, 34(2), pp. 256-263.
- 590 Hohenlohe, P.A., Bassham, S., Etter, P.D., Stiffler, N., Johnson, E.A., Cresko, W.A. (2010). Population
591 genomics of parallel adaptation in threespine stickleback using sequenced RAD tags. *PLoS Genetics*,
592 6(2), pp. e1000862.
- 593

- 594 Hoy, M.A. (1990). Genetic improvement of parasites and predators. FFTC-NARC International
595 Seminar on 'The use of parasitoids and predators to control agricultural pests', Tukuha Science City,
596 Ibaraki-ken, 305 Japan, October 2-7, 1989. pp.15
- 597 Huettel, M.D. (1976). Monitoring the quality of laboratory-reared insects: A biological and behavioral
598 perspective. *Environmental Entomology*, 5(5), pp. 807-814.
- 599 Jombart, T. (2008). adegenet: a R package for the multivariate analysis of genetic markers.
600 *Bioinformatics*, 24, pp. 1403-1405.
- 601 Joyner, K. and Gould, F. (1985). Developmental consequences of cannibalism in *Heliothis zea*
602 (Lepidoptera: Noctuidae). *Annals of the Entomological Society of America*, 78, pp. 24-28.
- 603 Kajitani, R., Toshimoto, K., Noguchi, H., Toyoda, A., Ogura, Y., Okuno, M., Yabana, M., Harada, M.,
604 Nagayasu, E., Maruyama, H., Kohara, Y., Fujiyama, A., Hayashi, T., and Itoh, T. (2014). Efficient *de*
605 *novo* assembly of highly heterozygous genomes from whole-genome shotgun short reads. *Genome*
606 *Research*, 24, pp. 1384-1395.
- 607 Keller, L.F., and Waller, D.M., (2002). Inbreeding effects in wild populations. *Trends in Ecology and*
608 *Evolution*, 17(5), pp. 230-241.
- 609 Kim, S.H., Cheng, K.M., Ritland, C., Ritland, K., and Silversides, F.G., (2007). Inbreeding in the
610 Japanese quail estimated by pedigree and microsatellite analyses. *Journal of Heredity*, 98(14), pp. 378-
611 381.
- 612 Li, R., Li, Y., Fang, X., Yang, H., Wang, J., Kristiansen, K., and Wang, J. (2009). SNP detection for
613 massively parallel whole-genome resequencing. *Genome Research*, 19, pp. 1124-1132.
- 614 Mackay, T.F.C., Richards, S., Stone, E.A., Barbadilla, A., Ayroles, J.F., Zhu, D., *et al.*, (2012) The
615 *Drosophila melanogaster* genetic reference panel. *Nature*, 482, pp. 173-178.
- 616 Magoc, T., and Salzberg, S. (2011). FLASH: Fast length adjustment of short reads to improve genome
617 assemblies. *Bioinformatics*, 27(21), pp. 2957-2963.
- 618 Margarido, G. R. A., Souza, A. P. and Garcia, A. A. F. (2007). OneMap: software for genetic mapping
619 in outcrossing species. *Hereditas*, 144, pp. 78-79.
- 620 Mita, K., Kasahara, M., Sasaki, S., Nagayasu, Y., Yamada, T., Kanamori, H., Namiki, N., Kitagawa, M.,
621 Yamashita, H., Yasukochi, Y., Kadono-Okuda, K., Yamamoto, K., Ajimura, M., Ravikumar, G.,
622 Shimomura, M., Nagamura, Y., Shin-i, T., Abe, H., Shimada, T., Morishita, S., and Sasaki, T. (2004)
623 The Genome Sequence of Silkworm, *Bombyx mori*. *DNA Research*, 11(1), pp. 27-35
- 624 Mukhopadhyay, J., Rangel, E.F., Ghosh, K., Munstermann, L.E. (1997). Patterns of genetic variability
625 in colonized strains of *Lutzomyia longipalpis* (Diptera: Psychodidae) and its consequences. *American*
626 *Journal of Tropical Medicine and Hygiene*, 57(2), pp. 216-221.
- 627 Munstermann, L.E. (1994). Unexpected genetic consequences of colonization and inbreeding: allozyme
628 tracking in Culicidae (Diptera). *Annals of the Entomological Society of America*, 87(2), pp. 157-164.
- 629 Nei M (1987) *Molecular evolutionary genetics*. New York: Columbia University Press.
- 630

- 631 Nickerson, D.A., Tobe, V.O., and Taylor, S.L. (1997). PolyPhred: automating the detection and
632 genotyping of single nucleotide substitutions using fluorescence-based resequencing. *Nucleic Acids*
633 *Research*, 25(14), pp. 2745-2751.
- 634 Nielsen, R., Paul, J.S., Albrechtsen, A., Song, Y.S. (2011). Genotype and SNP calling from next-
635 generation sequencing data. *Nature Reviews Genetics*, 12, pp. 443-451.
- 636 Norris, D.E., Shurtleff, A.C., Toure, Y.T., Lanzaro, G.C. (2001). Microsatellite DNA polymorphism and
637 heterozygosity among field and laboratory populations of *Anopheles gambiae* s.s. (Diptera: Culicidae).
638 *Journal of Medical Entomology*, 38(2), pp. 336-340.
- 639 Oppenheim, S.J., Gould, F., Hopper, K.R. (2012). The genetic architecture of a complex ecological
640 trait: host plant use in the specialist moth, *Heliothis subflexa*. *Evolution*, 66(11), pp. 3336-3351.
- 641 Paradis, E. (2010). pegas: an R package for population genetics with an integrated-modular approach.
642 *Bioinformatics*, 26, 419-420.
- 643 Paterson, A.H. (1996). Making genetic maps. A.H. Paterson (Ed.), *Genome Mapping In Plants*, R. G.
644 Landes Company, San Diego, California, pp. 23–29
- 645 Poland, J.A., Brown, P.H., Sorrells, M.E., Jannink, J.L. (2012). Development of high-density genetic
646 maps for barley and wheat using a novel two-enzyme genotyping-by-sequencing approach. *PLoS ONE*,
647 7(2), pp. e32253.
- 648 Pootakham, W., Jomchai, N., Ruang-areerate, P., Shearman, J. R., Sonthirod, C., Sangsrakru, D.
649 Tragoonrun, S., Tangphatsornruang, S., (2015). Genome-wide SNP discovery and identification of QTL
650 associated with agronomic traits in oil palm using genotyping-by-sequencing (GBS). *Genomics*,
651 105(5), pp. 288-295.
- 652 Pradeep, A. R., Chatterjee, S. N., & Nair, C. V. (2005). Genetic differentiation induced by selection in
653 an inbred population of the silkworm *Bombyx mori*, revealed by RAPD and ISSR marker systems.
654 *Journal of Applied Genetics*, 46(3), pp. 291.
- 655 R core team (2014). R: A language and environment for statistical computing. R Foundation for
656 Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- 657 Raulston, J.R. (1975). Tobacco budworm: observations on the laboratory adaptation of a wild strain.
658 *Annals of the Entomological Society of America*, 68, pp. 139-142.
- 659 Roush, R.T. (1986). Inbreeding depression and laboratory adaptation in *Heliothis virescens*
660 (Lepidoptera: Noctuidae). *Annals of the Entomological Society of America*, 79, pp. 583-587.
- 661 Shaw, K.L. (2000). Interspecific genetics of mate recognition: inheritance of female acoustic preference
662 in Hawaiian crickets. *Evolution*, 54(4), pp.1303-1312.
- 663 Sheck, A. L., and Gould, F. (1995). Genetic analysis of differences in oviposition preferences of
664 *Heliothis virescens* and *H. subflexa* (Lepidoptera: Noctuidae). *Environmental Entomology*, 24(2), pp.
665 341-347.
- 666 Sheck, A. L., and Gould, F. (1996). The genetic basis of differences in growth and behavior of
667 specialist and generalist herbivore species: selection on hybrids of *Heliothis virescens* and *Heliothis*


- 668 *subflexa* (Lepidoptera). *Evolution*, 50(2), pp. 831-841.
- 669 Sheck, A. L., Groot, A. T., Ward, C. M., Gemeno, C., Wang, J., Brownie, C., Schal, C., and Gould, F.
670 (2006). Genetics of sex pheromone blend differences between *Heliothis virescens* and *Heliothis*
671 *subflexa*: a chromosome mapping approach. *Journal of Evolutionary Biology*, 19(2), pp. 600-617.
- 672 Singh, R., Tan, S.G., Panandam, J.M., Rahman, R.A., Ooi, L.C.L., Low, E.L., Sharma, M., Jansen, J.,
673 and Cheah, S. (2009). Mapping quantitative trait loci (QTLs) for fatty acid composition in an
674 interspecific cross of oil palm. *BMC Plant Biology*, 9, 114.
- 675 Strimmer, K. (2008). fdrtool: a versatile R package for estimating local and tail area-based false
676 discovery rates. *Bioinformatics Applications*, 24(12), pp. 1461-1462.
- 677 Tajima, F. (1989) Statistical method for testing the neutral mutation hypothesis by DNA polymorphism.
678 *Genetics*, 123, pp. 585-595.
- 679 Taylor, M.F.J., Heckel, D.G., Brown, T.M., Kreitman, M.E., Black, G. (1993). Linkage of pyrethroid
680 resistance to a sodium channel locus in the tobacco budworm. *Insect Biochemistry and Molecular*
681 *Biology*, 23(7), pp. 763-775.
- 682 Tomaru, M., Doi, M., Higuchi, H., Oguma, Y. (2000). Courtship song recognition in the *Drosophila*
683 *melanogaster* complex: heterospecific songs make females receptive in *D. melanogaster*, but not in *D.*
684 *sechellia*. *Evolution*, 54(4), pp. 1286-1294.
- 685 Turissini, D.A., Gamez, S., White, B.J. (2014). Genome-wide patterns of polymorphism in an inbred
686 line of the African malaria mosquito, *Anopheles gambiae*. *Genome Biology and Evolution*, 6(11), pp.
687 3094-3104.
- 688 Van Os, H., Stam, P., Visser, R.G.F., Van Eck, H.J. (2005). RECORD: a novel method for ordering loci
689 on a genetic linkage map. *Theoretical and Applied Genetics*, 112, pp. 30-40.
- 690 Ward, M.D. (2009). Genetics of Sex Pheromones: Mapping Desaturase Genes in *Heliothis* Species
691 (Unpublished Master's Thesis). North Carolina State University, Raleigh, NC.
- 692 Watterson, G.A. (1975). On the number of segregating sites in genetical models without recombination.
693 *Theoretical Population Biology*, 7, pp. 256-276.
- 694 Xu, P., Xu, S., Wu, X., Tao, Y., Wang, B., Wang, S., Qin, D., Lu, Z., Li, G. (2014). Population genomic
695 analyses from low-coverage RAD-Seq data: a case study on the non-model cucurbit bottle gourd. *The*
696 *Plant Journal*, 77, pp. 430-442.
- 697 Young, J.R., Snow, J.W., Hamm, J.J., Perkins, W.D., Haile, D.G. (1975). Increasing the
698 competitiveness of laboratory-reared corn earworm by incorporation of indigenous moths from the area
699 of release. *Annals of the Entomological Society of America*, 68, pp. 40-42.

700
701 **Figures**
702




703 **Figure 1** – Mean numbers of unique alleles detected among 378 loci depend upon the number of
704 haplotypes sampled per population. For all but the inbred line (in green), as sample size increases, so
705 does the number of unique alleles detected per locus. Due to low survivorship in the inbred line, no
706 mean was computed for a haplotype sample size per locus of 24.


707
708



710 **Figure 2** – Total numbers of unique alleles detected in sib-mated (INT), Bt-selected (SEL), non-
711 selected (NS), and field-collected (FIELD) populations based upon random sampling of 18 haplotypes
712 per population per locus. Alleles were binned and color-coded according to the frequencies at which
713 they were present out of 18 total haplotypes. Numbers above each bar represent the total number of
714 unique alleles found per population out of 378 loci.



715 **Figure 3** – *Heliothis virescens* linkage map with a total length of 1919.5 cM. Centimorgan lengths are
716 below each linkage group. Each tick mark represents an individual marker that mapped to a particular
717 position in the linkage group. Red circles next to linkage groups 15 and 22 represent the positions of
718 the *HevCaLP*, and the *ABCC2*, respectively. The blue square represents the position of the delta-11-
719 desaturase on linkage group 16.
720



721 **Figure 4** – Nucleotide diversity per mapped marker site. Grey and blue circles represent nucleotide
722 diversity in non-selected (YDK), and Bt-selected (YHD2) populations, respectively. Red circles
723 represent nucleotide diversity in the inbred line following 10 generations of sib-mating.

724 **Tables**

725

726 **Table 1** - Population history, sample sizes (N) and ddRAD-seq read summary data. Filtered N refers to
727 the population sample size following removal of individuals with low read counts.

728

Population	Population History	Total N	Filtered N	Mean Number Reads per Individual
Inbred Line	10 generations of sib-mating	13	13	334689
YHD2	Bt-selected	43	41	346264
BENZ	Unselected	42	40	318636
YDK	Unselected	46	44	335728
LA2012	Field-collected	30	27	398336
TX2012	Field-collected	30	28	306164

730

731

732

733

734

735

736

737

738

739

740

741

742

743

744

745

746 **Table 2** – Multiple sets of consensus loci used to calculate population genetics parameters. Consensus sets of loci containing 125, 378, and
 747 583 loci are subsets of the largest set containing 1231 loci. The mean and maximum numbers of alleles per marker reported represent
 748 summary statistics for the entire multi-population dataset. The abbreviations Bt-sel and NS stand for Bt-selected and non-selected,
 749 respectively.

Number of Loci	Missing Genotypes (%)	Mean Number of Alleles per Marker	Max Number of Alleles per Marker	Fixed Loci (% of Total Examined)					
				Inbred line	Bt-Sel (YHD2)	NS (YDK)	NS (BENZ)	Field (LA)	Field (TX)
125	11.2	34	86	86.4	7.2	5.6	8.0	2.4	0
378	18.6	32	94	85.7	5.3	5.6	7.9	1.1	0.3
583	19.3	32	94	84.2	6.9	6.2	9.3	0.9	0.2
1231	29.5	29	94	81.3	7.3	7.5	10.9	1.4	0.9

751
 752
 753
 754
 755
 756

757 **Table 3** – Genome-wide nucleotide diversity values per 350 bp locus across populations. Values were generated by randomly sampling
 758 (without replacement) 18 haplotypes per population, then calculating π and S_K at each locus. Genome-wide values represent population-level
 759 π and S_K averaged across all loci. The abbreviations Bt-sel and NS stand for Bt-selected and non-selected, respectively.
 760

Population	Genome-wide π (2.5, 97.5% CIs)	Max π	Genome-wide S_K (2.5, 97.5% CIs)	Max S_K
Inbred line	0.0006 (0.0004, 0.0008)	0.013	0.15 (0.10, 0.19)	2.95
Bt-Sel (YHD2)	0.0040 (0.0035, 0.0043)	0.020	1.67 (1.54, 1.81)	6.53
NS (YDK)	0.0067 (0.0061, 0.0073)	0.028	1.66 (1.53, 1.80)	6.20
NS (BENZ)	0.0051 (0.0046, 0.0056)	0.026	1.35 (1.24, 1.47)	5.46
Field (LA)	0.0093 (0.0086, 0.0100)	0.034	5.23 (4.92, 5.55)	15.40
Field (TX)	0.0092 (0.0085, 0.0099)	0.033	5.02 (5.15, 5.33)	15.63

761

762 **Table 4** – Linkage group (LG) correspondence with *B. mori* chromosome (Chr). Linkage groups with an asterisk contained one or more
763 markers that aligned uniquely to an unmapped *B. mori* sequence. Where linkage groups contained markers that aligned to more than one *B.*
764 *mori* chromosome, italicized marker names correspond to the italicized *B. mori* chromosome.
765
766

LG	<i>B.mori</i> Chr	Number Markers Aligned to <i>B.mori</i> Chr	Total markers in LG	LG Length (cM)	Average Marker Spacing (cM)	Names of Markers Aligned to <i>B.mori</i> Chr
1	25	3	25	81.6	3.3	19754, 29329, 20282
2	8	6	23	52.3	2.3	22394, 27499, 21475, 17667, 17852, 25500
3*	17	3	24	74.5	3.1	376, 21931, 25112
4	-		7	35.6	5.1	
5*	4	5	36	107.3	3.0	2556, 29595, 22095, 8695, 15160
6	5	6	40	79.5	2.0	18575, 4268, 21041, 23679, 66, 5723
7	22	9	49	95.5	1.9	13710, 2328, 1443, 3654, 13880, 22897, 3123, 2430, 19858
8*	11	9	25	60.2	1.7	22584, 939, 3060, 18820, 23156, 17932, 202, 22290, 21173
9	23	2	7	42.4	6.1	4275, 1286
10	3	5	26	69.6	2.7	5015, 5329, 4265, 4545, 23262
11	9	8	37	97.8	2.6	19938, 6113, 16969, 19392, 1572, 89, 22588, 19343
12	-		18	89.5	5.0	
13	13	6	31	86.8	2.8	2074, 15962, 26388, 1193, 16684, 18825
14	19	1	21	47.0	2.2	14242
15	6	7	35	78.9	2.3	25943, 4579, 25083, 1752, 3385, 2784, 18178
16	23	4	20	55.9	2.8	2485, 16350, 806, 20599
17	7	2	14	42.5	3.0	29343, 343
18	26	1	14	57.9	4.1	1113
19	11, 22	2	4	9.5	2.4	3536, <i>16185</i>
20*	10	6	25	57.0	2.3	17820, 2978, 18970, 12277, 16316, 12984

768 **Table 4 continued.**

769

770

LG	<i>B.mori</i> Chr	Number Markers Aligned to <i>B.mori</i> Chr	Total markers in LG	LG Length (cM)	Average Marker Spacing (cM)	Names of Markers Aligned to <i>B.mori</i> Chr
21	21	3	21	68.4	3.3	26113, 20542, 7624
22	15	2	33	82.3	2.5	21411, 5773
23	14	1	17	65.2	3.8	4156
24			7	7.0	1.0	
25	12,1	4	53	110.8	2.0	20280, 9211,12382, 23604
26			7	51.0	7.3	
27	2	1	3	18.6	6.2	5781
28			6	25.3	4.2	
29	16	1	12	65.2	5.4	23481
30			6	32.4	5.4	
31			7	50.3	7.2	
32			3	6.8	2.3	
33	14, 28	2	6	18.1	3.0	6601, 20184

772

773

774 **Table 5** – Markers that retained polymorphism following 10 generations of inbreeding, and showed statistically significant ($\alpha = 0.05$)
775 signatures of balancing selection according to Tajima's D. N represents the number of individuals (of 13 total) that were genotyped per
776 locus. The q-value is derived from a bayesian false discovery rate methodology (Strimmer, 2008) that accounts for multiple comparisons.

Marker	LG	Position (cM)	N	π	Tajima's D	Unadjusted p-value	q-value
856	3	20	11	4.3×10^{-3}	2.2	0.028	0.043
19130	3	22	13	8.7×10^{-3}	2.7	0.006	0.038
2599	3	26	11	7.2×10^{-3}	2.5	0.011	0.042
6113	11	11	9	6.0×10^{-3}	2.4	0.017	0.043
19392	11	12	12	7.8×10^{-3}	2.1	0.006	0.050
20645	11	45	12	3.0×10^{-3}	2.1	0.037	0.050
13922	11	54	10	6.0×10^{-3}	2.5	0.014	0.042
21320	11	86	10	6.8×10^{-3}	2.1	0.032	0.049
68	14	29	11	5.8×10^{-3}	2.4	0.017	0.043
13550	14	34	11	1.5×10^{-2}	3.1	0.002	0.031
18240	15	48	10	1.0×10^{-2}	2.6	0.009	0.041
17301	15	50	11	4.3×10^{-3}	2.2	0.030	0.048
18676	15	59	12	4.5×10^{-3}	2.4	0.017	0.043
20150	15	61	11	6.9×10^{-3}	2.3	0.023	0.045
18178	15	61	11	5.9×10^{-3}	2.5	0.012	0.042
18240	16	32	10	4.3×10^{-3}	2.1	0.034	0.041
13777	18	27	10	2.4×10^{-2}	3.0	0.002	0.031
20662	20	0	11	4.5×10^{-3}	2.3	0.021	0.045
29612	23	36	8	1.2×10^{-2}	2.7	0.007	0.040
4851	26	1	12	8.0×10^{-3}	2.2	0.027	0.047

778 **Table 6** – Genomic regions that share signatures of balancing selection for the inbred and non-selected
779 (YDK) populations of *H. virescens*. Numbers in parenthesis following cM ranges indicate the numbers
780 of markers with significant Tajima's D values at the corrected significance threshold of $\alpha = 0.05$ within
781 the cM ranges.

782
783
784

Linkage Group	Range of Positions in cM	
	Inbred Line	Non-selected
3	20-26 (3)	22-26 (2)
11	11-86 (5)	14-86 (4)
14	29-34 (2)	30-31 (2)
15	48-61 (5)	61 (1)
23	36 (1)	22 (1)
26	1 (1)	1 (1)