

## Collinearity Analysis of Allotetraploid *Gossypium Tomentosum* and *Gossypium Darwinii*

Fang Liu<sup>1,2</sup>, Zhong L. Zhou<sup>2</sup>, Chun Y. Wang<sup>2</sup>, Yu H. Wang<sup>2</sup>, Xiao Y. Cai<sup>2</sup>, Xing X. Wang<sup>2</sup>, Kun B. Wang<sup>2\*</sup> & Zheng S. Zhang<sup>1\*</sup>

1 College of Agronomy and biotechnology, Southwest University, Chongqing 400715, People's Republic of China

2 State Key Laboratory of Cotton Biology, China / Cotton Research Institute, Chinese Academy of Agricultural Sciences, Anyang 455000, China

Correspondence authors: [wkbcri@163.com](mailto:wkbcri@163.com), [zhangzs@swu.edu.cn](mailto:zhangzs@swu.edu.cn)

**Abstract:** *G. tomentosum* and *G. darwinii* are wild allotetraploid cotton species, characterized with many excellent traits including finer fiber fineness, drought tolerance, Fusarium wilt and Verticillium wilt resistance. Based on construction of F<sub>2</sub> linkage groups of *G. hirsutum* × *G. tomentosum* and *G. hirsutum* × *G. darwinii*, two genetic linkage maps were compared. As a result we found a total of 7 inversion fragments on chr02, chr05, chr08, chr12, chr14, chr16 and chr25, 3 translocation fragments on chr05, chr14 and chr26. Further comparing the inversion and translocation fragments, we noticed four of seven markers orientation of *G. tomentosum* consistent with *G. hirsutum* or *G. raimondii*, one of seven inversion markers orientation of *G. darwinii* consistent with *G. hirsutum*; meanwhile one of three translocation marker orientation of *G. tomentosum* consistent with *G. raimondii*. The result indicate, compare *G. darwinii*, *G. tomentosum* has closer genetic relationship to *G. hirsutum*. This study will play an important role in understanding the genome structure of *G. tomentosum* and *G. darwinii*, and open the doors for further in-depth genome research such as fine mapping, tagging genes of interest from wild relatives and evolutionary study.

**Key words:** Cotton germplasm resource, Allotetraploid cotton, *Gossypium tomentosum*, *Gossypium darwinii*, linearity relationship

## Introduction

Cotton is a natural vegetable white fibrous agricultural product of great economic importance as raw material not only for textile industry but also for a wide variety of uses such as paper industry, home fixtures, medical supplies, chemicals and oil (Ensminger et al. 1990). Cotton (*Gossypium* spp) is one of the most expansively grown species around the globe in tropical and subtropical regions between 36° South latitude and 46° North latitude (Reller et al. 1997) of more than 80 countries ( Fryxell 1979; Smith 1995 ). *Gossypium* genus consists of 50 species (Fryxell 1992; Stewart 1995; Ma et al. 2008), which includes five allotetraploid species ( $2n=4x=52$ ;  $(AD)_1$  to  $(AD)_5$ ) and 45 diploid species ( $2n=2x=26$ , A through G and K) (Fryxell 1979; Stewart 1995; Brubaker et al. 1999; Zhang et al. 2005). The evolution of domesticated cotton involves various steps that can help scientists to have deep insight understanding of polyploidy role in diversification process and better investigation of unique traits, and it can further help geneticists to utilize germplasm from wild relatives. Allopolyploids provide a great tool to understand the evolutionary process of two diploid genomes which evolve simultaneously but different in genome size. Evolution and diversity studies of *Gossypium* genus provide the basic knowledge of morphological diversity of the genus and plant biology which can help in the better utilization of genetic resources (Wendel et al. 2009).

More than 30 genetic maps have already been published in cotton, and most of them are based on interspecific crosses of domesticated tetraploid species namely *G. hirsutum* and *G. barbadense* (Jiang et al. 1998; Zhang et al. 2002; Lacape et al. 2003; Nguyen et al. 2004; Rong et al. 2004; Guo et al. 2007; He et al. 2007; Lacape et al. 2009). The interspecific tetraploid genetic maps are useful in understanding genome structure and exploring the genetics basis of important agronomic characters and also provide the basis for finding new DNA markers for further high density maps (Guo et al. 2007; Zhang et al. 2008b; Yu et al. 2011).

Wild cotton has long been used as a genetic resource to introduce new traits to increase the potential of cotton cultivated species (Stewart 1995). *G. darwinii*, a wild allotetraploid species with  $(AD)_5$  genome, is closely related to *G. barbadense* but quite different from the cultivated *G. hirsutum*, and it has many excellent traits including finer fiber fineness, drought tolerance, Fusarium wilt and Verticillium wilt resistance. *G. tomentosum* is

endemic to Hawaiian Islands, and it has many unique agronomic traits such as insect-pest resistance, drought tolerance, salt tolerance, heat tolerance, nectarilessness and lint color. Many genetic studies like genomic and genetic structure/organization have been conducted on the two cultivated tetraploids, *G. hirsutum* and *G. barbadense*, however very little is known about genomic architecture, gene transfer or introgression of unique traits from other three tetraploids.

In the present study, the SSR genetic maps are developed from two F<sub>2</sub> populations, (*G. hirsutum* × *G. tomentosum*) (Chen et al. 2015) and (*G. hirsutum* × *G. darwinii*) (Kashif et al. 2015). The linearity relationship is compared between the two genetic maps, which will serve as an indispensable genomic resource for genome structure study, comparative genomic analysis, and fine mapping and map-based gene cloning of important traits.

### **Materials and methods**

Linkage map were reconstructed reference Chen et, al (2015). and Kashif et, al (2015). SSR markers uniform distribution on different chromosomes were selected and checked after downsizing. JoinMap 4.0 software was used for linkage analysis and map construction. Kosambi mapping function was used to convert recombination frequencies into map distances (centimorgan, cM). Mapchart 2.2 software was used to draw the genetic map.

Linkage map assignment was established by the common markers that were already anchored according to the previous publications (Wang et al. 2012; Li et al. 2014; Li et al. 2015; Zhang et al. 2015). Chromosomal nomenclature was used as mentioned by Guo et al. (2008), i.e. SSR loci anchored on chromosome (Chr.) 1 to 13 were designated to the A sub-genome (At), whereas loci confined to Chr. 14 to 26 were designated to the D sub-genome (Dt).

### **Results**

Comparing the linear relationship of two genetic linkage maps, we found that most SSR markers present collinearity between the two genetic linkage maps. Meanwhile, part of the non linear relationship appeared on the individual chromosomes between the genetic linkage maps of *G. hirsutum* × *G. tomentosum* and *G. hirsutum* × *G. darwinii*, which include 7 inversion fragments and 3 translocation fragments (Figure 1-13).

Seven inversions were found on seven different chromosomes between the genetic maps

of *G. tomentosum* and *G. darwinii* respectively: on chr02 of MON\_COT043-HAU2643-HAU0248-HAU2923, on chr05 of MON-DPL0384-TMB1586-BNL2448, on chr08 of CGR6748-HAU0709-BNL3257, on chr12 of MON\_CGR5158-NAU943-BNL1673, on chr14 of NAU3691-MON\_CGR5818-TMB1513, on chr16 of CM56-JESPR102-NAU3676-GH56 and on chr25 of MON\_CGR5665-NAU2388-BNL3103. Three translocations were found on 3 different chromosomes between the genetic maps of *G. tomentosum* and *G. darwinii* respectively: on chr05 of NAU2001-NAU1200-NAU3569 fragment parallel translocation with MUSB0592-NAU1127 fragment, on chr26 of MON-CGR5678-MON-DPL0391 fragment parallel translocation with MON-CGR6318- MON-CGR5802 fragment and on chr14 TMB0803-NAU2987 fragment parallel translocation with MON\_C2\_0079-HAU0551 fragment.

Referring to the published *G. Raimondii* genome sequencing data (D<sub>5</sub>, Wang et al. 2012) and the latest *G. hirsutum* genome sequencing data (AD<sub>1</sub>, Zhang et al. 2015), we compared the inversions and translocations on the genetic linkage map with the physical position and arrangement direction on the corresponding genome chromosome. The result showed that, among the inversions, the fragment orientation of *G. darwinii* on chr05 is consistent with the physical direction of *G. Raimondii* (D<sub>5</sub>) and *G. hirsutum* (AD<sub>1</sub>) genome sequence. The fragments orientation of *G. tomentosum* on chr08, chr12 and chr25 are consistent with the physical directions of *G. Raimondii* (D<sub>5</sub>). The fragments orientation of *G. tomentosum* on chr08, chr12 and chr16 are consistent with the physical directions of *G. hirsutum* (AD<sub>1</sub>). Among the translocation, just the fragment on chr26 corresponding physical position was found on *G. Raimondii* (D<sub>5</sub>), with the same direction as *G. tomentosum*.

## **Discussion**

### **Reciprocal translocations and Inversion**

In the study of *Gossypium* genetic linkage map construction, Researchers have repeatedly found in the reciprocal translocation between homologous chromosomes (Rong et al. 2004; He et al. 2007; Wang et al. 2006; John et al. 2012). Very few study focus on different cotton linear relationship between homologous chromosomes. In this study, use the same variety of *G. hirsutum*, we constructed allotetraploid genetic linkage maps of *G. hirsutum* × *G.*

*tomentosum* and *G. hirsutum* × *G. darwinii*. Through linear relationship analysis on homologous position, there are seven inversions (corresponding on chr02, chr05, chr08, chr12, chr14, chr16 and chr25) and three translocations (corresponding on chr05, chr14 and chr26) were found separately on the two genetic linkage maps. Compare the inversions and translocations location with the physical position of *G. hirsutum*, we found the same inversion phenomena appeared on the corresponding position of chr02, chr05, chr12 and chr14 in *G. hirsutum*, the same with translocation on the corresponding position of chr05 in *G. hirsutum*. For the first time we discovered inversion on the homologous chromosome fragments of chr02 and chr14, and for the first time we found out the translocations on chr05 and chr14. In this study we also found inversion and translocation appeared at the same fragment, which indicate it is genetic active region. Further research need to confirm inversion occur before or after translocation.

### **Genetic mapping coupled with physical alignment of genomic**

A lot of research has been done on the origin and evolution of allotetraploid cotton by previous researchers. Fryxell (1992) and Wendel (2009) support the idea that *G. tomentosum* and *G. hirsutum* has a closer relationship, however *G. darwinii* and *G. barbadense* has a closer relationship. Rarely research has done to the linear relationship of chromosome genetic linkage between *G. tomentosum* and *G. darwinii*. With the cotton genome sequencing successively completed, relationship between genetic linkage map and physical map study became feasible and convenient. Using F<sub>2</sub> population derived from interspecific crosses, we constructed two linkage maps of allotetraploid cotton *G. hirsutum* × *G. tomentosum* and *G. hirsutum* × *G. darwinii* separately. Referring to the latest published *G. hirsutum* genome sequencing data (Zhang et al. 2015), linear relationship of the two wild allotetraploid cotton genetic linkage maps was compared based on genome-wide SSR markers. The results show that the genetic linkage maps of *G. hirsutum* × *G. tomentosum* and *G. hirsutum* × *G. darwinii* had good linear relationship. Meanwhile, part of the non linear relationship appeared on the individual chromosomes between the two genetic linkage maps, including 7 inversion fragments and 3 translocation fragments. Compared with the sequence data of *G. hirsutum* (Zhang et al. 2015), we noticed among the 7 inversion fragments, just one markers orientation (chr05) which comes from *G. darwinii* has the same direction of physical position arrangement of

*G. hirsutum*, while four markers orientation (chr08, chr12, chr16 and chr25) which comes from *G. tomentosum* has the same direction of physical position arrangement of *G. hirsutum* or *G. raimondii*. Among the 3 translocation fragments, one marker orientation (chr26) which comes from *G. tomentosum* has the same direction of physical position arrangement of *G. raimondii*. The result indicate compare *G. darwinii*, *G. tomentosum* has closer genetic relationship to *G. hirsutum*.

**Acknowledgements** This work was co-supported by the National Science and Technology Support Plan of China (2013AA102601) and National Key Technology Support Program of China (2013BAD01B03).

### **Competing interests**

The authors declare that they have no competing interests.

### **Authors' contributions**

Kunbo Wang, Zhengsheng Zhang and Fang Liu participated in the experiments design. Fang Liu carried out most of the experiments and all computational analyses. Zhong-Li Zhou, Chun-Ying Wang and Yu-Hong Wang participated in part of groups construction, Xiao-Yan Cai, Xing-Xing Wang participated in part of mapping experiments. Fang Liu drafted the manuscript. Baohong Zhang, Kunbo Wang and Zhengsheng Zhang revised the manuscript. All authors read and approved the final manuscript.

### **References**

1. Ensminger, M. E., J. E. Oldfield, W. W. Heinemann, 1990 Excerpts with reference to cottonseed components. In: Ensminger Publishing Company, USA.
2. Reller, A., J. Gerstenberg, 1997 White gold, where do you go? Status quo and prospects of cotton usage. GAIA. 1, 35 - 51.
3. Fryxell, P. A., 1979 The natural history of the cotton tribe. Texas A&M University.
4. Smith, C. W., 1995 Cotton (*Gossypium hirsutum* L.). Chapter 6. In: Crop Production: Evolution, History, and Technology. John Wiley and Sons, Inc., New York, 287-349.
5. Fryxell, P. A., 1992 A revised taxonomic interpretation of *Gossypium* L. Malvaceae. Rheedea, 2, 108-165.
6. Stewart, J. M., 1995 Potential for crop improvement with exotic germplasm and genetic engineering.

In: G. Constable and N. Forrester (eds.). Challenging the future. Proceedings of the World Cotton Research Conference-1. CSIRO, Melbourne, 313-327.

7. Ma, X., B. L. Zhou, Y. Lu, W. Z. Guo, T. Z. Zhang, 2008 Simple sequence repeat genetic linkage maps of A-genome diploid cotton (*Gossypium arboreum*). Journal of Integrative Plant Biology, 50, 491-502.

8. Brubaker, C. L., F. M. Bourland, J. F. Wendel, 1999 The origin and domestication of cotton. In W. Smith and T. Cothren (eds.) Cotton: Origin, history, technology and production. John Wiley & Sons, Inc, 3-31.

9. Zhang, J. F., Y. Z. Lu, S. X. Yu, 2005 Cleaved AFLP (cAFLP), a modified amplified fragment length polymorphism analysis for cotton. Theoretical and Applied Genetics. 111, 1385-1395.

10. Wendel, J. F., C. Brubaker, I. Alvarez, R. Cronn, J. M. Stewart, 2009 Plant Genetics and Genomics: Crops and Models, Genetics and Genomics of Cotton: Evolution and Natural History of the Cotton Genus. Springer Science Business Media, LLC, 3, 3-22.

11. Jiang, C. X., R. J. Wright, K. M. El-Zik, A. H. Paterson, 1998 Polyploid formation created unique avenues for response to selection in *Gossypium* (cotton). Proc. Natl. Acad. Sci., 95, 4419-4424.

12. Zhang, J., W. Z. Guo, T. Z. Zhang, 2002 Molecular linkage map of allotetraploid cotton (*Gossypium hirsutum* L. x *Gossypium barbadense* L.) with a haploid population. Theor. Appl. Genet. 105:1166-1174.

13. Lacape, J. M., T. B. Nguyen, S. Thibivilliers, B. Bojinov, B. Courtois, R. G. Cantrell, B. Burr, B. Hau, 2003 A combined RFLP-SSR-AFLP map of tetraploid cotton based on a *Gossypium hirsutum* x *Gossypium barbadense* backcross population. Genome. 46, 612-626.

14. Nguyen, T. B., M. Giband, P. Brottier, A. M. Risterucci, J. M. Lacape, 2004 Wide coverage of the tetraploid cotton genome using newly developed microsatellite markers. Theor Appl Genet. 109: 167-175.

15. Rong, J. K., C. Abbey, J. E. Bowers, C. L. Brubaker, C. Chang, P. W. Chee, T. A. Delmonte, X. Ding, J. J. Garza, B. S. Marler, C. Park, G. J. Pierce, K. M. Rainey, V. Rastogi, K. Schulze, N. L. Tronlinde, J. F. Wendel, T. A. Wilkins, R. A. Wing, R. J. Wright, X. Zhao, L. Zhu, A. H. Paterson, 2004 A 3347-locus genetic recombination map of sequence-tagged sites reveals features of genome organization, transmission and evolution of cotton (*Gossypium*). Genetics. 166. 389-417.

16. Guo, W. Z., C. P. Cai, C. B. Wang, Z. G. Han, X. L. Song, K. Wang, X. Niu, C. Wang, K. Lu, B. Shi, T. Z. Zhang, 2007 A microsatellite-based, gene-rich linkage map reveals genome structure, function and evolution in *Gossypium*. Genetics. 176:527-541.

17. He, D. H., Z. X. Lin, X. L. Zhang, Y. C. Nie, X. P. Guo, Y. X. Zhang, W. Li, 2007 QTL mapping for economic traits based on a dense genetic map of cotton with PCR-based markers using the interspecific cross of *Gossypium hirsutum* x *Gossypium barbadense*. Euphytica. 153, 181-197.

18. Lacape, J. M., J. Jacobs, T. Arioli, R. Derijcker, N. Forestier-Chiron, D. Llewellyn, J. Jean, E.

- Thomas, C. Viot, 2009 A new interspecific, *Gossypium hirsutum* x *G. barbadense*, RIL population: towards a unified consensus linkage map of tetraploid cotton. *Theor. Appl. Genet.* 119: 281 - 292.
19. Zhang, Y. X., Z. X. Lin, Q. Z. Xia, M. G. Zhang, X. L. Zhang, 2008b Characteristics and analysis of simple sequence repeats in the cotton genome based on a linkage map constructed from a BC<sub>1</sub> population between *Gossypium hirsutum* and *G. barbadense*. *Genome.* 51, 534-546.
20. Yu, Y. D. J. Yuan, S. G. Liang, X. M. Li, X. Q. Wang, Z. X. Lin, X. L. Zhang, 2011 Genome structure of cotton revealed by a genome-wide SSR genetic map constructed from a BC<sub>1</sub> population between *Gossypium hirsutum* and *G. barbadense*. *BMC Genomics.* 12, 15.
21. Stewart, J. M., 1995 Potential for crop improvement with exotic germplasm and genetic engineering. In: G. Constable, Forrester N W. Challenging the future. Proceedings of the World Cotton Research Conference-1 Melbourne, Australia, CSIRO. 313-327.
22. Chen, H. D., R. K. Kashif, Z. L. Zhou, I. Kashif, X. X. Wang, X. Y. Cai, Wang, C. Y. Wang, Y. H. Wang, Y. Q. Li, F. Liu, K. B. Wang, 2015 A High-density SSR genetic map constructed from a F<sub>2</sub> population of *Gossypium hirsutum* and *Gossypium darwinii*. *Gene*: <http://dx.doi.org/10.1016/j.gene.2015.08.022>
23. Kashif, R. K., H. D. Chen, Z. L. Zhou, X. X. Wang, X. Y. Cai, Wang, C. Y. Wang, Y. H. Wang, Y. Q. Li, F. Liu, K. B. Wang, 2015 Genome wide SSR high density genetic map construction from an interspecific cross of *G. hirsutum* × *G. tomentosum*. *Genetic Molecular Research.* accepted.
24. Wang, K. B., Z. W. Wang, F. G. Li, W. W. Ye, J. Y. Wang, G. L. Song, Z. Yue, L. Cong, H. H. Shang, S. L. Zhu, C. S. Zou, Q. Li, Y. L. Yuan, C. R. Lu, H. L. Wei, C. Y. Gou, Z. Q. Zheng, Y. Yin, X. Y. Zhang, K. Liu, B. Wang, C. Song, N. Shi, J. K. Russell, G. P. Richard, Z. Y. John, Y. X. Zhu, J. Wang, & S. X. Yu, 2012 The draft genome of a diploid cotton *Gossypium raimondii*. *Nature Genetics.* 44, 1098-1103; doi: 10.1038/ng. 2371.
25. Li, F. G., K. B. Wang, Y. L. Yuan, G. L. Song, C. S. Zou, H. H. Shang, W. W. Ye, X. Y. Zhang, H. L. Wei, Z. F. Li, & S. X. Yu, 2014 Genome sequence of the cultivated cotton *Gossypium arboreum*. *Nature Genetics.* doi: 10.1038/ng.2987.
26. Li, F. G., G. Y. Fan, C. R. Lu, G. H. Xiao, C. S. Zou, J. K. Russell, Z. Y. Ma, H. H. Shang, X. F. Ma, J. Y. Wu, X. M. Liang, G. Huang, G. P. Richard, K. Liu, W. H. Yang, W. B. Chen, X. M. Du, C. C. Shi, Y. L. Yuan, W. W. Ye, X. Liu, X. Y. Zhang, W. Q. Liu, H. L. Wei, S. J. Wei, G. D. Huang, X. L. Zhang, S. J. Zhu, H. Zhang, F. M. Sun, X. F. Wang, J. Liang, J. H. Wang, Q. He, L. H. Huang, J. Wang, J. J. Cui, G. L. Song, K. B. Wang, X. Xu, Z. Y. John Y. X. Zhu, & S. X. Yu, 2015 Genome sequence of cultivated Upland cotton (*Gossypium hirsutum* TM-1) provides insights into genome evolution. *Nature biotechnology.* doi: 10.1038/nbt.3208.
27. Zhang, T. Z., Y. Hu, W. K. Jiang, L. Fang, X. Y. Guan, J. D. Chen, J. B. Zhang, A. S. Christopher, E. S. Brian, M. S. David, M. H. Amanda, Q. Wan, B. L. Liu, C. X. Liu, S. Wang, M. Q. Pan, Y. K. Wang, D. W. Wang, W. X. Ye, L. J. Chang, W. P. Zhang, Q. X. Song, C. K. Ryan, X. Y. Chen, D.

Elizabeth, J. L. Danny, G. P. Daniel, T. Peggy, C. J. Don, Q. Wang, X. Y. Xu, H. Zhang, H. T. Wu, L. Zhou, G. F. Mei, S. Q. Chen, Y. Tian, D. Xiang, X. H. Li, J. Ding, Q. Y. Zuo, L. Tao, Y. C. Liu, J. Li, Y. Lin, Y. Y. Hui, Z. S. Cao, C. P. Cai, X. F. Zhu, Z. Jiang, B. L. Zhou, W. Z. Guo, R. Q. Li, & Z. J. Chen, 2015 Sequencing of allotetraploid cotton (*Gossypium hirsutum* L. acc. TM-1) provides a resource for fiber improvement. *Nature biotechnology*. doi:10.1038/nbt.3208.

28. Guo, W. Z., C. P. Cai, C. B. Wang, L. Zhao, L. Wang, T. Z. Zhang, 2008 A preliminary analysis of genome structure and composition in *Gossypium hirsutum*. *BMC Genomics*. **9**, 314.

29. Wang, K., X. L. Song, Z. G. Han, W. Z. Guo, J. Z. Yu, et al. 2006 Complete assignment of the chromosomes of *Gossypium hirsutum* L. by translocation and fluorescence in situ hybridization mapping. *Theor. Appl. Genet*, 113, 73–80.

30. John, Z. Y., J. K. Russell, D. F. David, C. Jaemin, V. D. Allen, U. Mauricio, M. H. Steven, E. P. Alan, M. S. David, N. J. Johnie, S. Sukumar, P. K. Siva, R. S. Manali, V. H. William, & G. P. Richard, 2012 A High-Density Simple Sequence Repeat and Single Nucleotide Polymorphism Genetic Map of the Tetraploid Cotton Genome. *Genes Genomes Genetics*. doi: 10.1534/g3.111.001552.

Figure 1 to 13: genetic map and Collinearity comparing of *G. Tomentosum* and *G. Darwinii*.

Gd: *Gossypium Darwinii*



GT: *Gossypium Tomentosum*



Linkage group	Markers	Location on <i>G. darwinii</i> (genetic position, cM)	Location on <i>G. tomentosum</i> (genetic position)	Corresponding location on <i>G. raimondii</i> (physical position, Mb)	Corresponding location on <i>G. hirsutum</i> (physical position, Mb)	Events
chr02	MON_COT043-HAU2643-HAU0248-HAU2923	77.7-78.8-78.9-79.4	92.5-91.3-89.7-87.8			inversion
chr05	MON_DPL0384-TMB1586-BNL2448	144.1-146.1-147.8	86.3-84-82.6	21.4-21.6	37.2-58.2	inversion
chr08	MON_CGR6748-HAU1632-BNL3257	66.4-70.3-71.7	71.1-68.1-64.7	6.5-31.3-46.1	85.6-58.9	inversion
chr12	MON_CGR5158-NAU943-BNL1673	107.3-108-111	111.2-109.2-107.1	44.2-42.6	93.5-41.7	inversion
chr16	CM56-JESPR102-NAU3676-GH56	34-34.7-36.6-39	70.6-69.4-63.4-58.4		47.1-39.5	inversion
chr25	MON_CGR5665-NAU2388-BNL3103	18.6-20.9-23.3	29.8-29.4-28.6	42.9-40.6		inversion
chr05	NAU2001-NAU1200-NAU3569-MUSB0592-NAU1127	195.1-199.5-202.9-205.7-207	132.7-133.4-133.7-129-132.5			translocation
chr26	MON_CGR5678-MON_DPL0391-MON_CGR6318-MON_CGR5802	34.9-35.3-37.4-38.2	31.5-33-29.6-30	17.1-31.1-10.2-10.8		translocation
chr14	TMB0803-NAU2987-MON_C2_0079-HAU0551-NAU3691-MON_CGR5818-TMB1513	51.9-52.1-53.9-54.8-55-55.3-56.6	68-69.5-65.9-66.2-64.5-62.3-60.4			inversion & translocation


Table 1 Location comparing of genetic linkage fraction and genome physical position

chr01Gd(A1)

chr01Gt(A1)








chr04Gt(A4)

chr22Gd(D4)

chr22Gt(D4)

chr04Gd(A4)




## chr05Gd(A5)


## ch05Gt(A5)

## chr19Gd(D5)


## chr19Gt(D5)




chr06Gd(A6)




chr06Gt(A6)



chr25Gd(D6)




chr25Gt(D6)




## chr07Gt(A7)

## chr07Gd(A7)



## chr16Gd(D7)




## chr08Gd(A8)


## chr08Gt (A8)

## chr24Gd(D8)


## chr24Gt (D8)




chr09Gd(A9)




chr09Gt(A9)



chr23Gd(D9)



chr23Gt(D9)




## chr10Gd(A10)

## chr10Gt(A10)

## chr20Gd(D10)

## chr20Gt(D10)




## chr11Gt(A11)

## chr21Gd(D11)

## chr21Gt(D11)

## chr11Gd(A11)




## chr12Gd(A12)

## chr12Gt(A12)

## chr26Gd(D12)

## chr26Gt(D12)



## chr13Gd(A13)

0.0  
0.8  
6.8  
21.6  
22.8  
26.7  
29.2  
44.0  
47.3  
47.5  
47.8  
48.3  
49.6  
51.0  
51.4  
56.2  
57.3  
58.1  
58.7  
58.8  
59.8  
65.5  
68.9  
70.0  
70.3  
70.4  
70.9  
75.9  
76.9  
80.1  
82.8  
83.1  
85.9  
87.2  
88.2  
88.5  
88.9  
89.0  
89.8  
90.8  
91.0  
91.1  
91.2  
91.3  
91.4  
91.9  
92.0  
92.3  
92.6  
93.0  
93.5  
94.6  
95.1  
97.0  
99.6  
100.1  
101.7  
102.7  
105.2  
105.3  
106.1NAU3991  
HAU2977  
MON\_SHIN1512  
HAU3159  
HAU2752  
HAU0696  
HAU0534  
NAU2300  
MON\_CGR5554 MON\_CGR5331  
MON\_SHIN1462  
NAU5074  
HAU1997  
MUCS235  
HAU1182  
NAU944  
HAU0539  
HAU2857  
NAU3468  
NAU3017  
MON\_SHIN0412  
GH60a  
MON\_DPL0083  
GH592  
GH34  
MON\_CGR5827  
MON\_CER0005  
MON\_CGR5242  
MON\_COT009  
DPL0635  
MON\_DPL0535  
NAU1201  
MON\_DC40094  
MON\_DPL0553  
MON\_DC20120  
MON\_C2-0026C  
STV080  
NAU2458  
NAU3570  
GH331  
MON\_DC40441  
HAU3291  
MON\_CGR6359  
MON\_CGR6732  
MON\_SHIN1377  
SWU21985  
SWU10810  
MON\_DPL0572 HAU1813  
MON\_C2-0002Bb  
TMB0482  
CIR300  
MON\_DPL0754  
MON\_DC20119  
MON\_DC20047  
MON\_SHIN1598  
ICRC0114a  
ICRC0819  
HAU0429  
BNL1495  
NAU3948a  
BNL1421

## chr13Gt(A13)

0.0  
2.1  
2.9  
4.8  
5.9  
7.2  
8.6  
10.1  
19.2  
20.4  
22.4  
23.0  
24.1  
25.4  
27.6  
29.2  
30.6  
31.3  
31.7  
32.1  
33.7  
34.5  
35.2  
35.7  
36.5  
36.9  
37.8  
39.0  
40.3  
41.2  
41.6  
41.9  
42.4  
42.7  
43.1  
43.5  
44.5  
45.1  
46.7  
48.6  
50.5  
50.7  
51.1  
51.8  
52.4  
53.7  
58.5MON\_SHIN-1462  
MON\_CGR5021(a)  
NAU5074(a)  
HAU2331  
HAU1182  
HAU0539  
Gh497  
NAU3398  
MON\_DPL0083  
MON\_CGR5827  
MUSB0551  
Gh34  
MON\_CER0005  
TMB0856  
MON\_DPL0330  
BNL4029  
NAU5509(a)  
MON\_CGR6436  
NAU3307  
NBRI-0161  
Gh163  
MON\_DC20120  
MON\_DC30178  
NAU3570  
NAU1023(a)  
Gh215  
TMB1216  
MON\_DPL0406  
NBRI-0188  
TMB0403  
BNL2449  
CIR201  
MON\_DPL0274  
JESPR175  
CIR144  
MUSB0621  
BNL2652  
NAU1211  
MGHES-72  
MON\_SHIN-1598  
MUSS349  
NAU4102  
CIR342  
TMB0635  
BNL1421  
NAU4103(b)  
NAU3948(b)

## chr18Gd(D13)

0.0  
1.2  
1.9  
3.6  
10.5  
18.9  
20.0  
20.1  
24.0  
24.6  
24.8  
25.3  
28.1  
40.3  
40.8  
41.3  
48.1  
49.9  
51.4  
52.5  
53.1  
53.2  
53.3  
53.8  
56.9  
59.7  
60.1  
60.3  
60.8  
61.5  
67.7  
69.0  
69.2  
69.8  
70.3  
70.4  
76.1  
77.1  
78.7  
78.9  
80.0  
80.6  
80.7  
84.0  
85.1  
85.8  
86.3  
86.6  
87.7  
89.1  
89.6  
90.0  
90.1  
90.2  
91.4  
91.6  
96.6  
97.2  
98.2  
98.4  
98.6  
99.1  
106.3  
107.7  
110.8  
111.5  
116.3  
118.1  
121.6  
121.8  
122.1  
137.1  
138.4  
140.4  
140.5  
143.5NAU2980  
NAU3378  
HAU0933  
MUSB1135a  
JESPR246  
NAU3827  
HAU2276  
NAU3447 HAU1922  
NAU3080  
NAU5262  
NAU4861  
NAU3861  
HAU3099  
HAU1212  
CCRI1318  
MON\_C2-0104  
BNL2667  
CCRI283  
HAU1909  
MON\_SHIN1403  
MUS803  
CCRI294  
MUSB1166  
MON\_CGR5446  
MON\_DPL0424  
GH60b  
HAU1036  
MUCS405  
MUCS135  
MON\_CER0145  
MON\_SHIN1346  
NAU2443  
STV010  
MON\_DPL0420  
MON\_DPL0660  
MON\_DPL0741  
MON\_CGR6754  
MON\_COT114  
NAU3211  
MON\_CGR6231  
ICRC0465  
MON\_CGR5856  
HAU3061  
BNL1721  
NAU2138  
HAU0231  
MUS544  
BNL1079  
MON\_SHIN1530  
MON\_C2-0002Ba  
HAU2675  
MON\_CGR6787  
MON\_CGR5564  
BNL2652b  
GH653  
GH160  
BNL3558  
HAU0712  
STV097b  
NAU3203b  
HAU0693  
NAU3948b  
ICRC0066  
HAU1927  
HAU1880  
MON\_CGR5390  
MON\_DPL0286b  
BNL2544  
NAU3943  
BNL193  
HAU0596  
HAU2631b  
HAU2901a  
HAU0250b  
BNL243  
NAU3685

## chr18Gt(D13)

0.0  
3.1  
5.2  
6.9  
7.7  
11.2  
15.9  
23.7  
25.2  
32.2  
32.9  
34.2  
35.7  
37.0  
40.6  
42.2  
42.7  
43.0  
43.1  
43.4  
43.9  
44.3  
45.5  
46.6  
48.4  
49.7  
50.3  
51.1  
52.3  
53.2  
53.9  
54.6  
55.7  
56.8  
57.4  
58.0  
58.5  
59.1  
59.3  
59.6  
60.3  
60.9  
61.1  
61.6  
61.8  
62.1  
62.5  
62.7  
63.0  
63.2  
63.4  
63.8  
64.3  
64.7  
65.0  
65.3  
66.0  
66.5  
67.7  
72.3  
73.4  
80.3  
80.5  
81.4  
82.7  
84.7  
88.7HAU1355(b)  
NAU3827  
NAU3223  
MON\_CGR6750(a)  
MON\_DPL0375(a)  
NAU3861(a)  
NAU3130  
NAU3043  
JESPR178  
HAU2332(a)  
HAU1212  
NAU5074(b)  
MON\_CGR5021(b)  
CIR221  
MON\_SHIN-1403  
HAU1381  
HAU1909  
MUS803  
HAU3304  
NAU3589  
MON\_DC20165  
NAU3397  
MON\_DPL0424  
JESPR56(a)  
HAU1036  
SWU22280  
NAU5248  
MUSB0193  
Gh92  
NAU2647  
MON\_SHIN-1346  
NAU2443  
STV010  
TMB0068  
MON\_CGR6522  
Gh678  
MUSB1160  
MON\_DC40289  
BNL3445  
HAU3220  
NAU5509(b)  
DPL0049  
NAU5186  
MON\_SHIN-1530  
BNL1721  
BNL1079  
CM63  
Gh326  
NAU3816(b)  
SWU21957  
NBRI-0189  
NAU1023(b)  
Gh443(b)  
Gh160  
MON\_CGR5564  
Gh653  
SWU21879  
Gh153(a)  
NAU4103(a)  
NAU3948(a)  
BNL3558  
MON\_DPL0894  
NAU2697  
HAU1880  
MON\_CGR5390  
NBRI-0123  
MON\_CER0122

bioRxiv preprint doi: <https://doi.org/10.1101/031104>; this version posted November 10, 2015. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.