

Exploring miRNAs as the key to understand symptoms induced by ZIKA virus infection through a collaborative database.

Victor Satler Pylro^{1*#}; Francislton Silva de Oliveira^{1*}, Daniel Kumazawa Morais¹, Sara Cuadros Orellana¹, Fabiano Sviatopolk-Mirsky Pais¹, Julliane Dutra Medeiros¹, Juliana Assis Geraldo¹, Jack Gilbert^{2,3,4}, Angela Volpini¹, Gabriel da Rocha Fernandes^{1#}

¹Genomics and Computational Biology Group, René Rachou Research Center (CPqRR-FIOCRUZ), Belo Horizonte, MG, 30190-002, Brazil.

²Department of Ecology and Evolution, The University of Chicago, Chicago, Illinois, USA

³Argonne National Laboratory, Institute for Genomic and Systems Biology, Argonne, Illinois, USA

⁴Department of Surgery, The University of Chicago, Chicago, Illinois, USA

*These authors contributed equally to this work.

#Correspondence: gabriel.fernandes@cpqrr.fiocruz.br and victor.pylro@cpqrr.fiocruz.br

Zika virus (ZIKV) is an emerging mosquito-borne flavivirus, first isolated in 1947 from the serum of a pyrexial rhesus monkey caged in the Zika Forest (Uganda/Africa)¹. In 2007 ZIKV was reported to of been responsible for an outbreak of relatively mild disease, characterized by rash, arthralgia, and conjunctivitis on Yap Island, in the western Pacific Ocean². In the past year, ZIKV has been circulating in the Americas, probably introduced through Easter Island (Chile), by French Polynesians³. In early 2015, a new outbreak was recognized in northeast Brazil⁴, where concerns over its possible links with infant microcephaly have been discussed^{5,6}. Providing a definitive link between ZIKV infection and birth defects is still a big challenge⁷. MicroRNAs (miRNAs), are small noncoding RNAs that regulating post-transcriptional gene expression by translational repression, and play important roles in viral pathogenesis⁸ and brain development⁹. It is estimated that more than 60% of human protein-coding genes contain at least one conserved miRNA-binding site¹⁰. The potential for flavivirus-mediated miRNA signaling dysfunction in brain-tissue develop provides a compelling mechanism underlying perceived linked between ZIKV and microcephaly. Here, we provide strong evidences toward to understand the mechanism in which miRNAs can be linked to the “congenital Zika syndrome” symptoms. Moreover, following World Health Organization (WHO) recommendations¹¹, we have assembled a database that could help target mechanistic investigations of this possible relationship between ZIKV symptoms and miRNA mediated human gene expression, helping to foster potential targets for therapy.

Two hypotheses of how miRNAs can influence ZIKV/human-host interaction were raised to support our database. First, viruses can transcribe miRNAs that enable them to benefit from cellular and viral gene expression (e.g. Herpesvirus, Polyomavirus, Ascovirus, Baculovirus, Iridovirus, Adenovirus families)^{12,13}. RNA retrovirus miRNAs are encoded

through a RNA polymerase III (pol III) promoter, instead of being transcribed by pol II. Virus-encoded miRNAs can play a role in persistent infections through subtle modulation of gene expression, leading to prevention of host cell death, evasion of the host immune system and regulation of the latent-lytic switch¹⁴. Second, retrovirus genomes can directly interact with cellular miRNAs to enhance viral replication potential through a mechanism of innate antiviral immunity¹². In this sense, by recruiting/exploiting cellular miRNAs, an RNA virus can disturb normal host gene expression regulation, which can trigger different molecular diseases. To enable testing of these hypotheses, the *ZIKV collaborative database* (ZIKV-CDB) was assembled to allow for: (i) searching for predicted ZIKV miRNAs mimicking human miRNAs [searching criteria includes: “Gene name”, “Gene Symbol” or “Ensembl ID”] (hypothesis 1); and (ii) searching for human miRNAs with possible binding-sites to the ZIKV genomes (hypothesis 2).

The ZIKV-CDB comprises miRNAs predicted using HHMMiR¹⁵ for all complete ZIKV genomes currently available at the GenBank (February, 2016 - <http://www.ncbi.nlm.nih.gov>). Hairpin prediction was also performed for all *de novo* miRNAs using previously predicted RNA secondary structure¹⁶, and mature miRNAs were delineated with PHDcleav¹⁷. Finally, to test hypothesis 1, potential human genome (Ensembl GRCh37) target sites for the predicted ZIKV miRNAs were detected with miRanda¹⁸ using default parameters (minimum score=140; minimum energy=1). To test hypothesis 2, we retrieved all mature human miRNA sequences from miRBase Sequence Database (Release 21 – <http://www.mirbase.org>), and mapped these against the available ZIKV genomes using miRanda¹⁸ with default parameters, to keep only those miRNAs with a minimum complementarity to ZIKV genomes [at least with complementarity to the miRNA seed region (6-10 nt) of the miRNA]. The ZIKV-CDB can be accessed through a web interface at <http://zikadb.cpqrr.fiocruz.br>.

The ZIKV-CDB has enabled us to predict miRNAs from ZIKV that have high-confidence targets in the human genome (Table 1), as well as cellular (human) miRNAs able to interact with ZIKV genomes (Table 2). This knowledge base should help to delineate targets that have the potential to affect the expression of genes associated with microcephaly and other neurodevelopmental syndromes. Some of these targets include, the peroxisomal biogenesis factor 26 gene (PEX26), the fibroblast growth factor 2 (FGF2), the SET binding factor 1 (SBF1), the hook microtubule-tethering protein 3 (Hook3) and the pleckstrin homology and RhoGEF domain containing G4 (PLEKHG4) (Table 1). PLEKHG4 polymorphisms have been related to spinocerebellar ataxia¹⁹, a progressive-degenerative genetic disease. Also, Hook3 has been reported to interact with Pericentriolar Material 1 (PCM1) during brain development, and an imbalance in the Hook3-PCM1 interaction can cause premature depletion of the neural progenitor pool in the developing neocortex²⁰.

Finally, defects in the PEX26 gene can lead to a failure of protein import into the peroxisomal membrane or matrix, being the cause of several neuronal disorders, including Zellweger syndrome (ZWS), and neonatal adrenoleukodystrophy (NALD)^{21, 22}.

Interestingly, several human miRNAs known to exert an influence on the expression of genes with a functional role in neuronal development, were found to have sequence complementarity to regions in the ZIKV genomes (Table 2). The human hsa-miR-34a miRNA has, as one of its targets, the Cyclin-Dependent Kinase 6 (CDK6) gene, which is technically able to interact with the ZIKV genome. CDK6 is associated with the centrosome during mitosis, controlling the cell cycle division phases in neuron production²³. Mutation in CDK6 can lead to a deficient centrosomes division, which in turns can cause autosomal recessive primary microcephaly (MCHP)²³. There are seven well-know genes encoding centrosomal proteins that are involved in the autosomal recessive primary microcephaly (MCPH)²⁴, including the CDK5 Regulatory Subunit Associated Protein 2 (Cdk5rap2 or MCHP3) gene. We found a possible binding-site to the hsa-mir-324-3p, a cellular miRNA targeting Cdk5rap2 gene, in the ZIKV genomes. Equally, we found that ZIKV genomic regions can bind the hsa-mir-615-3p and hsa-miR-193b-3p human miRNAs, which target the WD Repeat Domain 62 (WDR62 or MCHP2), also related to MCHP when mutated. Remarkably, a hsa-mir-21-5p miRNA complementary site was found on the genomes of ZIKV from Brazil, Haiti, Martinique and French Polynesia, but not in those from Africa. This miRNA targets the MCHP4 gene, also linked to microcephaly cases. This geographic and hence historical accumulation of genomic differences may explain the recent rise of microcephaly, and also corroborates the predicted pathway of transmission from Africa, through Oceania, and into Central and South America.

To further support this, phylogenetic analysis was performed for all complete ZIKV genomes (Figure 1), which identified a cluster of strains isolated in Americas and Oceania (“recent” strains), and another with the African strains (“ancient” strains). A third group, including other two Flavivirus’ close related to ZIKV, was added as out-group. These difference were also supported by phylogenetic analysis of only the predicted miRNAs encoded by each ZIKV strain. Nine predicted miRNA types were identified with types 1-4 restricted to recent strains and 5-8 restricted to ancient strains. Type 9 was only found in strains from Central Africa Republic (Figure 2). The predicted miRNAs could target 14,745 human genes; 9,106 were specific to the ‘ancient’ strains, 2,840 specific to the ‘recent’ strains, and 2,789 were shared.

Recently, ZIKV was isolated from the brain tissue of a fetus diagnosed with microcephaly²⁵. However, the mechanism by which ZIKV alters neurophysiological development remains unknown, preventing therapeutic intervention. Our results provide new insights associated with the potential influence of miRNAs on the expression of human-

genes associated with the symptoms of “congenital ZIKA syndrome”. The ZIKV-CDB provides a crucial knowledge base to mitigate the impacts of this emerging health problem²⁶. ZIKV-CDB is an open-source and collaboration-based forum for sharing and identification of potential mechanistic and therapeutic targets. These data will guide experimental investigation to elucidate any association between ZIKV infection and neurobiological development in infants. The ZIKV-CDB will be constantly maintained and curated by the staff of the Genomics and Computational Biology Group, from FIOCRUZ/CPqRR (<http://www.cpqrr.fiocruz.br>).

References

1. Dick, G. W. A., Kitchen, S. F., Haddow, A. J. Zika Virus (I). Isolations and serological specificity. *Trans. R. Soc. Trop. Med. Hyg.* **46**, 509-520 (1952).
2. Lanciotti, R. S., et al. Genetic and serologic properties of Zika virus associated with an epidemic, Yap State, Micronesia, 2007. *Emerg. Infect. Dis.* **14**, 1232-1239 (2008).
3. Musso, D. Zika virus transmission from French Polynesia to Brazil. *Emerg. Infect. Dis.* **21**, 1897 (2015).
4. Campos, G. S., Bandeira, A. C. & Sardi, S. I. Zika Virus Outbreak, Bahia, Brazil. *Emerg. Infect. Dis.* **21**, 1885-1886 (2015).
5. Schuler-Faccini, L., et al. Possible association between Zika virus infection and microcephaly — Brazil, 2015. *Morb. Mortal. Wkly. Rep.* **65**, 59-62 (2016).
6. Butler, D. Zika virus: Brazil's surge in small-headed babies questioned by report. *Nature* **530**, 13–14 (2016).
7. Hayden, E. C. Proving Zika link to birth defects poses huge challenge. *Nature* **530**, 142–143 (2016).
8. Gragnani, L. et al., MicroRNA expression in hepatitis C virus-related malignancies: A brief review. *World J. Gastroenterol.* **21**, 8562-8568 (2015).
9. Im, H. & Paul, J. MicroRNAs in neuronal function and dysfunction Kenny. *Trends Neurosci.* **35**, 325-334 (2012).
10. Friedman, R. C., Farh, K. K., Burge, C. B. & Bartel, D. P. Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res.* **19**, 92–105 (2009).
11. Dye, C., Bartolomeos, K., Moorthy, V. Kieny, M. P Data sharing in public health emergencies: a call to researchers. *Bull WHO*, (2016). doi: 10.2471/BLT.16.170860
12. Skalsky, R. L. & Cullen, B. R. Viruses, microRNAs, and host interactions. *Annu. Rev. Microbiol.* **64**, 123-141 (2010).
13. Klase, Z. A., Sampey, G. C. & Kashanchi, F. Retrovirus infected cells contain viral microRNAs. *Retrovirology* **10**, 1-4 (2013).
14. Kincaid, R. P. & Sullivan, C. S. Virus-encoded microRNAs: An overview and a look to the future. *PLoS Pathog.* **8**, e1003018 (2012).
15. Kadri, S. Hinman, V., Benos, P. V. HHMMiR: efficient de novo prediction of microRNAs using hierarchical hidden Markov models. *BMC Bioinformatics* **10**, S35 (2009).
16. Lorenz R., et al. ViennaRNA Package 2.0. *Algorithms Mol. Biol.* **26**, 1-14 (2011).
17. Ahmed, F., Kaundal, R. & Raghava, G. P. S. PHDcleav: a SVM based method for predicting human Dicer cleavage sites using sequence and secondary structure of miRNA precursors. *BMC Bioinformatics* **14**, S9 (2013).

18. Enright, A.J. et al. MicroRNA targets in Drosophila. *Genome Biol.* **5**, R1.1-R1.14 (2003).
19. Gupta, M., et al. Plekhg4 is a novel Dbl family guanine nucleotide exchange factor protein for rho family GTPases. *J. Biol. Chem.* **288**, 14522-14530 (2013).
20. Xuecai, G., et al. Hook3 interacts with PCM1 to regulate pericentriolar material assembly and the timing of neurogenesis. *Neuron.* **65**, 191-203 (2010).
21. Steinberg, S. J., et al. Peroxisome biogenesis disorders. *Biochim. Biophys. Acta.* **1763**, 1733-1748 (2006).
22. Berger, J., Dorninger, F., Forss-Petter, S. & Kunze, M. Peroxisomes in brain development and function. *Biochim. Biophys. Acta.* **4889**, 426-427 (2015).
23. Hussain, M. S. et al. CDK6 associates with the centrosome during mitosis and is mutated in a large Pakistani family with primary microcephaly. *Hum. Mol. Gen.* **22**, 5199–5214 (2013).
24. Megraw, T. L., Sharkey, J. T., Nowakowski, R. S. Cdk5rap2 exposes the centrosomal root of microcephaly syndromes. *Trends Cell Biol.* **21**, 470-480, 2011.
25. Mlakar J, et al. Zika virus associated with microcephaly. *N Engl J Med.* (2016) doi: 10.1056/NEJMoa1600651
26. Whitty, C. J. M., et al. Providing incentives to share data early in health emergencies: the role of journal editors. *Lancet* **386**, 1797-1798 (2015).




Figure 1 – Phylogenetic analysis of all complete genomes of Zika Virus, available at GenBank (February, 2016). The GenBank accession number and the country of origin are indicated on the ZIKV branches for all strains, except for those from the out-group, where the name of the viruses is provided [Kedougou virus (NC_012533 and AY632540); Bagaza virus (AY632545 and NC_012534)]. The size of the full circles on the branches means percentage of bootstrap (100 replicates).




Figure 2 – Venn diagram showing the nine different types of miRNAs predicted to be encoded by all analyzed ZIKV genomes. Brazil (BR); Central African Republic (CF); French Polynesia (FP); Haiti (HT); Martinique (MQ); Puerto Rico (PR); Uganda (UG).

Table 1 – Examples of human targeting-genes to ZIKV predicted miRNAs.

Ensembl ID	Gene Symbol	Official Full Name	Cause/Effect	Reference PMID
ENSG00000215193	PEX26	peroxisomal biogenesis factor 26	Neonatal Adrenoleukodystrophy, Zellweger syndrome	17055079, 26686055
ENSG00000138685	FGF2	fibroblast growth factor 2 (basic)	Influence on cortical volume	24646670
ENSG00000100241	SBF1	SET binding factor 1	Microcephaly, mental retardation	21210780
ENSG00000168172	HOOK3	hook microtubule-tethering protein 3	Depletion of the neural progenitor pool in the developing neocortex	20152126
ENSG00000196155	PLEKHG4	pleckstrin homology and RhoGEF domain containing G4	Spinocerebellar ataxia	23572525

Table 2 – Examples of human miRNAs able to interact with ZIKV genomes.

miRNAs	Description*	Reference PMID
hsa-miR-34a	CDK6 target	23918663
hsa-mir-324-3p	Cdk5rap2/MCHP3 target	21632253 23726037
hsa-mir-15b-5p	MCPH1/BRIT1 target	12046007
hsa-mir-21-5p	MCPH4/CEP152 target	20598275
hsa-mir-335-5p	MCPH1/BRIT1 target	12046007
hsa-mir-615-3p	WDR62/MCPH2 target	20890278
hsa-miR-193b-3p	WDR62/MCPH2/ target	20890278

*All these genes have been previously related to microcephaly.