

1

2 **ABSTRACT**

3 SARS-CoV-2 employs the angiotensin-converting enzyme 2 (ACE2) receptor and
4 the transmembrane serine protease (TMPRSS2) to infect human lung cells. Previous studies
5 have suggested that different host genetic backgrounds in *ACE2* and *TMPRSS2* could
6 contribute to differences in the rate of infection or severity of COVID-19. Recent studies
7 also showed that variants in 15 genes related to type I interferon immunity to influenza
8 virus could predispose to life-threatening COVID-19 pneumonia. Additional genes
9 (*SLC6A20*, *LZTFL1*, *CCR9*, *FYCO1*, *CXCR6*, *XCRI*, *IL6*, *CTSL*, *ABO*, and *FURIN*) and
10 HLA alleles have also been implicated in response to infection with SARS-CoV-2.
11 Currently, Brazil has recorded the third-highest number of COVID-19 patients worldwide.
12 We aim to investigate the genetic variation present in COVID-19-related genes in the
13 Brazilian population. We analysed 27 candidate genes and HLA alleles in 954 admixed
14 Brazilian exomes. We used the information available in two public databases
15 (<http://www.bipmed.org> and <http://abraom.ib.usp.br/>), and additional exomes from
16 individuals born in southeast Brazil, the region with the highest number of COVID-19
17 patients in the country. Variant allele frequencies were compared with the 1000 Genomes
18 Project phase 3 (1KGP) and the gnomAD databases. We found 395 non-synonymous
19 variants; of these, 325 were also found in the 1000 Genome Project phase 3 (1KGP) and/or
20 gnomAD. Six of these variants were previously reported as putatively influencing the rate
21 of infection or clinical prognosis for COVID-19. The remaining 70 variants were identified
22 exclusively in the Brazilian sample, with a mean allele frequency of 0.0025. *In silico*
23 prediction of the impact in protein function revealed that three of these rare variants were
24 pathogenic. Furthermore, we identified HLA alleles that were previously associated with

1 COVID-19 response at loci DQB1 and DRB1. Our results showed genetic variability
2 common to other populations, but also rare and ultra-rare variants exclusively found in the
3 Brazilian population. These findings could potentially lead to differences in the rate of
4 infection or response to infection by SARS-CoV-2 and should be further investigated in
5 patients with the disease.

6

7

8 **Keywords:** population genomics, admixed population, COVID-19, SARS-Cov-2

9

1 **Introduction**

2 COVID-19 disease, caused by the SARS-CoV-2 coronavirus, is currently a
3 worldwide pandemic. To enter human lung cells, SARS-CoV-2 employs the spike protein,
4 which is primed by the host serine protease (TMPRSS2), followed by angiotensin-
5 converting enzyme 2 (ACE2) receptor binding, and proteolysis with activation of
6 membrane fusion within endosomes by cathepsin L (CTSL)¹⁻⁴. The main feature in SARS-
7 CoV-2 infection is pre-activation of the spike protein by FURIN inside the host cell, which
8 leads to increased SARS-CoV-2 spread into lung cells and increased virulence⁵. The rapid
9 SARS-CoV-2 infection leads to an exacerbated immune reaction, and a few studies have
10 shown that increased levels of IL-6 (an essential immune response mediator) are associated
11 with increased inflammatory response, respiratory failure, increased probability of
12 intubation, the presence of clinical complications, and higher mortality in patients with
13 COVID-19⁶⁻⁸. Additional studies found the enrichment of rare variants predicted to be loss-
14 of-function in genes related to type I interferon (IFN) immunity to influenza virus among
15 patients with life-threatening COVID-19 pneumonia (*TLR3*, *TICAM1*, *TRIF*, *UNC93B1*,
16 *TRAF3*, *TBK1*, *IRF3*, *NEMO*, *IKBKG*, *IFNAR1*, *IFNAR2*, *STAT1*, *STAT2*, *IRF7*, and
17 *IRF9*)⁹.

18 Specific variants in the *ACE2* and *TMPRSS2* genes have been reported among
19 diverse populations worldwide, suggesting that different host genetic backgrounds could
20 contribute to differences in COVID-19 infection and severity^{2,10}. Ellinghaus et al.¹¹
21 performed a genome-wide association study (GWAS) including Italian and Spanish patients
22 with confirmed COVID-19 and controls and identified six candidate genes associated with
23 COVID-19 response on chromosome (chr) 3p21.31 (*SLC6A20*, *LZTFL1*, *FYCO1*, *CXCR6*,

1 *XCRI*, *CCR9*), and one on chr 9q34.2, the locus for genes encoding for the *ABO* blood
2 group antigens. A subsequent, more extensive study replicated the association between the
3 locus on chr 3p21.31 and COVID-19. It revealed a COVID-19 risk core haplotype ranging
4 from 45,859,651bp to 45,909,024bp, which was inherited from Neanderthals and is
5 currently carried by approximately 50% of people in South Asia and about 16% of people
6 in Europe¹². Interestingly, no evidence of association was found for the previously
7 identified candidate genes that are potentially involved in the response to infection by
8 SARS-CoV-2: *ACE2*, *TMPRSS2*, *FURIN*, and *IL6*.

9 Furthermore, one significant factor modulating resistance or susceptibility to viral
10 infections is the human leukocyte antigens (HLAs). HLA polymorphism results from a set
11 of amino-acid substitutions in the peptide-binding groove of the HLA molecules that
12 produce variability in the peptide epitope binding-site and presentation to T cells, which
13 may protect against epidemic infection¹³. Thus, genetic variability in the HLA alleles could
14 influence the immune response in patients with COVID-19, modulating disease severity.
15 Indeed, *in silico* analysis found that HLA-B*46:01 had the fewest predicted binding sites
16 for SARS-CoV-2 peptides, and HLA-B*15:03 showed the greatest capacity to present
17 highly conserved shared SARS-CoV-2 peptides to immune cells¹⁴.

18 Brazil has reported the third-highest number of COVID-19 infections worldwide
19 (updated on September 28th 2020; <https://covid19.who.int/>;
20 <https://coronavirus.jhu.edu/map.html>), and the highest number of cases is concentrated in
21 the south-eastern region of the country (updated on September 28th 2020;
22 <https://covid.saude.gov.br/>). Brazilian individuals feature an admixed genome,
23 encompassing European, sub-Saharan African, and Native Americans as the three main
24 ancestry populations¹⁵⁻¹⁷, and the distribution of ancestry components varies remarkably

1 throughout the genome¹⁸. Furthermore, it has been demonstrated that a significant
2 proportion of genetic variability is still undiscovered in admixed Brazilians¹⁹ and that
3 genetic variability may lead to differential response to infection²⁰. Therefore, we aim to
4 investigate the genetic variation present in COVID-19 related genes in the Brazilian
5 population.

6

7 **Results**

8 *Exome analysis*

9 We found 7,172 variants among the candidate genes analysed (Supplementary Table
10 1). Of these, 395 variants putatively impact protein function, including 354 non-
11 synonymous variants, seven frameshift substitutions, three in-frame deletions, one in-frame
12 insertion, 12 stop gains, two start losses, and 16 splice site variants (Supplementary Table
13 1). Three hundred and twenty-five variants were also present in the gnomAD and/or 1000
14 Genome Project phase 3 (1KGP) databases, including 56 common variants, with an
15 alternative allele frequency (AAF) ≥ 0.01 and 269 rare variants (AAF < 0.01)
16 (Supplementary Data 1). Although AAF from the admixed Brazilian sample follows the
17 distribution from NFE/EUR, AFR, and AMR in gnomAD and 1KGP databases (Fig. 1), we
18 found differences in the AAF of these common and rare variants in the admixed Brazilian
19 sample compared to gnomAD²¹ and/or 1KGP²² databases shown in Fig. 1 and
20 Supplementary Data 1. Interestingly, we also observed some variability in the AAF among
21 samples from different Brazilian towns and the two public databases of genomic
22 information on the Brazilian population, BIPMed and ABraOM (Fig. 2).

1 More importantly, there were 70 variants which were exclusive to the Brazilian
2 sample, including 11 variants in genes related to type I INF immunity to influenza virus⁹,
3 six in candidate genes for COVID response identified by GWAS¹¹, and five related to
4 SARS-CoV-2 entry in lung cells and virus replication^{2,10}. These are rare or ultra-rare
5 variants, presenting a mean AF of 0.0025 (Supplementary Data 1). Among these, we found
6 one in the dataset from Belo Horizonte and two in the ABraOM database for *ACE2*
7 p.Arg219Cys; one in the dataset from Barretos and two in the ABraOM database for *ACE2*
8 p.Leu731Phe; and the *TMPRSS2* p.Val160Met variant was present in samples from all the
9 different Brazilian towns and the two public databases (BIPMed and ABraOM), with an
10 AAF ranging from 0.1333 in Belo Horizonte to 0.2931 in Campinas. Among the reported
11 variants in genes influencing type I INF immunity to influenza virus⁹, we found three
12 variants in the ABraOM database (one *TLR3* p.Pro554Ser, one *IFR3* p.Asn146Lys and one
13 *IRF7* p.Pro246Ser) (Supplementary Data1 and 2).

14 In addition, we identified five variants (rs35044562, rs34326463, rs35508621,
15 rs67959919, and rs35624553) which were previously described in the COVID-19 risk core
16 haplotype and inherited from Neanderthals¹². These were only present in samples from
17 Ribeirão Preto and the BIPMed dataset (rs34326463), Campinas (rs35044562, and
18 rs35508621), and the ABraOM dataset (rs35044562, rs35508621, rs67959919, and
19 rs35624553) (Table 1).

20

21 *In silico predictions*

22 We identified seven variants that were predicted to affect protein function for the 12
23 algorithms used: p.Phe249Ser, p.Gly164Val, and p.Leu25Pro in *the SLC6A20* gene;

1 p.Leu96Arg in *LZTFL1*; p.Tyr287Ser in *XCRI*; and p.Gly146Ser and p.Asn414Ser in the
2 *FURIN* gene (Table 2). Furthermore, the variant p.Gly146Ser in the *FURIN* gene was
3 predicted to destabilise the protein ($\Delta\Delta G$: -1.576 kcal/mol). We observed that the
4 p.Phe249Ser variant is present in samples from Barretos, the BIPMed dataset, gnomAD,
5 and 1KGP (NFE/EUR, AFR, AMR, and SAS populations), whereas the p.Gly164Val
6 variant is present in the ABraOM dataset, gnomAD, and 1KGP (NFE/EUR populations),
7 and the p.Gly146Ser variant is present in the ABraOM dataset, gnomAD, and 1KGP
8 (NFE/EUR, AFR, AMR, EAS, and SAS populations). Notably, four of the variants
9 predicted to be deleterious are found exclusively in admixed Brazilian individuals
10 (p.Leu25Pro in Barretos; p.Leu96Arg in the ABraOM dataset; p.Tyr287Ser in Belo
11 Horizonte; and p.Asn414Ser in the BIPMed dataset).

12 We did not find any predicted deleterious variants in *ACE2* and *TMPRSS2* based on
13 our 12 algorithm criteria. However, Hou et al.² applied only Polyphen2 and CADD scores
14 to variants in *ACE2* and *TMPRSS2* (Polyphen2 >0.96 and CADD >20 as the cut-off).
15 Therefore, only variants defined as ‘probably damaging’ by Polyphen2
16 (<http://genetics.bwh.harvard.edu/pph2/dokuwiki/overview>) and CADD (>20) were
17 included. We found 79 variants predicted to affect protein function, including the
18 p.Val160Met variant in *TMPRSS2* reported by Hou et al.², and the p.Pro554Ser variant in
19 *TLR3* previously reported by Zhang et al.⁹ (Supplementary Data 2).

20

21 *HLA analysis*

22 Overall, we identified 331 different HLA alleles in the admixed Brazilian samples.
23 Of these, three HLA alleles have been previously associated with COVID-19 response^{14,23}.

1 We compared the frequency of these HLA alleles in admixed Brazilians and in populations
2 that occupy the top 10 positions with most cases of COVID-19 and the five populations less
3 affected by the disease, including the United States, India, Russia, Colombia, Peru, Mexico,
4 Spain, Argentina, South Africa, Japan, Australia, South Korea, Vietnam, and Taiwan. The
5 frequency of these alleles is described in Supplementary Data 3. We noticed that the HLA-
6 B*46:01, HLA-B*27:07, HLA-B*15:27, and HLA-C*07:29 alleles were absent in the
7 Brazilian samples. The HLA-C*07:29 allele was also absent from other populations and is
8 present at a low frequency (AF = 0.0003) in the Indian population. HLA-B*15:27 was
9 identified in Vietnam, Taiwan, Japan, with an AF >0.01 and Spain with AF <0.0001. HLA-
10 B*27:07 was detected at a low frequency in India, Colombia, Spain, and South Africa. The
11 HLA-B*46:01 allele was detected in Russia, Mexico, Vietnam, Taiwan, and Japan.

12 Sixty-six Brazilian individuals (17.1%) presented the HLA-DQB1*06:02 allele
13 (AF=0.08938), 47 individuals (12.2%) carry the HLA-DRB1*15:01 allele (AF=0.06477)
14 and 32 individuals (8.29%) have both the HLA-DRB1*15:01 and HLA-DQB1*06:02
15 alleles. The population of other continents, except Oceania, also have these two HLA
16 alleles (HLA-DRB1*15:01 and HLA-DQB1*06:02) with an AF >0.01. Also, 15 Brazilian
17 individuals (3.88%) carry HLA-B*15:03 (AF = 0.02073), which is predicted to have the
18 greatest capacity to present SARS-CoV-2 peptides to immune cells¹⁴. This allele was not
19 found in the Asian population of Japan, South Korea, and Vietnam (Supplementary Data 3).

20

21 *In silico analysis of viral peptide-HLA class I and II binding affinity*

22 To verify the potential for an HLA allele type to produce an antiviral response to
23 SARS-CoV-2, we performed HLA binding affinity analyses to the SARS-CoV-2 proteome.

1 We tested 42 HLA-A proteins, 77 HLA-B, 38 HLA-C, 60 HLA-DP (DPA1/DPB1), 145
2 HLA-DQ (DQA1/DQB1), 46 HLA-DRB1, 4 DRB3, 2 DRB4, and 6 DRB5.

3 The SARS-CoV-2 proteome was presented by a diversity of HLA alleles from
4 classes I and II (Supplementary Table 2). The HLA proteins are predicted to bind a small
5 proportion of all possible SARS-CoV-2 derived peptides with high affinity (on average
6 0.5% for HLA class I and 2% for HLA class II). Also, we found a small proportion of weak
7 binders (on average, 1.5% for HLA class I and 8.2% for class II). Most of the HLA proteins
8 do not bind either Class I (on average >96%) or class II (on average >89%) molecules
9 (Supplementary data 4). Supplementary Data 5 shows a list of HLA strongest binders
10 (>300 peptides bound at high affinity) of SARS-CoV-2 peptides. These are found in loci
11 HLA-A, -B, -C, and DQ.

12

13 **Discussion**

14 Accessing the genomic sequences of the general population is relevant to identify
15 the genetic variability involved in the molecular mechanisms of infection²⁰. Also, it is
16 known that the admixed Brazilian population is underrepresented in large public
17 databases^{21,22}, and previous studies revealed variants present exclusively in Brazilian
18 individuals^{19,24}.

19 We studied 27 human COVID-19-related genes and the HLA region in two public
20 genomic databases of admixed Brazilians (BIPMed, www.bipmed.org¹⁹; ABraOM
21 <http://abraom.ib.usp.br>²⁴), and additional samples from individuals born in three different
22 towns of south-eastern Brazil. We reported the variants and HLA alleles found in these
23 samples and compared them with worldwide populations. We also reported variants

1 constituting the COVID-19 risk core haplotype on locus 3p21.31, described as being
2 inherited from Neanderthals¹².

3 Previous studies showed that the *ACE2*, *TMPRSS2*, *CTSL*, *FURIN*, and *IL6* genes,
4 as well as the HLA region, may be involved in SARS-CoV-2 infection^{1-5,10} and immune
5 response^{6-8,14,23,25}. Furthermore, variants on loci 3p21.31 and 9q34.2 (encompassing
6 *SLC6A20*, *LZTFL1*, *FYCO1*, *CXCR6*, *XCRI*, *CCR9*, and *ABO*) have been associated with
7 Spanish and Italian patients with COVID-19¹¹, and different variants were found to affect
8 the predisposition to life-threatening illness in patients with COVID-19 from different
9 ancestries⁹.

10 The analysis of genetic variability in candidate genes for specific populations can
11 help to identify individuals at a higher risk of infection or severe disease by constructing
12 risk haplotypes, which can also provide therapeutic targets for the development of more
13 effective treatments and the control of COVID-19^{2,10}. Thus, in addition to investigating
14 genetic variability in the 27 candidate genes, we extended our analysis to include HLA
15 alleles, which influence immunological response to many infectious agents (updated on
16 September 28th 2020; <https://covid19.who.int/>; <https://coronavirus.jhu.edu/map.html>). This
17 is the first comprehensive study of genetic variability of COVID-19 genes in admixed
18 individuals from Latin America, a hard-stricken population in the COVID-19 pandemic²⁶,
19 both in terms of the number of infected individuals and the severity of disease leading to
20 increased death rates. Indeed, in the USA, remarkable disparities of SARS-CoV-2 infection
21 by ethnicity have been shown, with Hispanic/Latino and African American individuals
22 presenting higher SARS-CoV-2 infection rates and risk mortality than ‘non-Hispanic
23 white’ Americans²⁷⁻²⁹. Therefore, by looking at population genomics data, one may gain
24 insights into disease-related variants, which could be disproportionally represented in

1 specific populations^{18,30-32}. Furthermore, by evaluating individuals with unknown
2 information on SARS-CoV-2 infection, one can achieve the random distribution of these
3 variants, allowing better estimates of the distribution of population allele frequencies.

4 We identified small differences in AF in the 395 candidate variants identified
5 among Brazilian samples, strengthening the hypothesis that different genetic backgrounds
6 could influence SARS-CoV-2 infection and behaviour in human host cells^{2,10}. Furthermore,
7 this study and previous works^{2,10} identified individuals who carry unique deleterious
8 variants, which may influence gene function and could potentially lead to different
9 responses to SARS-CoV-2 infection on an individual scale. However, the rather similar
10 distribution of AFs among Brazilians and their ancestry populations (NFE/EUR and AFR),
11 as well as other admixed Americans (AMR), and the fact that the unique variants identified
12 in the Brazilian population are rare or ultra-rare, indicates that the admixed Brazilian
13 genetic background is not sufficient to influence SARS-CoV-2 infection on a population
14 scale.

15 Zeberg and Pääbo¹² have shown that the major genetic risk factor for severe
16 COVID-19 was inherited from Neanderthals¹². This finding is important on a regional
17 scale, since 4% of admixed Americans analysed by Zeberg and Pääbo¹² (including 1533
18 Brazilian controls from the BRACOVID dataset) presented the core haplotype derived from
19 Neanderthals. Interestingly, Campinas, Ribeirão Preto, and the BIPMed dataset showed
20 only one risk allele, while Barretos and Belo Horizonte did not present any risk allele of the
21 Neanderthal's core haplotype reported. Therefore, if further studies demonstrate that the
22 Neanderthal-derived region confers a risk to COVID-19, this information should be
23 carefully evaluated in additional admixed Brazilian samples from different geographic
24 areas.

1 Currently, there is no consensus regarding a possible association of HLA alleles and
2 susceptibility to COVID-19. Ellinghaus et al.¹¹ did not find any evidence of an association
3 between HLA and COVID-19. On the other hand, HLA-DRB1*15:01, HLA-DQB1*06:02,
4 and HLA-B*27:07 alleles were associated with Italian patients affected by an extremely
5 severe or severe form of COVID-19²³, and an increased frequency of HLA-C*07:29 and
6 HLA-B*15:27 was detected in Chinese patients with COVID-19 in comparison to the
7 Chinese control population²⁵. Interestingly, the HLA-C*07:29 allele is absent from the
8 Brazilian admixed samples included in the present study and in all populations used in the
9 comparisons, except for individuals from India, where this allele was found at a low
10 frequency (0.0003). On the other hand, the HLA-B*15:27 allele was identified in
11 individuals from three Asian countries (Vietnam, Taiwan and Japan) with AF >0.01, and at
12 a low frequency in Spain (0.0001), but absent from Brazilian samples. The HLA-B*27:07
13 allele found in Italian individuals with a severe manifestation of COVID-19 was also
14 identified in India, Colombia, Spain, and South Africa, but not in populations from Asia
15 and Oceania (countries that are less affected by COVID-19) and from Brazil. In contrast,
16 the HLA-DQB1*06:02 is present in all populations surveyed in this study, including
17 Brazilian individuals (17.1%), with the exception of individuals from Australia. Also, the
18 HLA-DRB1*15:01 allele is present in all populations investigated in this study, including
19 Brazilian individuals (12.2%), but not in individuals from Australia and Peru. Interestingly,
20 8.29% of Brazilian individuals carry both the HLA-DRB1*15:01 and HLA-DQB1*06:02
21 alleles.

22 Furthermore, the HLA-A, -B, -C, and DQ loci show haplotypes that are strong
23 binders of SARS-CoV-2 peptides in the Brazilian samples, especially for the HLA-A locus
24 (20 alleles, Table 3). When comparing different populations, we found marked variability

1 in the frequency of the different HLA alleles putatively associated with the severe
2 manifestation of COVID-19, such as HLA-DRB1*15:01, HLA-DQB1*06:02, and HLA-
3 B*27:07 alleles²³. Overall, 10% of Brazilian individuals carried at least two of the alleles
4 associated with the severe manifestation of COVID-19. Interestingly, the same alleles were
5 absent from individuals from Australia. Variability in the frequency of HLA alleles
6 previously associated with COVID-19 highlights the importance of considering ethnic and
7 geographic origin when performing studies investigating the role of HLA alleles and
8 disease. Thus, it seems likely that different population-specific haplotypes may be
9 associated with an increased risk of severe disease in different populations.

10 In conclusion, we found rare variants in three COVID-19-related genes that are
11 present only in the Brazilian dataset and are predicted to affect protein function.
12 Furthermore, we identified HLA alleles previously associated with COVID-19
13 immunological response and 31 HLA alleles predicted as strong binders to SARS-CoV-2
14 peptides at loci -A, -B, -C, and DQ, which indicates the importance of further investigation
15 on the role of HLA haplotypes as modulators of response to infection to SARS-CoV-2.
16 Although the variants predicted to affect protein function in COVID-19-related genes are
17 rare in admixed Brazilians (varying from 0.0001 to 0.0032), these also emerge as
18 candidates for modulating response to infection by the SARS-CoV-2 in the Brazilian
19 population. Furthermore, our study suggests the utility of population genomic studies in the
20 context of precision health to stratify risk for infection disorders.

21

1 **Methods**

2 *Subjects*

3 We evaluated exomes from 257 individuals from the BIPMed dataset¹⁹, 609 from
4 the ABraOM dataset²⁴, and an additional 88 exomes from individuals born in three towns in
5 south-eastern Brazil: Barretos (N=30), Ribeirão Preto (N=30), located in the state of São
6 Paulo, and Belo Horizonte (N=28) the capital of the state of Minas Gerais. Among the
7 BIPMed individuals, 193 had information about their city of birth available. The HLA
8 region was sequenced in 386 individuals, including the 257 from BIPMed, the 88 additional
9 exomes, and an additional 41 individuals (22 from southeast Brazil). We signed terms of
10 data privacy to obtain permission to use the raw data from BIPMed and ABraOM public
11 databases and use raw data of the 88 exomes from Barretos, Ribeirão Preto, and Belo
12 Horizonte. This study was approved by the University of Campinas Research Ethics
13 Committee (UNICAMP, Campinas, São Paulo, Brazil). All methods were performed
14 according to the relevant guidelines and regulations.

15

16 *Exome analysis*

17 Whole exome data were stored in variant call format (VCF) files built-in
18 GRCh37/hg19 assembly. Gene regions were extracted by *vcftools*³³ based on the position
19 reported in Ensembl GRCh37 Release 101³⁴ (Supplementary Table 1). Variant
20 consequences were annotated from each gene region by ANNOVAR software (version
21 2019Oct24)³⁵, using the following flags: *-otherinfo* (to include Brazil AF); *-onetranscript*; *-*
22 *buildver hg19*; *-remove*; *-protocol refGene,gnomad211_exome; ALL.sites.2015_08;*
23 *EUR.sites.2015_08; AFR.sites.2015_08; AMR.sites.2015_08; EAS.sites.2015_08;*

1 *SAS.sites.2015_08*; *-operation g,f*; and *-nastring*. ANNOVAR software provides allele
2 frequency (AF) information from African/African-American (AFR/AFA), Latino/admixed
3 American (LAT/AMR), East Asian (EAS), non-Finish European (NFE), and South Asian
4 (SAS) populations from gnomAD exome dataset²¹, as well as sub-Saharan Africans (AFR),
5 Europeans (EUR), admixed Americans (AMR), east Asians (EAS), and south Asians (SAS)
6 from 1KGP phase 3 dataset²². In addition, we annotated variants which were not identified
7 by ANNOVAR using Variant Effect Prediction (VEP) algorithm³⁶, with the following
8 parameters: *--buffer_size 500*; *--canonical*; *--distance 5000*; *--species homo_sapiens*; *--*
9 *symbol*.

10 To evaluate whether regional variability is observed among Brazilian samples, we
11 separated the samples based on the city in which individuals were born, including 32
12 individuals from Campinas extracted from the BIPMed dataset.

13

14 *In silico prediction analysis*

15 To predict the impact on protein function of the non-synonymous variants
16 identified, we applied the following computer algorithms, which is currently recommended
17 by the ACMG/AMP guidelines: PANTHER³⁷, MutationTaster³⁸, Condel³⁹, PROVEAN⁴⁰,
18 PolyPhen2⁴¹, Sort Intolerant from tolerant (SIFT)⁴², Align Grantham Variation/ Grantham
19 Difference score (GVGD)⁴³, Combined Annotation Dependent Depletion (CADD)⁴⁴, PhD-
20 SNPg⁴⁵, Functional Analysis through Hidden Markov Models (FATHMM)⁴⁶, SNPs&GO⁴⁷,
21 and MutPred2 (<http://mutpred.mutdb.org>).

22 For Align-GVGD, we classified the variants based on the graded classifier with a
23 cut-off of C35 or higher for deleterious classification. For CADD, we used the PHRED-like
24 score with a cut-off of 20, below which the variants were classified as benign and otherwise

1 deleterious. For MutPred2, we considered a score threshold of 0.50 for pathogenicity. For
2 all other algorithms, we considered the classification provided as an output.

3 To access the impact of mutations on protein dynamics and stability, we used the
4 DynaMut server (<http://biosig.unimelb.edu.au/dynamut/>)⁴⁸. The server requires an input file
5 of protein structure in PDB format or by providing the four-letter accession code for any
6 entry on the Protein Data Bank database (PDB; <http://wwpdb.org>). The code for the *FURIN*
7 gene used was 5jxg. The other proteins are not available in the PDB database to be tested.

8

9 *HLA analysis*

10 We sequenced 11 HLA Loci (HLA-A, HLA-B, HLA-C, HLA-DRB1, HLA-DQB1,
11 HLA-DPB1, HLA-DQA1, HLA -DPA1, HLA-DRB3, HLA-DRB4, HLA-DRB5) in 298
12 samples using NGSgo[®] panels (GenDx, Utrecht, The Netherlands). The DNA libraries
13 were loaded onto a MiSeq Sequencer (Illumina Inc., San Diego, CA, USA), and the data
14 were analysed with the NGSengine v.2.16.2 software (GenDx, Utrecht, The Netherlands).
15 We determined the HLA alleles from the remaining 88 exomes using the HLA-HD (HLA
16 typing from High-quality Dictionary) tool v.1.3.0⁴⁹⁻⁵¹. The IPD-IMGT/HLA database
17 release 3.40.0⁵² was used as a reference. Even though we obtained results with six- and
18 eight-digit precision, we restricted the results to four-digit accuracy to compare with
19 published data. HLA allele frequencies were calculated by Arlequin v.3.5.2.2 software⁵³.

20

21 *In silico analysis of viral peptide-HLA class I and II binding affinity*

22 We performed *in silico* analysis of viral peptide-HLA class I and II binding affinity
23 across HLA proteins found in our population for the entire SARS-CoV-2 proteome. All
24 HLA-A, -B, -C alleles were selected to assess the peptide-binding affinity of their

1 corresponding proteins HLA-A, HLA-B, HLA-C, respectively. The HLA-DR is
2 represented by HLA-DRA/DRB1 dimer. Since HLA-DRA is considered monomorphic, we
3 just used the HLA-DRB1. The HLA-DP and DQ are represented by the HLA-DPA1/DPB1
4 dimer and HLA-DQA1/DQB1 dimer, respectively.

5 FASTA-formatted protein sequence data was retrieved from the National Center of
6 Biotechnology Information (NCBI) database
7 (<https://www.ncbi.nlm.nih.gov/genome/browse#!/overview/Sars-cov-2>). The follow eleven
8 protein viral product was used in the analyses: ORF1ab (YP_009724389.1), Surface
9 Glycoprotein (S) (YP_009724390.1), ORF3a (YP_009724391.1), Envelope (E)
10 (YP_009724392.1), Membrane Glycoprotein (M) (YP_009724393.1), ORF6
11 (YP_009724394.1), ORF7a (YP_009724395.1), ORF7b (YP_009725318.1), ORF8
12 (YP_009724396.1), Nucleocapsid (N) (YP_009724397.2), and ORF10 (YP_009725255.1).

13 We k-merised these sequences into 8- to 12-mers to assess HLA class I-peptide
14 binding affinity and into 15-mers to assess HLA class II binding affinity across the entire
15 proteome. Predictions for HLA were performed using different HLA alleles found in our
16 population with netMHCpan v4.1 for class I⁵⁴ and NetMHCIIpan - 3.2 for class II⁵⁵.

17

18 *HLA allele and haplotype frequencies of other populations*

19 HLA frequency data were obtained from the Allele Frequency Net Database
20 (<http://www.allelefrequencies.net/>)⁵⁶ for 10 distinct populations that are most and least
21 affected by COVID-19. We checked the HLA of the populations that occupy the top 10
22 positions (USA, India, Brazil, Russia, Colombia, Peru, Spain, Mexico, Argentina, South
23 Africa) and those that were less affected (Australia, Vietnam, Taiwan, Japan, and South

1 Korea) (accessed on 04/24/2020, <https://www.worldometers.info/coronavirus/>) according to
2 the availability of this data in the Allele Frequency Net Database.

3

4 **Data availability**

5 BIPMed raw dataset that supports this study's findings is available in EVA
6 repository/PRJEB39251, <https://www.ebi.ac.uk/eva/?eva-study=PRJEB39251>. ABraOM
7 raw dataset that supports the results of this study is available from ABraOM
8 (<http://abraom.ib.usp.br/>).

9

10 **Acknowledgments**

11 This work was supported by the Fundação de Amparo à Pesquisa do Estado de São
12 Paulo (FAPESP, grant number 2013/07559-3). RS was supported by FAPESP (grant
13 number 2019/08526-8). TKA is supported by FAPESP (grant number 2017/01900-6). IL-C
14 is supported by CNPq (grant number 311923/2019-4).

15

16 **Competing interests**

17 The authors declare no competing interests.

18

19 **Author contributions**

20 RS contributed with the study design, conceptualization, data acquisition, analysis, and
21 paper writing; TKA contributed with HLA sequencing, analysis, *in silico* prediction
22 analysis, and writing of the paper; MCG contributed with *in silico* prediction analysis and
23 paper writing; CSR contributed with public data acquisition and processing; MN and MZ

- 1 contribute with public data acquisition; LD and MACB contributed with Belo Horizonte
- 2 data acquisition and sample information; VLV contributed with Barretos data acquisition
- 3 and sample information; WAS contributes with Ribeirão Preto data acquisition and sample
- 4 information; ILC contributed with project conceptualization and served as principal
- 5 investigators. All authors reviewed the manuscript.

1 **References**

- 2 1 Shang, J. *et al.* Cell entry mechanisms of SARS-CoV-2. *Proceedings of the*
3 *National Academy of Sciences* **117**, 11727-11734, doi:10.1073/pnas.2003138117
4 (2020).
- 5 2 Hou, Y. *et al.* New insights into genetic susceptibility of COVID-19: an ACE2 and
6 TMPRSS2 polymorphism analysis. *BMC Medicine* **18**, 216, doi:10.1186/s12916-
7 020-01673-z (2020).
- 8 3 Lu, R. *et al.* Genomic characterisation and epidemiology of 2019 novel coronavirus:
9 implications for virus origins and receptor binding. *The Lancet* **395**, 565-574,
10 doi:10.1016/S0140-6736(20)30251-8 (2020).
- 11 4 Simmons, G. *et al.* Inhibitors of cathepsin L prevent severe acute respiratory
12 syndrome coronavirus entry. *Proceedings of the National Academy of Sciences of*
13 *the United States of America* **102**, 11876-11881, doi:10.1073/pnas.0505577102
14 (2005).
- 15 5 Hoffmann, M., Kleine-Weber, H. & Pöhlmann, S. A Multibasic cleavage site in the
16 spike protein of SARS-CoV-2 is essential for infection of human lung cells.
17 *Molecular Cell* **78**, 779-784.e775, doi:10.1016/j.molcel.2020.04.022 (2020).
- 18 6 Ye, Q., Wang, B. & Mao, J. The pathogenesis and treatment of the 'Cytokine Storm'
19 in COVID-19. *Journal of Infection* **80**, 607-613,
20 doi:<https://doi.org/10.1016/j.jinf.2020.03.037> (2020).
- 21 7 Herold, T. *et al.* Elevated levels of IL-6 and CRP predict the need for mechanical
22 ventilation in COVID-19. *Journal of Allergy and Clinical Immunology* **146**, 128-
23 136.e124, doi:10.1016/j.jaci.2020.05.008 (2020).

- 1 8 Aziz, M., Fatima, R. & Assaly, R. Elevated interleukin-6 and severe COVID-19: A
2 meta-analysis. *Journal of Medical Virology* **n/a**, doi:10.1002/jmv.25948 (2020).
- 3 9 Zhang, Q. *et al.* Inborn errors of type I IFN immunity in patients with life-
4 threatening COVID-19. *Science*, doi:10.1126/science.abd4570 (2020).
- 5 10 Cao, Y. *et al.* Comparative genetic analysis of the novel coronavirus (2019-
6 nCoV/SARS-CoV-2) receptor ACE2 in different populations. *Cell Discov* **6**, 11,
7 doi:10.1038/s41421-020-0147-1 (2020).
- 8 11 Ellinghaus, D. *et al.* Genome-wide association study of severe covid-19 with
9 respiratory failure. *New England Journal of Medicine*,
10 doi:10.1056/NEJMoa2020283 (2020).
- 11 12 Zeberg, H. & Pääbo, S. The major genetic risk factor for severe COVID-19 is
12 inherited from Neanderthals. *Nature*, doi:10.1038/s41586-020-2818-3 (2020).
- 13 13 Shi, Y. *et al.* COVID-19 infection: the perspectives on immune responses. *Cell*
14 *Death Differ* **27**, 1451-1454, doi:10.1038/s41418-020-0530-3 (2020).
- 15 14 Nguyen, A. *et al.* Human leukocyte antigen susceptibility map for severe acute
16 respiratory syndrome coronavirus 2. *Journal of Virology* **94**, e00510-00520,
17 doi:10.1128/JVI.00510-20 (2020).
- 18 15 Rodrigues de Moura, R., Coelho, A. V. C., de Queiroz Balbino, V., Crovella, S. &
19 Brandão, L. A. C. Meta-analysis of Brazilian genetic admixture and comparison
20 with other Latin America countries. *American Journal of Human Biology* **27**, 674-
21 680, doi:10.1002/ajhb.22714 (2015).
- 22 16 Kehdy, F. S. G. *et al.* Origin and dynamics of admixture in Brazilians and its effect
23 on the pattern of deleterious mutations. *Proceedings of the National Academy of*
24 *Sciences* **112**, 8696-8701, doi:10.1073/pnas.1504447112 (2015).

- 1 17 Lima-Costa, M. F. *et al.* Genomic ancestry and ethnoracial self-classification based
2 on 5,871 community-dwelling Brazilians (The Epigen Initiative). *Sci Rep* **5**, 9812,
3 doi:10.1038/srep09812 (2015).
- 4 18 Secolin, R. *et al.* Distribution of local ancestry and evidence of adaptation in
5 admixed populations. *Sci Rep* **9**, 13900, doi:10.1038/s41598-019-50362-2 (2019).
- 6 19 Rocha, C. S., Secolin, R., Rodrigues, M. R., Carvalho, B. S. & Lopes-Cendes, I.
7 The Brazilian Initiative on Precision Medicine (BIPMed): fostering genomic data-
8 sharing of underrepresented populations. *npj Genomic Medicine* **5**, 42,
9 doi:10.1038/s41525-020-00149-6 (2020).
- 10 20 Chapman, S. J. & Hill, A. V. Human genetic susceptibility to infectious disease. *Nat*
11 *Rev Genet* **13**, 175-188, doi:10.1038/nrg3114 (2012).
- 12 21 Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from
13 variation in 141,456 humans. *Nature* **581**, 434-443, doi:10.1038/s41586-020-2308-7
14 (2020).
- 15 22 Auton, A. *et al.* A global reference for human genetic variation. *Nature* **526**, 68-74,
16 doi:10.1038/nature15393 (2015).
- 17 23 Novelli, A. *et al.* HLA alleles frequencies and susceptibility to COVID-19 in a
18 group of 99 Italian patients. *HLA* **n/a**, doi:10.1111/tan.14047 (2020).
- 19 24 Naslavsky, M. S. *et al.* Exomic variants of an elderly cohort of Brazilians in the
20 ABraOM database. *Hum Mutat* **38**, 751-763, doi:10.1002/humu.23220 (2017).
- 21 25 Wang, W., Zhang, W., Zhang, J., He, J. & Zhu, F. Distribution of HLA allele
22 frequencies in 82 Chinese individuals with coronavirus disease-2019 (COVID-19).
23 *HLA* **96**, 194-196, doi:10.1111/tan.13941 (2020).

- 1 26 Dendrou, C. A., Petersen, J., Rossjohn, J. & Fugger, L. HLA variation and disease.
2 *Nat Rev Immunol* **18**, 325-339, doi:10.1038/nri.2017.143 (2018).
- 3 27 Alcendor, D. J. Racial disparities-associated COVID-19 mortality among minority
4 populations in the US. *J Clin Med* **9**, doi:10.3390/jcm9082442 (2020).
- 5 28 Kaufman, H. W., Niles, J. K. & Nash, D. B. Disparities in SARS-CoV-2 positivity
6 rates: Associations with race and ethnicity. *Popul Health Manag*,
7 doi:10.1089/pop.2020.0163 (2020).
- 8 29 Poulson, M. *et al.* National disparities in COVID-19 outcomes between black and
9 white Americans. *J Natl Med Assoc*, doi:10.1016/j.jnma.2020.07.009 (2020).
- 10 30 Deng, L., Ruiz-linares, A., Xu, S. & Wang, S. Ancestry variation and footprints of
11 natural selection along the genome in Latin American populations. *Scientific*
12 *Reports* **6**, 1-7, doi:10.1038/srep21766 (2016).
- 13 31 Cáceres, A. & González, J. R. Following the footprints of polymorphic inversions
14 on SNP data: from detection to association tests. *Nucleic acids research* **43**, e53-
15 e53, doi:10.1093/nar/gkv073 (2015).
- 16 32 Homburger, J. R. *et al.* Genomic insights into the ancestry and demographic history
17 of South America. *PLoS Genetics* **11**, doi:10.1371/journal.pgen.1005602 (2015).
- 18 33 Danecek, P. *et al.* The variant call format and VCF tools. *Bioinformatics* **27**, 2156-
19 2158, doi:10.1093/bioinformatics/btr330 (2011).
- 20 34 Yates, A. D. *et al.* Ensembl 2020. *Nucleic Acids Research* **48**, D682-D688,
21 doi:10.1093/nar/gkz966 (2020).
- 22 35 Wang, K., Li, M. & Hakonarson, H. ANNOVAR: functional annotation of genetic
23 variants from high-throughput sequencing data. *Nucleic acids research* **38**, e164-
24 e164, doi:10.1093/nar/gkq603 (2010).

- 1 36 McLaren, W. *et al.* The Ensembl variant effect predictor. *Genome Biol* **17**, 122,
2 doi:10.1186/s13059-016-0974-4 (2016).
- 3 37 Mi, H., Muruganujan, A. & Thomas, P. D. PANTHER in 2013: modelling the
4 evolution of gene function, and other gene attributes, in the context of phylogenetic
5 trees. *Nucleic Acids Res* **41**, D377-386, doi:10.1093/nar/gks1118 (2013).
- 6 38 Schwarz, J. M., Rödelberger, C., Schuelke, M. & Seelow, D. MutationTaster
7 evaluates disease-causing potential of sequence alterations. *Nat Methods* **7**, 575-
8 576, doi:10.1038/nmeth0810-575 (2010).
- 9 39 González-Pérez, A. & López-Bigas, N. Improving the assessment of the outcome of
10 non-synonymous SNVs with a consensus deleteriousness score, Condel. *Am J Hum*
11 *Genet* **88**, 440-449, doi:10.1016/j.ajhg.2011.03.004 (2011).
- 12 40 Choi, Y., Sims, G. E., Murphy, S., Miller, J. R. & Chan, A. P. Predicting the
13 functional effect of amino acid substitutions and indels. *PLoS One* **7**, e46688,
14 doi:10.1371/journal.pone.0046688 (2012).
- 15 41 Adzhubei, I., Jordan, D. M. & Sunyaev, S. R. Predicting functional effect of human
16 missense mutations using PolyPhen-2. *Curr Protoc Hum Genet* **Chapter 7**,
17 Unit7.20, doi:10.1002/0471142905.hg0720s76 (2013).
- 18 42 Sim, N. L. *et al.* SIFT web server: predicting effects of amino acid substitutions on
19 proteins. *Nucleic Acids Res* **40**, W452-457, doi:10.1093/nar/gks539 (2012).
- 20 43 Tavtigian, S. V. *et al.* Comprehensive statistical study of 452 BRCA1 missense
21 substitutions with classification of eight recurrent substitutions as neutral. *J Med*
22 *Genet* **43**, 295-305, doi:10.1136/jmg.2005.033878 (2006).

- 1 44 Rentzsch, P., Witten, D., Cooper, G. M., Shendure, J. & Kircher, M. CADD:
2 predicting the deleteriousness of variants throughout the human genome. *Nucleic
3 Acids Res* **47**, D886-D894, doi:10.1093/nar/gky1016 (2019).
- 4 45 Capriotti, E. & Fariselli, P. PhD-SNPg: a webserver and lightweight tool for scoring
5 single nucleotide variants. *Nucleic Acids Res* **45**, W247-W252,
6 doi:10.1093/nar/gkx369 (2017).
- 7 46 Shihab, H. A. *et al.* An integrative approach to predicting the functional effects of
8 non-coding and coding sequence variation. *Bioinformatics* **31**, 1536-1543,
9 doi:10.1093/bioinformatics/btv009 (2015).
- 10 47 Calabrese, R., Capriotti, E., Fariselli, P., Martelli, P. L. & Casadio, R. Functional
11 annotations improve the predictive score of human disease-related mutations in
12 proteins. *Hum Mutat* **30**, 1237-1244, doi:10.1002/humu.21047 (2009).
- 13 48 Rodrigues, C. H., Pires, D. E. & Ascher, D. B. DynaMut: predicting the impact of
14 mutations on protein conformation, flexibility and stability. *Nucleic Acids Res* **46**,
15 W350-W355, doi:10.1093/nar/gky300 (2018).
- 16 49 Kawaguchi, S. & Matsuda, F. High-definition genomic analysis of HLA genes via
17 comprehensive HLA allele genotyping. *Methods Mol Biol* **2131**, 31-38,
18 doi:10.1007/978-1-0716-0389-5_3 (2020).
- 19 50 Kawaguchi, S., Higasa, K., Yamada, R. & Matsuda, F. Comprehensive HLA typing
20 from a current allele database using next-generation sequencing data. *Methods Mol
21 Biol* **1802**, 225-233, doi:10.1007/978-1-4939-8546-3_16 (2018).
- 22 51 Kawaguchi, S., Higasa, K., Shimizu, M., Yamada, R. & Matsuda, F. HLA-HD: An
23 accurate HLA typing algorithm for next-generation sequencing data. *Hum Mutat* **38**,
24 788-797, doi:10.1002/humu.23230 (2017).

- 1 52 Robinson, J. *et al.* IPD-IMGT/HLA database. *Nucleic Acids Res* **48**, D948-D955,
2 doi:10.1093/nar/gkz950 (2020).
- 3 53 Excoffier, L. & Lischer, H. E. Arlequin suite ver 3.5: a new series of programs to
4 perform population genetics analyses under Linux and Windows. *Mol Ecol Resour*
5 **10**, 564-567, doi:10.1111/j.1755-0998.2010.02847.x (2010).
- 6 54 Reynisson, B., Alvarez, B., Paul, S., Peters, B. & Nielsen, M. NetMHCpan-4.1 and
7 NetMHCIIpan-4.0: improved predictions of MHC antigen presentation by
8 concurrent motif deconvolution and integration of MS MHC eluted ligand data.
9 *Nucleic Acids Res* **48**, W449-W454, doi:10.1093/nar/gkaa379 (2020).
- 10 55 Nielsen, M., Justesen, S., Lund, O., Lundegaard, C. & Buus, S. NetMHCIIpan-2.0 -
11 Improved pan-specific HLA-DR predictions using a novel concurrent alignment and
12 weight optimization training procedure. *Immunome Res* **6**, 9, doi:10.1186/1745-
13 7580-6-9 (2010).
- 14 56 Gonzalez-Galarza, F. F. *et al.* Allele frequency net database (AFND) 2020 update:
15 gold-standard data classification, open access genotype data and new query tools.
16 *Nucleic Acids Res* **48**, D783-D788, doi:10.1093/nar/gkz1029 (2020).
- 17

Tables

Table 1. Alternative allele frequency (AAF) of variants encompassing the COVID-19 risk core haplotype and the alleles present in Neanderthal samples.

dbSNP	Chr	Pos	ALT	Brazil AF	Risk allele*	Campinas	Barretos	Ribeirão Preto	Belo Horizonte	BIPMed	ABraOM
rs35044562	3	45909024	G	0.0011	G	0.0862	0.0000	0.0000	0.0000	0.0279	0.0311
rs34326463	3	45899651	G	0.0011	G	0.0000	0.0000	0.0357	0.0000	0.0000	0.0000
rs35508621	3	45880481	C	0.0011	C	0.0345	0.0000	0.0000	0.0000	0.0000	0.0039
rs67959919	3	45871908	A	0.0011	A	0.0000	0.0000	0.0000	0.0000	0.0000	0.0039
rs35624553	3	45867440	G	0.0262	G	0.0000	0.0000	0.0000	0.0000	0.0000	0.0039

*data extracted from Zeberg and Pääbo, 2020 (doi:10.1038/s41586-020-2818-3)

Table 2. Alternative allele frequency of deleterious variants according to 12 different prediction algorithms

Gene	Variant	Alternative Allele Frequency										
		Campinas	Barretos	Ribeirão Preto	Belo Horizonte	ABraOM	BIPMed	NFE	AFR	AMR	EAS	SAS
<i>SLC6A20</i>	Phe249Ser	0	0.0167	0	0	0.0008	0.0019	0.0005	0.0615	0.0003	0	0.0084
<i>SLC6A20</i>	Gly164Val	0	0	0	0	0.0008	0	0.0088	0	0	0	0
<i>SLC6A20</i>	Leu25Pro	0	0.0167	0	0	0	0	0	0	0	0	0
<i>LZTFL1</i>	Leu96Arg	0	0	0	0	0.0008	0	0	0	0	0	0
<i>XCR1</i>	Tyr287Ser	0	0	0	0.0167	0	0	0	0	0	0	0
<i>FURIN</i>	Gly146Ser	0	0	0	0	0.0008	0	0.0004	0.0002	0.0005	0.0544	0.0327
<i>FURIN</i>	Asn414Ser	0	0	0	0	0	0.0019	0	0	0	0	0

NFE=non-Finish European; AFR=sub-Saharan African/African American; AMR=admixed Americans/Latinos; EAS=east Asians; SAS=south Asians.

Table 3. List of HLA strongest binders (>300 peptides bound at high affinity) of SARS-CoV-2 peptides and frequency in the Brazilian sample.

HLA	HLA alleles	Allele frequency
HLA -A	A*01:01	0.10233
	A*11:01	0.04145
	A*11:67	0.00130
	A*23:01	0.03756
	A*23:17	0.00389
	A*24:02	0.10104
	A*24:03	0.00259
	A*24:05	0.00130
	A*25:01	0.00389
	A*26:01	0.02979
	A*26:02	0.00130
	A*26:08	0.00130
	A*29:01	0.00259
	A*29:02	0.04534
	A*29:119	0.00130
	A*30:02	0.02591
	A*30:04	0.00259
A*34:02	0.00777	
A*36:01	0.00389	
A*80:01	0.00130	
HLA-B	B*15:08	0.00130
	B*15:11	0.00130
HLA-C	C*03:02	0.00389
	C*07:02	0.05699
	C*07:50	0.00130
	C*14:02	0.02979
	C*14:03	0.00259
HLA-DQ	DQA1*0201-DQB1*0402	0.00259
	DQA1*0301-DQB1*0402	0.00130
	DQA1*0303-DQB1*0401	0.00130
	DQA1*0303-DQB1*0402	0.00130

Figures legend

Figure 1. Distribution of alternative allele frequency of common variants ($AAF \geq 0.01$) from samples of admixed Brazilians and worldwide public datasets. The x-axis shows the 56 variants found in common between the Brazil sample and the gnomAD and 1KGP dataset. (A) Comparison between Brazilians and gnomAD, and (B) including non-Finland Europeans (NFE), sub-Saharan Africans/African Americans (AFR) Venn diagrams that show the overlap between samples (A) and variants (B) in the WES and the SNP array datasets from BIPMed reference samples.

Figure 2. Distribution of alternative allele frequency of common variants ($AAF > 0.01$) separated by Brazilian cities. The x-axis shows the 56 variants found in common with gnomAD and 1KGP. This barplot also includes the two public Brazilian datasets (BIPMed and ABraOM) and the frequency of all samples combined (Brazil).

Figure 1




Figure 2

