

## **Prevalence and correlates of anemia among children aged 6-23 months in Wolaita Zone, Southern Ethiopia**

Mihiretu Alemayehu<sup>1\*</sup>, Mengistu Meskele<sup>1</sup>, Bereket Alemayehu<sup>2</sup>, Bereket Yakob<sup>3</sup>

<sup>1</sup>Wolaita Sodo University, College of Health Sciences and Medicine, School of Public Health: PO Box: 138, Ethiopia

<sup>2</sup>Wolaita Sodo University, College of Natural and Computational Sciences, Department of Biology, Biomedical Science division: PO Box: 138, Ethiopia

<sup>3</sup>Department of Global Health and Population, Harvard T.H. Chan School of Public Health: 665 Huntington Avenue, Boston, MA 02115

Mihiretu A: mihiretua@gmail.com (Corresponding author)

Mengistu M: mengistu77@gmail.com

Bereket A: bereketalemayehu@gmail.com

Bereket Y: berekbot@yahoo.com

### **Abstract**

**Background:** Anemia, the world's most common micro-nutrient deficiency disorder, can affect a person at any time and at all stages of life, although children aged 6 -23 months are particularly at higher risk. If left untreated, it adversely affects the health, cognitive development, school achievement, and work performance. However, little was investigated among young children in Sub-Saharan countries including Ethiopia. This research aimed to investigate its magnitude and correlates to address the gap and guide design of evidence based intervention.

**Methods:** A community-based cross-sectional study was conducted from May -June 2016 in rural districts of Wolaita Zone. Multi-stage sampling technique was applied to select 990 mother-child pairs. Socio-demography, health and nutritional characteristics were collected by

administering interview type questionnaire to mothers/care-givers. Blood samples were taken to diagnose anemia by using HemoCue device, and status was determined using cut-offs used for children aged 6-59 months. Hemoglobin concentration below 11.0 g/dl was considered anemic. Data were analyzed with Statav 14. Bivariate and multivariable logistic regressions were applied to identify candidate and predictor variables respectively. Statistical significance was determined at p-value < 0.05 at 95% confidence interval.

**Results:** The mean hemoglobin level of children was 10.44±1.3g/dl, and 65.7% of them were anemic. Among anemic children, 0.4% were severely anemic (<7.0g/dl), while 28.1% and 37.2% were mildly (10.0-10.9g/dl) and moderately (7.0-9.9g/dl) anemic, respectively. In the multivariable analysis, having maternal age of 35 years and above (AOR=1.96), being government employee (AOR=0.29), being merchant (AOR= 0.43) and ‘other’ occupation (AOR= 3.17) were correlated with anemia in children in rural Wolaita. Similarly, receiving anti-helminthic drugs (AOR= 0.39), being female child (AOR=1.76), consuming poor dietary diversity (AOR=1.40), and having moderate household food insecurity (AOR=1.72) were associated with anemia in rural Wolaita.

**Conclusion:** A large majority of children in the rural Wolaita were anemic and the need for proven public health interventions such as food diversification, provision of anti-helminthic drugs and ensuring household food security is crucial. In addition, educating women on nutrition and diet diversification, as well as helping them with alternative sources of income might be interventions in the study area.

**Keywords:** Anemia; Wolaita Zone; Under-two children; Infants; Children; Malnutrition; Iron deficiency anemia

## **Background**

Anemia is the world’s most common micro-nutrient deficiency disorder that affects more than 2 billion people globally [1]. It can affect a person at any time and at all stages of life. However, in most parts of the world, children aged 6–23months are at particularly higher risk. It primarily affects infants and young children because of their higher iron requirements related to growth, and women of childbearing age due to menstrual loss and pregnancy [2].

If left untreated, anemia can adversely affect the health, cognitive development, school achievement, and work performance of individuals. Low oxygenation of brain tissues, a consequence of anemia, may lead to impaired cognitive function, growth and psychomotor development in children. Infants, under 5-year-old children and pregnant women have greater susceptibility to anemia because of their increased iron requirements for rapid body growth and expansion of red blood cells [3]. During childhood period, it is strongly associated with poor health and physical development [4, 5], mild and moderate mental retardation [6], and poor motor development and its control [7]. Iron deficiency anemia leads to reduced academic achievement and work capacity which reduces the earning potential of individuals and hence damages national economic growth at large [5]. It also increases the risk of mortality and morbidity from infectious diseases [4, 8].

Childhood anemia is mostly caused by dietary iron deficiency, infectious and genetic diseases, and other nutrient deficiencies [9]. While more than half of the anemia burden in children is attributed to iron deficiency, only very small fraction is due to genetic causes [10]. In early childhood, bad feeding habits, especially during the weaning period as breast milk is replaced by foods that are poor in iron, vitamin B12 and folic acid [11].

In Ethiopia, evidence showed varying magnitudes of anemia ranging from 41% in Amhara region to 83% in Somali region [12]. Many studies showed that factors contributing to and affecting anemia in children varied from one geographical area to another [4, 6, 7, 10]. However, most of those studies were based on small sample size, conducted on urban dwellers, were facility-based, and lacked representativeness. Besides, no published research exists which uncovered the problem in the context of Woita Zone. Hence, the present study was conducted to fill the gap in a rural setting in Wolaita Zone with an aim of determining the magnitude of anemia and associated factors among children aged 6-23 months.

## **Methodology**

### **Study design and Study area**

A community based cross-sectional study was conducted among children aged 6-23 months residing in rural districts of Wolaita Zone, Southern Ethiopia, from May to June 2016. Wolaita Zone is one of the 13 administrative zones of southern region (SNNPR) which has 12 rural and 3

urban districts. The Zone was inhabited by over 1.8 million people in 2016 [13]. Wolaita Sodo, the capital of the Zone, is located at 6° 49' N latitude and 39° 47' E longitude, at an altitude of about 1900 meters above sea level. It is located at 330 km south-west of Addis Ababa, Ethiopia. The zone is characterized by its dense population. The majority (88.3%) of its population reside in rural districts whose major livelihood is agriculture. The major food crops cultivated in the zone are maize, sweet potato, *enset* (false banana), *teff* (*Eragrostis tef*), haricot bean, taro, sorghum, Irish potato, yam and cassava [14].

## Sampling

Multi-stage sampling technique was applied to select mother-child paired study population. Children aged 6-23 months were the source population for the study. Initially, four districts were randomly selected from the 12 rural districts. Damot Gale, Boloso Bombe, Humbo and Sodo Zuria districts were selected as study districts. Then, three kebeles (the lowest administrative unit of Ethiopia consisting of nearly 5000 population) were randomly selected from each of the selected districts, making the total number of kebeles included in the study 12. Finally, the study participants were selected through systematic sampling technique from each of the selected kebeles by probability proportional to size i.e., allocating the sample size with regard to the respective kebeles' population.

The total 993 sample size needed for this study was determined by the formula to estimate a single population proportion based on the following assumptions: 71.2% prevalence of anemia in Sub-Saharan Africa [15], 95% confidence interval, 5% margin of error, and 10% non-response rate and design effect of 3.

## Data collection tools and procedures

Data were collected through interviewer administered questionnaire prepared in English and translated to local language (Wolaita Dona). The questionnaire was developed by reviewing guidelines and related literatures [12, 15, 16, 17]. It consisted of demographic characteristics variables, household wealth indicators and anemia risk factors such as health service utilization, recent illnesses, and dietary practice of both mother and child.

**Anemia diagnosis:** Hemoglobin concentration was used to determine anemia status of the study participants by taking finger-prick blood sample. Hemoglobin level was analyzed onsite by using

HemoCue device (HemoCueHb 301), and values were adjusted for altitude using the UNICEF/WHO guideline. Anemia status was determined using cut-offs used for children aged 6-59 months. Hemoglobin concentration below 11.0 g/dl was considered anemic, whereas, hemoglobin concentration of 11.0 g/dl and above was considered as normal. Severity of anemia was categorized based on the UNICEF, UNU, WHO guideline as follows: children were categorized as mildly, moderately and severely anemic if their blood hemoglobin concentration is between 10.0-10.9g/dl, 7.0-9.9g/dl and < 7.0 g/dl, respectively. Maternal anemia (hemoglobin concentration below 12 g/dl) was diagnosed with the same procedure and device used for child anemia, followed by adjustment for pregnancy and altitude [16].

**Dietary assessment:** A 24-hour dietary recall method was used to assess dietary practice. Dietary Diversity Score of children was calculated by asking mothers/caregivers about the food items their children consumed in the past 24 hours preceding the survey. All food items consumed by the children in the last 24 hours preceding the survey were categorized into seven food groups as (1) grains, roots, and tubers, (2) legumes and nuts, (3) milk and milk products, (4) flesh foods, (5) eggs, (6) vitamin-A rich fruits and vegetables, and (7) other fruits and vegetables. Finally, the food groups consumed by the child were added together to obtain dietary diversity score [17].

**Food insecurity:** Food insecurity was measured by HFIAS (Household Food Insecurity Access Scale) tool developed by FANTA (Food and Nutrition Technical Assistant) project. The tool has nine questions asking household's about the three domains of food insecurity: feeling uncertainty of food supply, insufficient quality of food, and insufficient food intake and its physical consequences in the last month. The households participating in the study were categorized into the four levels of food-security (food secure, mildly food insecure, moderately food insecure and severely food insecure) based on the guideline's recommendation [18].

**Wealth index:** Household wealth index was constructed using household asset data through PCA (Principal Component Analysis) based on interview responses adopted from Ethiopian Demographic and Health Survey. The presence or absence of each household items such as plow oxen, table, Animal-drawn cart, chair, etc. were asked and their responses were coded as '0' for No and '1' for Yes. Finally, the common factor score for each household was produced for grouping households as lower, middle and higher wealth quantile households [19].

**Under-nutrition:** chronic energy deficiency (malnutrition) was assessed using WHO guideline. The WHO Anthro 2005 software was used to calculate Z-score. MUACZ cut-off- point of negative two (-2) was used to define under-nutrition[20].

**Data management and analysis:** Data were entered into Epi-Data software version 1.4.4.0 and analyzed with Stata software version 14 (College Station, Texas). Proportions, means and standard deviations (SD) were used to describe the study population by independent variables and anemia. Bivariate logistic regression was done using thirty independent variables to identify the candidate variables (p-value < 0.25) for multivariable regression. Finally, predictors of anemia were determined using multivariable logistic regression model among the selected eleven candidate variables. Multi co-linearity, interaction and mediation among independent variables were checked based on the assumptions such as tolerance, variance inflation factors (VIF), correlation coefficient of interaction, and others to assure the fitness of logistic regression model. The independent variables used for checking multi co-linearity, interaction and mediation were; age of child, age of mothers, , religion, occupation, family size, meal frequency of mother, meal frequency of child, initiation of complementary feeding, receiving anti-helminthic drug, sex of child, dietary diversity score of child (DDS) and household food insecurity access scale (HFIAS). The statistical significance was determined at p-value<0.05 at 95% confidence interval.

**Quality control:** Interviewers and laboratory technicians were trained for two days prior to data collection. A pilot study was done among 50 children who were selected from areas outside the actual study area. The data collection was regularly supervised by trained supervisors and the investigators.

**Ethical consideration:** The actual data collection was started after Wolaita Sodo University College of health sciences and medicine approved the study. Local administrative bodies (Damot Gale, Boloso Bombe, Humbo and Sodo Zuria district health offices) were also communicated and provided permission to undertake the study. Finally, written informed consent was obtained from the mothers/caregivers. Children diagnosed with anemia were counseled and those with hemoglobin concentration below 9.0 g/dl were referred to health facility for further treatment and follow up.

## Result

### Socio-demography

A total of 990 children were included in the survey making 99.7% response rate. The mean age of children was 14.96 months with a standard deviation of 5.37 months. Around 434(43.8%) of the children were females; 293(29.6%) were born with short birth interval (less than 2 years interval); 191(19.3%) were born from young mothers (mothers aged 15-24 years);and 60 (6.1%) were raised by single parents (Table 1).

**Table 1: Socio-demographic characteristics of children aged 6-23 months residing in rural districts of Wolaita zone, 2016 (N=990)**

Variable	Category	Number	Percent
Child age (in months)	6-11	336	33.9
	12-17	289	29.2
	18-23	365	36.9
Sex of child	Male	556	56.2
	Female	434	43.8
Birth interval	First born child	221	22.3
	below 24 month	293	29.6
	25-48 month	330	33.3
	above 48	146	14.8
Age of mother (in years)	15-24	191	19.3
	25-34	685	69.2
	35 and above	114	11.5
Marital status of mother	In a union	930	93.9
	Not in a union	60	6.1
Ethnicity of mother	Wolaita	963	97.3
	Others	27	2.7
Religion of mother	Protestant	857	86.6
	Orthodox	121	12.2
	Others	12	1.2
Occupation of mother	Housewife	881	89.0
	Government employee	36	3.6
	Merchant	33	3.3
	Others	40	4.0
Occupation of father	Farmer	624	63.0
	Government employee	111	11.2

	Merchant	175	17.7
	Others	80	8.1
Home distance from health facility (foot walk)	Below 30 min	423	42.7
	30-45 min	443	44.8
	Above 45 min	124	12.5
Family size	Below 5	307	31.0
	5-7	451	45.6
	Above 7	232	23.4

Others<sup>1</sup>– Catholic, Muslim, Adventist,

Others<sup>2</sup>– Daily laborers, potters, housemaid

### Health and nutrition characteristics of children

The majority (93.1%) of children was fully immunized and only 11% of them were given anti-helminthic drugs to treat intestinal parasites. A total of 409 (41.3%) children had at least one of the following symptoms within the past two weeks preceding the survey: cough, difficulty of breathing, fever, diarrhea (with/without blood), or visible parasite on stool. Based on the Mid-upper arm circumference (MUAC) measurement, 688(69.5%)of the children were well-nourished, whereas the rest 302(30.5%) had chronic energy deficiency.

The children had mean dietary diversity score of 3.4 with 1.5 standard deviation, and only 422 (42.6%) of them consumed more than half of the recommended seven food groups. Furthermore, more than one-third (36.3%) drank either tea or coffee in their usual meal (that contained tannins which reduce iron absorption). Nearly one out of ten children stopped breastfeeding at least one week before the survey (Table 2).

**Table 2: Health and nutrition characteristics of children in rural districts of Wolaita Zone, 2016 (N=990)**

Variable	Response Category	Number	Percent
Fully vaccinated	No	68	6.9
	Yes	922	93.1
Received anti-helminthes drug	No	881	89.0
	Yes	109	11.0
Sick in the past two weeks	No	581	58.7



	Yes	409	41.3
Ever caught malaria	No	642	64.8
	Yes	348	35.2
Usually drinks tea or coffee	No	631	63.7
	Yes	359	36.3
Bottle-feed	No	884	89.3
	Yes	106	10.7
Breast-feeding(currently)	No	110	11.1
	Yes	880	88.9
Complementary feeding frequency (Within 24 hours)	1-3	507	51.2
	4-5	440	44.4
	6-8	43	4.3
Breastfeeding frequency	1-6	194	19.6
	7-8	343	34.6
	Above 8	453	45.8
Initiation of complementary feeding	Before 6 month	90	9.1
	At 6 month	614	62.0
	After 6 month	286	28.9
Dietary diversity score	Below 4	568	57.4
	4 and above	422	42.6
Nutritional status (MUACZ)	≤-3	80	8.1
	Between -2 and -3	222	22.4
	Normal	688	69.5

Based on the household food insecurity access scale measurement (HFIAS), 628 (63.4%) of the children were born and lived in food-secure households; whereas the rest were born and lived in a household suffering from food insecurity. Four hundred and four (40.8%) mothers fully attended the four WHO recommended ANC visits, while only 86(8.7%) of them had no ANC visit. The prevalence of maternal anemia was 13.4%; and three-fourth (734) of them usually consumed fewer than four meals per day (Table 3).

**Table 3: Maternal health and nutrition in rural districts of Wolaita Zone, 2016 (N=990)**

Variable	Category	Number	Percent
Household food insecurity	Food secure	628	63.4
	Mildly food insecure	78	7.9
	Moderately food insecure	208	21.0
	Severely food insecure	76	7.7
Number of pregnancies	1-2	397	40.1
	3-4	298	30.1
	Above 4	295	29.8
Antenatal care follow-up	No visit	86	8.7
	1-3 visit	500	50.5
	4 and above	404	40.8
Place of delivery	Home	446	45.1
	Health facility	544	54.9
Maternal anemia	Yes	133	13.4
	No	857	86.6
Frequency of meals/day	Less than or equal to 3	734	74.1
	Above 3	256	25.9

### **Magnitude and severity of anemia**

The mean hemoglobin concentration of the children was 10.44 g/dl (95% C.I: 10.36, 10.52 g/dl) with standard deviation of 1.30 g/dl, and with a minimum and maximum hemoglobin concentration of 6.64 to 17.31g/dl. Overall, 650 (65.7%) of the children had anemia with hemoglobin level below 11g/dl (95% C.I: 62.6% - 68.6%). Most of them were mildly (10-10.9g/dl) and moderately (7-9.9g/dl) anemic accounting for 278 (28.1%) and 368 (37.2%), respectively; whereas only 4 (0.4%) of children were severely anemic with hemoglobin level below 7g/dl (Figure 1). The prevalence of anemia decreased as age increased i.e. from as high as 72.6% among infants aged 6-11 months to as low as 61.4% among young children aged 18-23 months (Figure 1).

## Factors associated with anemia

Bivariate logistic regression showed that age of child, age of mother, religion, occupation of mother, family size, child's meal frequency, initiation of complementary feeding, receiving anti-helminthic drugs, sex of child and food insecurity were associated with anemia in children ( $P < 0.05$ ).

However, in the multivariable analysis age of child, religion, family size and initiation of complementary feeding time were not associated with child anemia ( $p > 0.05$ ); whereas maternal age and occupation, child's anti-helminthic drug intake, sex, dietary diversity score, and household food insecurity were identified as predictors of child anemia in the multivariable analysis ( $p < 0.05$ ).

The multivariable logistic regression showed that children of mothers whose age was 35 years and above were two times more likely to be anemic as compared to children of mothers whose age was between 15-24 years, AOR=1.96 (95% C.I: 1.01, 3.85),  $p=0.049$ . Children of government employee mothers had 61% lower chance of being anemic as compared to children of housewife mothers, AOR=0.29 (95% C.I: 0.09, 0.47),  $p < 0.001$ . Similarly, children of merchant mothers had 57% lower chance of getting anemia than children of housewife mothers, AOR= 0.43 (95% C.I: 0.20, 0.92),  $p= 0.029$ . In contrary, children of mothers whose occupation was classified as 'other occupation' (daily laborers, potters and housemaid) had greater chance of being anemic than children of housewives, AOR= 3.17 (95% C.I: 1.35, 7.43),  $p= 0.008$ .

Receiving anti-helminthic drugs was found to be inversely related with childhood anemia. Accordingly, children who received anti-helminthic drugs had 61% lower chance of getting anemia than their counterparts, AOR= 0.39 (95% C.I: 0.24, 0.63),  $p < 0.001$ . Female children are also twice more likely to be anemic than male children, AOR= 1.76 (95% C.I: 1.30, 2.38),  $p < 0.001$ . Nutritional characteristics are also found to be predictive factors of childhood anemia. Accordingly, children who consumed poor dietary diversity (DDS < 4) were found to be more likely to be diagnosed anemic than their counterparts, AOR=1.40 (95% C.I: 1.03, 1.92),  $p= 0.034$ . Last but not least, children from moderately food insecure households were two times more likely to be anemic as compared to those from food secure households, AOR=1.72 (95% C.I: 1.07, 2.32),  $p= 0.023$  (Table 4).

**Table 4: Multivariable analysis of factors associated with anemia among children in rural districts of Wolaita zone, 2016 (N=990)**

Variable	Category			COR	95% CI for COR		AOR	95% CI for AOR		P value for AOR
					Lower	Upper		Lower	Upper	
		Anemic	Normal							
Age of child	6-11 month	244	92							
	12-17 month	182	107	0.64*	0.46	0.90	0.77	0.52	1.12	0.175
	18-23 month	224	141	0.60**	0.44	0.82	0.76	0.52	1.11	0.154
Age of mother	15-24	113	78							
	25-34	447	238	1.30	0.93	1.80	1.52	0.98	2.36	0.061
	35 and above	90	24	2.59**	1.52	4.42	1.96*	1.01	3.85	0.049
Religion	Protestant	571	286							
	Orthodox	67	54	0.62*	0.42	0.91	0.66	0.43	1.02	0.060
	Others <sup>1</sup>	12	0	--	--	--	--	--	--	--
Occupation of mother	Housewife	592	289							
	Government employee	10	26	0.19**	0.09	0.39	0.21**	0.09	0.47	0.000
	Merchant	16	17	0.46*	0.23	0.92	0.43*	0.20	0.92	0.029
	Others <sup>2</sup>	32	8	1.95	0.89	4.29	3.17**	1.35	7.43	0.008
Family size	Below 5	195	112							
	5-7	280	171	0.94	0.70	1.27	0.81	0.55	1.21	0.311
	8 and above	175	57	1.76**	1.21	2.57	1.27	0.79	2.04	0.320
Child's meal	<4	394	113							
	4-5	241	199	0.35**	0.26	0.46	0.38**	0.27	0.52	0.000

frequency	6-8	15	28	0.15**	0.08	0.30	0.23**	0.10	0.49	0.000
Initiation of complementary feeding	< 6 month	68	22							
	At 6 month	374	240	0.50**	0.30	0.84	0.98	0.55	1.73	0.947
	> 6 month	208	78	0.86	0.50	1.49	0.96	0.53	1.75	0.903
Received anti-helminthic drug	No	609	272							
	Yes	41	68	0.27**	0.18	0.41	0.39**	0.24	0.63	0.000
Sex of child	Male	343	213							
	Female	307	127	1.50**	1.15	1.96	1.76**	1.30	2.38	0.000
Dietary diversity score	Below 4	376	192	0.95	0.73	1.23	1.40*	1.03	1.92	0.034
	4 and above	274	148							
Food insecurity	Food secure	396	232							
	Mildly insecure	48	30	0.94	0.58	1.52	0.93	0.54	1.62	0.809
	Moderately insecure	152	56	1.59**	1.12	2.25	1.57*	1.07	2.32	0.023
	Severely insecure	54	22	1.44	0.85	2.42	1.10	0.62	1.94	0.748

\*- P<0.05   \*\*- P<0.01

Others<sup>1</sup>– Catholic, Muslim, Adventist,

Others<sup>2</sup>–Daily laborers, potters, housemaid

## Discussion

Anemia remains one of the major health problems that result in grave health outcomes in developing countries despite progresses seen in nutrition interventions. The Global Burden of Diseases study [21] shows that anemia in children is one of the most common causes of child death in Ethiopia, and continues to be a major public health problem. Similarly, our study showed that two-third of the young children in rural Wolaita had anemia standing as a severe

public health problem (above the WHO cut-off point of 40%) [22]. On the other hand, studies elsewhere in Ethiopia [23] reported similar findings although severe anemia was found to be lower in the study area (0.4%) as compared to Ethiopia's national rate among young children (3.5%) [12].

UNICEF and WHO recommend adequate breastfeeding, iron supplementation and fortification, and nutrition education for mothers [24] in order to curve health loss (high morbidity and mortality) due to iron deficiency anemia in children. Ethiopia's nutrition strategy puts weight on the above nutrition intervention with the vision of ensuring adequate micronutrient intake for all children and its citizens [25]. Implementation of the national nutrition strategy with focus to young children will be vital in preventing and treating anemia in rural Wolaita.

The present study showed that maternal age, type of occupation, child's anti-helminthic drug intake, sex of child, dietary diversity score and household food insecurity were associated with child anemia.

Studies have shown that consuming diverse food prevents anemia although this is a difficult option for households in developing countries where there is recurrent food insecurity problem. In conformity to this study, several studies [23, 26, 27] have shown children who had poor dietary diversity score had a higher chance of having anemia. Similarly, WHO report has shown that feeding children diverse foods increases the bioavailability of micronutrients including iron, and this is one of the recommended practices [28]. For instance, eating iron rich animal source food items such as flesh meat, organ meat, poultry; and non-animal source food items such as legumes and green leafy vegetables increase the bioavailability of iron and enables micronutrient demands for children. Consumption of fruits, vegetables, and tubers that are good sources of vitamins A and C, and folic acid enhances the absorption and utilization of iron [24].

De-worming intestinal parasites had also a positive impact on prevention of anemia among children [24]. Similarly, in this study, children in rural Wolaita who had undergone de-worming had lower odds of having iron deficiency anemia. A report of study conducted in 25 Sub-Saharan countries stated that de-worming and iron intake for more than six months prevented anemia in children [15]. De-worming is an essential public health intervention as intestinal parasites especially hookworm infection results in intestinal blood loss which in turn, contributes to anemia.

Household food security had also a significant role on preventing anemia among children [24]. In this study, we found that food insecurity was associated with the development of anemia, which was higher among children of moderately food insecure households. Studies elsewhere showed similar finding [29], and it is important to address food insecurity problems in rural areas to prevent anemia in children of such areas. In the absence of such measures, it will be difficult to achieve global and national targets of micronutrient problems. The efforts to ensure and assist local people with food security may need implementation of programs funded by government and non-governmental organizations. This could be an important means of intervention if such programs are implemented according to the standard [24] and prioritize people with severe or moderate food insecurity problems. Nevertheless, this study did not attempt to identify households supported by such programs.

Sex of the child was also one of the factors which influenced blood hemoglobin level of children. Many studies found the correlation between sex and hemoglobin level [29, 30, 31, 32]. According to those studies, the role of sex varies from place to place which gives different gender value for the sex of child with respect to the culture of the community. In the present study, female children were more likely to have anemia than males, although precise cause could not be spelled out. Some of the reasons could be high prevalence of sex bias (negatively affecting the feminine gender), late initiation of complementary feeding for female children and the community's belief of greater need of care for male children than females [33]. Based on the finding of this study and other similar studies, female children are not favored to get iron rich food, which increases the risk of iron deficiency anemia. Therefore, it is important to educate the rural community about the importance of providing iron rich food for all children regardless of sex. Further studies may be required to validate the role of gender in iron deficiency anemia in Ethiopian context.

This study also showed that maternal age and occupation were correlated with children's hemoglobin level. The risk of children's anemia increased with the age of their mothers. In line with this finding, a study in Northeastern Brazil also showed similar trends of iron deficiency anemia [34]. On the other hand, parental occupation was also one of significant predictors of children's hemoglobin level [27]. In this study, mothers who made relatively better incomes i.e. government employee and merchant mothers had a lower chance of having anemic children as compared to housewives. This indicates that anemia is significantly prevalent among families

with less educated parents and low income, indicating the role of social inequality in the development of anemia [35]. Therefore, it will be important to ensure women with lower income either providing them opportunities to make more income or supplemental food for their children.

## **Conclusion and recommendation**

Based on the World Health Organization cut-off point, anemia is found to be a severe public health problem among young children residing in rural districts of Wolaita zone. Nearly two third of children aged 6-23 months were diagnosed to be anemic having hemoglobin level below 11g/dl. Poor dietary diversity, female sex of child, failure to take anti-helminthic drug, household food insecurity, maternal age and occupation were significantly associated with child anemia.

A large majority of children in the rural Wolaita were anemic, and the need for proven public health intervention such as food diversification, provision of anti-helminthic drugs and ensuring household food security is crucial. In addition, nutrition education and diet diversification through provision of alternative sources of income for women might be useful interventions.

## **List of abbreviations**

ANC: Antenatal Care

AOR: Adjusted Odds Ratio

C.I: Confidence Interval

COR: Crude Odds Ratio

EDHS: Ethiopian Demographic and Health Survey

FANTA: Food and nutrition technical assistant

Hb: Hemoglobin

HFIAS: Household Food Insecurity Access Scale

MUACZ: Mid Upper Arm Circumference Z-score



PCA: Principal Component Analysis

SD: Standard Deviation

SNNPR: Southern Nations, Nationalities and Peoples Region

UNICEF: United Nations Children's Fund

WHO: World Health Organization

## **Declarations**

### **Ethical approval and consent to participate**

The actual data collection was started after Wolaita Sodo University College of Health Sciences and Medicine approved the study. Local administrative bodies were also communicated about the study and permission was obtained ahead of the study. Finally, informed written consent was obtained from the mothers/caregivers. Children diagnosed with anemia were counseled and those with hemoglobin concentration below 9.0 g/dl were referred to health facility for further treatment and follow up.

### **Consent for publication**

Consent to publish this manuscript from the participants was deemed not applicable since the manuscript does not contain any individual person data.

### **Availability of data and material**

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

### **Competing interest**

All authors declared that they have no competing interests.

### **Funding**

This research was funded by Wolaita Sodo University. The university supported data collection process.

### **Authors' contribution**

MA: conceived the study and designed the experiments. MA and MM: obtained ethical clearance, and supervised data collection. MA, BA and BY Performed statistical analysis, interpreted the data and revised the article. All authors read and approved the final manuscript.

### **Acknowledgements**

We would like to acknowledge Wolaita Sodo University for financial support for field work. We also extend our gratitude to respective district administrative bodies, data collectors, data clerk and study participants for their cooperation and technical assistance.

### **Authors' Information**

MA has Masters of Public Health (MPH) in reproductive Health specialty and works at Wolaita Sodo University as assistant professor. MM has Masters of Public Health (MPH) in reproductive Health specialty and works at Wolaita Sodo University as assistant professor. BA has Masters of Science (MSc) in Biomedical Sciences and works as Associate professor at Wolaita Sodo University. BY has PhD in Public Health, and he is a Visiting Scientist at the Department of Global Health and Population, Harvard T.H Chan School of Public Health.

### **References**

1. WHO. Micronutrient deficiency: Battling iron deficiency anemia: the challenge. 2004. Available from: <http://www.who.int/nut/ida.htm>, accessed on December, 2016.
2. Lozoff B. Iron deficiency and child development. *Food Nutr Bull.* 2007; 28:S560–71.
3. Walter T, de Andraca I, Chadud P, Perales CG. Iron deficiency anemia: adverse effects on infant psychomotor development. *Pediatrics.* 1989; 84(1): 7-17.
4. McLean E, Cogswell M, Egli I, Wojdyla D, De Benoist B. Worldwide prevalence of anemia, WHO Vitamin and Mineral Nutrition Information System, 1993–2005. *Public Health Nutrition.* 2009;12(4):444–454. doi:10.1017/s1368980008002401.

5. Victora CG, Adair L, Fall C, et al. Maternal and child undernutrition: consequences for adult health and human capital. *The Lancet*. 2008;371(9609):340–357. doi: 10.1016/s0140-6736(07)61692-4.
6. Shafir T, Angulo-Barroso R, Su J, Jacobson S W, Lozoff B. Iron deficiency anemia in infancy and reach and grasp development. *Infant Behavior and Development*. 2009;32(4):366–375. doi: 10.1016/j.infbeh.2009.06.002.
7. Cardoso M A, Scopel K K G, Muniz P T, Villamor E, Ferreira MU. Underlying factors associated with anemia in amazonian children: a population-based, cross-sectional study. *PLoS ONE*. 2012;7(5) doi: 10.1371/journal.pone.0036341.e36341.
8. Pollitt E. Early iron deficiency anemia and later mental retardation. *The American Journal of Clinical Nutrition*. 1999;69(1):4–5.
9. Simon B, Willis A, Rachel P, Benson E, Siân EC, et al. Epidemiology of Plasmodium-Helminth Co-Infection in Africa: Populations at risk, potential impact on anemia, and prospects for combining control. *Am J Trop Med Hyg*. 2007; 77(6): 88–98.
10. Stoltzfus R, Mullany L, Black RE. Iron deficiency anemia. In: Ezzati M, Lopez A, Rodgers A, Murray CJL, eds. *Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors*. Geneva, World Health Organization, 2004, 163–210.
11. Torres MA, Sato K, Queiroz SS. Anemia in children under 2 years in basic health care units in the State of São Paulo, Brazil. *Rev Saúde Pública*. 1994; 28(4): 290-4.
12. Central Statistical Agency [Ethiopia] and ICF International. *Ethiopia Demographic and Health Survey, 2016*. Addis Ababa EaC, Maryland, USA: Central Statistical Agency and ICF International.
13. *Summary and statistical report of the 2007 population and housing census, Federal democratic republic of Ethiopia, 2008*.
14. *Prospects for integrating food and food production in Wolaita Sodo, Ethiopia*: available at <http://www.fao.org/wairdocs/ILRI/x5519B/x5519b19.htm> accessed by Nov. 2016
15. Calistus W, Shiro T, Fabian E, Koji K. Prenatal anemia control and anemia in children aged 6-23 months in Sub-Saharan Africa. *Matern Child Nutr*. 2016; 1–10. Doi 10.1111/mcn12375.

16. Iron deficiency anemia assessment, prevention, and control: A guide for program managers. UNICEF, UNU, WHO, 2001 (WHO/NHD/01.2). Available at <http://www.who.int/reproductive-health/docs/anaemia.pdf> accessed by Feb 2016.
17. World Health Organization, Indicators for Assessing Infant and Young Child Feeding Practices, WHO Press, Geneva, Switzerland, 2010.
18. Jennifer C, Anne S, Paula B. Household Food Insecurity Access Scale (HFIAS) for Measurement of Household Food Access: Indicator Guide. Washington, D.C.: Food and Nutrition Technical Assistance Project: 2007. Available at [http://www.fao.org/fileadmin/user\\_upload/eufao-fsi4dm/doc-training/hfias.pdf](http://www.fao.org/fileadmin/user_upload/eufao-fsi4dm/doc-training/hfias.pdf). Accessed by June 2016.
19. Central Statistical Agency [Ethiopia] and ICF International. 2012. Ethiopia Demographic and Health Survey 2011. Addis Ababa EaC, Maryland, USA: Central Statistical Agency and ICF International.
20. World Health Organization, “Physical status: the use and interpretation of anthropometry: report of a WHO expert committee,” WHO Technical Report Series 854, World Health Organization, Geneva, Switzerland, 1995, [http://www.who.int/childgrowth/publications/physical\\_status/en/](http://www.who.int/childgrowth/publications/physical_status/en/).
21. Awoke M, Tilahun NH, Kebede D, Gizachew AT, Amare D. National mortality burden due to communicable, non-communicable, and other diseases in Ethiopia, 1990–2015: findings from the Global Burden of Disease Study. *Population Health Metrics*: 2015; 15(29): DOI: 10.1186/s12963-017-0145-1
22. World Health Organization. Iron deficiency anemia; assessment, prevention and control. A guide for program managers Geneva: World Health Organization; 2005. Available from: [www.who.int/nutrition/publications/en/ida\\_assessment\\_prevention\\_control](http://www.who.int/nutrition/publications/en/ida_assessment_prevention_control).
23. Woldie H, Kebede Y, Tariku A. Factors Associated with Anemia among Children Aged 6-23 Months Attending Growth Monitoring at Tsitsika Health Center, Wag-Himra Zone, Northeast Ethiopia. *J NutrMetab*. 2015; 2015: 928632. doi: 10.1155/2015/928632.0
24. Iron deficiency anemia assessment, prevention, and control: A guide for program managers. UNICEF, UNU, WHO, 2001 (WHO/NHD/01.2). Available at <http://www.who.int/reproductive-health/docs/anaemia.pdf> accessed by Feb 2016.

25. The federal Democratic Republic of Ethiopia. National Nutrition Strategy. Available at [http://extranet.who.int/nutrition/gina/sites/default/files/ETH%202008%20National%20Nutrition%20Strategy\\_1.pdf](http://extranet.who.int/nutrition/gina/sites/default/files/ETH%202008%20National%20Nutrition%20Strategy_1.pdf) accessed on June 2017
26. Mahama S. Relationship between Dietary Diversity and Haematological Status of Children aged 6-59 months in Northern Ghana: *Nessa J Nutritional Health Sciences*. 2016;1(1); 1-23
27. Desalegn A, Mossie A, Gedefaw L. Nutritional Iron Deficiency Anemia: Magnitude and Its Predictors among School Age Children, Southwest Ethiopia: A Community Based Cross-Sectional Study. *PLoS ONE*. 2014; 9(12): e114059. doi:10.1371/journal.pone.0114059
28. World Health Organization and Working Group on Infant and Young Child Feeding Indicators, *Developing and Validating Simple Indicators of Dietary Quality and Energy Intake of Infants and Young Child in Developing Countries*, World Health Organization, Washington, DC, USA, 2007.
29. Sant-R, James B, Sumithra M, Anita S, Vijay B, et al. Determinants of anemia among young children in rural India. *Pediatrics*. 2010; 126(1):e140. doi: 10.1542/peds.2009-3108.
30. Michael OF, Frank OF, Michael OA. Factors Associated with Hemoglobin Prevalence among Ghanaian Children Aged 6 – 59 months. *Journal of Biology, Agriculture and Health care*; 2014. 4(2); 132-140.
31. Monica MO, Pedro IC, Malaquias B, Ann A. Prevalence of anemia in children 6–59 months old in the state of Pernambuco, Brazil. *Pan Am J Public Health* 2001; 10(2): 101-107.
32. Chitra B, Nisha N, Jesni KJ, Jesina B, Nazar A, et al. Study on Prevalence of Anemia among School Children in a Rural Community Setup. *Sch. Acad. J. Pharm*. 2014; 3(6): 423-426.
33. Agumasie S, Gezahegn T, Alemayehu B. Complementary feeding practice of mothers and associated factors in HiwotFana Hospital, Eastern Ethiopia. *Pan Afr Med J*. 2014; 18: 143. Doi: 10.11604/pamj.2014.18.143.3496.
34. Luciana PL, Malaquias BF, Pedro IC, José NF, Monica MO. Prevalence of anemia and associated factors in children aged 6-59 months in Pernambuco, Northeastern Brazil. *Rev. Saúde Pública*. 2011; 45(3): 457-466

35. Malkanthi R, Silva K, Jayasinghe-Mudalige U. Risk factors associated with high prevalence of anemia among children under 5 years of age in paddy-farming households in Sri Lanka. *Food and Nutrition Bulletin*. 2010; 31(4): 475-482.




Figure 1: Magnitude and severity of anemia among children aged 6-23 months in rural districts of Wolaita Zone, 2016 (N=990)