

Chemogenetic inhibition of the amygdala modulates emotional behavior expression in infant rhesus monkeys

Abbreviated title: Infant Chemogenetic Amygdala Inhibition

*Jessica Raper¹; Lauren Murphy^{1,2}; Rebecca Richardson¹; Zoe Romm³; Zsafia Kovacs-Balint¹; Christa Payne⁴; Adriana Galvan^{1,5}

1 Yerkes National Primate Research Center, Emory University, Atlanta, GA 30329

2 Department of Psychology, Emory University, Atlanta, GA 30329

3 Drexel University, Philadelphia, PA 19104

4 Marcus Autism Center, Atlanta, GA 30329

5 Department of Neurology, School of Medicine, Emory University, Atlanta, GA 30329

***Corresponding Author:** Dr. Jessica Raper, Yerkes National Primate Research Center, 954 Gatewood Rd NE, Atlanta, GA 30329. Email: jraper@emory.edu

Number of Pages 34

Number of Figures 5

Number of Tables 1

Conflicts of Interest: None, authors have no conflicts of interest to report.

Acknowledgements: Authors would like to thank Dr. Jocelyne Bachevalier for assistance with viral vector surgery, as well as Jean-Francois Pare and Susan Jenkins for expert technical help. Thanks to Dr. Katalin Gothard for the use of her conspecific video stimuli, as well as Shanice Wilson, B.S., Madi Seaver, Noa Shapiro-Fanklin, and Sahrudh Dharanedra for drawing areas of interests on eyetracking stimulus videos. Many thanks to Drs. Ami Klin and Warren Jones for supporting this research. Thanks to Dr. Thomas Vanderford and Oliva Delmas for conducting the neutralizing antibody assay.

Funding Sources: National Institute of Mental Health (P50-MH081756). Yerkes National Primate Research Centre is supported by NIH/Office of the Director P51-OD011132.

Author Contributions: JR, LM, and CP Designed Research; JR, LM, RR, ZR, and AG Performed Research; JR, AG, and ZK-B Analyzed Data; JR, LM, CP, and AG Wrote Paper

Abstract

Manipulation of neuronal activity during the early postnatal period in monkeys has been largely limited to permanent lesion studies, which can be impacted by developmental plasticity leading to reorganization and compensation from other brain structures that can interfere with the interpretations of results. Chemogenetic tools, such as DREADDs (designer receptors exclusively activated by designer drugs), can transiently and reversibly activate or inactivate brain structures, avoiding the pitfalls of permanent lesions to better address important developmental neuroscience questions. We demonstrate that inhibitory DREADDs in the amygdala can be used to manipulate socioemotional behavior in infant monkeys. Two infant rhesus monkeys (1 male, 1 female) received AAV5-hSyn-HA-hM4Di-IRES-mCitrine injections bilaterally in the amygdala at 9 months of age. DREADD activation after systemic administration of either clozapine-N-oxide or low dose clozapine resulted in decreased freezing and anxiety on the human intruder paradigm and changed the looking patterns on a socioemotional attention eyetracking task, compared to vehicle administration. The DREADD-induced behaviors were reminiscent of, but not identical to, those seen after permanent lesions of the amygdala in infant monkeys, such that early amygdala lesions produce a more extensive array of behavioral changes in response to the human intruder task that were not seen with DREADD-evoked inhibition of this region. Our results support the notion that early permanent damage leads to brain reorganization manifesting in a broader impact on behavior. The current study provides a proof-of-principle that DREADDs can be used in young infant monkeys to transiently and reversibly manipulate behavior.

Statement of Significance

Many neurodevelopmental disorders exhibit abnormal structural or functional amygdala development and alterations in socioemotional behavior. To date, developmental neuroscience studies have relied on permanent lesions techniques to investigate how atypical amygdala development impacts socioemotional behaviors, which may not adequately recapitulate the role of amygdala dysfunction in the manifestation of aberrant behavior. The present study sought to demonstrate that the DREADDs (designer receptors exclusively activated by designer drugs) chemogenetic tool could transiently inhibit amygdala activity in infant monkeys resulting in alterations in socioemotional behavior. This proof-of-principle study supports the use of chemogenetics for developmental neuroscience research, providing an opportunity to broaden our understanding of how changes in neuronal activity across early postnatal development influences behavior and clinical symptoms.

Introduction

The amygdala plays an important role in social behavior and emotional responses to threats in the environment (Aggleton, 2000). This knowledge has largely come from research on adult humans and animals. Yet, the amygdala has a protracted postnatal development (Payne et al., 2010; Chareyron et al., 2012; Hunsaker et al., 2014), in which growth of the amygdala appears to coincide with refinement of social and threat detecting skills (Mendelson, 1982; Mendelson et al., 1982.; Kalin et al., 1991). Investigations of how amygdala development contributes to these skills have largely utilized permanent lesion techniques, finding that damage to the amygdala during the early postnatal period contributes to life-long changes in social, emotional, and neuroendocrine function (Bauman et al., 2004; Bliss-Moreau et al., 2010; Bliss-Moreau et al., 2011a; Bliss-Moreau et al., 2011b; Raper et al., 2013b; Raper et al., 2014b; Raper et al., 2014a; Stephens et al., 2015; Bliss-Moreau et al., 2017; Moadab et al., 2017). Given the potential for neural plasticity during this postnatal period, there is considerable opportunity for reorganization and compensation from other brain structures. In fact, several studies have demonstrated how lesions early in life contribute to such reorganization (Machado et al., 2008; Raper et al., 2014b; Grayson et al., 2017; Payne et al., 2017). In order to avoid the pitfalls of permanent lesions and better address these questions we need tools that transiently activate or inactivate the amygdala during early development.

Pharmacological inactivation techniques have been used to manipulate amygdala activity and influence behavior (Wellman et al., 2005; Wellman et al., 2016). However, these techniques require the surgical placement of skull-mounted chambers for intracerebral drug infusions, which are not ideal for developmental research on infant monkeys with rapidly growing brains and bodies. An alternative approach that would not require permanently mounted hardware nor rely on repeated intracerebral injections is greatly needed for developmental research.

Chemogenetic techniques provide powerful tools for behavioral neuroscience because they allow remote manipulation of neuronal activity (Sternson and Roth, 2014). Designer receptors exclusively activated by designer drugs (DREADDs) are the most commonly used chemogenetic tool. DREADDs involves the use of artificial G protein-coupled receptors that are synthetic variants of muscarinic receptors (Armbruster et al., 2007; Farrell and Roth, 2013; Roth, 2016), are not activated by acetylcholine (the endogenous ligand of muscarinic receptors) and are not constitutively active. Instead, DREADDs are exclusively activated with high efficacy by exogenous ligands. The available DREADDs include receptors that are coupled to Gi/o, Gq/11 and Gs proteins termed M4Di, M3Dq and GsD, respectively (Wess et al., 2013; Roth, 2016), providing tools that can either increase or decrease the activity of transduced neurons, depending on the intracellular pathway activated. The expression of DREADDs can be targeted to specific neurons using viral transfection (Urban and Roth, 2015; Roth, 2016). The most commonly used ligand to activate DREADDs is clozapine-N-oxide (CNO), although recent reports have shown that CNO can be reverse metabolized into clozapine (Raper et al., 2017; Allen et al., 2018). Thus, an alternative method to active the DREADDs is to use clozapine itself, at low doses that will only preferentially activate the DREADDs (Gomez et al., 2017).

DREADDs have been widely used in rodent models, but reports in nonhuman primates (NHPs) have been limited. To date, only four publications have reported DREADDs manipulation of behavior in NHPs (Eldridge et al., 2016; Grayson et al., 2016; Nagai et al., 2016; Upright et al., 2018). While these studies provide critical information demonstrating the ability to use chemogenetic tools in NHPs to manipulate neuronal activity and impact behavior, they have all focused on adult monkeys. Considering its advantages over permanent lesions and other transient inactivation techniques (i.e. reversibility and minimal invasiveness), chemogenetics hold their greatest promise for developmental neuroscience research. The current study provides proof-of-principle that chemogenetic tools can be used to manipulate amygdala activity

in infant monkeys.

Materials and Methods

Subjects:

Two infant Indian rhesus macaques (*Macaca mulatta*), one male and one female, were used in this study. The animals were raised by their mothers in large social groups at the Yerkes National Primate Research Center (YNPRC) Field Station (Lawrenceville, GA) until 7 months of age, when they were transferred to indoor pair-housing at the YNPRC Main Station (Atlanta, GA) for the duration of the study. The Animal Care and Use Committee of Emory University approved all procedures, which were performed in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals. Figure 1 outlines the study design, behavioral testing, and timing of ligand administrations.

Screening for AAV Neutralizing antibodies: Nonhuman primates' native immunity has been shown to impact transduction in other CNS tissues (i.e. eye (Kotterman et al., 2015)) and neutralizing antibody response to multiple virus solution intracerebral injections have been directly linked to weak transduction in neurons (Mendoza et al., 2017). However, it is currently unknown whether native immunity to AAVs can impact transduction of neuronal tissue. Out of an abundance of caution, infant monkeys were screened for neutralizing antibodies to AAV5 prior to surgery to reduce the possibility that such antibodies might interfere with viral transduction.

A 2ml blood samples were collected in tubes containing EDTA (3.5 mg), centrifuged at 2500 rcf for 15 min, and plasma was pipetted off for assay by the Virology Core at the YNPRC (Atlanta, GA). AAV5 neutralization titers were determined by 50% infection inhibition assay (IC50) on HeLa cells (ATCC, Manassas, VA) freshly infected with Adenovirus serotype 5 (Ad5, ATCC,

Manassas, VA). Briefly, plasma samples were heat inactivated for 1 hour at 56°C and diluted to 1:5, 1:20, 1:80, and 1:240. Plasma dilutions were then mixed with AAV5 bearing a luciferase expression construct (AAV5-luc, University of North Carolina Vector Core, Chapel Hill, NC) and incubated for 1 hour at 37°C. After initial incubation, plasma/AAV5-luc mixtures were added to the Ad5-infected HeLa cells and incubated at 37°C in 5% CO₂ incubator for 48 hours. Luciferase expression was determined using the Bright-Glo Luciferase Assay system (Promega, Madison, WI) according to the manufacturer's protocol, and measured on a BioTek Gen5 luminometer. IC50 was calculated relative to negative serum controls. Results revealed that neutralizing antibody titers for AAV5 were below the level of detection in the male and 1:20 for the female.

Surgery:

At 9 months of age, animals underwent DREADDs transduction surgery. Injection of inhibitory DREADDs bilaterally into the amygdala followed procedures similar to those described in previous studies of amygdala lesions in monkeys (Raper et al., 2014b). Briefly, on the day of surgery, the infants were removed from their cagemate, sedated with ketamine hydrochloride (1 mg/kg) and maintained with isoflurane (1–2% to effect). Their head was secured in a nonferromagnetic stereotaxic apparatus, and T1-weighted MRI sequences were acquired using a Siemens 3.0T/90 cm whole-body scanner and a 3 inch circular surface coil. A high resolution T1-weighted scan [spin-echo sequence, echo time (TE) = 11 ms, repetition time (TR) = 450 ms, contiguous 1 mm section, 12 cm field-of-view (FOV), 256 X 256 matrix] obtained in the coronal plane was used to determine the coordinates of injection sites in the amygdala. An additional T1-weighted MRI was acquired at 4 months post-surgery in the context of other studies (not reported here), following these same procedures.

All surgical procedures were performed under aseptic conditions, an intravenous drip (0.45% dextrose/0.9% NaCl) was placed to maintain normal hydration, and vital signs (heart rate,

respirations, blood pressure, expired CO₂) were monitored throughout the surgery. Nolvasan solution was used to disinfect the scalp and a local anesthetic (Bupivacaine 0.25% concentration, 1.5 ml) was injected subcutaneously along the midline to reduce the pain during skin incision. After the skin and underlying connective tissue were gently displaced laterally, two small craniotomies were made in front of bregma and above the amygdalae, and the dura was cut and retracted to expose the brain. Animals received viral vector injections of AAV5-hSyn-HA-hM4Di-IRES-mCitrine (6×10^{12} vg/ml; Duke Viral Vector Core, Duke University, Durham, NC) administered in eight sites within the center of each amygdala using 10 μ l Hamilton syringe. Needles were lowered simultaneously in both hemispheres, 5 μ l were injected at each site for a total of 40 μ l manually injected at a rate of 0.4 μ l/min. After each injection, a 3 min waiting period was allotted to minimize viral spread during needle retractions.

At the completion of the surgical procedures, the dura was closed with silk sutures, the bone opening was covered with Surgicel NU-KNIT (absorbable hemostat), and connective tissues and skin were closed. The animal was removed from anesthesia, and placed in a temperature controlled incubator ventilated with oxygen until full recovery from anesthesia. All animals received banamine (1 mg/kg for 3 d), dexamethasone (0.5 mg/kg for 3 d), and antibiotic (rocephin, 25 mg/kg for 7 d) after surgery to prevent pain, edema, and infection, respectively. Behavioral testing began 5 weeks after surgery to allow animals to recover and provide adequate time for DREADD expression.

Administration of Ligands and Plasma Assays:

All ligands were administered through subcutaneous injection in the interstitial space between the shoulder blades of the animals. CNO was administered at 10mg/kg body weight, whereas clozapine was administered at 0.1 μ g/kg body weight. CNO powder (NIMH C-929, RITI International) was stored at -20 °C, protected from light. CNO solutions were freshly prepared

on each experimental day, by dissolving the drug in 100% dimethyl sulfoxide (DMSO, Sigma-Aldrich, St. Louis MO, USA) and then diluting with calcium and magnesium-free phosphate-buffered saline (PBS, Corning, Corning, NY, USA) to a final concentration of 10 mg/mL in 15% DMSO. To prevent degradation due to light exposure, vials and syringes that contained CNO solution were wrapped in aluminum foil. Clozapine powder (Tocris, Minneapolis, MN) was stored at room temperature, protected from light. Clozapine solutions were also freshly prepared on each day of experiment, using the same dilution methods described for CNO. Percentages of DMSO and PBS remained constant across all injections to avoid any potential confound that might arise from differences in DMSO concentrations per injection. Vehicle injections consisted of 85% PBS and 15% DMSO.

Animals were trained for awake blood collection from the saphenous vein, following established protocols at the YNPRC (Raper et al., 2014b). On behavioral testing days, blood samples were collected at various times after ligand administration to examine plasma concentrations of the ligand during behavioral tasks. For CNO administration, blood was collected at 20, 60, 90 and 135min post-injection. For clozapine administration, blood was collected at 2, 40, and 60 minutes post-injection. All blood samples (1-2ml) were collected in prechilled 2 ml tubes containing EDTA (3.5 mg) and immediately placed on ice. Samples were centrifuged at 2500 rcf for 15 min in a refrigerated centrifuge (at 4 °C). Plasma was pipetted off and stored at -80 °C until assayed.

All plasma samples were assayed for CNO and clozapine by Precera Bioscience (Franklin, TN) using previously published techniques (Raper et al., 2017). Briefly, liquid chromatography–tandem mass spectroscopy (LC-MS/MS) analyses were performed via reverse phase chromatography using a Shimadzu Nexera X2 UPLC (Columbia, MD) coupled with a QTrap 5500 (Sciex, Framingham, MA, USA). For plasma samples, 20 µL aliquots of each

standard, quality control sample, plasma sample, blank, and double blank were transferred to a labeled 96-well plate. Next, 120 μ L of 50 ng/mL internal standard (carbamazepine in acetonitrile) was added to standards, quality controls, plasma samples, and blanks, while 120 μ L of acetonitrile was added to the double blank, then the plate was centrifuged for 5 min at 3000 rcf. Then, 75 μ L of supernatant was transferred into a labeled 96-well plate, and 75 μ L of water was added to all samples. Finally, the plate was heat-sealed for analysis. The calibration range was 1–5000 ng/mL for CNO and 1–1000 ng/mL for clozapine.

Histology and Visualization of DREADDs:

Perfusion and initial preparation of tissue: At the end of the behavioral experiments, the animals received an overdose of pentobarbital, and were transcardially perfused with Ringer's solution, followed by 4% paraformaldehyde and 0.1% glutaraldehyde in phosphate buffer (PB, 0.2 M, pH 7.4). After removing the brain from the skull, we collected vibratome coronal sections (60 μ m) in cold phosphate-buffered saline (PBS, 0.01M, pH 7.4), that were stored at -20° C in an anti-freeze solution (30% ethylene glycol/30% glycerol in PB), until further processing.

Staining of Nissl substance was done to identify the localization of the injection tracks. To reveal the localization of hM4Di, we used antibodies against the hemagglutinin (HA) tag, which is fused to hM4Di. We selected brain sections containing or adjacent to the virus solution injection tracks, pretreated with 1% normal goat serum, 1% bovine serum albumin and 0.3% Triton X-100, and then incubated for 24 hours in solutions containing either HA-tag antibodies (raised in rabbit, clone C29F4, 1:400, cat # 3724, Cell Signaling Technology, Danvers, MA. As controls, some sections were processed in solutions without the primary antibodies. Incubation in the primary antibodies was followed by 2-hour incubation in anti-rabbit secondary biotinylated antibodies (1:200, Vector Laboratories, Burlingame, CA), and then in avidin-biotin-peroxidase complex (ABC) solution (1:200; Vectastain standard kit, Vector) for 90 min. The sections were

then placed in 0.025% 3-3'-diaminobenzidine tetrahydrochloride (DAB, Sigma-Aldrich, St. Louis, MO), 0.01M Imidazole (Fisher Scientific, Pittsburgh, PA) and 0.006% H₂O₂ for 10 min. The sections were mounted on slides, cover-slipped, and digitized with an Aperio Scanscope CS system (Aperio Technologies, Vista, CA).

To estimate the extent of the brain regions expressing the hM4Di, we used a series of HA-Tag sections (0.5 mm apart) that encompassed the transfected amygdaloid regions. In digital images (2X), we delineated regions with HA-positive cell bodies, and calculated the area using ImageJ (<http://imagej.nih.gov/ij/>).

Behavioral Assessments:

Human Intruder Paradigm: The human intruder paradigm was chosen because it has been shown to be a robust task for detecting differences in emotional responses of nonhuman primates after adult or neonatal amygdala lesions (Kalin et al., 2004; Machado and Bachevalier, 2008; Raper et al., 2013b; Raper et al., 2013a). Animals' responses to the human intruder were assessed under CNO prior to transduction, then again under CNO at 1- and 3-months post-transduction, or under vehicle conditions at 2- and 6-months post-transduction. Clozapine activation of the DREADDs was tested at 5-months post-transduction (Figure 1). On test days when CNO was administered, behavioral testing began approximately 80-90 minutes after injection based on the peak of CNO in cerebrospinal fluid in adult rhesus monkeys (Raper et al., 2017). On test days when clozapine was administered, testing began approximately 5-10 minutes after injection because clozapine readily crosses the blood brain barrier (Baldessarini et al., 1993). During testing, monkeys were separated from their cagemate, transported to a testing room, and transferred to a stainless-steel testing cage (53 X 53 X 55 cm) with one wall made of clear plastic to allow for unobstructed video recording. The human intruder paradigm consisted of three conditions (alone, profile, stare) presented in the same order to all monkeys

for a total duration of 30 min. The experimenter wore a rubber mask depicting a different male face at each test, such that both monkeys saw the same novel human intruder at each time point in the study. Other attempts to ensure novelty of the test throughout the six sessions, included the experimenter wearing a different colored lab jacket and either different curtains hung around the room or using a new room to make the environment less familiar. The human intruder test session started with the monkey remaining alone in the cage for 9 min (alone condition) to acclimatize to the environment and obtain a baseline level of behavior. Then the intruder entered the room and sat 2 m from the test cage, presenting his/her profile to the animal for 9 min (profile condition or no eye contact condition in other experiments). The intruder then left the room while the monkey remained in the cage for a 3 min period, then the intruder re-entered the room, and sat 2 m from the cage, making direct eye contact for 9 min (stare condition). Emotional behavior responses to the intruder were assessed using Observer XT 14 software (Noldus Inc., Wageningen, The Netherlands) and a detailed ethogram (Table 1). One experimenter (JR) with a high degree of both intra-rater reliability (Cohen's $\kappa = 0.98$) and inter-rater reliability (Cohen's $\kappa = 0.86$) with other trained experimenters at YNPRC coded all of the videotapes. The experimenter was blind to the animals' ligand administration while coding the videos.

Socioemotional Attention Task: Damage to the amygdala has been shown to alter attention toward social cues as well as threats in the environment (Aggleton, 2000; Adolphs et al., 2005; Feinstein et al., 2011; Gothard et al., 2018; Payne and Bachevalier, 2019). To measure changes in attention to social and innate aversive stimuli, subjects' eye movements were monitored using a Tobii T60/T120 eye tracker (Tobii Technology, Mountain View, CA). Eye tracking technology uses noninvasive infrared light reflections on the subjects' cornea and retina to measure looking behavior. These reflections were calibrated using a 5-point calibration paradigm at the start of each testing session to control for subtle changes in distance and head

position. Subjects were acclimated to an infant nonhuman primate chair (Crist Instrument Co., Hagerstown, MD) and positioned 22" from the eye tracker monitor on which videos were displayed. During test sessions, an experimenter running the eye tracker was visually separated from the monkey by a curtain. The experimenter could monitor whether the tracker was detecting the eyes, and a gaze-trail was displayed over a view of the presented stimuli on the experimenter's laptop.

Each test session consisted of a novel set of 54 10-second videos containing Social, Aversive, or Neutral images. Since subjects were not restricted to facing the monitor, the videos were presented twice making the testing session 20 minutes. Social videos contained clips of an unfamiliar monkey looking toward the camera expressing either a threatening, affiliative, or neutral expression (Mosher et al., 2011; Mosher et al., 2014). Aversive videos contained clips of snakes or spiders, which monkeys innately fear (Mineka et al., 1980; Mineka, 1987; Izquierdo et al., 2005; Chudasama et al., 2009). Neutral videos contained clips of enrichment food items (e.g. oranges, grapes, popcorn) or novel items (e.g. train, flower, hot air balloon). Animals were tested on this socioemotional attention eye tracking task under CNO at 3.5 months, under vehicle conditions at 3.5, 4, and 5.5 months, or under clozapine at 5 months post-transduction (Figure 1). Timing between ligand administration and the beginning of behavioral testing was the same as reported for the human intruder task.

Gaze information was recorded at a rate of 60Hz on a Windows laptop running Tobii Studio software 3.2.2 (Tobii Technology, Mountain View, CA). Several areas of interest (AOIs) were drawn on all video stimuli prior to data collection using the Tobii Studio software. For Social videos, regions were defined as eyes (bridge of nose to brow), mouth (nose to chin), face (eyes + mouth), body (outline of monkey body), and background (total viewable movie area). For Aversive and Neutral videos, regions were defined as foreground (outline of object) and

background (total viewable movie area). The primary output measure was fixation duration for each pre-drawn AOI. Fixation duration for each AOI was transformed according to the proportion of time spent in the containing AOI. For example, the average fixation duration in the mouth AOI was divided by the average fixation duration of the entire face AOI (which contains the mouth AOI). The average proportion of fixation duration at each time point was then averaged across the two presentations of the stimuli and for each category (Social, Aversive, Neutral).

Statistical Analyses:

Pharmacokinetic parameters (i.e. area under the curve (AUC)) for CNO and clozapine were determined using Microsoft Office Excel (Microsoft Corporation, Redmond, WA) and results were graphed using GraphPad Prism 7.02 (GraphPad Software Inc., La Jolla, CA).

To determine whether CNO or clozapine activation of inhibitory DREADDs modulated emotional behavior on the human intruder task or looking patterns on the socioemotional attention task, we used an analysis of variance (ANOVA) with ligand (CNO/clozapine, Vehicle/CNO prior to transduction) as the between subjects factor. All behavioral data were analyzed using SPSS 24 for Windows (IBM Corporation, Armonk, NY), significance was set at $p < 0.05$, and effect sizes were calculated using partial eta squared (η_p^2).

Results

Plasma Concentrations of Ligands:

Although CNO is the most widely used ligand for activating the DREADDs, it has been shown to have poor penetration of the blood brain barrier and can be metabolized into clozapine (Gomez et al., 2017; Raper et al., 2017; Allen et al., 2018). Importantly, clozapine has a high affinity for the DREADDs and has been shown to preferentially activate these receptors after low dose

administration (Armbruster et al., 2007; Gomez et al., 2017). To help interpret DREADD mediated behavioral effects, we collected plasma samples before and after behavioral testing sessions with CNO or low dose clozapine administrations. The two infant monkeys had similar pharmacokinetic profiles for 10mg/kg doses of CNO (Figure 2A). Infants plasma concentrations of CNO were high throughout sample collections with an average AUC value of 171.66 ng/ml·h. As observed previously in adults (Raper et al., 2017), infants also readily metabolized CNO into clozapine, which was detectable starting at 20minutes post-injection with an average AUC value of 2.03ng/ml·h. Since developmental changes in physiology can impact drug pharmacokinetics and pharmacodynamics (Milsap and Jusko, 1994; Lu and Rosenbaum, 2014), it is important to note that the conversion ratio of CNO to clozapine in infant monkeys (1.1%) was similar to that of adult monkeys (1.3%; (Raper et al., 2017)). Low dose clozapine (0.1mg/kg) administration also exhibited a similar pharmacokinetic profile between the two infant monkeys, however clozapine was not detectable until the second sample collection at 40 minutes post-injection, and plasma concentrations of clozapine were below levels of converted clozapine after 10mg/kg CNO administration (Figure 2B).

Structural MRI Scans and Post-mortem Verification of DREADD Expression:

While the pre-surgical T1-weighted MRI scan did not reveal any structural abnormalities in the monkeys (not shown), a second scan performed at 4 months post-surgery, showed a reduction in the volume of the left amygdala in the infant female subject (Figure 3B). Since the female's right amygdala and both hemispheres in the male monkey looked normal in the post-surgical MRI scan, we presume that the reduced volume was due to a unilateral lesion of the amygdala likely caused by a hemorrhage or other ischemic event associated with the intracerebral injection surgery, rather than as a result of the DREADD expression. The unilateral lesion was confirmed during histology (see inset Figure 3B).

To identify the areas with robust DREADD expression, we used antibodies against the HA-tag and revealed these antibodies using biotinylated antibodies and immunoperoxidase. Figure 3C-E show representative images of brain sections close to the viral vector injection sites in the two animals used in this study. During tissue processing, we found considerable damage in the amygdala region in the left hemisphere of the female subject (see inset, Figure 3B), most likely related to the anomaly identified in the MRI scan. As a consequence, we were unable to evaluate the DREADD expression in this hemisphere. In the other 3 hemispheres, expression of DREADDs (visualized using HA-tag immunoreactivity) was found in numerous neurons in the basolateral amygdala, surrounding the injection sites (Figure 3A). The HA-positive area extended approximately 3 X 4 X 4 mm in the antero-posterior, dorsoventral and lateromedial planes, respectively. In higher magnification images, we observed that the labeling was confined, for the most part, to the cell bodies and proximal dendrites of HA-positive neurons (see insets, Figure 3C-E).

Modulation of Emotional Reactivity:

CNO administration did not impact behavior prior to DREADD transduction since both infants exhibited high levels of freezing, hostile, and anxious behaviors on the human intruder task (see Figure 4). By contrast, inhibition of the amygdala with either CNO or clozapine administration lead to decreased freezing during the profile condition of the human intruder task as compared to baseline or vehicle administration ($F[1,11] = 12.98$, $p = 0.005$, $\eta_p^2 = 0.57$; Figure 4A).

DREADD inhibition of the amygdala did not impact the expression of hostile behaviors during the stare condition ($F[1,11] = 0.28$, $p = 0.61$, $\eta_p^2 = 0.03$; Figure 4B) or total vocalizations across all conditions ($F[1,11] = 0.65$, $p = 0.44$, $\eta_p^2 = 0.06$; Figure 4C). However, the expression of anxious behaviors during the Stare condition decreased during amygdala DREADD inhibition ($F[1,11] = 4.99$, $p = 0.049$, $\eta_p^2 = 0.33$; Figure 4D).

Modulation of Socioemotional Attention:

DREADD activation with either CNO or clozapine administration lead to increased attention toward the conspecifics mouth in social stimulus videos as compared to vehicle administration ($F[1,9] = 12.71$, $p = 0.007$, $\eta_p^2 = 0.61$; Figure 5A), but did not impact the amount of time spent looking toward the eyes ($F[1,9] = 0.55$, $p = 0.48$, $\eta_p^2 = 0.06$; Figure 5B). Amygdala inhibition also increased attention toward Aversive stimuli ($F[1,9] = 18.70$, $p = 0.003$, $\eta_p^2 = 0.70$; Figure 5C), but did not impact attention to Neutral stimuli ($F[1,9] = 1.18$, $p = 0.31$, $\eta_p^2 = 0.13$; Figure 5D).

Discussion

This study provides proof-of-principle that chemogenetic tools can be used in infant nonhuman primates to modulate neuronal activity resulting in transient and reversible changes in behavior. To date nonhuman primate chemogenetic studies have focused on adults (Eldridge et al., 2016; Grayson et al., 2016; Nagai et al., 2016; Upright et al., 2018), and considering the reversibility and low invasiveness of this technique, it holds great promise for developmental studies in which more invasive techniques cannot be employed. Our study incorporated the use of two amygdala dependent behavioral tasks and separate investigations of the ability of both CNO and low dose clozapine to activate the DREADD receptors. We have shown that, in infant monkeys, reductive metabolism of CNO to clozapine is similar to adult monkeys, and that low doses of clozapine are enough to elicit the DREADD-mediated behavioral effects. We have also demonstrated that amygdala neurons can be transduced to express DREADDs after virus solution injections, that the expression is stable for at least 7 months, and that activation of DREADDs in the neurons modulates some anxiety-related behaviors. Although this study has technical limitations (see below), our results provide essential pilot data demonstrating that chemogenetics can be used to investigate developmental behavioral neuroscience research questions.

For this study, we chose two tasks that exhibit robust effects after amygdala damage, the human intruder paradigm (Kalin et al., 2004; Machado and Bachevalier, 2008; Raper et al., 2013b; Raper et al., 2013a) and socioemotional attention task (Adolphs et al., 1994; Adolphs et al., 2005; Chudasama et al., 2009; Machado et al., 2009; Machado et al., 2011; Hadj-Bouziane et al., 2012; Gothard et al., 2018). Given that CNO can be reductively metabolized into the psychoactive compound clozapine (Gomez et al., 2017; Raper et al., 2017; Allen et al., 2018), it was important to establish that CNO administration alone would not impact behavior prior to introducing DREADDs into the amygdala. We demonstrate that a 10mg/kg CNO administration did not impact behavior on the human intruder task prior to DREADDs transduction (Figure 4). With these appropriate control measures in place, we found that activation of hM4Di DREADDs in the amygdala results in behavioral alterations similar to those seen after both permanent lesion or transient inactivation (Kalin et al., 2001; Kalin et al., 2004; Machado and Bachevalier, 2008; Chudasama et al., 2009; Machado et al., 2009; Bliss-Moreau et al., 2011b; Raper et al., 2013b; Raper et al., 2013a; Dal Monte et al., 2015; Wellman et al., 2016). After the viral DREADD transduction, CNO or low dose clozapine administration reliably decreased freezing and anxiety on the human intruder task. Importantly, low dose clozapine appeared to be equally effective at decreasing freezing and anxiety as compared to CNO. Considering the differences in plasma clozapine levels after CNO or low dose clozapine administration, the current behavioral effects support the finding that very low doses of clozapine are highly effective for DREADD activation (Gomez et al., 2017). Decreased freezing and anxiety in the human intruder task have been reported following amygdala lesions in nonhuman primates during both adulthood (Kalin et al., 2004; Machado and Bachevalier, 2008) or infancy (Raper et al., 2013b; Raper et al., 2013a). In contrast to results obtained after neonatal amygdala lesions on the human intruder task (Raper et al., 2013b; Raper et al., 2013a), the present study in infant monkeys did not exhibit changes in vocalizations and hostile behavior expression after DREADD-mediated amygdala inactivation. The differences between the previous lesion studies

and the current study support previous findings of brain reorganization after early permanent damage (Machado et al., 2008; Raper et al., 2014b; Grayson et al., 2017; Payne et al., 2017) resulting in more extensive behavioral changes.

Amygdala damage has been shown to alter social behavior and attention to social cues (Adolphs et al., 1994; Bauman et al., 2004; Spezio et al., 2007; Raper et al., 2014a; Dal Monte et al., 2015; Wellman et al., 2016; Bliss-Moreau et al., 2017). Lesions of the amygdala have also been shown to decrease fear reactivity and increase visual interest toward aversive stimuli, such as snakes (Kalin et al., 2004; Chudasama et al., 2009; Machado et al., 2009; Bliss-Moreau et al., 2010; Bliss-Moreau et al., 2011a; Bliss-Moreau et al., 2011b; Feinstein et al., 2011). Using an eye tracking task, we examined how DREADD inhibition of the amygdala altered attention toward social and aversive stimuli. We found that amygdala inhibition with either CNO or clozapine administration resulted in increased looking toward the mouth of novel conspecifics in social stimulus videos. Our findings of increased looking toward the mouth of social videos is similar to results obtained after amygdala damage in adult humans and nonhuman primates (Adolphs et al., 1994; Spezio et al., 2007; Hadj-Bouziane et al., 2012; Dal Monte et al., 2015). A recent study found that adult monkeys with neonatal amygdala lesions looked equally to the mouth and eyes of a conspecific, whereas controls attended more to the eyes (Payne and Bachevalaier, 2019). Those results are similar to the current study, demonstrating that amygdala inhibition only increased the percent looking toward the mouth and did not alter looking at the eyes of conspecifics. In addition, DREADD-mediated amygdala inhibition resulted in increased looking at aversive videos containing snakes and spiders, similar to previous lesion studies (Kalin et al., 2004; Izquierdo et al., 2005; Chudasama et al., 2009; Machado et al., 2009; Bliss-Moreau et al., 2010; Bliss-Moreau et al., 2011a; Bliss-Moreau et al., 2011b). Overall, the similarity between the current results and previous lesion research suggest successful chemogenetic inhibition of the amygdala in infant monkeys.

There are limitations to this study. First, only two infant monkeys were used in this study. However, the within subjects repeated design of the study in which the same behaviors were measured several times under vehicle or CNO/clozapine does provide greater power due to reducing individual variability in behavior that comes from not having a separate control group for comparison. Second, we only conducted a pre-surgery test of CNO on the human intruder task and not the socioemotional attention eyetracking task. Although, it is possible that CNO/clozapine could have impacted social attention, this is unlikely considering a recent study demonstrating that low dose clozapine does not impact social behavior or working memory in rats (Ilg et al., 2018). However, future studies should include testing in naïve animals of the DREADD actuators in all relevant behavioral tasks. Furthermore, although clozapine at the low doses used here preferentially activated DREADDs, we cannot completely rule out contributions of other neurotransmitter receptors, therefore, future studies should investigate the use of novel ligands that will not convert to clozapine and be more selective for the DREADDs (Bonaventura et al., 2018). Third, there was an unintended amygdala lesion in the infant female subject. We cannot rule out the possibility that the lesion was caused by the introduction of the artificial DREADD receptors. However, this is unlikely since the damage occurred only in one subject and was localized in one hemisphere. Instead, it was most likely due to a hemorrhage or other ischemic event that occurred as a consequence of the intracerebral injections. This event highlights the importance of conducting a post-surgical MRI. Future studies should consider doing a T2-weighted Fluid Attenuated Inverse Recovery (FLAIR) MRI sequence within 5-10 days after surgery which could help to detect edema from a hemorrhage or ischemic event (Nemanic et al., 2002). The extent to which the unilateral lesion impacted behavioral results is unclear, although previous studies have shown that unilateral amygdala lesioned monkeys behave similarly to controls (Kalin et al., 2004; Raper et al., 2013b). Therefore, regardless of whether the bilateral silencing of the amygdala was mediated by DREADDs (infant male

subject) or achieved by a combination of unilateral lesion and DREADD activation (infant female subject), the transient and reversible effects on behavior were similar between both subjects (see Figures 4 and 5). Despite its limitations, the current study is the first to demonstrate the promise for using chemogenetic tools in young infant nonhuman primates to address developmental behavioral neuroscience questions.

In summary, transient and reversible inhibition of the amygdala is possible in infant monkeys using chemogenetics, specifically hM4Di DREADDs. This study provides critical pilot data demonstrating that chemogenetics can be used in young animals to investigate developmental neuroscience questions that could not previously be investigated due to the limitations of permanent lesions or invasiveness of pharmacological inactivation studies.

References

- Adolphs R, Tranel D, Damasio H, Damasio A (1994) Impaired recognition of emotion in facial expressions following bilateral damage to the human amygdala. *Nature* 372:669-672.
- Adolphs R, Gosselin F, Buchanan TW, Tranel D, Schyns P, Damasio AR (2005) A mechanism for impaired fear recognition after amygdala damage. *Nature* 433:68-72.
- Aggleton JP (2000) *The Amygdala. A Functional Analysis*, 2nd Edition. New York, NY.: Oxford University Press.
- Allen DC, Carlson TL, Xiong Y, Jin J, Grant KA, Cuzon Carlson VC (2018) A comparative study of the pharmacokinetics of clozapine N-oxide and clozapine N-oxide hydrochloride salt in rhesus macaques. *J Pharmacol Exp Ther* online ahead of print.
- Armbruster BN, Li X, Pausch MH, Herlitze S, Roth BL (2007) Evolving the lock to fit the key to create a family of G protein-coupled receptors potently activated by an inert ligand. *Proc Natl Acad Sci USA* 104:5163-5168.
- Baldessarini RJ, Centorrino F, Flood JG, Volpicelli SA, Huston-Lyons D, Cohen BM (1993) Tissue concentrations of clozapine and its metabolites in the rat. *Neuropsychopharmacology* 9:117-124.
- Bauman MD, Lavenex P, Mason WA, Capitanio JP, Amaral DG (2004) The development of social behavior following neonatal amygdala lesions in rhesus monkeys. *J Cogn Neurosci* 16:1388-1411.
- Bliss-Moreau E, Bauman MD, Amaral DG (2011a) Neonatal amygdala lesions result in globally blunted affect in adult rhesus macaques. *Behav Neurosci* 125:848-858.
- Bliss-Moreau E, Moadab G, Santistevan A, Amaral DG (2017) The effects of neonatal amygdala or hippocampus lesions on adult social behavior. *Behav Brain Res* 322:123-137.

Bliss-Moreau E, Toscano JE, Bauman MD, Mason WA, Amaral DG (2010) Neonatal amygdala or hippocampus lesions influence responsiveness to objects. *Dev Psychobiol* 52:487-503.

Bliss-Moreau E, Toscano JE, Bauman MD, Mason WA, Amaral DG (2011b) Neonatal amygdala lesions alter responsiveness to objects in juvenile macaques. *Neuroscience* 178:123-132.

Bonaventura J et al. (2018) Chemogenetic ligands for translational neurotherapeutics. *bioRxiv* doi10.1101/487637.

Chareyron LJ, Lavenex PB, Amaral DG, Lavenex P (2012) Postnatal development of the amygdala: A stereological study in macaque monkeys. *J Comp Neurol* 520:1965-1984.

Chudasama Y, Izquierdo A, Murray EA (2009) Distinct contributions of the amygdala and hippocampus to fear expression. *Eur J Neurosci* 30:2327-2337.

Dal Monte O, Costa VD, Noble PL, Murray EA, Averbeck BB (2015) Amygdala lesions in rhesus macaques decrease attention to threat. *Nat Commun* 6.

Eldridge MA, Lerchner W, Saunders RC, Kaneko H, Krausz KW, Gonzalez FJ, Ji B, Higuchi M, Minamimoto T, Richmond BJ (2016) Chemogenetic disconnection of monkey orbitofrontal and rhinal cortex reversibly disrupts reward value. *Nat Neurosci* 19:37-39.

Farrell MS, Roth BL (2013) Pharmacosynthetics: Reimagining the pharmacogenetic approach. *Brain Res* 1511:6-20.

Feinstein JS, Adolphs R, Damasio A, Tranel D (2011) The Human Amygdala and the Induction and Experience of Fear. *Current Biology* 21:34-38.

Gomez JL, Bonaventura J, Lesniak W, Mathews WB, Sysa-Shah P, Rodriguez LA, Ellis RJ, Richie CT, Harvey BK, Dannals RF, Pomper MG, Bonci A, Michaelides M (2017) Chemogenetics revealed: DREADD occupancy and activation via converted clozapine. *Science* 357:503-507.

- Gothard KM, Mosher CP, Zimmerman PE, Putnam PT, Morrow JK, Fuglevand AJ (2018) New perspectives on the neurophysiology of primate amygdala emerging from the study of naturalistic social behaviors. *Wiley Interdiscip Rev Cogn Sci* 9.
- Grayson DS, Bliss-Moreau E, Bennett J, Lavenex P, Amaral DG (2017) Neural Reorganization Due to Neonatal Amygdala Lesions in the Rhesus Monkey: Changes in Morphology and Network Structure. *Cereb Cortex* 27:3240-3253.
- Grayson DS, Bliss-Moreau E, Machado CJ, Bennett J, Shen K, Grant KA, Fair DA, Amaral DG (2016) The Rhesus Monkey Connectome Predicts Disrupted Functional Networks Resulting from Pharmacogenetic Inactivation of the Amygdala. *Neuron* 91:453-466.
- Hadj-Bouziane F, Liu N, Bell AH, Gothard KM, Luh WM, Tootell RB, Murray EA, Ungerleider LG (2012) Amygdala lesions disrupt modulation of functional MRI activity evoked by facial expression in the monkey inferior temporal cortex. *Proc Natl Acad Sci USA* 109:E3640-E3648.
- Hunsaker MR, Scott JA, Bauman MD, Schumann CM, Amaral DG (2014) Postnatal development of the hippocampus in the Rhesus macaque (*Macaca mulatta*): a longitudinal magnetic resonance imaging study. *Hippocampus* 24:794-807.
- Ilg AK, Enkel T, Bartsch D, Böhner F (2018) Behavioral Effects of Acute Systemic Low-Dose Clozapine in Wild-Type Rats: Implications for the Use of DREADDs in Behavioral Neuroscience. *Front Behav Neurosci* 12.
- Izquierdo A, Suda RK, Murray EA (2005) Comparison of the effects of bilateral orbital prefrontal cortex lesions and amygdala lesions on emotional responses in rhesus monkeys. *J Neurosci* 25:8534-8542.
- Kalin N, Shelton SE, Davidson RJ (2004) The role of the central nucleus of the amygdala in mediating fear and anxiety in primate. *J Neurosci* 24:5506-5515.

- Kalin N, Shelton SE, Davidson RJ, Kelley AE (2001) The primate amygdala mediates acute fear but not the behavioral and physiological components of anxious temperament. . J Neurosci 21:2067-2074.
- Kalin NH, Shelton SE, Takahashi LK (1991) Defensive behaviors in infant rhesus monkeys: Ontogeny and context-dependent selective expression. Child Development 62.
- Kotterman MA, Yin L, Strazzeri JM, Flannery JG, Merigan WH, Schaffer DV (2015) Antibody neutralization poses a barrier to intravitreal adeno-associated viral vector gene delivery to non-human primates. Gene Ther 22:116-126.
- Lu H, Rosenbaum S (2014) Developmental pharmacokinetics in pediatric populations. J Pediatr Pharmacol Ther 19:262-276.
- Machado CJ, Bachevalier J (2008) Behavioral and hormonal reactivity to threat: effects of selective amygdala, hippocampal or orbital frontal lesions in monkeys. Psychoneuroendocrinology 33:926-941.
- Machado CJ, Kazama AM, Bachevalier J (2009) Impact of amygdala, orbital frontal, or hippocampal lesions on threat avoidance and emotional reactivity in nonhuman primates. Emotion 9:147-163.
- Machado CJ, Bliss-Moreau E, Platt ML, Amaral DG (2011) Social and nonsocial content differentially modulates visual attention and autonomic arousal in Rhesus macaques. PLoS One 6:e26598.
- Machado CJ, Snyder AZ, Cherry SR, Lavenex P, Amaral DG (2008) Effects of neonatal amygdala or hippocampus lesions on resting brain metabolism in the macaque monkey: a microPET imaging study. Neuroimage 39:832-846.
- Mendelson MJ (1982) Clinical examination of visual and social responses in infant rhesus monkeys. Dev Psychol 18:659-664.
- Mendelson MJ, Haith MM, Goldman-Rakic PS (1982.) Face scanning and responsiveness to social cues in infant rhesus monkeys. Dev Psychol 18:222—228.

- Mendoza SD, El-Shamayleh Y, Horwitz GD (2017) AAV-mediated delivery of optogenetic constructs to the macaque brain triggers humoral immune responses. *J Neurophysiol* 117:2004-2013.
- Milsap RL, Jusko WJ (1994) Pharmacokinetics in the infant. *Environ Health Perspect* 102:107-110.
- Mineka S (1987) A primate model of phobic fears. In: *Theoretical foundations of behavior therapy* (Eysenck HJ, Martin I, eds), pp 81–111. New York: Plenum.
- Mineka S, Keir R, Price V (1980) Fear of snakes in wild- and laboratoryreared rhesus monkeys (*Macaca mulatta*). *Anim Learn Behav* 8:653– 663.
- Moadab G, Bliss-Moreau E, Bauman MD, Amaral DG (2017) Early amygdala or hippocampus damage influences adolescent female social behavior during group formation. *Behav Neurosci* 131:68-82.
- Mosher CM, Zimmerman PE, Gothard KM (2011) Videos of conspecifics elicit interactive looking patterns and facial expressions in monkeys. *Behav Neurosci* 25:639-652.
- Mosher CM, Zimmerman PE, Gothard KM (2014) Neurons in the monkey amygdala detect eye-contact during naturalistic social interactions. *Current Biology* 24:2459-2464.
- Nagai Y et al. (2016) PET imaging-guided chemogenetic silencing reveals a critical role of primate rostromedial caudate in reward evaluation. *Nat Commun* 7:13605.
- Nemanic S, Alvarado MC, Price RE, Jackson EF, Bachevalier J (2002) Assessment of locus and extent of neurotoxic lesions in monkeys using neuroimaging techniques: a replication *J Neurosci Methods* 121:199-209.
- Payne C, Bachevalier J (2019) Early amygdala damage alters the way rhesus macaques process species-specific audio-visual vocalizations. *Behav Neurosci* In Press.
- Payne C, Cirilli L, Bachevalier J (2017) An MRI study of the corpus callosum in monkeys: Developmental trajectories and effects of neonatal hippocampal and amygdala lesions. *Dev Psychobiol* 59:495-506.

- Payne C, Machado CJ, Bliwise NG, Bachevalier J (2010) Maturation of the hippocampal formation and amygdala in *Macaca mulatta*: a volumetric magnetic resonance imaging study. *Hippocampus* 20:922-935.
- Raper J, Wilson ME, Sanchez M, Machado C, Bachevalier J (2013a) Pervasive alterations of emotional and neuroendocrine responses to an acute stressor after neonatal amygdala lesions in rhesus monkeys. *Psychoneuroendocrinology* 38:1021-1035.
- Raper J, Stephens SBZ, Sanchez M, Bachevalier J, Wallen K (2014a) Neonatal amygdala lesions alter mother-infant interactions in rhesus monkeys living in a species-typical social environment. *Dev Psychobiol* 56:1711-1722.
- Raper J, Wallen K, Sanchez MM, Stephens SB, Henry A, Villareal T, Bachevalier J (2013b) Sex-dependent role of the amygdala in the development of emotional and neuroendocrine reactivity to threatening stimuli in infant and juvenile rhesus monkeys. *Horm Behav* 63:646-658.
- Raper J, Stephens SBZ, Henry A, Villarreal T, Bachevalier J, Wallen K, Sanchez MM (2014b) Neonatal amygdala lesions lead to increased activity of brain CRF systems and hypothalamic-pituitary-adrenal axis of juvenile rhesus monkeys. *J Neurosci* 34:11452-11460.
- Raper J, Morrison RD, Daniels JS, Howell L, Bachevalier J, Wichmann T, Galvan A (2017) Metabolism and Distribution of Clozapine-N-oxide: Implications for Nonhuman Primate Chemogenetics. *ACS Chem Neurosci* 8:1570-1576.
- Roth BL (2016) DREADDs for Neuroscientists. *Neuron* 89:683-694.
- Spezio ML, Huang PY, Castelli F, Adolphs R (2007) Amygdala damage impairs eye contact during conversations with real people. *J Neurosci* 27:3994-3997.
- Stephens SBZ, Raper J, Bachevalier J, Wallen K (2015) Neonatal amygdala lesions advance pubertal timing in female rhesus macaques. *Psychoneuroendocrinology* 51:307-317.

Sternson SM, Roth BL (2014) Chemogenetic tools to interrogate brain functions. *Annu Rev Neurosci* 37:387-407.

Upright NA, Brookshire SW, Schnebelen W, Damatac CG, Hof PR, Browning PGF, Croxson PL, Rudebeck PH, Baxter MG (2018) Behavioral Effect of Chemogenetic Inhibition Is Directly Related to Receptor Transduction Levels in Rhesus Monkeys. *J Neurosci* 38:7969-7975.

Urban DJ, Roth BL (2015) DREADDs (designer receptors exclusively activated by designer drugs): chemogenetic tools with therapeutic utility. *Annu Rev Pharmacol Toxicol* 55:399-417.

Wellman LL, Gale K, Malkova L (2005) GABAA-mediated inhibition of basolateral amygdala blocks reward devaluation in macaques. *J Neurosci* 25:4577-4586.

Wellman LL, Forcelli PA, Aguilar BL, Malkova L (2016) Bidirectional Control of Social Behavior by Activity within Basolateral and Central Amygdala of Primates. *J Neurosci* 36:8746-8756.

Wess J, Nakajima K, Jain S (2013) Novel designer receptors to probe GPCR signaling and physiology. *Trends Pharmacol Sci* 34:385-392.

Table 1. Behavioral Ethogram for Human Intruder

Category and Specific Behaviors	Measurement	Brief Descriptions
Freeze	Duration	Rigid, motionless posture except slight head movement
Hostile	Cumulative Frequency	
Threat bark	Frequency	Low pitch, high intensity, rasping, guttural
Threat	Frequency	Any of the following: open mouth (no teeth exposed), head-bobbing, or ear flapping
Cage aggression	Frequency	Vigorously slaps, shakes, or slams body against cage
Lunge	Frequency	A quick, jerky movement toward the stimulus
Vocalizations	Cumulative Frequency	
Coo	Frequency	Clear soft pitch and intensity, sounds like “ooooh”
Scream	Frequency	High pitch, high intensity screech or loud chirp
Anxiety	Cumulative Frequency	
Scratch	Frequency	Rapid scratch of body with hands or feet
Body shake	Frequency	Shake of the whole body or just head and shoulders region
Tooth grind ¹	Frequency	Repetitive, audible rubbing of upper and lower teeth
Yawn	Frequency	Open mouth widely, exposing teeth
Affiliative²	Cumulative Frequency	
Lipsmack	Frequency	Rapid movement of pursed lips, accompanied by a smacking sound
Present	Frequency	Rigid posture (knees locked) with tail elevated and rump oriented toward the stimulus object
Fearful²	Cumulative Frequency	
Withdrawal	Frequency	Quick, jerky motion away from the stimulus object (jump back)
Grimace	Frequency	Refracted lips, exposed clenched teeth
Stereotypies²	Cumulative Duration	
Pacing	Duration	Repetitive motor pattern around the test cage
Motor stereotypy	Duration	Repetitive, abnormal voluntary or involuntary motor patterns (e.g. swinging, twirling, flipping)

List of all behaviors scored, how they are measured and a brief definition. 1. Behavior for which total duration was also measured. 2. Behavior that was rarely or never seen in the current study.




Figure 1. Schematic of the study design. Two infant rhesus monkeys received bilateral injections of AAV-hM4Di DREADD into the amygdala at 9 months of age, followed by 6 months of testing with either Vehicle (Veh), clozapine-n-oxide (CNO, 10mg/kg), or clozapine (0.1mg/kg). Prior to DREADD transduction, infants were tested on the Human Intruder paradigm with CNO to rule out any impact of the drug on the behavior in naïve animals. Magnetic resonance imaging (MRI) was conducted immediately prior to surgery to aid with amygdala injection localization and again at 4 months post-surgery in relation to another study not reported here. Monkeys were sacrificed at 7 months post-surgery to determine DREADD expression with histology.




Figure 2. Time-concentration profiles of CNO and clozapine in infant monkeys. Plasma concentrations of CNO (yellow line) and its metabolite clozapine (green line) following SC administration of CNO at 10mg/kg (a) and clozapine at 0.1mg/kg (b) in the infant male (blue square) and female (pink circle) monkeys.




Figure 3. Expression of DREADDs after virus injections in the amygdala. A) Injection sites (each site marked by a red dot) in the amygdala, identified in Nissl stains. All sites identified from both animals were pooled for the figure. B) T1-weighted MRI scan at 4 months post-surgery for the female and the male subjects. In the left hemisphere of the female, a reduction in the amygdala was observed, which was later confirmed in the Nissl section (inset). C-E) HA immunoperoxidase labeling of regions of the amygdala. Areas in the rectangles in the left image in each panel are shown at higher magnifications at right, the insets show higher magnifications of the HA(hM4Di)-positive cell bodies. Scale bars in C apply to D and E. AP indicates the approximate antero-posterior plane, from the interaural line, abbreviations: AB, anterobasal amygdala; AC, anterior commissure; B, basal amygdala; CE, central amygdala; GPe, external segment of the globus pallidus, L, lateral amygdala; Put, putamen. magnifications of the HA(hM4Di)-positive cell bodies.




Figure 4. Behavioral responses on the Human Intruder paradigm with and without DREADD inhibition of the amygdala. Freezing (A), hostile (B), vocalizations (C), and anxiety (D) behaviors measured after CNO (yellow bars), clozapine (green bars), vehicle (Veh, open bars), or CNO prior to transduction (CNO Prior, striped bar). Individual scores for the infant male is represented by blue squares and infant female is represented by pink circles. * indicates a significant difference from vehicle and CNO Prior ($p < 0.05$).




Figure 5. Looking behavior during the Socioemotional Attention task with and without DREADD inhibition of the amygdala. Percent looking to the mouth (A) or looking to the eyes (B) of Social videos, looking at the Aversive stimulus (C), and looking at the Neutral stimulus (D) after CNO (yellow bars), clozapine (green bars), or vehicle (Veh, open bars). Individual scores for the infant male is represented by blue squares and infant female is represented by pink circles. * indicates a significant difference from vehicle ($p < 0.05$).