

The magnitude of neonatal mortality and its predictors in Ethiopia: a systematic review and meta-analysis.

Yared Asmare^{1*}, Wondimeneh Shibabaw¹, Tadesse Yirga², Abate Dargie¹, Tesfa Dejenie Habtewold³

¹College of Health Science, Debre Birhan University, Debre Birhan, Ethiopia

²College of Health Science Debre Markos University, Debre Markos, Ethiopia

^{1,3}Department of Epidemiology, University Medical Centre Groningen, University of Groningen, Groningen, The Netherlands.

Yared Asmare (YA): yaredasmare123@gmail.com

Tadesse Yirga (TY): tadesseyirga680@gmail.com

Wondimeneh Shibabaw (WS): wshibabaw2015@gmail.com

Abate Dargie (AD): abiarse999@gmail.com

Tesfa Dejenie (TD): tesfadej2003@gmail.com

*corresponding author

Email: yaredasmare123@gmail.com ,

P.O.Box 445, Addis Ababa, Ethiopia

Abstracts

Background: Although neonatal death is a global burden, it is the highest in Sub Saharan Africa countries such as Ethiopia. This study was aimed to provide pooled national prevalence and predictors of neonatal mortality in Ethiopia.

Objective: To assess the pooled prevalence and predictors of neonatal mortality in Ethiopia.

Search Strategy: global databases were systematically explored. Systematically searched using the following databases: Boolean operator, Cochrane library, PubMed, EMBASE, HINARI, and

Google Scholar. Selection, screening, reviewing and data extraction was done by two reviewers independently using Microsoft excel spread sheet. The modified Newcastle–Ottawa Scale (NOS) and the Joanna Briggs Institute Prevalence Critical Appraisal tools were used to assess the quality of evidence

Selection criteria: All studies conducted in Ethiopia and reporting the prevalence and predictors of neonatal mortality were included

Data Collection and Analysis: Data were extracted using a Microsoft Excel spreadsheet software and imported into STATA Version 14 s for further analysis. The pooled effect size with 95% confidence interval of neonatal mortality rate was determined using a weighted inverse variance random-effects model. Publication bias was checked using funnel plots, Egger’s and bagger’s regression test. Heterogeneity also checked by Higgins’s method. A random effects meta-analysis model was computed to estimate the pooled effect size (i.e. prevalence and odds ratio). Moreover, subgroup analysis based on region, sample size and study design were done.

Results: After reviewing 88 studies, 12 studies fulfilled the inclusion criteria and were included in the meta-analysis. The pooled national prevalence of neonatal mortality in Ethiopia was 16.3% (95% CI: 11.9, 20.7, $I^2=88.6\%$). The subgroup analysis indicated that the highest prevalence was observed in Amhara region with a prevalence of 20.3% (95% CI: 9.6, 31.1, $I^2=98.8$) followed by Oromia, 18.8% (95%CI: 11.9,49.4, $I^2=99.5$). Gestational age AOR,1.14 (95% CI: 0.94 ,1.3), neonatal sepsis (OR:1.2(95% CI: 0.8, 1.5), respiratory distros (OR: 1.2(95% CI: 0.8, 1.5) and place of residency (OR:1.93 (95% CI:1.1,2.7) were the most important predictor.

Conclusions: neonatal mortality in Ethiopia was significantly decreased than the national report. There was evidence that neonatal sepsis, gestational age, respiratory distress were the significant predictors. We strongly recommended that health care workers should give a priority for the identified predictors.

Keywords: incidence, prevalence, neonatal mortality, Ethiopia

Background

The neonatal period is the first four weeks of a child's life in which changes are very rapid and many critical events can occur in this period [1, 2]. Survival of new born babies had improved significantly through enhanced and specialized care. But, still it is the main reason of under-five death and risk of lifelong sequel [3-5]. Globally, the neonatal mortality rate (NMR) is declined from 49 % to 19%, but slower than under five mortality rate (dropped by 60%) .Of all death of under-five , 40% were attributed with new born death [5] in which close to 1 and 2 million deaths occur on the day of birth, and in the first week of life respectively [6]. A review of 20 studies also indicated that the total NMR greatly varied between developed (4 to 46%) and developing (0.2 to 64.4%) countries [7]. Despite this, NMR shares the highest proportion of under-five mortality worldwide. [8-14]. According to the report of United Nations international child education and fund(UNICEF) and Ethiopian Demographic and Health Survey (EDHS) , the neonatal mortality rate accounts 23% and 29%/1000 live births in Ethiopia [5, 15].

In Ethiopia, different studies have been conducted to assess the prevalence of neonatal mortality and associated factors. The findings of these uneven studies documented that there was a great variability in the prevalence of neonatal mortality across the regions of the country. Concerning predictors, these studies revealed that different maternal, neonatal and health service related factors influenced neonatal mortality; maternal residency [[16, 17]], history of antenatal, gestational age ,neonatal sepsis, RDS, asphyxia were some of the factors associated with neonatal mortality [18, 19].Globally, there are different policies, strategies, and programs which work on prevention and care of neonate [20, 21]. However, it is sustained as the most cause of under-five mortality worldwide [11, 22-25].Despite different strategies, health policies (like HSDP IV 2010-2015) and interventions were implemented to prevent and care neonatal death, the rate of neonatal mortality is high. Additionally, Ethiopia achieved millennium development goal in reducing child mortality by two-third, but neonatal mortality is still high. Several studies were conducted and showed important results, including prevalence of mortality and its major predictors in the study area. However, there is variation in the prevalence and associated factors of studies based on region, sample size and the study design that have been used.as far as our knowledge, there has been no any

systematic review and meta-analysis of studies reporting on the neonatal mortality and its predictors .So that we believe synthesizing these studies may fill the gap in the literature and provide stronger evidence for policy-making as there is an increasing recognition of systematic review and meta-analysis findings in the policy-making process. Thus, further meta-analysis and synthesis of the available evidence is needed now. Hence, this study aimed to synthesize nationally available evidence on the prevalence of neonatal mortality and the association with different variables in Ethiopia. The findings of this study will be used as an input to policy makers and program planners working in the area of neonatal health.

Methods

Searching strategies

The PRISMA guidelines protocol was used to write the systematic review [22]. The reviewer follows PRISMA systematic review protocol as a reporting guideline (for the PRISMA checklist, eligible studies for the study were selected in terms of titles alone, abstracts, and then full-text articles, based on inclusion criteria. Cochrane library, PubMed, EMBASE, HINARI, and Google Scholar was systematically searched for articles. The studies were accessed using the following search terms: “neonatal mortality”, “predictors”, “neonatal death”, “newborn”, “prevalence”, “neonatal mortality” and “Ethiopia”. The search terms were used individually and in combination using “AND” and “OR” Boolean operators. In addition, after identification of included studies, cross-references were searched to identify more eligible studies. The search was guided by PECO: Population - neonates (age < 28 days); occurrence of death within 28 days after delivery.

Inclusion and exclusion criteria

Studies with the following major criteria were considered for inclusion. Observational (i.e. cross-sectional, case-control and cohort) studies in Ethiopia, which reports the prevalence and predictors of neonatal mortality were included. Articles published in English language was considered as further inclusion criteria. On the other hand, studies which did not report the outcome and articles without full-text were excluded. Corresponding authors were approached by email at least twice to access full-texts.

Screening and data extraction

Two reviewers (YA and WS) screened titles and abstracts against the inclusion criteria-. Then, the full-texts of articles were accessed and independent assessment was carried out by two reviewers based on the predetermined inclusion and exclusion criteria. Discrepancies between the reviewers were resolved through discussion and common consensus of all investigators. Data were extracted from the included papers by AD, TD and TY independently extracted data from a random sample of 20% of papers to check consistency; consequently, there was no differences.

Assessment of study quality; Structured data abstraction form was constructed in Microsoft excel. In each abstract and full text of the article, which was considered to be appropriate; a special emphasis was given for clearness of objective, data about the study area, study design, year of publication, study population, sample, size respondent rate, prevalence/incidence of neonatal death and other useful variables were recorded (table 1). The Joanna Briggs Institute Prevalence Critical Appraisal Tool for use in systematic review for prevalence study was used for critical appraisal of studies [26]. Moreover, methodological and other quality of each article was assessed based on a modified version of the Newcastle-Ottawa Scale for cross-sectional study adapted from Modesti et al[27].

Data synthesis and statistical analysis

Data were extracted using a Microsoft Excel spreadsheet software and imported into STATA Version 14 software for further analysis. The pooled effect size with 95% confidence interval of national neonatal mortality rate was determined using a weighted inverse variance random-effects model [28]. Heterogeneity across the studies was assessed using I^2 statistic where 25, 50 and 75% representing low, moderate and high heterogeneity respectively[29]. A Funnel plot ,beggars and Egger's regression test were used to check publication bias[30]. Moreover, subgroup analysis based on study area (region), study design and sample size were done. Log odds ratios were used to examine the association between mortality and its major predictors.

Results

Characteristics of included studies

The abstract search resulted in 136985 PUBMED references when we search with "neonatal mortality) OR Ethiopia in which the report is not limited to the study area. When farther advanced searches are attempted the search, strategy retrieved 88 potential articles, of which 12 full text articles that fulfill the eligible criteria with a total sample size of 12397 neonates were included in the final analysis for the current systematic review and meta-analysis (Fig. 1). Information about authors, publication year, population, study area, region, study design, outcome and main results from the selected articles were extracted and summarize on table. the overall respondent rate was between eighty-three to hundred percent.




Fig. 1: PRISMA flow diagram for showing screening and selection process of studies

Features of Studies. All studies were done in Ethiopia and published in an indexed journal. Eight of them were cohort (both retrospective and prospective) and the remaining three were cross sectional studies. Detailed screening and selection process were shown in Figure 1.

Study characteristics

The studies were conducted in Amhara [19, 31-34], Tigray [35, 36], Southern Nations, Nationalities and Peoples (SNNP) [16], Oromia [37, 38] and Somali [39] region. Eight were cohort and three were cross sectional studies. The sample size of the studies ranging from 485 to 3,604.

Table 1: Descriptive summary of 12 studies included in the meta-analysis of the prevalence of neonatal mortality.

| Authors name | year | Area | region | study design | Total sample | Response rate | Prevalence |
|--|------|----------|--------|-----------------|--------------|---------------|------------|
| Gizaw et al. % [16] | 2013 | Butajira | SNNPR | Cohort | 1055 | 83.9 | 26% |
| Andargie et al. [34] | 2013 | Gondar | Amhara | Cross-sectional | 1752 | 100 | 5% |
| Mengesha et al. [35] | 2016 | Tigray | Tigray | Cohort | 1162 | 99.14 | 6% |
| Yismaw et al. [31] | 2019 | Gondar | Amhara | Cohort | 516 | 100 | 29% |
| Yismaw and Ta-rekegn% [32] | 2018 | Gondar | Amhara | Cross-sectional | 516 | 100 | 28.8% |
| Demisse et al. % [33] | 2017 | Gondar | Amhara | Cross-sectional | 769 | 100 | 14.3% |
| Debelew et al. [37] | 2014 | Jimma | Oromia | Cohort | 3604 | 96.1 | 3.2% |
| Farah et al. % [39] | 2018 | Somali | Somali | Cohort | 792 | 100 | 5.7% |
| Yehuala and Teka [19] | 2015 | Gondar | Amhara | Cohort | 485 | 100 | 25% |
| Wesenu et al. [38] | 2017 | Jimma | Oromia | Cohort | 490 | 100 | 34.5% |
| Mengesha et al. [36] | 2016 | Mekelle | Tigray | Cohort | 1162 | 99.04 | 5.8% |
| Orsido et al.[18] | 2019 | Weliata | SNNPR | cohort | 964 | 100 | 16.5 |

prevalence of neonatal mortality

In the current systematic review and meta-analysis, the pooled prevalence estimates of neonatal mortality were described by forest plot (Fig. 2). The pooled prevalence of neonatal mortality from the random effect's method was found to be 16.3% (95% CI; 12.1–20.6).




Figure 2: Forest plot of the pooled prevalence of neonatal mortality.

Investigation of heterogeneity and publication bias

Consequently, the analysis showed a substantial heterogeneity of egger test ($p < 0.001$) and I^2 statistics ($I^2 = 83\%$) for neonatal mortality. The discrepancy in the pooled estimates of the prevalence was adjusted through subgroup analysis based on the region, where the study conducted and based on the types of study design. A sensitivity analysis was conducted to check the stability of summary estimate. The funnel plot was found to be asymmetric (fig 3) and Egger's and bagger's test was found to be significant publication bias at p value ≤ 0.001 which confirms that there is a publication bias.




Fig.3: Funnel plot to show the distribution of 12 studies.

Meta-regression analysis

Based on the subgroup-analysis result, the highest prevalence (20.3%; 95% CI:9.6 -31.1) was reported in Amhara region with regard to sample size, the prevalence of neonatal mortality was higher in studies having a sample size < 800, 22.7% (95%CI:12.8, 32.7) compared to those having a sample size \geq 800,8.9 (95%CI:4.9, 12.9) (Table 3).

Table 3: Meta-regression analysis of neonatal sepsis with heterogeneity of neonatal mortality.

| Heterogeneity source | Coefficients | Std. Err. | P-value |
|----------------------|----------------|--------------|-------------|
| Publication Year | -1.243 | 4.738 | 0.64 |
| Sample size | 0.01800 | 0.290 | 0.86 |

Predictors of neonatal mortality

Gestational age

Seven studies[[17](#), [31](#), [33](#), [36-38](#), [40](#)], examined the association between gestational age and neonatal mortality. The pooled odds ratio was 1.32 (95% CI: 1.07 ,1.58), $I^2= 49\%$). Neonates born

as term had 14% lower chance of death compared with preterm neonates (Fig. 4). Begg's ($p=0.37$) and Egger's ($p=0.7$) tests did not reveal significant publication bias.




Figure 4: the pooled odds ratio of the association between GA and neonatal mortality.

Residency

Six studies [31-33, 36, 37, 41] reported the association between place of residency and neonatal mortality. The pooled odds ratio was 1.93 (95% CI: 1.1, 2.7; $I^2=3\%$) (fig 5). Neonates who were born in rural area were found to be 98% higher risk of mortality than their counterpart although not statistically significant. Begg's ($p=0.99$) and Egger's tests ($p=0.663$) showed that there was no significant publication bias.




Figure 5: the pooled odds ratio of the association between residency and neonatal mortality.

Respiratory distros syndromes

According to these finding, those neonates having RDS were also associated with neonatal mortality (fig 6). Neonates who had RDS were nearly one and half times more likely to die as compared to those who did not have RD (OR= 1.2,95% CI: 0.8, 1.5). We observed that a no any

heterogeneity across the studies ($I^2 = 4.43\%$ (d.f. = 5) with no any publication bias with begger ($p = 1.000$) and egger ($p = 0.138$) tests.




Figure 6: the pooled odds ratio of the association between RDS and neonatal mortality.

Neonatal sepsis

To assess the association of neonatal sepsis with neonatal mortality, we have include five studies [33, 36-38, 41], Patients who had neonatal sepsis had almost one and half times higher chance of neonatal death compared to those patients without sepsis OR: 1.23(95% CI:1.1, 1.4) (Figure 7). The heterogeneity test showed extreme evidence of variation across studies ($I^2 = 90\%$, $P =$

0.001. Moreover, the result of Egger's test to examine publication bias showed no statistically significant evidence of publication bias (I² = 0.806) and (egger P=0.511).




Figure7: Pooled odds ratio of the association between neonatal sepsis and neonatal mortality

Discussion

This systematic review and meta-analysis revealed the national burden of neonatal mortality. We also sought, gestational age, neonatal sepsis and residence significantly predicted neonatal mortality but not respiratory distress syndrome. Our study showed that the national prevalence of neonatal mortality was 16.3%. This is in line with a study findings in Cameroon, South Africa, South Sudan and Mauritanian [15, 42] [40]. However, our finding was lower than the UNICEF national 2016 report[3, 5] and other national reports in Africa[43, 44], Europe, USA, Central and West Asia [45-47]. This marked difference might be attributed to difference in methodology, sample size, study period and geographic area. For example, the UNICEF report includes all

regions and cities of Ethiopia but the current study includes studies only from five regions. Additionally, some of the previous reports limited study area instead of nationwide reports. Moreover, the difference in the study period could be a relevant factor as standard of care and treatment modalities changes over time. Additionally, though, neonatal health care services in Ethiopia have been remaining less consolidated, systematic efforts have been gaining momentum since relatively recent. Furthermore, the NICU service in our countries is not well included Level three (subspecialty) NICU all over the countries.

The subgroup analysis indicated that the highest prevalence of mortality was observed in Amhara region (20.3%) whereas the lowest prevalence was observed in Somalia (5.7%). The possible reason might be that in Amhara region five studies were included compared with other regions. Additionally, the sampled population included in Amhara region were higher than other regions. These studies reported that, having neonatal sepsis was significantly increased the risk of mortality. Those who had neonatal sepsis were nearly **two** times increase risk of mortality as compared to those who did not have sepsis. This finding is supported by a result in developed and developing country [19] [48] [7], [49], [50] [51], ([52], [53]. The possible reason might be due to newborns have many physiologic challenges when adapting to the extrauterine environment which might contributed to common problem like immature immunity, RDS, neurologic, cardiovascular, hematologic, nutritional, gastrointestinal and poor thermoregulation which farther increase risk of sepsis and mortality. We also found that neonates living in rural areas were more vulnerable to death than their urban counterparts. Neonates from rural households were nearly two times more likely to die as compared to their counterparts. This finding is similar to a study conducted in Washington, Louisiana, and Tennessee [54]. This variance could be due to the fact that rural residents are still relatively disadvantaged in terms of infrastructures, knowledge and awareness difference, distance from the site. Another possible reason could be that people living in rural areas tend to be poorer than their urban counterparts' area, a factor known to have an impact on the neonatal outcome. GA is also another important determinant of neonatal mortality in our meta-analysis. Accordingly, neonates born as preterm were almost one and half times more likely to die than those term neonates [49-57] [55], [56], [52], [53], [50], [7] [51], [49, 57]. RDS are an important determinant of neonatal mortality although not statistically significant in our meta-analysis. Neonates who had RDS were one times more likely to die as compared to those who did not have RD. Consistent results have been recorded in our countries and other studies [32,34,38,45].

This might be due to that Babies with RDS don't have a protein called surfactant that keeps small air sacs in the lungs from collapsing which increase the risk of neonatal mortality

This systematic review and meta-analysis were conducted to assess the pooled prevalence of neonatal mortality and its predictors in Ethiopia. Conducting this type of study will act as an input for program planners and policy makers working in the area of neonatal care and also indicates the quality of health care and the welfare of the society. This study is the first meta-analysis in the study area with novel findings. Even though this analysis has delivered valued evidence regarding the level of neonatal mortality and its predictors, there were some limitations, which we address below: This meta-analysis represented only studies reported from five regions of the country. Therefore, the regions may be under-represented due to the limited number of studies included.

Conclusion: Despite the variation across regions, in Ethiopia, at least one out of ten newborn had risk of death before their first birth date. Neonatal sepsis and residence were an important predictor of neonatal mortality. Therefore, based our findings, we recommend particular emphasis shall be given to the rural communities. Additionally, the government should have to strengthen any service related with reducing the neonatal mortality like expanding NICU all over the country and particular emphasis should be given for those neonate's diagnosis as preterm birth, neonatal sepsis and RDS.

Abbreviations

CI: Confidence Interval, OR: Odds Ratio, PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses, RDS: respiratory distros syndromes: Southern Nations, Nationalities, and Peoples, WHO: World Health Organization.

Declarations

Authors' contributions

YA Conception of research protocol, study design, literature review, data extraction, data analysis, interpretation and drafting the manuscript. WS, AD ,TD and TY : data extraction, quality assessment, data analysis and reviewing the manuscript. All authors have read and approved the manuscript.

Ethics approval and consent to participate

Not applicable.

Availability of data and materials

The data analyzed during the current systematic review and meta-analysis is available from the corresponding author on reasonable request.

Funding

Not applicable

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

1. **WHO. RECOMMENDED DEFINITIONS, TERMINOLOGY AND FORMAT FOR STATISTICAL TABLES RELATED TO THE PERINATAL PERIOD AND USE OF A NEW CERTIFICATE FOR CAUSE OF PERINATAL DEATHS: Modifications Recommended by FIGO as Amended October 14,1976:. Acta Obstet Gynecol Scand: . 1977;56:247-53.**
2. **Robert M. Kliegman BFS, Joseph W. St Geme III, Nina F. Schor, Richard E. Behrman. Nelson textbook of pediatrics. 20th. ed. Canada : Robert M. Kliegman; © 2016. Chapter 93, Overview of Mortality and Morbidity; p. 789-793.**
3. You D, Hug L, Ejdemyr S, Beise J: **Levels and trends in child mortality. Estimates developed by the UN Inter-agency Group for Child Mortality Estimation (IGME). Report 2015.** 2015.
4. Horbar JD, Carpenter JH, Badger GJ, Kenny MJ, Soll RF, Morrow KA, Buzas JS: **Mortality and neonatal morbidity among infants 501 to 1500 grams from 2000 to 2009.** *Pediatrics* 2012, **129**(6):1019-1026.
5. **Central Statistical Agency (CSA) [Ethiopia] and ICF. 2016. Ethiopia Demographic and Health Survey 2016. Addis Ababa, Ethiopia, and Rockville, Maryland, USA: CSA and ICF. .**
6. Unicef: **Committing to child survival: a promise renewed.** In.: eSocialSciences; 2015.
7. Chow S, Chow R, Popovic M, Lam M, Popovic M, Merrick J, Margalit RNS, Lam H, Milakovic M, Chow E: **A selected review of the mortality rates of neonatal intensive care units.** *Frontiers in public health* 2015, **3**.
8. Hannah H Chang JL, Hannah Blencowe, Catherine Y Spong, Christopher P Howson, Sarah Cairns-Smith, Eve M Lackritz, Shoo K Lee, Elizabeth Mason, Andrew C Serazin, Salimah Walani, Joe Leigh Simpson, Joy E Lawn: **Preventing preterm births: analysis of trends and potential**

- reductions with interventions in 39 countries with very high human development index.** *Lancet* 2013, **381**(223):12.
9. Gebreslasie. K: **Preterm Birth and Associated Factors among Mothers Who Gave Birth in Gondar Town Health Institutions.** *Advances in Nursing: Hindawi Publishing cooperation* 2016.
 10. Bekele I DTaDK: **Prevalence of Preterm Birth and its Associated Factors among Mothers Delivered in Jimma University Specialized Teaching and Referral Hospital, Jimma Zone, Oromia Regional State, South West Ethiopia:.** *J Women's Health Care:* 2017, **6**(1).
 11. WHO.: **Born Too Soon The Global Action Report on Preterm Birth.** In. Geneva; 2012: 128.
 12. WHO: **Revised Fact Sheet: Preterm births.** In. Geneva; 2016.
 13. Stay Beck DW, Lale Say, Ana Pilar Betran, Mario Meriardi, Jennifer Harris Requejo, Craig Rubens, Ramkumar Menon & Paul FA Van Look: **The worldwide incidence of preterm birth: a systematic review of maternal mortality and morbidity.** *Bull World Health Organ* 2010, **88**(31-38):8.
 14. Akilew Awoke Adane^{1*} TAA, Leta Gedefaw Ararsa², Bikes Destaw Bitew³ and Berihun Megabiaw Zeleke.: **Adverse birth outcomes among deliveries at Gondar University Hospital, Northwest Ethiopia:.** *BMC Pregnancy and Childbirth:* 2014, **14**(90).
 15. **UNICEF:A fair chance for every child: The State of The World's Children 2016, developed by UNICEF country and regional offices and headquarters divisions,UNCIF <data.unicef.org>. ISBN: 978-92-806-4838-6.**
 16. Gizaw M, Molla M, Mekonnen W: **Trends and risk factors for neonatal mortality in Butajira District, South Central Ethiopia,(1987-2008): a prospective cohort study.** *BMC pregnancy and childbirth* 2014, **14**(1):64.
 17. Kokeb M, Desta T: **Institution Based prospective cross-sectional study on patterns of neonatal morbidity at Gondar University Hospital Neonatal Unit, North-West Ethiopia.** *Ethiopian journal of health sciences* 2016, **26**(1):73-79.
 18. Orsido TT, Asseffa NA, Berheto TM: **Predictors of Neonatal mortality in Neonatal intensive care unit at referral Hospital in Southern Ethiopia: a retrospective cohort study.** *BMC Pregnancy and Childbirth* 2019, **19**(1):83.
 19. Yehuala S, Tekla Z: **Survival Analysis of Premature Infants Admitted to Neonatal Intensive Care Unit (NICU) in Northwest Ethiopia using Semi-Parametric Frailty Model.** *Journal of Biometrics & Biostatistics* 2015, **6**(1):1.
 20. United nations: **Transforming our world: the 2030 Agenda for Sustainable Development,.** 2015.
 21. Child. EWE: **THE GLOBAL STRATEGY FOR WOMEN'S, CHILDREN'S AND ADOLESCENTS' HEALTH (2016-2030):.** In. Geneva; 2015.
 22. March of Dimes WHO: **The Global Action Report on Preterm Birth.** In. Geneva, Switzerland; 2016.
 23. Kliegman RIN, Waldo E. (Waldo Emerson): **Nelson Textbook Of Pediatrics. Chapter 91: Prematurity and Intrauterine restriction** In., edn. United states of America: Elsevier; 2011.
 24. Joy E Lawn MGG, Toni M Nunes, Craig E Rubens, Cynthia Stanton, and the GAPPS Review Group: **Global report on preterm birth and stillbirth (1 of 7): definitions, description of the burden and opportunities to improve data.** *BMC Pregnancy and Childbirth* 2010, **10**:22.
 25. Mengesha HG, Sahle BW: **Cause of neonatal deaths in Northern Ethiopia: a prospective cohort study.** *BMC Public Health* 2017, **17**(1):62.
 26. **Munn Z, Moola S, Riitano D, Lisy K: The development of a critical appraisal tool for use in systematic reviews addressing questions of prevalence.** *International journal of health policy and management* 2014, **3**(3):123.

27. **Modesti PA, Reboldi G, Cappuccio FP, Agyemang C, Remuzzi G, Rapi S, Perruolo E, Parati G: Panethnic differences in blood pressure in Europe: a systematic review and meta-analysis. PLoS One 2016, 11(1):e0147601.**
28. **DerSimonian R, Kacker R. Random-effects model for metaanalysis of clinical trials: an update. Contemp Clin Trials. 2007;28(2):105–14.**
29. **Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. BMJ Br Med J. 2003;327(7414):557.**
30. **Peters JL, Sutton AJ, Jones DR, Abrams KR, Rushton L. Comparison of two methods to detect publication bias in meta-analysis. JAMA. 2006;295(6):676–80.**
31. **Yismaw AE, Gelagay AA, Sisay MM: Survival and predictors among preterm neonates admitted at University of Gondar comprehensive specialized hospital neonatal intensive care unit, Northwest Ethiopia. Italian journal of pediatrics 2019, 45(1):4.**
32. **Yismaw AE, Tarekegn AA: Proportion and factors of death among preterm neonates admitted in University of Gondar comprehensive specialized hospital neonatal intensive care unit, Northwest Ethiopia. BMC research notes 2018, 11(1):867.**
33. **Demisse AG, Alemu F, Gizaw MA, Tigabu Z: Patterns of admission and factors associated with neonatal mortality among neonates admitted to the neonatal intensive care unit of University of Gondar Hospital, Northwest Ethiopia. Pediatric health, medicine and therapeutics 2017, 8:57.**
34. **Andargie G, Berhane Y, Worku A, Kebede Y: Predictors of perinatal mortality in rural population of Northwest Ethiopia: a prospective longitudinal study. BMC public health 2013, 13(1):168.**
35. **Mengesha HG, Lerebo WT, Kidanemariam A, Gebrezgiabher G, Berhane Y: Pre-term and post-term births: predictors and implications on neonatal mortality in Northern Ethiopia. BMC nursing 2016, 15(1):48.**
36. **Mengesha HG, Wuneh AD, Lerebo WT, Tekle TH: Survival of neonates and predictors of their mortality in Tigray region, Northern Ethiopia: prospective cohort study. BMC pregnancy and childbirth 2016, 16(1):202.**
37. **Debelew GT, Afework MF, Yalew AW: Determinants and causes of neonatal mortality in Jimma zone, southwest Ethiopia: a multilevel analysis of prospective follow up study. PLoS One 2014, 9(9):e107184.**
38. **Wesenu M, Kulkarni S, Tilahun T: Modeling Determinants of Time-To-Death in Premature Infants Admitted to Neonatal Intensive Care Unit in Jimma University Specialized Hospital. Annals of Data Science 2017, 4(3):361-381.**
39. **Farah AE, Abbas AH, Ahmed AT: Trends of admission and predictors of neonatal mortality: A hospital based retrospective cohort study in Somali region of Ethiopia. PloS one 2018, 13(9):e0203314.**
40. **Roro EM, Sisay MM, Sibley LM: Determinants of perinatal mortality among cohorts of pregnant women in three districts of North Showa zone, Oromia Region, Ethiopia: Community based nested case control study. BMC public health 2018, 18(1):888.**
41. **Kolola T, Ekubay M, Tesfa E, Morka W: Determinants of Neonatal Mortality in North Shoa Zone, Amhara Regional State, Ethiopia. PloS one 2016, 11(10):e0164472.**
42. **United Nations Inter-agency Group for Child Mortality Estimation (UN IGME), 'Levels & Trends in Child Mortality: Report 2017, Estimates Developed by the UN Inter-agency Group for Child Mortality Estimation', United Nations Children's Fund, New York, 2017.**
43. **Bako B, Idrisa A, Garba M, Pius S, Obetta H: Determinants of neonatal survival following preterm delivery at the University of Maiduguri Teaching Hospital, Maiduguri, Nigeria. Tropical Journal of Obstetrics and Gynaecology 2017, 34(1):39.**

44. Kalimba EM: **Survival of extremely low birth weight infants at Charlotte Maxeke Johannesburg Academic Hospital**. 2014.
45. Ibrahimou B, Kodali S, Salihu H: **Survival of Preterm Singleton Deliveries: A Population-Based Retrospective Study**. *Advances in Epidemiology* 2015, **2015**.
46. Basiri B, Ashari FE, Shokouhi M, Sabzehei MK: **Neonatal Mortality and its Main Determinants in Premature Infants Hospitalized in Neonatal Intensive Care Unit in Fatemeh Hospital, Hamadan, Iran**. *Journal of Comprehensive Pediatrics* 2015, **6**(3).
47. Ghorbani F, Heidarzadeh M, Dastgiri S, Ghazi M, Rahkar Farshi M: **Survival of Premature and Low Birth Weight Infants: A Multicenter, Prospective, Cohort Study in Iran**. *Iranian Journal of Neonatology IJN* 2017, **8**(1):16-22.
48. Wesenu M, Kulkarni S, Tilahun T: **Modeling Determinants of Time-To-Death in Premature Infants Admitted to Neonatal Intensive Care Unit in Jimma University Specialized Hospital**. *Annals of Data Science* 2017:1-21.
49. Field D, Boyle E, Draper ES, Evans A, Johnson S, Khan K, Manktelow B, Marlow N, Petrou S, Pritchard C: **Towards reducing variations in infant mortality and morbidity: a population-based approach**. *Programme Grants for Applied Research* 2016, **4**(1):1-218.
50. Jacob J, Kamitsuka M, Clark RH, Kelleher AS, Spitzer AR: **Etiologies of NICU deaths**. *Pediatrics* 2014:peds. 2014-2967.
51. Shariati MK, Karimi Z, Rezaiejad M, Basiri A, Torkestani F, Gargari SS: **Perinatal complications associated with preterm deliveries at 24 to 33 weeks and 6 days gestation (2011-2012): A hospital-based retrospective study**. *Iranian journal of reproductive medicine* 2015, **13**(11):697.
52. Ntock FN, Angong FTE, Tochie JN, Mbuagbaw L, Temgoua MN, Ndombo PK, Ekei QM: **A cohort analysis of neonatal hospital mortality rate and predictors of neonatal mortality in a sub-urban hospital of Cameroon**. *Italian journal of pediatrics* 2017, **43**(1):52.
53. Khan U, Saeed MH, Bangash H, Kifayat A: **THE MORBIDITY AND MORTALITY PATTERNS OF PRETERM INFANTS AT A TERTIARY CARE HOSPITAL IN PESHAWAR: A MEDICAL RECORD REVIEW**. *KJMS* 2015, **8**(2):154.
54. Geronimus AT: **The effects of race, residence, and prenatal care on the relationship of maternal age to neonatal mortality**. *American Journal of Public Health* 1986, **76**(12):1416-1421.
55. Ballot DE, Chirwa T: **Factors Associated with Survival to Discharge of Newborns in a Middle-Income Country**. In: *Epidemiology of Communicable and Non-Communicable Diseases- Attributes of Lifestyle and Nature on Humankind*. edn.: InTech; 2016.
56. Haghighi L, Nojomi M, Mohabbatian B, Najmi Z: **Survival predictors of preterm neonates: Hospital based study in Iran (2010-2011)**. *Iranian journal of reproductive medicine* 2013, **11**(12):957.
57. Behrman RE, Koops BL, Morgan LJ, Battaglia FC: **Neonatal mortality risk in relation to birth weight and gestational age: update**. *The Journal of pediatrics* 1982, **101**(6):969-977.