

A Drug Combination Approach Targeting Both Growing Bacteria and Dormant Persisters Eradicate Persistent *Staphylococcus aureus* Biofilm Infection

Rebecca Yee^a, Yuting Yuan^a, Andreina Tarff^b, Cory Brayton^c, Naina Gour^d, Jie Feng^a, Wanliang Shi^a, and Ying Zhang^a

^aDepartment of Molecular Microbiology and Immunology, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD

^bThe Wilmer Eye Institute, Johns Hopkins University School of Medicine, Baltimore, MD

^cDepartment of Comparative Medicine, Johns Hopkins University School of Medicine, Baltimore, MD

^dThe Solomon H. Snyder Department of Neuroscience and the Center for Sensory Biology, Johns Hopkins University School of Medicine, Baltimore, MD

Abstract

Staphylococcus aureus can cause a variety of infections, many of which involve biofilm infections. Inside biofilms, growing and non-growing bacteria such as persisters co-exist, making it challenging to completely eradicate a persistent and recurrent infection with current treatments. Despite the clinical relevance, most of the current antibiotic treatments mainly kill the growing bacteria and have poor activity against non-growing persister bacteria and thus have limited effect on treating persistent infections including biofilm infections. We previously proposed a Yin-Yang model using a drug combination approach targeting both growing bacteria and persister bacteria for more effective clearance of persistent infections. Here, as a proof of principle, we showed that combining drugs that have high activity against growing forms, such as vancomycin or meropenem, with drugs that have robust anti-persister activity, such as clinafloxacin and oritavancin, could completely eradicate *S. aureus* biofilm bacteria in vitro. In contrast, single or two drugs including the current treatment for persistent *S. aureus* infection doxycycline plus rifampin failed to kill all biofilm bacteria in vitro. We then developed a chronic persistent skin infection mouse model with biofilm-seeded bacterial inocula demonstrating that biofilm bacteria caused more severe and persistent skin lesions than log phase *S. aureus* bacteria. More importantly, we found that the drug combination which eradicated biofilm bacteria in vitro is more efficacious than current treatments and completely eradicated *S. aureus* biofilm infection in mice. The complete eradication of biofilm bacteria is attributed to the unique high anti-persister activity of clinafloxacin, which could not be replaced by other fluoroquinolones such as moxifloxacin, levofloxacin or ciprofloxacin. Our study is the first to demonstrate that the combination of meropenem, daptomycin, plus clinafloxacin completely cleared the persistent infection, healed the lesions, and had less inflammation, while mice treated with doxycycline plus rifampin, the current clinically recommended treatment for chronic tissue infection, failed to do so. We also compared our persister drug combination with other approaches for treating persistent infections including gentamicin+fructose and ADEP4+rifampin in the *S. aureus* biofilm infection mouse model. Neither gentamicin+fructose nor ADEP4+rifampin could eradicate or cure the persistent biofilm infection in mice. In contrast, our drug combination regimen with persister drug clinafloxacin plus meropenem and daptomycin completely eradicated and cured the persistent biofilm infection in 7 days. An unexpected observation is that ADEP4 treatment group developed worsened skin lesions and caused more extensive pathology than the untreated control mice. Our study demonstrates an important treatment principle for persistent infections by targeting both growing and non-growing heterogeneous bacterial populations utilizing persister drugs for more effective eradication of persistent and biofilm infections. Our findings may have implications for improved treatment of many other persistent infections in general.

16 Introduction

17
18 Methicillin-resistant *Staphylococcus aureus* (MRSA) strains are highly prevalent in healthcare and community-
19 acquired staphylococcal infections¹. The mortality rate associated with MRSA infection is as high as 40%^{2,3}.
20 As an opportunistic pathogen, *S. aureus* is the most common cause of skin infections and can also cause chronic
21 infections such as endocarditis, osteomyelitis, and prosthetic joint infections⁴⁻⁶. In particular, indwelling
22 devices are conducive to biofilm formation, complicating treatment and leading to prolonged infections.
23 Globally, persistent and chronic infections are a huge burden to public health as they increase the length of
24 hospital stay, cause relapse, cost of treatment, and risk of death by at least three-folds⁷.

25
26 Bacteria in biofilms are more tolerant to antibiotics compared to planktonic cells⁸. Studies have shown that
27 antibiotics do indeed penetrate the biofilm but they do not always kill the bacteria, suggesting that tolerance to
28 treatment is not due to impaired antibiotic penetration or genetic resistance^{9,10}, but due to dormant non-growing
29 or slowing growing persister bacteria. Bacteria inside the biofilm are quite heterogeneous as some cells grow
30 slowly, which are representative of stationary phase bacteria, while others form dormant persister cells due to
31 the high cell density, nutrient and oxygen limiting environment inside the biofilm matrix¹¹.

32
33 First described in 1942, Hobby et al. found that while 99% of *S. aureus* cells were killed by penicillin about 1%
34 of residual metabolically quiescent or dormant cells called persister cells were not killed¹². The persister cells
35 were not resistant to penicillin and hence, did not undergo genetic changes but were phenotypic variants that
36 became tolerant to antibiotics¹³. Similarly, a clinical observation was also made as penicillin failed to clear
37 chronic infections due to the presence of persister cells found in patients¹³. While the mechanisms of *S. aureus*
38 persistence were largely unknown for a long time, recent studies have shown that pathways involved in quorum
39 sensing, pigmentation production, and metabolic processes such as oxidative phosphorylation, glycolysis,
40 amino acid and energy metabolism¹⁴⁻¹⁸ are responsible. Understanding the pathways of persistence would
41 facilitate development of novel drugs and therapeutic approaches to more effectively eradicate persistent
42 bacterial infections.

43
44 Despite the observation of persister bacteria from 1940s and their implications in causing prolonged treatment
45 and post-treatment relapse, the importance of persister bacteria in clinical setting has been ignored largely
46 because no persister drugs have been found that can shorten treatment duration and reduce relapse in clinically
47 relevant persistent infections. The importance of persister drug to more effectively cure persistent infections is
48 only recognized recently in the case of tuberculosis persister drug pyrazinamide (PZA) which shortens the
49 tuberculosis treatment from 9-12 months to 6 months after its inclusion in a drug combination setting¹⁹. PZA's
50 activity in killing persisters, unlike the other drugs used to treat tuberculosis, is crucial in developing a shorter
51 treatment¹⁹⁻²². A drug like PZA validates an important principle of use of a persister drug in combination with
52 other drugs targeting both persisters and growing cells in formulating an effective therapy for chronic persistent
53 infections²³⁻²⁵. More recently, a similar approach has been used to identify effective drug combinations to
54 eradicate biofilm-like structures consisting of heterogeneous cells of *Borrelia burgdorferi* *in vitro*²⁶.

55
56 Using this approach, in a recent study aimed at identifying drugs targeting non- growing persisters, we used
57 stationary phase culture of *S. aureus* as a drug screen model and identified several drugs such as clinafloxacin
58 and tosufloxacin with high activity against *S. aureus* persisters²⁷. However, their activities alone and in drug
59 combinations in killing biofilms have not been evaluated *in vitro* or in related infections caused by *S. aureus* *in vivo*.
60 In this study, we developed drug combinations that can more effectively eradicate *S. aureus* biofilms by
61 formulating drug combinations that have high activities against growing bacteria and non-growing persisters in
62 a biofilm model *in vitro* initially. Then, we established a persistent skin infection mouse model for *S. aureus*
63 using "biofilm seeding" and evaluated drug combinations in clearing the biofilm infection in this persistent skin
64 infection model. Here, we show that combining meropenem and daptomycin targeting growing bacteria, with
65 clinafloxacin targeting persister bacteria led to complete eradication of *S. aureus* biofilm not only *in vitro* but
66 more importantly also *in vivo* in a murine model of persistent skin infection, whereas other approaches for

37 treating persistent infections such as aminoglycoside plus sugar or ADEP4 plus rifampin and the currently
38 recommended drug combination treatment without persister drugs failed to do so.

39 40 **Results**

41 *Commonly used treatments for MRSA have poor activity against biofilms in vitro*

42
43
44 While vancomycin is highly effective in killing MRSA bacteria in vitro, monotherapy with vancomycin may
45 not be the most effective in clearing chronic infections with *S. aureus*. For conditions such as osteomyelitis and
46 prosthetic joint infections, treatment with vancomycin as a monotherapy or drug combination for at least 6
47 weeks are recommended. Drug combination such as doxycycline + rifampin for up to 10 days, vancomycin +
48 gentamicin + rifampin for at least 6 weeks are recommended to treat chronic infections such as recurrent tissue
49 infections and endocarditis on prosthetic valves, respectively (The Johns Hopkins Antibiotics Guide). We first
50 evaluated the activity of the above drugs in killing biofilm bacteria *in vitro* using traditional bacterial cell counts
51 (Fig. 1A), viability assessment by SYBR Green I/Propidium Iodide staining that has been developed to screen
52 for drugs targeting borrelia persister bacteria²⁸ (Fig. 1B), and staining of absolute biofilm (Fig. 1C-G). We
53 found that such clinically used combinations are not completely effective against biofilms. After 4-day
54 treatment, biofilm bacteria were not completely eradicated by any of current treatments with vancomycin alone,
55 or doxycycline + rifampin or vancomycin+gentamicin+rifampin as shown by significant numbers of bacteria
56 remaining (Fig. 1). However, it is worth noting that vancomycin+gentamicin+rifampin was more active than
57 vancomycin alone or doxycycline + rifampin in killing biofilm bacteria (Fig. 1A).

58 59 *Identification of drug combinations with strong anti-biofilm activity*

60
61 To address the clinical unmet need of better treatments against persistent infections, we hypothesize that a drug
62 combination that includes drugs that act on growing bacteria such as cell wall (e.g. vancomycin, meropenem) or
63 cell membrane inhibitors (e.g. daptomycin) plus a drug that acts on persister bacteria will be a more potent drug
64 combination in eradicating biofilm bacteria. Previous studies from our lab identified tosufloxacin and
65 clinafloxacin as having strong anti-persister activity against *S. aureus*²⁷. In order to identify a potent
66 combination, we tested various drug combinations that include drugs against both growing bacteria and non-
67 growing persisters in an *in vitro* biofilm model. Biofilms of *S. aureus* strain USA300, a common circulating
68 strain of community acquired-MRSA (CA-MRSA), were grown in 96-well microtiter plates to allow biofilm
69 formation on the bottom of the wells²⁹. While we previously showed that tosufloxacin had robust activity
70 against *S. aureus* persister cells, drug combination of vancomycin/meropenem + daptomycin + tosufloxacin
71 achieved only partial eradication, with 10⁵ CFU/ml in biofilms remaining after treatment (Fig. 2A-B). In
72 contrast, combination of vancomycin/meropenem + daptomycin + clinafloxacin showed absolute eradication of
73 biofilms after 4-day treatment as shown by 0 CFU and a live/dead ratio below the limit of detection (Fig. 2A-
74 B). Although we used the same molar concentration of each drug (50 μM of each drug) in our drug screen for
75 comparison of relative drug activity, to evaluate the activity of the combination in a more clinically relevant
76 manner, we treated the biofilms with the drugs at their C_{max} concentrations (Table 1). Our findings with C_{max}
77 drug concentrations were confirmatory as the combination of vancomycin/meropenem + daptomycin +
78 clinafloxacin still achieved complete eradication while our no treatment control and the clinically used
79 combination of doxycycline + rifampin could not. The clearance of biofilms was confirmed by both CFU counts
80 and viability staining (Fig. 2C-D).

81
82 We then tested the potential of the drug combination of meropenem + daptomycin + clinafloxacin to eradicate
83 biofilm bacteria from different MRSA *S. aureus* strains, including other CA-MRSA clinical isolates CA-409,
84 CA-127, and hospital-acquired MRSA strain GA-656. Complete eradication (0 CFU/ml) and undetectable
85 levels of live cells (under the limit of detection) were found for all of the MRSA strains tested after 4 days of
86 treating biofilms with our combination meropenem + daptomycin + clinafloxacin (Fig. 2E-F).

87 88 *Unique anti-persister activity of clinafloxacin that could not be replaced by other fluoroquinolone drugs*

19
20 Clinafloxacin is a member of the fluoroquinolone class of antibiotics which inhibits DNA replication by binding
21 to DNA gyrase. As our results suggest (Fig. 1), clinafloxacin is a powerful anti-persister drug. We then wanted
22 to rank the anti-biofilm activity of different fluoroquinolones to determine whether the robust anti-biofilm
23 activity of clinafloxacin used in combination is unique to the drug itself or can be replaced by other members of
24 fluoroquinolone antibiotics. To do so, we used the *S. aureus* Newman strain due to its susceptibility to many
25 fluoroquinolones as we wanted to eliminate any confounding factors due to inherent drug resistance. While
26 other fluoroquinolones such as ciprofloxacin, levofloxacin, and moxifloxacin had certain anti-persister or anti-
27 biofilm activity when used in combination with meropenem and daptomycin after 4-days of treatment, the drug
28 combination with clinafloxacin was indeed the most active and was the only combination that achieved
29 complete sterilization. By contrast, biofilms treated with combinations consisting of other quinolones still
30 harbored 10^4 - 10^8 CFU/ml. When used in combination, the activity of the quinolones from strongest to weakest
31 as ranked by both viability assessment and viable cell counts is as follows: clinafloxacin, ciprofloxacin,
32 moxifloxacin, and levofloxacin (Table 2). Hence, clinafloxacin has unique potent activity against persisters
33 compared to other fluoroquinolone counterparts.

34 *Anti-persister activity of oritavancin and dalbavancin*

35
36 Thus far, our data suggest that inclusion of a drug with great anti-persister activity can be beneficial in killing
37 biofilm bacteria (Fig. 1, Fig. 2, and Table 2). To identify other potential anti-persister candidates, we turned to
38 the new generation of lipoglycopeptides such as oritavancin and dalbavancin. These drugs have multiple
39 mechanisms of action: inhibition of transglycosylation, transpeptidation, and cell membrane disruption, a
40 property of persister drugs¹⁹. We first tested the activity of oritavancin and dalbavancin in killing *S. aureus*
41 persisters in comparison with its parent counterpart vancomycin, and the results revealed oritavancin was the
42 best in killing persisters among the three drugs (Fig. 3A). After 6-day drug exposure, oritavancin killed 10^6
43 CFU/ml of persisters as compared with dalbavancin or vancomycin which killed only about 10^2 CFU/ml.

44
45 Since oritavancin showed strong anti-persister activity, we next evaluated oritavancin's activity in drug
46 combinations. After replacing clinafloxacin with oritavancin, we observed that the combination of meropenem
47 + daptomycin + oritavancin exhibited partial activity against biofilms, a decrease of 10^5 CFU/ml, which is much
48 better than the activity achieved by treatment with single drugs or two-drug combinations, but still inferior to
49 the clinafloxacin combination (Fig. 3B).

50
51 Due to oritavancin's strong activity against growing phase *S. aureus* (MIC of 0.03 mg/L)³⁰ and its dual
52 mechanism of action that mimics cell wall + cell membrane inhibitors in our drug combination, we tested
53 oritavancin in place of meropenem and daptomycin. Surprisingly, the combination of oritavancin +
54 clinafloxacin was also able to achieve complete eradication of biofilms suggesting that oritavancin can replace
55 the component in our drug combination that targets actively growing bacteria (Fig. 3C). It is also important to
56 note that single drug of oritavancin cannot kill biofilms (no change in CFU after 4 day treatment) which further
57 validates the importance of drug combinations in biofilm bacteria.

58
59 To compare the activity of the three combinations tested thus far with clinafloxacin, we performed a time-
60 course kill experiment which revealed that oritavancin + clinafloxacin can kill all biofilms by 2-day treatment
61 whereas it took 4 days for meropenem/vancomycin + daptomycin + clinafloxacin to eradicate the biofilm
62 bacteria (Fig. 3D). Overall, our data suggest that inclusion of an anti-persister drug in a drug combination to
63 treat biofilms is paramount and these combinations possess better activity than current clinically used regimens
64 based on our in vitro studies.

65
66 *The drug combination meropenem + daptomycin + clinafloxacin eradicated biofilm infections in the mouse skin*
67 *persistent infection model*

10 Given the robust activity of our drug combinations in eradicating biofilms *in vitro*, we were interested to know
11 if our combination can also eradicate persistent infections *in-vivo*. To evaluate the efficacy of our drug
12 combinations in treating the persistent skin infection, we chose to infect mice with biofilm bacteria from *S.*
13 *aureus* strain USA300, a clinical strain most representative to causing persistent infections in a host. We
14 allowed the infection to develop for 7 days, followed by 7 day treatment with different regimens (Fig. 4A).
15 Previously, we have shown that mice infected with biofilm bacteria developed more chronic skin lesions³¹.
16 Administration of the combination of doxycycline + rifampin (a control group as a clinically used treatment) or
17 drug combination vancomycin + daptomycin + clinafloxacin decreased the bacteria load (about 1-log of bacteria)
18 but did not clear the infection (Fig. 4B). Other treatments which supposedly eradicate chronic *S. aureus*
19 infections such as ADEP4+rifampin³² or fructose+gentamicin³³ did not show sterilizing activity in our biofilm
20 infection model and instead, had increased lesion size and inflammation (Fig. 4C). Remarkably, our
21 combination of meropenem + daptomycin + clinafloxacin cleared the infection completely, decreased the size
22 of lesions, and reduced histopathology scores, and healed the lesions completely (Fig. 4D-G).

14 Because our *in-vivo* experiments were done using an MRSA strain USA300, we wanted to infect mice with a
15 methicillin susceptible *S. aureus* Newman strain (MSSA) to ensure that our drug combination is effective for
16 other *S. aureus* strains. In addition, we wanted to further confirm that clinafloxacin's activity, compared to other
17 quinolones, is superior despite the difference in background sensitivity of the bacterial strain. Despite
18 moxifloxacin and clinafloxacin having the same MIC for the Newman strain, the combination of meropenem +
19 daptomycin + moxifloxacin was not effective in clearing the biofilm infection whereas the combination of
20 meropenem + daptomycin + clinafloxacin indeed cleared the infection completely (Fig. 4H). This indicates
21 clinafloxacin combination works for both MRSA strain and MSSA strain and its unique sterilizing activity
22 cannot be replaced by moxifloxacin.

24 *Eradication of biofilm bacteria correlates with resolution of tissue inflammation and immunopathology*

26 Skin infections caused by *S. aureus* are cleared by neutrophil recruitment driven by IL-17 production³⁴. To
27 evaluate any potential immunopathology consequences of our treatment, we measured the levels of IL-17 and
28 proinflammatory cytokine IL-1 at the infection site and performed histology. Skin tissues of mice treated with
29 our drug combination of meropenem + daptomycin + clinafloxacin produced the lowest amount of IL-17 (Fig.
30 5A) and IL-1 (Fig. 5B) and showed the least amount of gross inflammation (Fig. 5C) compared to other control
31 treatments. These data support our hypothesis that a drug combination targeting both growing (e.g. meropenem,
32 daptomycin) and persister cells (e.g. clinafloxacin) is essential in clearing chronic infections *in-vivo* such as a
33 persistent skin infection caused by *S. aureus* biofilm.

35 **Discussion**

37 Numerous studies have documented how resilient biofilms and biofilm infections are to antibiotic treatments³⁵⁻
38 ³⁷. Since persister cells that are embedded in the biofilm are mostly responsible for recalcitrance of biofilms to
39 antibiotic treatments, many attempts have been made to identify novel treatments and synthetic compounds that
40 kill bacterial persisters^{38,39}. Some approaches include resuscitating or altering the metabolic status of persisters
41 ^{40,41} or enhancing the activity of aminoglycoside antibiotic with sugars³³, or activating protease by ADEP4 plus
42 rifampin³². Although these new therapeutic approaches showed promising results *in vitro* and in some cases, *in-*
43 *vivo*,^{32,38,42} not all treatments achieved sterilizing activity and their utility in more persistent biofilm infections
44 remains to be confirmed. The animal models used either rely on immunosuppressant agents or a short term
45 infection that do not reflect true persistent infections clinically. Here, we established a more relevant persistent
46 infection mouse model with biofilm inocula that mimic the human infections without the use of
47 immunosuppressant agents. Previous studies have mostly used log phase bacteria as inocula for infection in
48 animal models, however, in this study, we showed that biofilm inocula produced a more severe lesion and more
49 persistent infection than the log phase bacteria³¹. This biofilm-inocula model could serve as a useful model for
50 evaluating treatment regimens against biofilm infections *in vivo* in general. Importantly, we were able to show
51 that single drugs or even two drug combinations can this study, to identify more effective regimens to treat

52 chronic *S. aureus* infections, we first identified several drug combinations that are more active in killing
53 biofilms *in vitro* than currently recommended regimens (e.g. vancomycin alone, doxycycline + rifampin, and
54 vancomycin + gentamicin + rifampin) used clinically. Then, we confirmed the potent activity of the
55 combination meropenem + daptomycin + clinafloxacin in our newly established chronic, skin infection mouse
56 model.

57
58 Previously, we identified both clinafloxacin and tosufloxacin as having robust activity against *S. aureus*
59 persisters²⁷. However, in our drug combination studies, clinafloxacin used in combination displayed greater
60 activity against biofilms than tosufloxacin. This could potentially be explained by the genetic background and
61 inherent antibiotic resistance of the strain tested. Previously, tosufloxacin was identified to have great persister
62 activity in the background of the Newman strain²⁷, a methicillin-sensitive *S. aureus* strain whereas our biofilm
63 experiments conducted here used USA300, a MRSA strain. The ability between these strains to form biofilms
64 may also be a contributing factor⁴³. Nonetheless, despite being unable to achieve complete eradication, it must
65 be noted that inclusion of any anti-persister drug in combination (clinafloxacin, tosufloxacin, or oritavancin)
66 with drugs that can kill growing bacteria can kill more bacteria in biofilms than currently approved regimens,
67 confirming the importance of targeting the heterogeneous of bacterial populations in developing more effective
68 treatments.

69
70 We showed that meropenem + daptomycin + clinafloxacin achieved sterilizing activity in a chronic, skin
71 infection model in mice. The strong activity of meropenem or daptomycin against *S. aureus* growing bacteria is
72 indisputable as most *S. aureus* strains have a relatively low MIC to these drugs^{44,45}. The inclusion of drugs with
73 such strong activity against active bacteria allows for rapid killing of growing bacteria in the population. To kill
74 non-growing biofilm persisters, clinafloxacin is crucial in the combination. As shown here, the MICs for
75 clinafloxacin for both the MSSA Newman strain and MRSA USA300 strain were both under 0.25 µg/ml. Other
76 studies have shown that MICs for streptococci are from 0.06-0.12 µg/ml and for enterococci is 0.5 µg/ml. Both
77 oral and intravenous formulations have been developed^{46,47}. Although not commonly used, clinafloxacin
78 administration drastically improved the condition of a cystic fibrosis patient who had a chronic *Burkholderia*
79 *cenoepecia* infection and was not responding to different antibiotic treatment⁴⁸. A human trial with patients
80 having native or prosthetic valve endocarditis also showed that clinafloxacin was an effective treatment⁴⁹. As a
81 quinolone, clinafloxacin inhibits bacterial DNA gyrase and topoisomerase IV but not all quinolones have anti-
82 persister activity (Table 1). Comparing the chemical structure of clinafloxacin to the other quinolones that have
83 weak anti-persister activity (ciprofloxacin, tosufloxacin, moxifloxacin, and levofloxacin), a chloride group
84 attached to the benzene ring appears to be unique to only clinafloxacin (Fig. 6). Further studies to explore the
85 mechanism of clinafloxacin's unique ability to kill persisters requires further investigation.

86
87 While meropenem and daptomycin are our agents directed at killing growing bacteria, it is important to note
88 that these drugs also have some activity against persisters. Meropenem used in combination with polymyxin B
89 has been shown to eradicate persisters in *Acinetobacter baumannii* strains⁵⁰. Similarly, daptomycin has been
90 shown to be active against *S. aureus* biofilms found on implants⁵¹. Daptomycin in combination with
91 doxycycline and cefoperazone or cefuroxime have been shown to kill biofilm-like microcolonies of *B.*
92 *burgdorferi*^{26,52}. Its mechanism of action in disrupting membrane structure and rapid depolarization of the
93 membrane may impact the viability of persisters and thus, play an important role in the combination⁵³.

94
95 It is important to note that the chronic infection status of our mice is a key component to our disease model. The
96 phenotype of a more severe disease caused by persistent forms is an important observation³¹ and our model
97 could potentially better mimic chronic infections in humans. While Conlon et al. also used stationary phase
98 inocula (10⁶) to infect their mice and caused a deep-seated infection, the infection was allowed to develop for
99 only 24 hours before treatment which cannot be a persistent infection; and the mice were made neutropenic³², a
100 condition that may not apply to a majority of patients suffering from chronic *S. aureus* infections. Such
101 differences in the animal models may explain why ADEP4 + rifampin which was claimed to have sterilizing
102 activity by Conlon et al. failed to eradicate the persistent infection in our model established with biofilm inocula
103 and infection allowed to develop for 1 week before treatment. The combination of an aminoglycoside + sugar

was shown to be effective in an *E. coli* urinary tract infection model³³ but unfortunately this approach was not effective in our biofilm infection model either. Allison *et al.* showed that gentamicin + fructose reduced 1.5 fold of *S. aureus* biofilms *in vitro* after 4 hours of treatment³³ but was not tested in animals. In our study here, we showed that mice treated for 7 days with gentamicin + fructose still harbored 10⁵ CFU/ml in skin tissues and showed an increase in lesion size despite the treatment. In both these cases, the discrepancy could be due to differences in the disease model, as ours is a more persistent biofilm skin infection model established with biofilm inocula and would be expected to be more difficult to cure than the other studies that did not use biofilm inocula for the infection.

Our combination of meropenem + daptomycin + clinafloxacin showed sterilizing activity in mice after one week based on the concentrations of drugs and dosing regimens commonly found in literature (Table 2). Higher, yet safe, concentrations of drugs should be tested to see whether or not a shorter treatment period can achieve similar eradication, which would be beneficial for patients undergoing therapy. Further PK/PD studies of our drug combination are needed. Moreover, meropenem, daptomycin, and clinafloxacin are intravenous drugs and not convenient to administer. Future studies to develop oral regimens as effective as the identified combinations are needed for more convenient administration. Our *in vitro* data suggested that oritavancin used in combination with clinafloxacin had robust activity against biofilms, killing all the bacteria in the biofilms (10⁷ CFU) after a short treatment of 2 days. The administration of oritavancin is a single 1200-mg dose given in a slow, 3 hour infusion, which may also be of interest for patients due to the ease of administration and long half-life. Hence, preclinical studies in mice to test oritavancin's activity in chronic infections need to be performed carefully.

Currently used regimens for treating persistent infections are lengthy and the inability to clear the bacteria in a timely fashion may also increase the chances of developing antibiotic resistance. A drug combination that has both activities against growing and persister cells as proposed in the Yin-Yang model²⁵ have promising potential in developing a more effective therapy for treating chronic persistent infections. This study validates this Yin-Yang model²⁵ and emphasizes the importance of persister drugs like clinafloxacin in eradicating a persistent infection. This treatment algorithm takes into account the heterogeneous population of bacterial cells that exists upon encountering stress. With this principle in mind, this study reports novel drug combinations that are effective in killing *S. aureus* biofilms and treating chronic infections. We established a chronic skin infection mouse model that more appropriately mimics human chronic disease. Then, we developed a triple drug combination of meropenem (or vancomycin) + daptomycin + clinafloxacin or two drug combination of oritavancin + clinafloxacin that can achieve sterilizing activity *in vitro*. We also show that administration of meropenem + daptomycin + clinafloxacin allowed the mice with chronic skin infections to completely clear the bacterial load, heal lesions completely, and show reduced pathology and inflammation. Our approach of combining drugs targeting both growing and non-growing bacteria with persister drugs to completely eradicate biofilm infections may have implications for developing better treatments against other persistent infections by other bacterial pathogens, fungi, and even cancer.

Methods

Culture media, antibiotics, and chemicals

Staphylococcus aureus strains Newman, USA300, CA-409, CA-127, and GA-656, were obtained from American Type Tissue Collections (Manassas, VA, USA), and cultivated in tryptic soy broth (TSB) and tryptic soy agar (TSA) (Becton Dickinson Franklin Lakes, NJ, USA) at 37°C. Vancomycin, gentamicin, rifampicin, levofloxacin, ciprofloxacin, moxifloxacin, and oritavancin were obtained from Sigma-Aldrich Co. (St. Louis, MO, USA). Daptomycin, meropenem, tosufloxacin, and clinafloxacin were obtained from AK Scientific, Inc. (Union City, CA, USA). Stock solutions were prepared in the laboratory, filter-sterilized and used at indicated concentrations.

Microtiter plate biofilm assays

55

56 *S. aureus* strains grown overnight in TSB were diluted 1:100 in TSB. Then, 100- μ l aliquots of each diluted
57 culture were placed into a 96-well flat-bottom microtiter plate and statically incubated for 24 h at 37°C²⁹.
58 Planktonic cells were removed and discarded from the microtiter plates. Drugs at the indicated concentrations
59 were then added to the biofilms attached to the bottom of the microtiter plate at a total volume of 100 μ l in TSB.
60 To determine the cell and biofilm density, the supernatant was removed from the well and the biofilms were
61 washed twice with PBS (1X). To enumerate bacterial cell counts, the biofilms in the wells were resuspended in
62 TSB and scraped with a pipette tip before serial dilution and plating. To assess cell viability using the ratio of
63 green:red fluorescence to determine the ratio of live:dead cells, respectively, the biofilms were stained with
64 SYBR Green I/Propidium Iodide dyes as described^{28,54}. Briefly, SYBR Green I (10,000 \times stock) was mixed
65 with PI (20 mM) in distilled H₂O at a ratio of 1:3, respectively. The SYBR Green I/PI staining mix was added
66 to each sample at a ratio of 1:10 (10 μ l of dye for 100 μ l of sample). Upon incubation at room temperature in
67 the dark for 20 min, the green and red fluorescence intensity was detected using a Synergy H1 microplate reader
68 by BioTek Instruments (Winooski, VT, USA) at excitation wavelength of 485 nm and 538 nm and 612 nm for
69 green and red emission, respectively. To visualize biofilm biomass, biofilms were stained with crystal violet
70 (0.1%) for 15 minutes at room temperature. Excess dyes were washed with water and the biofilms were left to
71 air dry. Images were recorded using Keyence BZ-X710 microscope and were processed using BZ-X Analyzer
72 software provided by Keyence (Osaka, Japan).

73 *Mouse skin infection model*

74 Female Swiss-Webster mice of 6 weeks of age were obtained from Charles River. They were housed 3 to 5 per
75 cage under BSL-2 housing conditions. All animal procedures were approved by the Johns Hopkins University
76 Animal Care and Use Committee. *S. aureus* strain USA300 and strain Newman were used in the mouse
77 infection experiments. Mice were anesthetized and then shaved to remove a patch of fur of approximately 3 cm
78 by 2 cm. Bacteria of indicated inoculum size and age were subcutaneously injected into the mice. For log phase
79 inoculum, bacteria grown overnight were diluted 1:100 in TSB and grown for 2 hrs at 37°C with shaking at 220
80 RPM. For stationary phase inoculum, overnight cultures of bacteria grown at 37°C were used. For preparation
81 of biofilm inoculum, biofilms were first grown in microtiter plates as described previously, and then
82 resuspended and scraped up with a pipette tip. Quantification of all inoculum was performed by serial dilution
83 and plating. Treatment was started after 1 week infection with different drugs and drug combinations. For
84 details on drugs, drug dosage and route of administration, please refer to Table 1. Skin lesion sizes were
85 measured at the indicated time points using a caliper. Mice were euthanized after 1 week post-treatment and
86 skin tissues were removed, homogenized, and serially diluted for bacterial counting on TSA plates.

87 *Histopathology*

88 Skin tissues were dissected, laid flat, and fixed for 24 hrs with neutral buffered formalin. Tissues were
89 embedded in paraffin, cut into 5- μ m sections, and mounted on glass slides. Tissue sections were stained with
90 hematoxylin and eosin for histopathological scoring. Tissue sections were evaluated for lesion crust formation,
91 ulcer formation, hyperplasia, inflammation, gross size, and bacterial count and were assigned a score on a 0–3
92 scale (0 = none, 1 = mild, 2 = moderate, and 3 = severe). The cumulative pathology score represented the sum
93 of each individual pathology parameter. Scoring was performed by an observer in consultation with a veterinary
94 pathologist. Representative images were taken using a Keyence BZ-X710 Microscope.

95 *Statistical analyses*

96 Statistical analyses were performed using two-tailed Student's *t*-test and two-way ANOVAs where appropriate.
97 Mean differences were considered statistically significant if *p* value was <0.05. All experiments were performed
98 in triplicates. Analyses were performed using GraphPad Prism and Microsoft Office Excel.

99

100

11

12 References

- 13 1 DeLeo, F. R., Otto, M., Kreiswirth, B. N. & Chambers, H. F. Community-associated methicillin-resistant
14 *Staphylococcus aureus*. *Lancet* **375**, 1557-1568, doi:10.1016/s0140-6736(09)61999-1 (2010).
- 15 2 Selton-Suty, C. *et al.* Preeminence of *Staphylococcus aureus* in infective endocarditis: a 1-year
16 population-based survey. *Clinical infectious diseases : an official publication of the Infectious Diseases*
17 *Society of America* **54**, 1230-1239, doi:10.1093/cid/cis199 (2012).
- 18 3 Bae, I. G. *et al.* Heterogeneous vancomycin-intermediate susceptibility phenotype in bloodstream
19 methicillin-resistant *Staphylococcus aureus* isolates from an international cohort of patients with
20 infective endocarditis: prevalence, genotype, and clinical significance. *The Journal of infectious*
21 *diseases* **200**, 1355-1366, doi:10.1086/606027 (2009).
- 22 4 Chan, L. C. *et al.* Innate Immune Memory Contributes to Host Defense against Recurrent Skin and Skin
23 Structure Infections Caused by Methicillin-Resistant *Staphylococcus aureus*. *Infection and immunity* **85**,
24 doi:10.1128/iai.00876-16 (2017).
- 25 5 David, M. Z. & Daum, R. S. Community-associated methicillin-resistant *Staphylococcus aureus*:
26 epidemiology and clinical consequences of an emerging epidemic. *Clinical microbiology reviews* **23**,
27 616-687, doi:10.1128/cmr.00081-09 (2010).
- 28 6 Dryden, M. S. Complicated skin and soft tissue infection. *The Journal of antimicrobial chemotherapy* **65**
29 **Suppl 3**, iii35-44, doi:10.1093/jac/dkq302 (2010).
- 30 7 Noskin, G. A. *et al.* The burden of *Staphylococcus aureus* infections on hospitals in the United States:
31 an analysis of the 2000 and 2001 Nationwide Inpatient Sample Database. *Archives of internal medicine*
32 **165**, 1756-1761, doi:10.1001/archinte.165.15.1756 (2005).
- 33 8 de Oliveira, A. *et al.* Antimicrobial Resistance Profile of Planktonic and Biofilm Cells of *Staphylococcus*
34 *aureus* and Coagulase-Negative Staphylococci. *International journal of molecular sciences* **17**,
35 doi:10.3390/ijms17091423 (2016).
- 36 9 Stewart, P. S., Davison, W. M. & Steenbergen, J. N. Daptomycin rapidly penetrates a *Staphylococcus*
37 *epidermidis* biofilm. *Antimicrobial agents and chemotherapy* **53**, 3505-3507, doi:10.1128/aac.01728-08
38 (2009).
- 39 10 Kirker, K. R., Fisher, S. T. & James, G. A. Potency and penetration of telavancin in staphylococcal
40 biofilms. *International journal of antimicrobial agents* **46**, 451-455, doi:10.1016/j.ijantimicag.2015.05.022
41 (2015).
- 42 11 Kavanaugh, J. S. & Horswill, A. R. Impact of Environmental Cues on Staphylococcal Quorum Sensing
43 and Biofilm Development. *The Journal of biological chemistry* **291**, 12556-12564,
44 doi:10.1074/jbc.R116.722710 (2016).
- 45 12 Hobby, G. L., Meyer, K. & Chaffee, E. Observations on the mechanism of action of penicillin. *Proc Soc*
46 *Exp Biol NY* **50**, 281-285 (1942).
- 47 13 Bigger, J. W. Treatment of staphylococcal infections with penicillin by intermittent sterilisation. *Lancet*
48 **244**, 4 (1944).
- 49 14 Wang, W. *et al.* Transposon Mutagenesis Identifies Novel Genes Associated with *Staphylococcus*
50 *aureus* Persister Formation. *Frontiers in microbiology* **6**, 1437, doi:10.3389/fmicb.2015.01437 (2015).
- 51 15 Conlon, B. P., Rowe, S. E., Gandt, A. B., Nuxoll, A. S. & Donegan, N. P. Persister formation in
52 *Staphylococcus aureus* is associated with ATP depletion. **1**, 16051, doi:10.1038/nmicrobiol.2016.51
53 (2016).
- 54 16 Yee, R., Cui, P., Shi, W., Feng, J. & Zhang, Y. Genetic Screen Reveals the Role of Purine Metabolism
55 in *Staphylococcus aureus* Persistence to Rifampicin. *Antibiotics* **4**, 627 (2015).
- 56 17 Xu, T. *et al.* The Agr Quorum Sensing System Represses Persister Formation through Regulation of
Phenol Soluble Modulins in *Staphylococcus aureus*. *Frontiers in microbiology* **8**, 2189,
doi:10.3389/fmicb.2017.02189 (2017).
- 18 Sahukhal, G. S., Pandey, S. & Elasri, M. O. msaABCR operon is involved in persister cell formation in
Staphylococcus aureus. *BMC microbiology* **17**, 218, doi:10.1186/s12866-017-1129-9 (2017).
- 19 Zhang, Y. & Mitchison, D. The curious characteristics of pyrazinamide: a review. *The international*
journal of tuberculosis and lung disease : the official journal of the International Union against
Tuberculosis and Lung Disease **7**, 6-21 (2003).
- 20 Zhang, Y., Wade, M. M., Scorpio, A., Zhang, H. & Sun, Z. Mode of action of pyrazinamide: disruption of
Mycobacterium tuberculosis membrane transport and energetics by pyrazinoic acid. *The Journal of*
antimicrobial chemotherapy **52**, 790-795, doi:10.1093/jac/dkg446 (2003).

- 57 21 Shi, W. *et al.* Pyrazinamide inhibits trans-translation in *Mycobacterium tuberculosis*. *Science (New York, N. Y.)* **333**, 1630-1632, doi:10.1126/science.1208813 (2011).
- 58
- 59 22 Zhang, S. *et al.* Mutations in panD encoding aspartate decarboxylase are associated with pyrazinamide resistance in *Mycobacterium tuberculosis*. *Emerging microbes & infections* **2**, e34, doi:10.1038/emi.2013.38 (2013).
- 60
- 61
- 62 23 Coates, A. R. & Hu, Y. Targeting non-multiplying organisms as a way to develop novel antimicrobials. *Trends Pharmacol Sci* **29**, 143-150, doi:S0165-6147(08)00018-7 [pii] 10.1016/j.tips.2007.12.001 (2008).
- 63
- 64 24 Zhang, Y., Shi, W., Zhang, W. & Mitchison, D. Mechanisms of Pyrazinamide Action and Resistance. *Microbiology spectrum* **2**, 1-12, doi:10.1128/microbiolspec.MGM2-0023-2013 (2013).
- 65
- 66 25 Zhang, Y. Persisters, Persistent Infections and the Yin-Yang Model. *Emerging Microbes and Infection* **3**, 10 (2014).
- 67
- 68
- 69 26 Feng, J., Weitner, M., Shi, W., Zhang, S. & Zhang, Y. Eradication of Biofilm-Like Microcolony Structures of *Borrelia burgdorferi* by Daunomycin and Daptomycin but not Mitomycin C in Combination with Doxycycline and Cefuroxime. *Frontiers in microbiology* **7**, 62, doi:10.3389/fmicb.2016.00062 (2016).
- 70
- 71
- 72 27 Niu, H., Peng Cui, Rebecca Yee, Wanliang Shi, Shuo Zhang, Jie Feng, David Sullivan, Wenhong Zhang, Bingdong Zhu, Ying Zhang. A Clinical Drug Library Screen Identifies Tosufloxacin as Being Highly Active against *Staphylococcus aureus* Persisters. *Antibiotics* **4**, 329-336 (2015).
- 73
- 74
- 75 28 Feng, J., Wang, T., Zhang, S., Shi, W. & Zhang, Y. An optimized SYBR Green I/PI assay for rapid viability assessment and antibiotic susceptibility testing for *Borrelia burgdorferi*. *PLoS One* **9**, e111809, doi:10.1371/journal.pone.0111809 (2014).
- 76
- 77
- 78 29 O'Toole, G. A. Microtiter dish biofilm formation assay. *Journal of visualized experiments : JoVE*, doi:10.3791/2437 (2011).
- 79
- 80 30 Pfaller, M. A., Sader, H. S., Flamm, R. K., Castanheira, M. & Mendes, R. E. Oritavancin in vitro activity against gram-positive organisms from European and United States medical centers: results from the SENTRY Antimicrobial Surveillance Program for 2010-2014. *Diagnostic microbiology and infectious disease* **91**, 199-204, doi:10.1016/j.diagmicrobio.2018.01.029 (2018).
- 81
- 82
- 83 31 Yee, R., Yuan, Y., Brayton, C., Tarff, A., Feng, J., Shi, W., Behrens, A., Zhang, Y. Biofilm/Persister/Stationary Phase Bacteria Cause More Severe Disease Than Log Phase Bacteria II Infection with Persister Forms of *Staphylococcus aureus* Causes a Chronic Persistent Skin Infection with More Severe Lesion that Takes Longer to Heal and is not Eradicated by the Current Recommended Treatment in Mice. *bioRxiv* 476465, doi:https://doi.org/10.1101/476465 (2018).
- 84
- 85
- 86 32 Conlon, B. P. *et al.* Activated ClpP kills persisters and eradicates a chronic biofilm infection. *Nature* **503**, 365-370, doi:10.1038/nature12790 (2013).
- 87
- 88
- 89 33 Allison, K. R., Brynildsen, M. P. & Collins, J. J. Metabolite-enabled eradication of bacterial persisters by aminoglycosides. *Nature* **473**, 216-220, doi:10.1038/nature10069 (2011).
- 90
- 91
- 92 34 Cho, J. S. *et al.* IL-17 is essential for host defense against cutaneous *Staphylococcus aureus* infection in mice. *The Journal of clinical investigation* **120**, 1762-1773, doi:10.1172/jci40891 (2010).
- 93
- 94
- 95 35 Tong, S. Y., Davis, J. S., Eichenberger, E., Holland, T. L. & Fowler, V. G., Jr. *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations, and management. *Clinical microbiology reviews* **28**, 603-661, doi:10.1128/cmr.00134-14 (2015).
- 96
- 97
- 98 36 Hughes, G. & Webber, M. A. Novel approaches to the treatment of bacterial biofilm infections. *British journal of pharmacology* **174**, 2237-2246, doi:10.1111/bph.13706 (2017).
- 99
- 100 37 Romling, U. & Balsalobre, C. Biofilm infections, their resilience to therapy and innovative treatment strategies. *Journal of internal medicine* **272**, 541-561, doi:10.1111/joim.12004 (2012).
- 101
- 102 38 Kim, W. *et al.* A new class of synthetic retinoid antibiotics effective against bacterial persisters. *Nature* **556**, 103-107, doi:10.1038/nature26157 (2018).
- 103
- 104 39 Mohamed, M. F., Abdelkhalek, A. & Seleem, M. N. Evaluation of short synthetic antimicrobial peptides for treatment of drug-resistant and intracellular *Staphylococcus aureus*. *Scientific reports* **6**, 29707, doi:10.1038/srep29707 (2016).
- 105
- 106 40 Zhang, Y., Yew, W. W. & Barer, M. R. Targeting persisters for tuberculosis control. *Antimicrobial agents and chemotherapy* **56**, 2223-2230, doi:10.1128/aac.06288-11 (2012).
- 107
- 108 41 Joers, A., Kaldalu, N. & Tenson, T. The frequency of persisters in *Escherichia coli* reflects the kinetics of awakening from dormancy. *Journal of bacteriology* **192**, 3379-3384, doi:10.1128/jb.00056-10 (2010).
- 109
- 110

- 11 42 Mandell, J. B. *et al.* Elimination of Antibiotic Resistant Surgical Implant Biofilms Using an Engineered
12 Cationic Amphipathic Peptide WLBU2. *Scientific reports* **7**, 18098, doi:10.1038/s41598-017-17780-6
13 (2017).
- 14 43 Cue, D. *et al.* SaeRS-dependent inhibition of biofilm formation in *Staphylococcus aureus* Newman.
15 *PLoS one* **10**, e0123027, doi:10.1371/journal.pone.0123027 (2015).
- 16 44 Humphries, R. M., Pollett, S. & Sakoulas, G. A current perspective on daptomycin for the clinical
17 microbiologist. *Clinical microbiology reviews* **26**, 759-780, doi:10.1128/cmr.00030-13 (2013).
- 18 45 CLSI. (2016).
- 19 46 Barrett, M. S., Jones, R. N., Erwin, M. E., Johnson, D. M. & Briggs, B. M. Antimicrobial activity
20 evaluations of two new quinolones, PD127391 (CI-960 and AM-1091) and PD131628. *Diagnostic*
21 *microbiology and infectious disease* **14**, 389-401 (1991).
- 22 47 Cohen, M. A., Yoder, S. L., Huband, M. D., Roland, G. E. & Courtney, C. L. In vitro and in vivo activities
23 of clinafloxacin, CI-990 (PD 131112), and PD 138312 versus enterococci. *Antimicrobial agents and*
24 *chemotherapy* **39**, 2123-2127 (1995).
- 25 48 Balwan, A., Nicolau, D. P., Wungwattana, M., Zuckerman, J. B. & Waters, V. Clinafloxacin for
26 Treatment of *Burkholderia cenocepacia* Infection in a Cystic Fibrosis Patient. *Antimicrobial agents and*
27 *chemotherapy* **60**, 1-5, doi:10.1128/aac.01428-15 (2016).
- 28 49 Levine, D. P., Holley, H. P., Eiseman, I., Willcox, P. & Tack, K. Clinafloxacin for the treatment of
29 bacterial endocarditis. *Clinical infectious diseases : an official publication of the Infectious Diseases*
30 *Society of America* **38**, 620-631, doi:10.1086/381670 (2004).
- 31 50 Gallo, S. W., Ferreira, C. A. S. & de Oliveira, S. D. Combination of polymyxin B and meropenem
32 eradicates persister cells from *Acinetobacter baumannii* strains in exponential growth. *Journal of*
33 *medical microbiology* **66**, 1257-1260, doi:10.1099/jmm.0.000542 (2017).
- 34 51 Carpenter, C. F. & Chambers, H. F. Daptomycin: another novel agent for treating infections due to
35 drug-resistant gram-positive pathogens. *Clinical infectious diseases : an official publication of the*
36 *Infectious Diseases Society of America* **38**, 994-1000, doi:10.1086/383472 (2004).
- 37 52 Feng, J., Auwaerter, P. G. & Zhang, Y. Drug Combinations against *Borrelia burgdorferi* Persists In
38 Vitro: Eradication Achieved by Using Daptomycin, Cefoperazone and Doxycycline. *PLoS One* **10**,
39 e0117207, doi:10.1371/journal.pone.0117207 (2015).
- 40 53 Pogliano, J., Pogliano, N. & Silverman, J. A. Daptomycin-mediated reorganization of membrane
41 architecture causes mislocalization of essential cell division proteins. *Journal of bacteriology* **194**, 4494-
42 4504, doi:10.1128/jb.00011-12 (2012).
- 43 54 Feng, J. *et al.* A Rapid Growth-Independent Antibiotic Resistance Detection Test by SYBR
44 Green/Propidium Iodide Viability Assay. *Frontiers in medicine* **5**, 127, doi:10.3389/fmed.2018.00127
45 (2018).
- 46
47
48
49
50
51
52
53
54




Figure 1. Clinically recommended treatments for chronic *S. aureus* infections only partially killed biofilm bacteria *in vitro*.

Treatments (all 50 μ M) of vancomycin alone, doxycycline+rifampin, and the combination of vancomycin+gentamicin+rifampin for 4 days were evaluated for biofilm killing by CFU enumeration (A) and viability staining using SYBR Green I/PI (B). Images of biofilm biomass (C-F) and quantification of absolute biofilm biomass (G) after respective antibiotic treatments. Vancomycin, Van; Doxycycline, Dox; Rifampin, Rif; Gentamicin, Gen. Student's *t*-test, * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$.




Figure 2. Identification of drug combinations that kill MRSA biofilms. Various drug combinations consisting of drugs (at 50 μ M) highly active against growing phases and persister cells were tested and evaluated for their anti-biofilm activity by SYBR Green I/PI viability staining (A) and CFU enumeration (B). Drug combinations with sterilizing activity against USA300 biofilms were tested at

clinically achievable concentrations (Cmax). (E,F) Validation of meropenem + daptomycin + clinafloxacin in killing biofilms of various clinical isolates of MRSA (C,D). Vancomycin, Van; Meropenem, Mer; Daptomycin, Dap; Tosufloxacin, Tosu; Clinafloxacin, Clina; Doxycycline, Dox; Rifampin, Rif. Student's *t*-test, * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$.




Figure 3. Evaluation of oritavancin in killing biofilms as a single drug or in combination. (A) Comparing novel lipoglycopeptides oritavancin and dalbavancin to vancomycin in their activity to kill persisters (at 50 μ M). Evaluating oritavancin in (B) killing

persists in combination with meropenem+daptomycin and the agent (C) killing growing phase bacteria in combination with clinafloxacin (at Cmax concentrations). (D) Time-kill curve of biofilms comparing the top drug combinations candidates (at Cmax concentrations). Meropenem, Mer; Daptomycin, Dap; Oritavancin, Orita; Clinafloxacin, Clina; Doxycycline, Dox; Rifampin, Rif. Two-way ANOVA, and Student's *t*-test, * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$.




Figure 4. Validation of drug combinations in chronic skin infection model. (A) Study design of mouse treatment studies. (B) Bacterial load in the skin lesions. (C) Changes in lesion sizes and histology of skin tissues of mice infected with USA300 and treated

7-days with drug combinations and respective controls were measured. Histopathology of uninfected mice (D), infected mice receiving 7-days of no treatment (E), or treated with doxycycline + rifampin (F), or treated with meropenem + daptomycin + clinafloxacin (G) was analyzed. (H) Bacterial loads in the skin tissues of mice infected with MSSA Newman strain and treated with 7-days with drug combinations or control treatments were enumerated. Images were taken at 200X magnification. Meropenem, Mer; Daptomycin, Dap; Tosufloxacin, Tosu; Clinafloxacin, Clina; Doxycycline, Dox; Rifampin, Rif, Gentamicin, Gen. Blue arrows indicate crust formation, red brackets indicate hyperplasia and cellular infiltration; Student's *t*-test, * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$.




Figure 5. Meropenem + Daptomycin + Clinafloxacin reduced immune response of infected skin tissues. (A) IL-17, (B) IL-1, and (C) Gross pathology of skin tissues of mice treated 7-days with drug combinations and respective controls (in mm). Meropenem, Mer; Daptomycin, Dap; Tosufloxacin, Tosu; Clinafloxacin, Clina; Doxycycline, Dox; Rifampin, Rif, Gentamicin, Gen




Figure 6. Chemical structures of the fluoroquinolones tested. The chloride group (circled) is unique to clinafloxacin.

Table 1. Drug Dosage, Scheduling, and Administration

Drug	Dosage	Route	Times Treated	Cmax tested (clinically achievable concentrations)
Vancomycin	110 mg/kg	Intraperitoneally	Twice/daily	20 µg/ml
Daptomycin	50 mg/kg	Intraperitoneally	Once/daily	80 µg/ml
Meropenem	50 mg/kg	Intraperitoneally	Once/daily	20 µg/ml
Clinafloxacin	50 mg/kg	Intraperitoneally	Once/daily	2 µg/ml
Doxycycline	100 mg/kg	Oral	Twice/daily	5 µg/ml
Rifampin	10 mg/kg	Oral	Twice/daily	5 µg/ml
Moxifloxacin	100 mg/kg	Oral	Once/daily	4 µg/ml
ADEP4	25 mg/kg and 35 mg/kg	Intraperitoneally	Twice/daily	
Rifampin (for ADEP4 combination)	30 mg/kg	Intraperitoneally	Once/daily	
Gentamicin	20 mg/kg	Intraperitoneally	Once/daily	
Fructose	1.5 g/kg	Intraperitoneally	Once/daily	
Oritavancin	---	---	---	5 µg/ml
Ciprofloxacin	---	---	---	10 µg/ml
Levofloxacin	---	---	---	10 µg/ml

Table 2. Ranking of Fluoroquinolones for Their Activity in Killing Biofilm Bacteria

Treatment (Cmax)	Log (CFU/ml)	Live/Dead Ratio	Ranking of combination (1= best, 4= worst)
None	9.23 ± 0.4	11.95 ± 0.1	n/a
Meropenem	7.34 ± 0.7	8.11 ± 0.04	n/a
Daptomycin	9.34 ± 0.3	7.92 ± 0.2	n/a
Meropenem+Daptomycin	6.77 ± 0.6	4.52 ± 0.03	n/a
Mer+Dap+Clina	0	1.03 ± 0.07 T	1
Mer+Dap+Cipro	3.87 ± 0.3	1.34 ± 0.08 T	2
Mer+Dap+Moxi	4.99 ± 0.6	1.24 ± 0.06 T	3
Mer+Dap+Levo	8.91 ± 0.5	7.19 ± 0.1 T	4

T = Limit of detection has been reached for the SYBR Green I/PI viability assay

Meropenem, Mer; Daptomycin, Dap; Clinafloxacin, Clina; Ciprofloxacin, Cipro; Moxifloxacin, Moxi; Levofloxacin, Levo

n/a= none available