

1 Transcriptional changes suggest a major 2 involvement of Gibberellins in *Trifolium* 3 *pratense* regrowth after mowing

4 Short title: Gibberellin influences regrowth in red clover

5 Authors: Denise Brigitte Herbert (1)*, Thomas Gross (1)*, Oliver Rupp (2), Annette Becker (1)

6 * These authors contributed equally to this work

7 Affiliations:

8 (1) Justus Liebig University, Institute of Botany, Heinrich-Buff-Ring 38, D-35392 Giessen, Germany

9 (2) Justus Liebig University, Department of Bioinformatics and Systems Biology, Heinrich-Buff-Ring
10 26-32, D-35392, Giessen, Germany

11 Corresponding author: Denise Herbert (Denise.Herbert@bio.uni-giessen.de)

12 **Abstract**

13 Red clover (*Trifolium pratense*) is used worldwide as a fodder plant due its high nutritional value. In
14 response to mowing, red clover exhibits specific morphological traits to compensate the loss of
15 biomass. The morphological reaction is well described, but knowledge of the underlying molecular
16 mechanisms are still lacking. Here we characterize the molecular genetic response to mowing of red
17 clover by using comparative transcriptomics in greenhouse conditions and agriculturally used field.
18 The analysis of mown and control plants revealed candidate genes possibly regulating crucial steps
19 of the genetic network governing the regrowth reaction. In addition, multiple identified gibberellic

20 acid (GA) related genes suggest a major role for GA in establishing the regrowth morphology of red
21 clover. Mown red clover plants showing this regrowth morphology were partially “rescued” by
22 exogenous GA application, demonstrating the influence of GA during regrowth. Our findings provide
23 insights into the physiological and genetic processes of mowing red clover, to serve as a base for red
24 clover yield improvement.

25

26 **Introduction**

27 *Trifolium pratense* (red clover) is an important worldwide forage crop and thus of great economic
28 interest. This perennial plant offers several advantages like a high protein content and soil improving
29 characteristics, which can reduce the use of artificial nitrogen application and can enhance intake in
30 livestock. Well-known disadvantages of red clover include poor persistence under several land use
31 scenarios, like grazing or cutting [1–3]. *T. pratense* is a member of the Fabaceae (or legumes), which
32 are, due to their economic value, among the most examined families in the plant kingdom with
33 genome sequences available for species like *Medicago truncatula* (barrel clover) [4], *Lotus japonicus*
34 (birdsfoot trefoil) [5], *Glycine max* (soy) [6], *Phaseolus vulgaris* (common bean) [7], *Cicer arietinum*
35 (chickpea) [8], *Vigna unguiculata* (cowpea) [9], *Trifolium subterraneum* (subterranean clover) [10]
36 *Trifolium medium* (zigzag clover) [11], and *T. pratense* (red clover) [12,13].

37 Facing today’s challenges such as an increased demand on food production in an era of global
38 climate change together with the aim to solve these problems in an environmental friendly and
39 sustainable way requires improvement of forage crops like *T. pratense* [14,15]. *T. pratense* breeding
40 aims to offer genotypes with improved key agronomic traits (dry matter yield, high quality,
41 resistance to diseases and abiotic/biotic stress, persistency, [16]), while improving its regrowth
42 ability [2,17]. Unfortunately, the morphological investigations of several *T. pratense* populations
43 showed a correlation of persistency with non-favorable traits, like small plant size and prostrate

44 growth habit [18]. Moreover, most *T. pratense* cultivars or accessions are locally adapted and
45 require their specific local conditions to show the favored traits [19,20], which decreases the
46 stability for individual traits in breeding efforts [21]. *T. pratense* exhibits significant intraspecific
47 variation due to high intrapopulation genetic diversity, thus, persistence and performance in
48 response to mowing or cutting, depends on the variety, as well as developmental stage at the
49 moment of damage [22–25].

50 Persistency can be defined as a sustained forage yield over several growing periods [26] and is a
51 complex trait influenced by a variety of abiotic and biotic factors, and the regrowth ability of a plant
52 [27]. Plants with high regrowth ability can survive more frequent and intense biomass loss and could
53 be therefore more persistent. Decapitation or biomass loss due to herbivory or mowing triggers a
54 complex reaction affected by environmental conditions, plant morphology, architecture,
55 developmental stage and genotype [22]. After decapitation, the first stress response in other
56 legumes like *Medicago sativa* and *Pisum sativum* involves the production of phytohormones:
57 cytokinin, auxin, and strigolactones [28–30]. In addition, the mobilization of energy reserves is
58 activated [31]. Phenotypic plasticity of plant architecture in combination with alterations of
59 hormone concentrations can be observed in *P. sativum* (pea) and *T. pratense* after decapitation
60 [25,30,32]. However, the molecular processes allowing plants to thrive even after an enormous loss
61 of biomass remain still unclear, even in *Arabidopsis thaliana* [33,34].

62 Here, we compare the transcriptomes of mown (cut) vs. unmown (uncut) *T. pratense* plants from
63 two different field locations on the Biodiversity Exploratory “Heinich-Dün” [35] and greenhouse
64 grown plants. Our field samples were subjected to standard agricultural treatment and we can thus
65 discriminate transcriptional changes caused by abiotic factors and biotic interactions in the field
66 from those regulating regrowth. We present the identification and *in silico* characterization of
67 putative developmental regulators differentially expressed in the regrowth phase after mowing in
68 the field and in the greenhouse that may contribute to the regrowth response of *T. pratense* and

69 demonstrate that gibberellic acid (GA) is a major regulator of specific aspects of the regrowth
70 morphology in red clover.

71

72 **Material and Methods**

73 **Plant growth conditions, GA treatment, tissue sampling, and RNA extraction, cDNA** 74 **library construction and RNA-Seq**

75

76 Plant material for RNA-Seq was collected from three locations (fields and greenhouse, Fig. 1 A and
77 table S1, thereby one field location includes two neighboring field sites). Field plant tissue for RNA-
78 Seq was sampled on 11.06.2014 within the area of the Biodiversity Exploratory “Hainich-Dün” [35],
79 located in Thuringia, Germany. Material was sampled on four neighboring sites; two mown pastures
80 and two meadows that were not mown (FaM, FaNM, FbM, FbNM). For the greenhouse samples,
81 seeds of regional *T. pratense* populations (from a region covering mainly Thuringia, Saxony, Saxony-
82 Anhalt, Thuringian Forest and Uckermark, Germany) were obtained from the Rieger Hofmann seed
83 company (Blaufelden, Germany). Plants were grown in 23 °C with 16 h of light in pots of 12 cm
84 diameter. Plants in the greenhouse were watered daily and compound fertilizer (8’8’6’+) was given
85 every ten days. After 122 days after sowing, half of the plants were cut to 5 cm (GM and GNM).
86 Material from mown plants was sampled approximately 14 days after mowing/cutting, to avoid
87 sequencing of the transcripts related to the first stress response [36]. After collection, the samples
88 were snap frozen in liquid nitrogen. For each site and the greenhouse two biological replicates of
89 four pooled plants (shoot and leaf material) each were collected.

90

91 Fig. 1: Overview of sampling locations and classification of DEGs. A: Overview of the sampling
92 locations for the plant material. Names of the fields belonging to the Biodiversity Exploratory or
93 greenhouse populations are shown, as well as the conditions (mown/cut and not mown/uncut, HG
94 15 and HG 42; HG13 and HG 08). Distances between sampling locations in the field have been
95 estimated. B-D: Classification of DEGs with a $|\log_2\text{FoldChange}| < 2$. Percentage share of each class to
96 the corresponding gene list is shown in bar charts B: Classes of DEGs from field a mown vs. unmown.
97 C: Classes of DE field b mown vs. not mown. D: Classes of DE contigs of mown plants grown in the
98 greenhouse vs. unmown plants. E-F: Shared genes between the different treatments and locations.
99 The Venn diagrams show the number of shared upregulated gene within the “mown” samples (E)
100 and the number of shared genes within the “not mown” samples (F), blue circles indicate
101 greenhouse data, green field a and red field b. G: Number of genes belonging to the class
102 “phytohormones” within the DEG list of field (a and b) and greenhouse transcriptomes. The pie chart
103 shows the number of the different plant phytohormones (abscisic acid, ABA; auxin, AUX; genes
104 common between the auxin and cytokinin pathway, AUX/CK; cytokinin, CK; ethylene, ET;
105 gibberellins, GA; jasmonic acid; JA; salicylic acid, SA).

106

107 RNA was extracted using NucleoSpin® RNA Plant Kit (Macherey-Nagel GmbH & Co. KG, Düren,
108 Germany) according to the manufacturer’s instructions. Preparation of the cDNA libraries and the
109 strand-specific sequencing was conducted by Eurofins Genomics (Ebersberg, Germany). The RNAs of
110 four individuals were pooled for each RNA-Seq library and sequenced on an Illumina HiSeq2000
111 platform with chemistry v3.0, creating 2x 100 bp paired end reads.

112 In order to assess the effect of GA during the regrowth reaction of *T. pratense*, 14 red clover plants
113 were mown as described in [25]. Of these plants, seven were used as control plants and seven plants
114 were sprayed with 100 μM GA₃ (Duchefa Biochemie B.V, Haarlem, The Netherlands) once per week
115 as described in [37]. Different morphological characters (leaf number, length/width of leaflets,

116 petiole length, number of inflorescences, and number of main shoots) were measured for four
117 weeks.

118

119 **Assembly of reference transcriptome and annotation**

120

121 The raw-read-quality of the RNA-Seq data was analyzed with FastQC (available online at:
122 <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>). Illumina adapter and low quality
123 regions were trimmed using Trimmomatic [38] with ILLUMINACLIP, SLIDINGWINDOW:5:20 and
124 MINLEN:50 options. Quality trimmed reads were pooled and digitally normalized [39]. Multiple *de*
125 *novo* assemblies were computed using Trinity [40] and Oases [41] with all odd k-mer parameters
126 between 19 and 85. In addition, a genome guided assembly was performed using Trinity using the
127 draft genome of *T. pratense* 1.0 (GCA_000583005.2) [12,42]. The resulting contigs were screened for
128 potential coding sequences (CDS) using TransDecoder (<https://transdecoder.github.io/>). The
129 EvidentialGene pipeline ([http://arthropods.eugenis.org/EvidentialGene/about/EvidentialGene](http://arthropods.eugenis.org/EvidentialGene/about/EvidentialGene_trassembly_pipe.html)
130 [_trassembly_pipe.html](http://arthropods.eugenis.org/EvidentialGene/about/EvidentialGene_trassembly_pipe.html)) was used to merge and filter the contigs based on the TransDecoder CDS
131 prediction. Completeness of the final contig was confirmed by computing the mapping-rate of the
132 non-normalized reads to the contigs. The raw sequence reads can be found at NCBI: PRJNA561285.

133 The contigs were uploaded to the “Sequence Analysis and Management System” (SAMS) [43] for
134 functional annotation with the SwissProt [44], TrEMBL [45] and Phytozome [46] (e-value cutoff of
135 1e-5) databases. Additionally, attributes like gene name or functional description were extracted
136 from the blast hits. Contigs were mapped to the *T. pratense* reference genome using gmap [47]. All
137 non-Viridiplantae contigs were discarded. Transcription factors were identified using a blastp search
138 of the protein sequences against the plant transcription factor database Potsdam (PlnTFDB) (48 [48],

139 version 3.0, <http://plntfdb.bio.uni-potsdam.de/v3.0/>) protein database with an e-value cutoff of 1e-
140 20. The files contain the functional annotation description of all transcripts e-Appendix (Table S11).

141

142 **Differential gene expression analysis, enrichment analysis, and classification of** 143 **differentially expressed genes**

144

145 Read counts for each contig of the final assembly in each sample were computed using RSEM [49]
146 with bowtie mapping. To identify differentially expressed *T. pratense* genes (DEG) a pairwise
147 comparison of all treatments was performed using the DESeq2 [50] tool with $FDR \leq 0.01$ and
148 $|\log_2 \text{FoldChange}| \geq 2$ between FaM and FaNM, FbM and FbNM; GM and GNM respectively. The top
149 20 DEG were determined for each comparison based on the expression strength (\log_2 fold change).
150 Homologues in the next closest species and *A. thaliana* for each *T. pratense* candidate gene were
151 searched based on the *T. pratense* genome sequence deposited in Phytozome [46]. TPM (transcript
152 per million) values were calculated to estimate contig expression level (Wagner et al 2012).

153 We used the description and gene names obtained from TrEMBLE and SwissProt to search the
154 UniProt [51], NCBI [52] and TAIR [53] databases to obtain further information (Table S8). Raw reads
155 that were assembled to contigs, exhibiting a gene structure (ORF) and attained a putative annotation
156 referred to below as genes.

157

158 **Blast2Go Analysis of *T. pratense* genomes**

159

160 Two local BLAST searches [54] with word-size of 3, e-value of 1.0E-3 and HSP length cutoff of 33
161 were performed against the PlnTFDB using Blast2GO [55]. Only the blast hits with the highest

162 similarity were used for further comparisons (number of BLAST hits = 1), sequences with similarity
163 below 50% and an e-value higher than 1.0e-4 were omitted. The Blast2GO output was compared
164 with an in-house python3-script utilizing NumPy (<https://numpy.org/>), Pandas
165 (<https://pandas.pydata.org/>) and Seaborn (<https://seaborn.pydata.org/>) applying the list of
166 transcription factors (TF) downloaded from PlnTFDB (<http://plntfdb.bio.uni-potsdam.de/v3.0/>) to
167 the blast output and furthermore visualizing the generated datasets. We searched Uniprot database
168 hits for development and phytohormone related genes. Subsequently, gene IDs of gibberellin genes
169 we searched for matches within our annotated *T. pratense* transcriptomes. Matches were filtered
170 based on TPM values and classified based on biosynthesis, catabolism activation/repression or
171 signaling/response, corresponding expression patterns within the transcriptome have been
172 identified additionally.

173

174

175 **Results**

176 **RNA-Seq results, *de novo* assembly, and functional description of contigs**

177

178 The RNA-Seq produced a total number of short reads between 44.7 and 58.1 million for each library
179 with two exceptions (table S2) totaling 608,041,012 raw reads. The *de novo* assembly of the
180 reference transcriptome of *T. pratense* produced 44,643 contigs, of which 41,505 contigs were
181 annotated and 29,781 contigs were identified as plant specific. The minimum length of the contigs
182 was 124 bp, the maximum length 1171.31 bp (Table S3). After the *de novo* assembly of the *T.*
183 *pratense* transcriptome, each individual library was mapped back against the reference

184 transcriptome individually, to determine the overall alignment rate, which was between 77.85 % and
185 90.32 % (Table S4).

186 63 % of the 44,643 contigs could be mapped to a known locus of the *T. pratense* genome annotation
187 [12,42]. 32 % could be mapped to an unknown locus of the *T. pratense* genome and 5 % could not be
188 mapped to the *T. pratense* genome (Fig. S1). All plant-specific contigs were annotated with several
189 databases (Table S5). To further verify the quality of our replicates, we identified the transcripts
190 shared by the two replicates. We identified TPM values for each transcript and discarded transcripts
191 with TPM values <1. Then we compared the transcripts of each library with each other and
192 calculated the percentage of this number compared with the total number of transcripts within each
193 library. The percentage of transcripts shared between the two replicates is between 90 % and 94 %
194 for all treatments/localities, suggesting that the RNA-Seq data are reproducible (Table S6).

195 **Specific transcriptional regulator families are differentially expressed during the** 196 **regrowth process**

197 We were firstly interested to identify transcriptional regulators initiating and maintaining the
198 regrowth morphology and mapped the transcriptome to the PlnTFDB to identify these
199 transcriptional regulators. All members of a specific transcriptional regulator family (TRF) were *in*
200 *silico* identified in the transcriptome and their expression was compared between mown and
201 unmown plants (Fig. S2). Only those TRFs are shown for which at least 10% of the members showed
202 significantly differential expression between mown and unmown conditions (Fig. 2).

203

204 Fig. 2: Differentially expressed TRF members in mown and not mown *T. pratense* plants. The y-axis
205 shows the number of expressed contigs (TPM value over 5 TPM) that are members of the specific
206 TRF. Names of the transcriptomes and TRFs are given on the x-axis. Expression of transcription factor
207 members was compared in a pairwise manner (GM vs GNM, FaM vs FaNM, FbM vs FbNM). Shown

208 are only those plant TRFs in which at least one of the comparisons resulted in a difference of more
209 than 10% of the contigs significantly upregulated in either the mown or the unmown condition
210 (orange bars).

211

212 17 TRFs were identified of which at least 10% of the members showed differential expression in the
213 mown versus unmown comparisons (Fig. 2): ABI3VP1, AP2-EREBP, C₂C₂-Dof, C₂C₂-GATA, GRAS, HSF,
214 LOB, MADS, mTERF, MYB, NAC, PHD, SBP, SNF2, TCP, TRAF, WRKY.

215 Two TRFs show expression activation upon mowing: a significant number of WRKY transcripts are
216 up-regulated in mown plants regardless of the provenance. MADS-box transcripts were found
217 upregulated as well, but only in the field-derived transcriptomes. Generally, only five of the 17 TRFs
218 analyzed here showed significant changes in expression towards mowing in the greenhouse-derived
219 plants suggesting that they react less strongly towards mowing than the field-derived plants. Six
220 TRFs (AP2-EREBP, MYB, NAC, PHD, SBP, and TCP) show transcriptional changes in reaction to mowing
221 only in field location a and three TRFs (mTERF, SNF2, TRAF) show this only in field location b
222 suggesting that combination of biotic and abiotic factors with mowing differ between the two field
223 locations.

224 Notably, only the C₂C₂-GATA TRF reacts towards mowing under greenhouse but not under field-
225 conditions suggesting that transcriptional changes in reaction to other biotic and abiotic factors may
226 overlay the regrowth reaction. Taken together, the TRF analysis shows that the reaction towards
227 mowing induces transcriptional changes in only a subset of TRFs, suggesting that those play a major
228 role in relieving the stress biomass loss and regrowth.

229

230 **Differentially expressed genes analysis reveals diverse subsets of genes involved**
231 **in regrowth influenced by location and environmental conditions**

232

233 To identify gene expression responses underlying the regrowth response after mowing digital gene
234 expression analysis was performed comparing FaM vs FaNM; FbM vs FbNM; GM vs GNM to identify
235 DETs (Table S12) from mown plants. Interestingly, using the log fold change 2, the number of DEG is
236 rather similar in all comparisons, ranging from 119 (Gm vs. GNM) to 142 (FaM vs. FaNM) (Table 1).

237

238 **Table 1 Table shows the numbers of differentially expressed transcripts (contigs) between libraries with changes above**
239 **logfold 2. Up- or down regulation for each comparison is shown.**

Analysis	total DEG	Number of transcripts up regulated (library)	Number of transcripts down regulated (library)
GM vs. GNM	119	54 (greenhouse mown)	65 (greenhouse not mown)
FaM vs. FaNM	142	49 (mown)	93 (not mown)
FbM vs. FbNM	122	59 (mown)	63 (not mown)

240

241 We were then interested to identify developmental processes in greater detail that are required for
242 the regrowth process. Thus, the results of the DEG analysis were restructured such that the DEG
243 were grouped in 16 descriptive classes by database and literature mining (Table S7 and Table S8).
244 Those classes describe major functional groups and serve to identify the potential role of a gene.

245 The results of the top 20 DEG showed that the greenhouse plants displayed more DEG involved in
246 regrowth processes and less genes related to environmental conditions when compared with field
247 plants. Most likely, the greenhouse grown plants displayed the regrowth reaction more prominently,

248 as they grew under less stressful conditions than the field grown plants, for which more stress
249 related DEG were observed (Fig. 1 B-D /Table 2-4).

250 **Table 2: Top twenty differentially expressed genes of GM vs. GNM analysis. The table shows the transcript name, log2 fold change of the corresponding transcript, the library in which the**
 251 **transcript is upregulated (pattern), gene name based on *T. pratense* genome annotation, corresponding Phytozome description, gene name and species name of the next homologues and**
 252 ***A. thaliana* gene name and locus name based on information available on Tair.**

ID	Patt ern	Contig ID	log2 Fold Chan ge	Class (basis of classification)	Gene name <i>T. Pratense</i>	Next homologue gene name	Next homologue species name	<i>A. thaliana</i> gene name	<i>A. thaliana</i> locus name
1	GHN M	tdn_99733	-9.5	Growth (<i>M. truncatula</i>)	Tp57577_TGAC_v2_mRNA4544.v2	Medtr4g029550.1	<i>M. truncatula</i>	-	-
2	GHN M	k41_54584	-6.3	Biotic stress (<i>T.pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA28349.v2	Medtr5g073620.1	<i>M. truncatula</i>	ATEXO70B1	AT5G58430
3	GHN M	tdn_92791	-5.5	Abiotic/biotic stress (<i>T.pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA20498.v2	Medtr1g041150.1	<i>M. truncatula</i>	ATCPK1	AT5G04870
4	GHN M	k41_130218	-5.5	-	-	-	-	-	-
5	GHN M	tdn_53091	-4.8	Phytohormone (<i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA39912.v2	Medtr4g010250.1	<i>M. truncatula</i>	-	AT5G20190
6	GHN M	tgg_43136	-4.4	Transcription (<i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA29629.v2	Medtr4g098630.1	<i>M. truncatula</i>	ANAC071	AT4G17980
7	GHN	tdn_141837	-4.3	Abiotic stress (<i>T.pratense</i> , <i>M. truncatula</i> , <i>A.</i>	Tp57577_TGAC_v2_mRNA760.v2	Medtr2g022700.1	<i>M. truncatula</i>	ATGPT2	AT1G61800

	M			<i>thaliana</i>)					
8	GHN	tdn_40997	-4.2	Abiotic stress (<i>T.pratense</i> , <i>M. truncatula</i> , <i>A.</i>	Tp57577_TGAC_v2_mRNA25718.v2	Medtr4g130540.1	<i>M. truncatula</i>	<i>HSP70B</i>	AT1G16030
	M			<i>thaliana</i>)					
9	GHN	k71_5292	-4.1	Biotic stress (<i>T.pratense</i> , <i>M. truncatula</i>)	Tp57577_TGAC_v2_mRNA23166.v2	Medtr0163s0020.	<i>M. truncatula</i>	<i>LECRK-IX.1</i>	AT5G10530
	M					1			
10	GHN	k59_6358	-3.9	Growth (<i>T.pratense</i> , <i>M. truncatula</i> , <i>A.</i>	Tp57577_TGAC_v2_mRNA12337.v2	Medtr3g435430.1	<i>M. truncatula</i>	<i>ATEXP15</i>	AT2G03090
	M			<i>thaliana</i>)					
11	GHM	tdn_86219	8.0	Biotic stress (<i>T.pratense</i> , <i>M. truncatula</i> , <i>A.</i>	Tp57577_TGAC_v2_mRNA29036.v2	Medtr4g066210.1	<i>M. truncatula</i>	<i>BGLU12</i>	AT5G42260
				<i>thaliana</i>)					
12	GHM	k23_115785	8.0	Abiotic stress (<i>T.pratense</i> , <i>M. truncatula</i> , <i>A.</i>	Tp57577_TGAC_v2_mRNA22071.v2	Glyma.01G00100	<i>G. max</i>	-	AT5G58110
				<i>thaliana</i>)		0.1			
13	GHM	tdn_91159	8.1	Biotic stress (<i>T.pratense</i> , <i>M. truncatula</i> , <i>A.</i>	Tp57577_TGAC_v2_mRNA7745.v2	Medtr4g035870.1	<i>M. truncatula</i>	-	AT5G62360
				<i>thaliana</i>)					
14	GHM	k65_43517	8.3	Phytohormone (<i>T.pratense</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA6281.v2	Medtr1g082750.1	<i>M. truncatula</i>	<i>ATAMI1</i>	AT1G08980
15	GHM	tgg_18067	8.4	-	Tp57577_TGAC_v2_mRNA32019.v2	-	-	-	-
16	GHM	k61_38813	9.0	-	-	-	-	-	-
17	GHM	k49_82496	9.0	Abiotic/biotic stress (<i>G. max</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA37976.v2	Glyma.06G26880	<i>G. max</i>	-	AT4G04790
						0.1			

18	GHM	k67_38815	9.1	Biotic stress (<i>T.pratense</i>)	Tp57577_TGAC_v2_mRNA41666.v2	Medtr0062s0020.	<i>M. truncatula</i>	-	-
									1
19	GHM	k45_11164	9.6	Transcription (<i>T.pratense</i>)	Tp57577_TGAC_v2_mRNA29953.v2	Medtr3g092510.1	<i>M. truncatula</i>	<i>ATRBP37</i>	AT4G10610
20	GHM	tdn_25484	9.6	Growth (<i>Phaseolus vulgaris</i>)	Tp57577_TGAC_v2_mRNA13093.v2	Phvul.006G03380	<i>Phaseolus vulgaris</i>	-	-
									0.1

253

254

255 **Table 3: Top twenty differentially expressed genes of FaM vs. FaNM analysis. The table shows the transcript name, log2 fold change of the corresponding transcript, the library in which**
 256 **the transcript is upregulated (pattern), gene name based on *T. pratense* genome annotation, corresponding Phytozome description, gene name and species name of the next homologues**
 257 **and *A. thaliana* gene name and locus name based on information available on Tair.**

ID	Pattern	Contig ID	log2F	Class (basis of classification)	Gene name <i>T. pratense</i>	Next homologue gen name	Next homologue species name	<i>A. thaliana</i> gene name	<i>A. thaliana</i> locus name
1	TPNM2	k33_17052	-9,0	Biotic stress (<i>T. pratense, M. truncatula, A. thaliana</i>)	Tp57577_TGAC_v2_mRNA21474.v2	Medtr4g079440.1	<i>M. truncatula</i>	na	AT1G06260
2	TPNM2	k43_11179	-8,8	Biotic stress (<i>M. truncatula</i>)	Tp57577_TGAC_v2_mRNA26333.v2	Medtr8g101900.1	<i>M. truncatula</i>	CCOAOMT7	AT4G26220
									2
3	TPNM2	tdn_34568	-8,6	-	Tp57577_TGAC_v2_mRNA9104.v2	Glyma.13G06180	<i>G. max</i>	-	AT5G39530
									0.1
4	TPNM2	tdn_49640	-8,6	-	-	-	-	-	-

5	TPNM2	tdn_58745	-8,5	Biotic stress (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA20190.v2	Medtr8g075200.1	<i>M. truncatula</i>	-	AT1G75900
6	TPNM2	tdn_47209	-8,5	Growth (<i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA10703.v2	Medtr1g053315.1	<i>M. truncatula</i>	-	AT1G03390
7	TPNM2	tdn_48478	-8,4	Biotic stress (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA19516.v2	Medtr2g099020.1	<i>M. truncatula</i>	-	AT3G59510
8	TPNM2	k41_17597	-8,4	Growth stress (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA8526.v2	Medtr1g036490.1	<i>M. truncatula</i>	ATCOMT, ATOMT1	AT5G54160
9	TPNM2	k51_82581	-8,2	Growth (<i>T. pratense</i>)	Tp57577_TGAC_v2_mRNA23127.v2	Medtr2g436480.1	<i>M. truncatula</i>	KCS21	AT5G49070
10	TPNM2	tdn_82424	-8,1	Growth (<i>T. pratense</i>)	Tp57577_TGAC_v2_mRNA17103.v2	Medtr2g013740.1	<i>M. truncatula</i>	KCS10	AT2G26250
11	TPM2	k49_380	7,5	Development (<i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA37185.v2	SapurV1A.0885s0 040.1	<i>Salix purpurea</i>	DAYSLEEPER	AT3G42170
12	TPM2	tdn_49869	7,6	-	-	-	-	-	-
13	TPM2	tdn_54983	7,7	-	-	-	-	-	-
14	TPM2	k37_9029	7,8	-	-	-	-	-	-
15	TPM2	k45_6120	8,4	-	Tp57577_TGAC_v2_mRNA2166.v2	Medtr2g007510.1	<i>M. truncatula</i>	-	-
16	TPM2	k71_23808	8,4	Development (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA14131.v2	Medtr1g021320.1	<i>M. truncatula</i>	-	AT4G33280
17	TPM2	k59_3541	8,4	Development (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA34193.v2	Medtr4g089030.1	<i>M. truncatula</i>	CYP71A26	AT3G48270
18	TPM2	k59_360	8,6	Metabolism (<i>Linum usitatissimum</i> , <i>T. pratense</i>)	Tp57577_TGAC_v2_mRNA21875.v2	Lus10012445	<i>Linum usitatissimum</i>	-	AT1G50020
19	TPM2	k53_38903	9,0	Abiotic stress (<i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA37328.v2	Medtr8g063190.1	<i>M. truncatula</i>	PRIN2	AT1G10522

20 TPM2 tdn_12997 9,6 - Tp57577_TGAC_v2_mRNA9318.v2 Medtr7g062280.1 *M. truncatula* - AT5G01140
8

258

259 **Table 4: Top twenty differentially expressed genes of FbM vs. FbNM analysis. The table shows the transcript name, log2 fold change of the corresponding transcript, the library in which**
260 **the transcript is upregulated (pattern), gene name based on *T. pratense* genome annotation, corresponding Phytozome description, gene name and species name of the next homologues**
261 **and *A. thaliana* gene name and locus name based on information available on Tair.**

ID	Patter n	Contig ID	log2 Fold	Class (basis of classification)	Gene name <i>T. pratense</i>	Next homologue gen name	next homologue species name	<i>A. thaliana</i> gene name	<i>A. thaliana</i> locus name
1	TPNM 3	tdn_100726	-9,4	Biotic stress (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA24659.v2	Medtr4g094772.1	<i>M. truncatula</i>	<i>CYP81D</i>	AT4G37340
2	TPNM 3	tgg_49631	-8,0	Biotic stress (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA37846.v2	Medtr6g034470.1	<i>M. truncatula</i>	-	AT2G34930
3	TPNM 3	tdn_152262	-7,9	-	-	-	-	-	-
4	TPNM 3	tdn_56712	-7,9	Biotic stress (<i>T. pratense</i> , <i>M. truncatula</i>)	Tp57577_TGAC_v2_mRNA30556.v2	Medtr8g027540.1	<i>M. truncatula</i>	-	-
5	TPNM 3	tdn_87762	-7,9	Biotic stress (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA10533.v2	Medtr7g451400.1	<i>M. truncatula</i>	<i>ATMCP1B</i> , <i>ATMCPB1</i>	AT1G02170

6	TPNM	tdn_86129	-7,1	General cell functions (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA10207.v2	Glyma.11G15450 0.1	<i>G. max</i>	<i>RPB5E</i>	AT3G54490
7	TPNM	k55_46241	-6,9	Growth (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA31452.v2	Medtr4g128150.1	<i>M. truncatula</i>	<i>histone 4</i>	AT2G28740
8	TPNM	tdn_55533	-6,2	Abiotic stress (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA39263.v2	Medtr5g007790.1	<i>M. truncatula</i>	<i>ATCRM1</i> , <i>ATXPO1</i>	AT5G17020
9	TPNM	tgg_51443	-4,7	Growth (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA37076.v2	Medtr5g019580.2	<i>M. truncatula</i>	<i>UGT72E2</i>	AT5G66690
10	TPNM	tdn_136706	-4,7	-	-	-	-	-	-
11	TPM1	tdn_140636	8,8	General cell functions (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA28209.v2	Medtr8g005980.1	<i>M. truncatula</i>	<i>C-NAD-MDH2</i>	AT5G43330
12	TPM1	tdn_154158	8,9	General cell functions (<i>T. pratense</i> , <i>M. truncatula</i>)	Tp57577_TGAC_v2_mRNA39482.v2	Medtr3g114970.2	<i>M. truncatula</i>	-	AT5G55150
13	TPM1	tdn_65187	9,1	Transposon (<i>T. pratense</i> , <i>Prunus persica</i> A. <i>thaliana</i>)	Tp57577_TGAC_v2_mRNA30115.v2	Prupe.4G011200. 1	<i>Prunus persica</i>	-	AT4G29090
14	TPM1	tdn_100956	9,2	Metabolism (<i>T. pratense</i> , <i>Capsella rubella</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA9542.v2	Carubv10008027 m	<i>Capsella rubella</i>	<i>AHA2</i>	AT4G30190
15	TPM1	k63_21505	9,3	Biotic stress (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA19467.v2	Medtr3g022400.1	<i>M. truncatula</i>	-	AT3G14470

16	TPM1	tdn_142681	9,3	Secondary metabolite biosynthesis (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA15473.v2	Medtr8g074550.1	<i>M. truncatula</i>	-	AT2G18570
17	TPM1	k45_6120	9,6	-	Tp57577_TGAC_v2_mRNA2166.v2	Medtr2g007510.1	<i>M. truncatula</i>	-	-
18	TPM1	tdn_52922	10,1	-	Tp57577_TGAC_v2_mRNA41271.v2	mrna20290.1- v1.0-hybrid	<i>Fragaria vesca</i>	-	AT1G21280
19	TPM1	tdn_65185	10,9	-	-	-	-	-	-
20	TPM1	tdn_109277	11,7	Transcription (<i>T. pratense</i> , <i>M. truncatula</i> , <i>A. thaliana</i>)	Tp57577_TGAC_v2_mRNA29560.v2	Medtr5g028610.1	<i>M. truncatula</i>	-	AT3G14460

262

263

264

265

266 Several functional groups show a similar pattern in the mown vs. unmown plants of all three
267 locations: more genes related to biotic stress processes and metabolism were upregulated in the
268 unmown locations (Fig. 1 B-D). In mown plants, more genes related to signaling and transposons
269 were upregulated. Only a single functional group (growth) shows similar patterns in only the field
270 locations suggesting that plants in the two field locations cope with very different habitat conditions
271 and stress factors.

272 The photosynthesis- and phytohormone-related genes of field a show a similar pattern to the
273 greenhouse plants as do the development- and signaling related genes. Genes related to
274 development, general cell functions and transcription have similar patterns between field b and the
275 greenhouse grown plants, such that more transcription - and development-related genes are
276 upregulated in mown plants. DEG related to symbiosis were found upregulated in unmown plants
277 grown in the greenhouse, even though these plants were fertilized. And unexpectedly, senescence-
278 related genes are upregulated in mown plants of field A. However, because our analysis cannot
279 discriminate between activating and repressing factors of senescence, we cannot conclude from our
280 data if the mown plants have activated or repressed their senescence program.

281 The largest group of differentially expressed genes is the one related to biotic stress with up to 38%
282 differentially expressed genes in one location (field b, Fig. 1 C). This suggests that different biotic
283 stresses act upon the mown vs. unmown plants. A similar phenomenon can be observed for growth
284 related processes, where up to 24% genes were upregulated in the mown and unmown plants
285 indicating that different growth programs are active in mown vs. unmown plants.

286 Taken together we can state that mown plants in all locations change their regulatory programs
287 upon mowing to cope with different biotic factors suggesting that they massively change their
288 metabolism and signaling processes. Further, transposons are more active in mown plants. Apart

289 from these conclusions, the molecular answer to substantial biomass loss differs between all three
290 locations.

291 To find similarly regulated genes between the treatments and/or locations, a Venn diagram was
292 generated to compare the number of shared significantly DEG within the “mown” samples and the
293 “not mown” samples (Fig. 1 E-F, Table S9). Within the “mown” samples we detected no overlap
294 between the groups with the exception of four genes that are differentially expressed and
295 upregulated in “mown” condition and are shared between the two field transcriptomes (FbM and
296 FaM (Fig. 1 E). Within the “not mown” samples also four genes are shared between the field
297 transcriptomes (FbNM and FaNM)) and one is shared between the field b and the greenhouse (Fig. 1
298 F). No genes are shared between all three samples, neither in the “mown” treatment, nor in the “not
299 mown” treatment. The genes that were shared between the transcriptomes belong to the main
300 classes “growth”, “phytohormone”, “general cell functions”, “biotic stress”, “development” and
301 “transcription” (Table S9).

302 Two of the genes could not be annotated. The annotated genes include for example genes
303 tdn_60472 (shared between FaM/FbM, class: phytohormone), that was found to be the homolog of
304 the *A. thaliana* locus AT1G75750, describing a GA-responsive GASA1 protein homolog. Another *A.*
305 *thaliana* homolog was identified, Chitinase A (ATCHIA), shared between FaNM/FbNM (tdn_129843,
306 class: biotic stress). In addition one gene was found, with a *T. pratense* annotation but no further
307 description or homologs to *A. thaliana* (k45_6120, shared between FaM/FbM). This suggests that
308 the molecular mechanisms directing regrowth overlaid by other processes, such as stress response
309 which have a more dramatic impact on the number of DEG than growth processes have. The shared
310 genes between the field conditions and the almost complete absence of shared genes between field
311 and greenhouse indicates that the growth conditions in the field are more like each other, even
312 when the fields are far apart from each other than any field to a greenhouse.

313

314 **Gibberellins are major players in the regrowth reaction**

315

316 As phytohormones play a major role in the regulation of development and stress response, we
317 identified DEGs related to phytohormone synthesis, homeostasis, transport, and signaling within all
318 transcriptome comparisons (Table 1). DEG links for all major classes of phytohormones were
319 identified, except for strigolactone. DEGs association to four phytohormones was most abundant:
320 abscisic acid (ABA, 8 DEGs), gibberellins (GA, 8 DEGs), salicylic acid (SA, 6 DEGs), and auxin (AUX, 5
321 DEGs) (Fig. 1 G). While ABA and SA are mainly involved in response to biotic and abiotic stresses, and
322 AUX is known to play a major role in growth and development, we identified GA as a novel candidate
323 phytohormone for regrowth response.

324 To learn more about the role of GA in the regrowth response, we identified 32 GA-related genes out
325 of 151 within the transcriptomes of the greenhouse and the field grown plants, matching our
326 selection criteria (TPM <5, involved in GA biosynthesis, signaling, GA responsive genes or catabolism,
327 displaying certain expression patterns Fig. 3 A) and classified them according to their function in the
328 GA biosynthesis and signaling processes (Table S13). Ranges of expression strength were calculated
329 and color coded to compare expression patterns (Fig. 3 A).

330

331 Fig. 3: Analysis of GA related contigs and regrowth processes. A) Differentially expressed GA-related
332 contigs within the *T. pratense* transcriptomes. Ranges of expression were calculated (0-39.99% blue
333 (low expression), 40-59.99% grey (neutral expression), 60-100% yellow (high expression)) according
334 to their TPM values. On the left, the gene names of the *T. pratense*'s closest *A. thaliana* homologs
335 are given if available. B) and C) show morphological changes in leaves after GA treatment. B) leaflet
336 area in cm², C) length of petioles in cm. The graphs show average values for each sampling date and
337 95% confidence interval. GA treated plants, blue; control plants, orange.

338

339 Five genes predicted to be involved in GA biosynthesis and signaling show similar expression
340 differences between mown and unmown plants from at least two of the three locations:
341 *tdn_142825* (AT2G46590) and the homologs of *ZHD1*, *GID1B*, *MPT1*, and *MPT3*. Further, homologs
342 of three GA responsive genes (*tdn_75969* (*MYB 44*, AT5G67300), *tdn_157683* (Os04g0670200,
343 AT1G47128), and *XERICO* and homologs of two GA catabolism genes (*GA2OX1*, *GA2OX8*) react
344 towards mowing. Interestingly, we also find many differences in expression between the field sites
345 in the GA related genes suggesting fundamental differences in the living conditions between the two
346 field sites that also impact regrowth after biomass loss.

347 When considering only the greenhouse grown plants, homologs of GA biosynthesis genes were not
348 differentially regulated, but genes most likely involved in GA signaling such as *MPT1*, *MPT3*, and
349 *SOC1* are down regulated in mown plants. Five homologs of GA responsive genes are upregulated in
350 the mown plants, while three, among them *XERICO*, are down regulated. Further, of the three
351 *GA2OXIDASE8* homologs encoded by the *T. pratense* genome, two are differentially expressed, one
352 up- and the other down regulated upon mowing. Also, one *GA2OXIDASE1* homolog is down
353 regulated in mown plants.

354 In summary, this suggests a highly dynamic response of several GA-related genes to mowing *or T.*
355 *pratense*. Interestingly, we were unable to identify larger changes in the GA biosynthesis pathway of
356 the greenhouse plants, but in GA catabolism genes, suggesting that GA availability in response to
357 mowing is regulated by catabolism and signaling rather than by biosynthesis. Further, we identified
358 two contrastingly regulated sets of genes acting in the mowing response.

359 **GA treatment after mowing induces specific changes to the regrowth response**

360

361 Identification of several GA related genes that changed expression suggested an involvement of GA
362 in the regulation of development after mowing. We were interested to corroborate this hypothesis
363 experimentally and treated *T. pratense* plants with GA after mowing. Weekly GA application during
364 the regrowth process led to significant and specific changes in morphology (Fig. 3 B, C). Previous
365 work suggested that regrowing plants produce smaller and rounder leaflets with shorter petioles
366 than uncut plants [25]. Number of leaves, shoots and inflorescences, leaf area and the roundness of
367 leaflets were measured (Fig. 3 B, C, Suppl. Fig. 3). The first visible effects of GA treatment were
368 recognized after 1.5 weeks, showing a significant difference in leaflet area between GA treated and
369 control plants. Later it was observed that the petioles of treated plants were in average twice as long
370 as petioles of untreated plants (16.7 ± 1.9 cm and 8 ± 1.2 cm, respectively). GA leaflets were with 4.7
371 ± 0.9 cm² almost double the size than those of untreated plants (2.4 ± 0.6 cm²). However, GA treated
372 plants grew only 30% more total leaf area than control plants, because the control plants had more
373 leaves than GA treated plants (Fig. S3 A, B, F, and G). Other morphological traits such as number of
374 inflorescences, leaves, and shoots remained unaffected by the GA treatment. In summary, mown
375 plants normally produce leaves with shorter petioles, restrict their leaflet area and their leaves
376 become rounder. GA treatment partially alleviated these developmental changes such that the
377 mown, GA treated plants produced larger leaves with longer petioles while the leaf shape was
378 unaffected by GA treatment.

379

380 **Discussion**

381 **RNA-Seq and assembly**

382

383 The *de novo* assembly in combination with a reference-based approach for the annotation led to
384 44643 contigs of which 29781 could be annotated as plant-specific (Fig. S1). With the prior *de novo*
385 assembly it was possible to attain 4051 additional contigs that could be not found within the
386 genome of *T. pratense* 1.0 (GCA_000583005.2) [12,42]. The estimated genome size of *T. pratense* is
387 ~440 Mbp [28]. The *T. pratense* transcriptome data in study was ~55 Mbp in size, corresponding to
388 ~12.5% transcribed regions in the *T. pratense* genome, which is within the range of previously
389 published transcriptomes (~10% (42 Mbp) [56]). Interestingly, we found plant-specific, previously
390 unreported contigs suggesting that the *T. pratense* genome might need improvement in terms of
391 sequencing coverage and protein coding sequence annotation.

392 **Cell walls are remodeled after mowing**

393

394 Our data analysis shows that several plant TRFs are predominantly involved in the regrowth reaction
395 (Fig. 2 and S2). After massive biomass loss like mowing inflicts on *T. pratense*, plants firstly need to
396 seal wounded tissues. Several transcriptional regulators are known to play a role in the tissue
397 reunion processes were identified in *Solanum lycopersicum*, *Cucumis sativus*, and *A. thaliana*
398 (reviewed in [36]). Homologs of these genes were also identified to be differentially regulated in the
399 *T. pratense* transcriptome after mowing, such as several members of the Auxin Response Factor
400 (ARF) family or the No Apical Meristem (NAM) family member *ANAC071*. [57] suggested that high
401 levels of AUX induce the expression of *ANAC071* via ARF6 and ARF8 (in the upper part of incised
402 stems), at the same time reduced AUX level directly after the cutting activate the expression of
403 *RAP2.6L*. In addition auxin signaling via ARF6 and ARF8 influences JA synthesis, via the activation of
404 *DAD1*, thus together with LOX2 increases *RAP2.6L* expression during tissue reunion in *A. thaliana*
405 [57]. Further it was demonstrated that *ANAC071* can as a transcription factor initiate the expression
406 of members of xyloglucan endotransglucosylase/hydrolases family (XTH20 and XTH19) which
407 recombine hemicellulose chains to drives the cell proliferation during tissue reunion [58].

408 Interestingly, we were able to identify all members of the cell wall remodeling pathway mentioned
409 above, displaying distinct expression pattern with some of them upregulated in mown plants
410 including for example *XTH32* (k69_7012, upregulated in FbM, tdn_94651, upregulated in GM, FaM
411 and FbM), *XTH6* (tdn_91763, upregulated in GM), *XTH8* (k71_5058, upregulated in GM, FbM), *XTH9*
412 (tdn_113578, upregulated in GM), *XTHA* (tdn_87930, upregulated in GM), *LOX2* (tdn_156279,
413 upregulated FbM), and *ARF8* (tdn_156886 upregulated in GM, tdn_156890 upregulated in GM)
414 (Table S10, S12 and S13). This is suggesting that the early steps in the regrowth reaction are
415 conserved in core eudicots and that the cell wall remodeling processes continue at least two weeks
416 after mowing.

417 **Biotic and abiotic stresses contribute to differential gene expression**

418

419 RNA-Seq experiments create a large amount of raw data which requires significant downstream
420 analysis to provide a biologically meaningful dataset. We thus compared those 20 genes, that
421 differed most strongly in their expression between the different treatments and locations (see table
422 4-6 and Fig. 1 B-D). These comparisons revealed that the mown greenhouse plants show the highest
423 percentage of genes possibly involved in regrowth processes. Contrasting, the field transcriptomes
424 display patterns of abiotic and biotic stress reactions. Comparisons of the top 20 DEG of the
425 unmown field transcriptomes showed that plants grown on field a and b face biotic stress more than
426 abiotic stress. One of the upregulated genes in field a is a chitinase homolog suggesting that those
427 plants are under attack of fungi and/or insects. Follow-up analyses to correlate environmental
428 conditions, biotic and abiotic stresses monitored within the Biodiversity Exploratories with
429 differential gene expression at the two field locations would be an interesting project but are
430 beyond the scope of this work. In contrast, the top 20 DE transcripts of the greenhouse plants
431 include phytohormone- and transcription-related genes, but also a high proportion of biotic and
432 abiotic stress-related genes. This suggests that also these plants have to cope with stresses, but to a

433 lesser extent. Thus, their regrowth reaction is more visible within the top 20 DEG. Generally, the
434 non-mown plants show a much higher number of upregulated biotic stress-related genes during a
435 phase in their life when senescence commences and they become more susceptible to pathogen
436 attacks. The mown plants during their regrowth phase are not senescing and their younger organs
437 seem be less affected by pathogens.

438

439

440 **GA related genes influence regrowth of *T. pratense***

441 GAs are involved in multiple aspects of plant development like cell elongation, flowering time
442 regulation, and seed germination. Consequently, genes encoding for proteins involved in the
443 synthesis, perception, and catabolism of the various GAs influence plant form. The RNA-Seq data
444 presented showed a high abundance of GA associated genes (Fig. 1 G and 3) which might explain the
445 morphological changes to mowing, such as rounder leaves, temporary dwarf-like appearance, and
446 higher cumulative biomass production in mown plants [25]. Two of the genes, expressed higher in
447 mown plants than in control clover plants are *GA20OX1* and *GA20OX2*. They are key enzymes of GA
448 synthesis by producing precursors of the active GA forms (reviewed in [59]) and a deficiency in their
449 activity is correlated with a dwarfed growth phenotype in *A. thaliana* and *O. sativa* [60,61]. These
450 mutants show also slow down cell division and expansion rate [62]. The upregulation of *GA20OX2* in
451 mown plants may meet the enhanced demand of active GA to promote and sustain the regrowth to
452 increase cell division and elongation. Similar to an increased GA synthesis, the expression of genes
453 involved in reception of GA, such as the ortholog of *AtGID1B* is enhanced in mown plants. In *A.*
454 *thaliana*, loss-of-function mutants of this GA receptor show a dwarfed phenotype [63] and over-
455 expression of *GID1* in *Medicago sativa* promotes biomass accumulation and leaf roundness [64].

456 In contrast to these GA responsive genes, GA2OX2 and GA2OX8 are involved in GA catabolism [65]
457 and high levels of GA are known to activate the expression of degrading enzymes [66]. Both genes
458 are expressed higher in not mown control plants than in mown plants and they inactivate the
459 bioactive GAs GA2OX over-expression results in stunted plants and delayed flowering time [67,68].
460 While mown clover plants show a higher expression of genes associated with growth activation, the
461 ABA biosynthesis regulator XERICO is higher expressed in unmown plants (Table S13). It is a direct
462 target of the DELLA protein RGL2 and in addition negatively regulated by GID1B [69,70]. Thus,
463 XERICO might restrict GA mediated growth to confer the drought adaptation of the not mown plants
464 as more water is lost through the high leaf biomass.

465 In summary, mowing seems to trigger differential gene expression of GA activating enzymes and
466 catabolic enzymes suggesting a dynamic GA response, but the gene expression patterns were not
467 informative in respect to the consequences for the phenotype. When analyzing the morphological
468 effects of GA application to mown plants (Fig. 3 A-C) we could show that external GA application
469 lead to the disappearance of specific traits typical of the mowing response. Mown plants develop
470 shorter petioles and produce a smaller leaf size area [25], but when treated with GA, leaves and
471 petioles grow up the size seen in unmown plants.

472 The growth promoting abilities of GAs by cell expansion and proliferation via stimulating the
473 degradation of growth-repressing DELLA proteins are well established [62]. The length increase of
474 petioles in GA treated mown plants is in line with reported data from non-mown *Pisum sativum*
475 (pea) plants, but in those, leaf sizes remained unchanged after GA treatment [71], suggesting a more
476 specific role for GA in the regrowth reaction after biomass loss. Moreover, it was shown in *A.*
477 *thaliana* previously, that elevated GA concentrations enhance cell division rates in the distal end of
478 leaves (reviewed in [72]). If these results are transferred to *T. pratense* GA treatment should result in
479 longer leaflets after GA treatment of mown plants. However, the leaf shape did not change, only the

480 size increased suggesting a regrowth-specific shift of growth pattern which is unaffected by GA but
481 similar to leaf shape of juvenile plants [25].

482 Interestingly, GA treatment of mown *T. pratense* plants does not generally lead to stronger
483 longitudinal growth as leaves retained the round shape characteristic for untreated mown plants.
484 These regrowth-specific characteristics can also be found in other species, for example in *A.*
485 *thaliana*, *Fragaria ananassa*, *Duchesnea indica* and *G. max* GA treatment causes elongated petioles
486 and increased leaf sizes and a more erect growth habit [73–76]. This proposes a new method to
487 increase the accumulation of biomass, suitable for animal fodder. Previous experiments with the
488 grasses *Leymus chinensis* and *Lolium perenne* showed GA action to be limited by N fertilization
489 [77,78]. Red clover, living in symbiosis with nitrogen fixing bacteria, is not dependent on additional N
490 fertilization and can produce high-protein content biomass without fertilizer on poor soils.

491

492

493 **Acknowledgements**

494 We thank Andrea Weisert for excellent technical help and Dietmar Haffer for skillfully raising the
495 plants. We also thank Volker Wissemann and Birgit Gemeinholzer for continuous discussions on the
496 project. We thank Hermann Finke for his help during visualization.

497

498 **References**

499 1. Isobe S, Klimenko I, Ivashuta S, Gau M, Kozlov NN. First RFLP linkage map of red clover (
500 *Trifolium pratense* L.) based on cDNA probes and its transferability to other red clover
501 germplasm. *Theor Appl Genet.* 2003; 108: 105–112. doi: 10.1007/s00122-003-1412-z.

- 502 2. Isobe S, Sawai A, Yamaguchi H, Gau M, Uchiyama K. Breeding potential of the backcross
503 progenies of a hybrid between *Trifolium medium* × *T. pratense* to *T. pratense*. *Can. J. Plant Sci.*
504 2002; 82: 395–399. doi: 10.4141/P01-034.
- 505 3. Eriksen J, Askegaard M, Sjøgaard K. Complementary effects of red clover inclusion in ryegrass-
506 white clover swards for grazing and cutting. *Grass Forage Sci.* 2014; 69: 241–250.
507 doi: 10.1111/gfs.12025.
- 508 4. Young ND, Debellé F, Oldroyd GED, Geurts R, Cannon SB, Udvardi MK, et al. The *Medicago*
509 genome provides insight into the evolution of rhizobial symbioses. *Nature.* 2011; 480: 520–524.
510 doi: 10.1038/nature10625.
- 511 5. Sato S, Nakamura Y, Kaneko T, Asamizu E, Kato T, Nakao M, et al. Genome Structure of the
512 Legume, *Lotus japonicus*. *Theor Appl Genet.* 2008; 15: 227–239. doi: 10.1093/dnares/dsn008.
- 513 6. Schmutz J, Cannon SB, Schlueter J, Ma J, Mitros T, Nelson W, et al. Genome sequence of the
514 palaeopolyploid soybean. *Nature.* 2010; 463: 178–183. doi: 10.1038/nature08670.
- 515 7. Schmutz J, McClean PE, Mamidi S, Wu GA, Cannon SB, Grimwood J, et al. A reference genome
516 for common bean and genome-wide analysis of dual domestications. *Nat Genet.* 2014; 46: 707–
517 713. doi: 10.1038/ng.3008.
- 518 8. Varshney RK, Song C, Saxena RK, Azam S, Yu S, Sharpe AG, et al. Draft genome sequence of
519 chickpea (*Cicer arietinum*) provides a resource for trait improvement. *Nat Biotechnol.* 2013; 31:
520 240–246. doi: 10.1038/nbt.2491.
- 521 9. Lonardi S, Muñoz-Amatriaín M, Liang Q, Shu S, Wanamaker SI, Lo S, et al. The genome of
522 cowpea (*Vigna unguiculata* L. Walp.). *Plant J.* 2019; 98: 767–782. doi: 10.1111/tpj.14349.
- 523 10. Kaur P, Bayer PE, Milec Z, Vrána J, Yuan Y, Appels R, et al. An advanced reference genome of
524 *Trifolium subterraneum* L. reveals genes related to agronomic performance. *Plant Biotechnol J.*
525 2017; 15: 1034–1046. doi: 10.1111/pbi.12697.

- 526 11. Dluhošová J, Ištváněk J, Nedělník J, Řepková J. Red Clover (*Trifolium pratense*) and Zigzag Clover
527 (*T. medium*) - A Picture of Genomic Similarities and Differences. *Front Plant Sci.* 2018; 9: 724.
528 doi: 10.3389/fpls.2018.00724.
- 529 12. Ištváněk J, Jaros M, Krenek A, Řepková J. Genome assembly and annotation for red clover
530 (*Trifolium pratense*; Fabaceae). *Am J Bot.* 2014; 101: 327–337. doi: 10.3732/ajb.1300340.
- 531 13. Vega JJ de, Ayling S, Hegarty M, Kudrna D, Goicoechea JL, Ergon Å, et al. Red clover (*Trifolium*
532 *pratense* L.) draft genome provides a platform for trait improvement. *Sci Rep.* 2015; 5: 17394.
533 doi: 10.1038/srep17394.
- 534 14. Jahufer MZZ, Ford JL, Widdup KH, Harris C, Cousins G, Ayres JF, et al. Improving white clover for
535 Australasia. *Crop Pasture Sci.* 2012; 63: 739. doi: 10.1071/CP12142.
- 536 15. Barrett BA, Faville MJ, Nichols SN, Simpson WR, Bryan GT, Conner AJ. Breaking through the feed
537 barrier: options for improving forage genetics. *Anim. Prod. Sci.* 2015; 55: 883.
538 doi: 10.1071/AN14833.
- 539 16. Řepková J, Nedělník J. Modern Methods for Genetic Improvement of *Trifolium pratense*. *Czech*
540 *Journal of Genetics & Plant Breeding.* 2014: 92–99.
- 541 17. Řepková J, Nedělník J. Modern methods for genetic improvement of *Trifolium pratense*. *Czech*
542 *J. Genet. Plant Breed.* 2014; 50: 92–99. doi: 10.17221/139/2013-CJGPB.
- 543 18. Dias PMB, Julier B, Sampoux J-P, Barre P, Dall'Agnol M. Genetic diversity in red clover (*Trifolium*
544 *pratense* L.) revealed by morphological and microsatellite (SSR) markers. *Euphytica.* 2008; 160:
545 189–205. doi: 10.1007/s10681-007-9534-z.
- 546 19. Annicchiarico P, Proietti S. White clover selected for enhanced competitive ability widens the
547 compatibility with grasses and favours the optimization of legume content and forage yield in
548 mown clover-grass mixtures. *Grass Forage Sci.* 2010; 140: no-no. doi: 10.1111/j.1365-
549 2494.2010.00749.x.

- 550 20. Ford JL, Barrett BA. Improving red clover persistence under grazing. Proceedings of the NZ
551 Grassland Association. 2011; 73: 119–124.
- 552 21. Naydenova G, Hristova T, Aleksiev Y. Objectives and approaches in the breeding of perennial
553 legumes for use in temporary pastures. Bio Anim Husb. 2013; 29: 233–250.
554 doi: 10.2298/BAH1302233N.
- 555 22. Tiffin P. Mechanisms of tolerance to herbivore damage: what do we know. Evolutionary
556 Ecology. 2000; 14: 523–536. doi: 10.1023/A:1010881317261.
- 557 23. Diaz S, Lavorel S, McIntyre SUE, Falczuk V, Casanoves F, Milchunas DG, et al. Plant trait
558 responses to grazing ? a global synthesis. Global Change Biol. 2007; 13: 313–341.
559 doi: 10.1111/j.1365-2486.2006.01288.x.
- 560 24. van Minnebruggen A, Roldán-Ruiz I, van Bockstaele E, Haesaert G, Cnops G. The relationship
561 between architectural characteristics and regrowth in *Trifolium pratense* (red clover). Grass
562 Forage Sci. 2015; 70: 507–518. doi: 10.1111/gfs.12138.
- 563 25. Herbert DB, Ekschmitt K, Wissemann V, Becker A. Cutting reduces variation in biomass
564 production of forage crops and allows low-performers to catch up: A case study of *Trifolium*
565 *pratense* L. (red clover). Plant Biol (Stuttg). 2018; 20: 465–473. doi: 10.1111/plb.12695.
- 566 26. Conaghan P, Casler MD. A theoretical and practical analysis of the optimum breeding system
567 for perennial ryegrass. Irish Journal of Agricultural and Food Research. 2011; 50: 47–63.
- 568 27. Ortega F, Parra L, Quiroz A. Breeding red clover for improved persistence in Chile: a review.
569 Crop Pasture Sci. 2014; 65: 1138. doi: 10.1071/CP13323.
- 570 28. Sato S, Isobe S, Asamizu E, Ohmido N, Kataoka R, Nakamura Y, et al. Comprehensive structural
571 analysis of the genome of red clover (*Trifolium pratense* L.). DNA Res. 2005; 12: 301–364.
572 doi: 10.1093/dnares/dsi018.

- 573 29. Shimizu-Sato S, Tanaka M, Mori H. Auxin-cytokinin interactions in the control of shoot
574 branching. *Plant Mol Biol*. 2009; 69: 429–435. doi: 10.1007/s11103-008-9416-3.
- 575 30. Stafstrom J. Influence of Bud Position and Plant Ontogeny on the Morphology of Branch Shoots
576 in Pea (*Pisum sativum* L. cv. Alaska). *Annals of Botany*. 1995; 76: 343–348.
577 doi: 10.1006/anbo.1995.1106.
- 578 31. Briske DD, Richards JH. Plant responses to defoliation: a physiological, morphological and
579 demographic evaluation. In: Bedunah DJ, Sosebee RE, editors. *Wildland plants. Physiological
580 ecology and developmental morphology*. 1st ed. Denver, Colo.: Society for Range Management;
581 1995. pp. 635–710.
- 582 32. Kotova LM, Kotov AA, Kara AN. Changes in Phytohormone Status in Stems and Roots after
583 Decapitation of Pea Seedlings. *Russian Journal of Plant Physiology*. 2004; 51: 107–111.
584 doi: 10.1023/B:RUPP.0000011309.47328.23.
- 585 33. Li S, Strid Å. Anthocyanin accumulation and changes in CHS and PR-5 gene expression in
586 *Arabidopsis thaliana* after removal of the inflorescence stem (decapitation). *Plant Physiology
587 and Biochemistry*. 2005; 43: 521–525. doi: 10.1016/j.plaphy.2005.05.004.
- 588 34. Scholes DR, Wszalek AE, Paige KN. Regrowth patterns and rosette attributes contribute to the
589 differential compensatory responses of *Arabidopsis thaliana* genotypes to apical damage. *Plant
590 Biol (Stuttg)*. 2016; 18: 239–248. doi: 10.1111/plb.12404.
- 591 35. Fischer M, Bossdorf O, Gockel S, Hänsel F, Hemp A, Hessenmöller D, et al. Implementing large-
592 scale and long-term functional biodiversity research: The Biodiversity Exploratories. *Basic and
593 Applied Ecology*. 2010; 11: 473–485. doi: 10.1016/j.baae.2010.07.009.
- 594 36. Asahina M, Satoh S. Molecular and physiological mechanisms regulating tissue reunion in
595 incised plant tissues. *J Plant Res*. 2015; 128: 381–388. doi: 10.1007/s10265-015-0705-z.

- 596 37. Blazquez MA, Green R, Nilsson O, Sussman MR, Weigel D. Gibberellins promote flowering of
597 arabidopsis by activating the LEAFY promoter. *Plant Cell*. 1998; 10: 791–800.
598 doi: 10.1105/tpc.10.5.791.
- 599 38. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
600 *Bioinformatics*. 2014; 30: 2114–2120. doi: 10.1093/bioinformatics/btu170.
- 601 39. Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, et al. De novo transcript
602 sequence reconstruction from RNA-seq using the Trinity platform for reference generation and
603 analysis. *Nat Protoc*. 2013; 8: 1494–1512. doi: 10.1038/nprot.2013.084.
- 604 40. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length
605 transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol*. 2011;
606 29: 644–652. doi: 10.1038/nbt.1883.
- 607 41. Schulz MH, Zerbino DR, Vingron M, Birney E. Oases: robust de novo RNA-seq assembly across
608 the dynamic range of expression levels. *Bioinformatics*. 2012; 28: 1086–1092.
609 doi: 10.1093/bioinformatics/bts094.
- 610 42. Ištváněk J, Dluhošová J, Dluhoš P, Pátková L, Nedělník J, Řepková J. Gene Classification and
611 Mining of Molecular Markers Useful in Red Clover (*Trifolium pratense*) Breeding. *Front Plant*
612 *Sci*. 2017; 8. doi: 10.3389/fpls.2017.00367.
- 613 43. Bekel T, Henckel K, Küster H, Meyer F, Mittard Runte V, Neuweger H, et al. The Sequence
614 Analysis and Management System – SAMS-2.0: Data management and sequence analysis
615 adapted to changing requirements from traditional sanger sequencing to ultrafast sequencing
616 technologies. *Journal of Biotechnology*. 2009; 140: 3–12. doi: 10.1016/j.jbiotec.2009.01.006.
- 617 44. Boutet E, Lieberherr D, Tognolli M, Schneider M, Bairoch A. UniProtKB/Swiss-Prot. *Methods*
618 *Mol Biol*. 2007; 406: 89–112.

- 619 45. Bairoch A, Apweiler R. The SWISS-PROT protein sequence data bank and its supplement
620 TrEMBL. *Nucleic Acids Res.* 1997; 25: 31–36. doi: 10.1093/nar/25.1.31.
- 621 46. Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, et al. Phytozome: a comparative
622 platform for green plant genomics. *Nucleic Acids Res.* 2012; 40: D1178-86.
623 doi: 10.1093/nar/gkr944.
- 624 47. Wu TD, Watanabe CK. GMAP: a genomic mapping and alignment program for mRNA and EST
625 sequences. *Bioinformatics.* 2005; 21: 1859–1875. doi: 10.1093/bioinformatics/bti310.
- 626 48. Pérez-Rodríguez P, Riaño-Pachón DM, Corrêa LGG, Rensing SA, Kersten B, Mueller-Roeber B.
627 PlnTFDB: updated content and new features of the plant transcription factor database. *Nucleic*
628 *Acids Res.* 2010; 38: D822-7. doi: 10.1093/nar/gkp805.
- 629 49. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or without a
630 reference genome. *BMC Bioinformatics.* 2011; 12: 323. doi: 10.1186/1471-2105-12-323.
- 631 50. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq
632 data with DESeq2. *Genome Biol.* 2014; 15: 550. doi: 10.1186/s13059-014-0550-8.
- 633 51. The UniProt Consortium. UniProt: the universal protein knowledgebase. *Nucleic Acids Res.*
634 2016; 45: D158-69. doi: 10.1093/nar/gkw1099.
- 635 52. NCBI Resource Coordinators. Database resources of the National Center for Biotechnology
636 Information. *Nucleic Acids Res.* 2016; 44: D7-19. doi: 10.1093/nar/gkv1290.
- 637 53. Berardini TZ, Reiser L, Li D, Mezheritsky Y, Muller R, Strait E, et al. The Arabidopsis Information
638 Resource: Making and Mining the ‘Gold Standard’ Annotated Reference Plant Genome.
639 *Genesis.* 2015; 53: 474–485. doi: 10.1002/dvg.22877.
- 640 54. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *Journal*
641 *of Molecular Biology.* 1990; 215: 403–410. doi: 10.1016/S0022-2836(05)80360-2.

- 642 55. Götz S, García-Gómez JM, Terol J, Williams TD, Nagaraj SH, Nueda MJ, et al. High-throughput
643 functional annotation and data mining with the Blast2GO suite. *Nucleic Acids Res.* 2008; 36:
644 3420–3435. doi: 10.1093/nar/gkn176.
- 645 56. Yates SA, Swain MT, Hegarty MJ, Chernukin I, Lowe M, Allison GG, et al. De novo assembly of
646 red clover transcriptome based on RNA-Seq data provides insight into drought response, gene
647 discovery and marker identification. *BMC Genomics.* 2014; 15: 453. doi: 10.1186/1471-2164-
648 15-453.
- 649 57. Pitaksaringkarn W, Ishiguro S, Asahina M, Satoh S. ARF6 and ARF8 contribute to tissue reunion
650 in incised *Arabidopsis* inflorescence stems. *Plant Biotechnology.* 2014; 31: 49–53.
651 doi: 10.5511/plantbiotechnology.13.1028b.
- 652 58. Pitaksaringkarn W, Matsuoka K, Asahina M, Miura K, Sage-Ono K, Ono M, et al. XTH20 and
653 XTH19 regulated by ANAC071 under auxin flow are involved in cell proliferation in incised
654 *Arabidopsis* inflorescence stems. *Plant J.* 2014; 80: 604–614. doi: 10.1111/tpj.12654.
- 655 59. Salazar-Cerezo S, Martínez-Montiel N, García-Sánchez J, Pérez-y-Terrón R, Martínez-Contreras
656 RD. Gibberellin biosynthesis and metabolism: A convergent route for plants, fungi and bacteria.
657 *Microbiological Research.* 2018; 208: 85–98. doi: 10.1016/j.micres.2018.01.010.
- 658 60. Rieu I, Ruiz-Rivero O, Fernandez-Garcia N, Griffiths J, Powers SJ, Gong F, et al. The gibberellin
659 biosynthetic genes AtGA20ox1 and AtGA20ox2 act, partially redundantly, to promote growth
660 and development throughout the *Arabidopsis* life cycle. *Plant J.* 2008; 53: 488–504.
661 doi: 10.1111/j.1365-313X.2007.03356.x.
- 662 61. Spielmeier W, Ellis MH, Chandler PM. Semidwarf (*sd-1*), "green revolution" rice, contains a
663 defective gibberellin 20-oxidase gene. *Proc Natl Acad Sci U S A.* 2002; 99: 9043–9048.
664 doi: 10.1073/pnas.132266399.

- 665 62. Achard P, Gusti A, Cheminant S, Alioua M, Dhondt S, Coppens F, et al. Gibberellin Signaling
666 Controls Cell Proliferation Rate in Arabidopsis. *Current Biology*. 2009; 19: 1188–1193.
667 doi: 10.1016/j.cub.2009.05.059.
- 668 63. Iuchi S, Suzuki H, Kim Y-C, Iuchi A, Kuromori T, Ueguchi-Tanaka M, et al. Multiple loss-of-
669 function of Arabidopsis gibberellin receptor AtGID1s completely shuts down a gibberellin signal.
670 *Plant J*. 2007; 50: 958–966. doi: 10.1111/j.1365-313X.2007.03098.x.
- 671 64. Wang X, Li J, Ban L, Wu Y, Wu X, Wang Y, et al. Functional characterization of a gibberellin
672 receptor and its application in alfalfa biomass improvement. *Sci Rep*. 2017; 7: 41296.
673 doi: 10.1038/srep41296.
- 674 65. Thomas SG, Phillips AL, Hedden P. Molecular cloning and functional expression of gibberellin 2-
675 oxidases, multifunctional enzymes involved in gibberellin deactivation. *Proc Natl Acad Sci U S A*.
676 1999; 96: 4698–4703. doi: 10.1073/pnas.96.8.4698.
- 677 66. Yamaguchi S. Gibberellin metabolism and its regulation. *Annu Rev Plant Biol*. 2008; 59: 225–
678 251. doi: 10.1146/annurev.arplant.59.032607.092804.
- 679 67. Curtis IS, Hanada A, Yamaguchi S, Kamiya Y. Modification of plant architecture through the
680 expression of GA 2-oxidase under the control of an estrogen inducible promoter in Arabidopsis
681 thaliana L. *Planta*. 2005; 222: 957–967. doi: 10.1007/s00425-005-0037-7.
- 682 68. Sakamoto T, Kobayashi M, Itoh H, Tagiri A, Kayano T, Tanaka H, et al. Expression of a gibberellin
683 2-oxidase gene around the shoot apex is related to phase transition in rice. *Plant Physiol*. 2001;
684 125: 1508–1516. doi: 10.1104/pp.125.3.1508.
- 685 69. Ariizumi T, Hauvermale AL, Nelson SK, Hanada A, Yamaguchi S, Steber CM. Lifting della
686 repression of Arabidopsis seed germination by nonproteolytic gibberellin signaling. *Plant*
687 *Physiol*. 2013; 162: 2125–2139. doi: 10.1104/pp.113.219451.

- 688 70. Zentella R, Zhang Z-L, Park M, Thomas SG, Endo A, Murase K, et al. Global analysis of della
689 direct targets in early gibberellin signaling in Arabidopsis. *Plant Cell*. 2007; 19: 3037–3057.
690 doi: 10.1105/tpc.107.054999.
- 691 71. DeMason DA, Chetty VJ. Interactions between GA, auxin, and UNI expression controlling shoot
692 ontogeny, leaf morphogenesis, and auxin response in *Pisum sativum* (Fabaceae): or how the
693 uni-tac mutant is rescued. *Am J Bot*. 2011; 98: 775–791. doi: 10.3732/ajb.1000358.
- 694 72. Nelissen H, Gonzalez N, Inzé D. Leaf growth in dicots and monocots: so different yet so alike.
695 *Curr Opin Plant Biol*. 2016; 33: 72–76. doi: 10.1016/j.pbi.2016.06.009.
- 696 73. Guttridge CG, Thombsen PA. The Effect of Gibberellins on Growth and Flowering of *Fragaria*
697 and *Duchesnea*. *J Exp Bot*. 1964; 15: 631–646. doi: 10.1093/jxb/15.3.631.
- 698 74. Leite VM, Rosolem CA, Rodrigues JD. Gibberellin and cytokinin effects on soybean growth. *Sci*.
699 *agric. (Piracicaba, Braz.)*. 2003; 60: 537–541. doi: 10.1590/S0103-90162003000300019.
- 700 75. Tsukaya H, Kozuka T, Kim G-T. Genetic control of petiole length in *Arabidopsis thaliana*. *Plant*
701 *Cell Physiol*. 2002; 43: 1221–1228. doi: 10.1093/pcp/pcf147.
- 702 76. Hisamatsu T, King RW, Helliwell CA, Koshioka M. The involvement of gibberellin 20-oxidase
703 genes in phytochrome-regulated petiole elongation of *Arabidopsis*. *Plant Physiol*. 2005; 138:
704 1106–1116. doi: 10.1104/pp.104.059055.
- 705 77. Cai Y, Shao L, Li X, Liu G, Chen S. Gibberellin stimulates regrowth after defoliation of sheepgrass
706 (*Leymus chinensis*) by regulating expression of fructan-related genes. *J Plant Res*. 2016; 129:
707 935–944. doi: 10.1007/s10265-016-0832-1.
- 708 78. Morvan-Bertrand A, Ernstsen A, Lindgard B, Koshioka M, Le Saos J, Boucaud J, et al.
709 Endogenous gibberellins in *Lolium perenne* and influence of defoliation on their contents in
710 elongating leaf bases and in leaf sheaths. *Physiol Plant*. 2001; 111: 225–231.
711 doi: 10.1034/j.1399-3054.2001.1110214.x.

712 **Supplement Figures**


713 Figure S1: Annotation Overview: A: Distribution of transcripts that could be mapped to the *T.*
714 *pratense* genome, to a known locus and were annotated with *T. pratense* genome identifier. B:
715 Distribution of transcripts that could be mapped to an unknown *T. pratense* gene locus. C:
716 Distribution of transcripts that could not be mapped o the *T. pratense* genome. D: Distribution of
717 transcripts of whole transcriptome representing all 12 libraries.

718 Figure S2 Differential expression of putative transcription factors of *T. pratense*. The Y axis denotes
719 the number of expressed TF family members, the x axis shows the treatments and TF families.
720 Orange bars indicate that >10% of the TF members are differentially expressed between the
721 treatments, the red bars indicates that >5% are differentially expressed.

722 Figure S3: Plant architectural characteristics and growth habit of GA treated plants. A-E Measured,
723 counted or calculated plant characteristics during phenotypic monitoring experiments. GA treated
724 plants, blue; control plants, orange. Graphs show average values and 95% confidence intervals. Time
725 is shown in weeks. Growth habit of control plants (left side) vs GA treated plants (right side), after
726 approximately 2 weeks of GA treatment and regrowth (F), and after 4 weeks (G).


727

A




bioRxiv preprint doi: <https://doi.org/10.1101/775841>; this version posted September 19, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.


B




C




D



E



F



G




Figure 1





Figure 2






Figure 3