1

Interferon-responsive genes are targeted during the establishment of human cytomegalovirus latency

- 4
- 5 Elizabeth G. Elder¹, Benjamin A. Krishna^{1,2}, James Williamson^{1,3}, Eleanor Y. Lim¹, Emma Poole¹,
- 6 George X. Sedikides¹, Mark Wills¹, Christine M. O'Connor², Paul J. Lehner^{1,3}, John Sinclair^{1,*}

7 Author affiliations

- 8 1. Department of Medicine, University of Cambridge, UK
- 9 2. Lerner Research Institute, Cleveland Clinic, Cleveland, Ohio, USA
- 10 3. Cambridge Institute for Medical Research, University of Cambridge, UK
- 11 *Corresponding author, js152@cam.ac.uk

12 Abstract

Human cytomegalovirus (HCMV) latency is an active process which remodels the latently infected 13 14 cell to optimise latent carriage and reactivation. This is achieved, in part, through the expression of 15 viral genes, including the G-protein coupled receptor US28. Here, we use an unbiased proteomic screen to assess changes in host proteins induced by US28, revealing that interferon-inducible genes 16 17 are downregulated by US28. We validate that MHC Class II and two PYHIN proteins, MNDA and 18 IFI16, are downregulated during experimental latency in primary human CD14⁺ monocytes. We find that IFI16 is targeted rapidly during the establishment of latency in a US28-dependent manner, but 19 20 only in undifferentiated myeloid cells, a natural site of latent carriage. Finally, by overexpressing 21 IFI16, we show that IFI16 can activate the viral major immediate early promoter and immediate early

2

22 gene expression during latency via NF-KB, a function which explains why downregulation of IFI16

23 during latency is advantageous for the virus.

24 Importance

Human cytomegalovirus (HCMV) is a ubiquitous herpesvirus which infects 50-100% of humans
worldwide. HCMV causes a lifelong subclinical infection in immunocompetent individuals, but is a
serious cause of mortality and morbidity in the immunocompromised and in neonates. In particular,
reactivation of HCMV in the transplant setting is a major cause of transplant failure and related
disease. Therefore, a molecular understanding of HCMV latency and reactivation could provide
insights into potential ways to target the latent viral reservoir in at-risk patient populations.

31

32 Introduction

33 Lifelong persistence of Human cytomegalovirus (HCMV) is underpinned by viral latency and 34 reactivation. Following primary infection, the ubiquitous betaherpesvirus HCMV establishes latency in cell types including early myeloid lineage cells ^{1–4}. Viral genome is maintained in these cells in the 35 36 relative absence of immediate early (IE) gene expression or production of infectious virions. 37 Reactivation of HCMV is associated with differentiation of myeloid lineage cells to mature dendritic 38 cells and macrophages; as such, reactivation events are thought to occur sporadically throughout the lifetime of the host ^{5–8}. In immunocompetent individuals, both primary infection and reactivation 39 events are well-controlled by a broad and robust immune response⁹. However, HCMV reactivation is 40 41 a major cause of morbidity and mortality in immunocompromised patients, including stem cell and organ transplant recipients ^{10,11}. 42

A key hallmark of latency is the relative suppression of IE gene expression ^{1,2,12-14}, which is controlled
by the major immediate early promoter ¹⁵⁻¹⁷, and the subsequent lack of infectious virion
production. The establishment of latency via MIEP repression in early myeloid lineage cells requires

46	both host and viral factors ¹⁸ . One viral factor that suppresses MIEP activity is the G-protein coupled
47	receptor (GPCR) US28, a virally encoded chemokine receptor homologue, which is expressed de
48	<i>novo</i> during latency as well as being delivered to cells with the incoming virion $^{19-24}$ and this
49	incoming viral US28 is functional ²⁵ . US28 modulates the signalling pathways of early myeloid cells; it
50	attenuates MAP kinases, NF-KB, and c-fos, whilst activating STAT3 and iNOS ^{21,23,25} . All of these
51	contribute to the repression of MIEP activity. This US28-mediated signalling is so critical to latency
52	that US28-deleted viruses, or the loss of G-protein coupling by the US28 mutant R129A, result in lytic
53	infection of undifferentiated myeloid cells ^{19,21,23,25} . Furthermore, when examined, these US28-
54	mediated effects on cell signalling did not occur during lytic infection or in permissive cells ²¹ ,
55	implying that US28 represses the MIEP during latency but does not impair reactivation following
56	cellular differentiation. This is reflective of the cell type-specific nature of US28-mediated signalling
57	21,26
58	Since US28 can modulate all these pathways and control the MIEP, we hypothesised that US28
59	would also cause changes in host protein expression. Here, we perform a proteomic screen
60	comparing host cell protein abundance in myelomonocytic THP-1 cells expressing wild-type (WT)
61	US28 or the US28-R129A signalling mutant. We find that the expression of many host proteins are
62	decreased in the presence of US28-WT compared with the US28-R129A mutant and a large
63	proportion of these proteins are interferon inducible. In particular, the two Pyrin and HIN domain
64	(PYHIN) family proteins, Myeloid Cell Nuclear Differentiation Antigen (MNDA) and Gamma-
65	Interferon-Inducible Protein 16 (IFI16), are downregulated by US28, as well as MHC Class II proteins.
66	IFI16 is associated with the nuclear sensing of herpesvirus DNA ^{27–31} and control of herpesvirus gene
67	expression ³²⁻³⁹ , and also represses viral transcription during HIV latency ⁴⁰ , but the effects of IFI16
68	on HCMV latent infection is unknown. Downregulation of HLA-DR/MHC Class II is important for the
69	evasion of CD4 ⁺ T cell responses to latently infected myeloid cells 41 , while antiviral effects of MNDA
70	have yet to be reported.

We have validated the downregulation of IF116, MNDA, and HLA-DR in US28-expressing THP-1 cells
and during experimental latency in primary CD14⁺ monocytes. We find that HCMV downregulates
IF116 within the first 24 hours of infection of myeloid cells in a US28-dependent manner, but this
effect is lost in differentiated dendritic cells. We propose that downregulation of IF116 is beneficial to
the establishment of latency because overexpression of IF116 drives MIEP activity and IE gene
expression via NF-KB. By targeting the downregulation of IF116, US28 actively promotes the
establishment of latency in early myeloid lineage cells.

78

79 Materials and Methods

80 Cells

81 All cells were maintained at 37 °C in a 5% CO₂ atmosphere. THP-1 cells (ECACC 88081201) were 82 cultured in RPMI-1640 media (Sigma) supplemented with 10% heat-inactivated fetal bovine serum 83 (FBS; PAN Biotech), 100 U/mL penicillin and 100 µg/mL streptomycin (Sigma), and 0.05 mM 2mercaptoethanol (Gibco). Kasumi-3 cells (ATCC® CRL-2725) were cultured in RPMI-1640 media 84 85 (Sigma) supplemented with 20% heat-inactivated fetal bovine serum (FBS; PAN Biotech), 100 U/mL 86 penicillin and 100 µg/mL streptomycin (Sigma). During infections, THP-1 and Kasumi-3 cells were cultured in a low-serum (1%) version of this media for a minimum of 24 hours prior to inoculation, 87 and maintained in this low-serum media throughout the infection. MIEP-eGFP reporter THP-1 cells ⁴² 88 89 were a gift from M Van Loock, Johnson & Johnson. RPE-1 cells (ATCC[®] CRL-4000[™]) and Human 90 foreskin fibroblasts (Hff1; ATCC[®] SCRC-1041[™]) were maintained in DMEM (Sigma) supplemented 91 with 10% heat-inactivated FBS and 100 U/mL penicillin and 100 µg/mL streptomycin. 293T cells 92 (ECACC 12022001) were maintained in DMEM (Sigma) supplemented with 10% heat-inactivated FBS 93 but without penicillin or streptomycin. Phorbol 12-myristate 13-acetate (PMA, Sigma) was used to 94 induce myeloid cell differentiation at a concentration of 20 ng/mL.

95	Primary CD14 ⁺ monocytes were isolated from apheresis cones (NHS Blood & Transplant Service) or
96	from peripheral blood taken from healthy volunteers as previously described 43 . Briefly, CD14 $^{+}$
97	monocytes were isolated from total PBMC by magnetic-activated cell sorting (MACS) using CD14 $^{\scriptscriptstyle +}$
98	microbeads (Miltenyi Biotech). The monocytes were plated on tissue culture dishes (Corning) or
99	slides (Ibidi), or kept in suspension in X-Vivo 15 media (Lonza) supplemented with 2 mM L-glutamine.
100	Mature dendritic cells were generated by treating CD14 $^{\scriptscriptstyle +}$ monocytes with granulocyte-macrophage
101	colony-stimulating factor (GM-CSF, Miltenyi, 1000U/mL) and interleukin-4 (IL-4, Miltenyi, 1000U/mL)
102	for 5 days before addition of lipopolysaccharide (LPS, Invivogen, 50 ng/mL) for 2 further days.
103	Primary human CD34 ⁺ hematopoietic progenitor cells, isolated from adult bone marrow, were
104	purchased from STEMCELL Technologies and cultured in X-Vivo 15 media (Lonza).
105	Generation of lentiviruses and retroviruses
106	The lentiviral vectors encoding US28 from the VHL/E strain of HCMV have been described previously

107 ²¹; US28 is expressed in these vectors via the SFFV promoter. The lentiviral vectors pHRSIN UbEm and pHRsin SV40blast were a kind gift from D. van den Boomen, University of Cambridge, and were 108 based upon a previously published lentiviral system^{44,45}. Briefly, expression of the gene of interest is 109 110 also driven by the Spleen Focus-Forming Virus (SFFV) promoter, and the selectable markers Emerald 111 and blasticidin resistance are driven by the Ubiquitin promoter (UbEm) and the SV40 promoter (SV40blast), respectively. The sequence encoding US28 from the VHL/E strain of HCMV was cloned 112 113 into pHRSIN UbEm using the EcoR1 and Spe1 sites using the following primers: US28 FW 5' GCACGAATTCCATATGACGCCGACGACGAC AND RV 5' CTGCACTAGTTTACGGTATAATTTGTGAGAC. The 114 115 sequence encoding IFI16 was cloned into pHRsin SV40blast using the BamHI and NotI sites using the 116 following primers: IFI16 FW 5' GATTGCGGCCGCATGGGAAAAAAATACAAGAACATTGTTC and RV 5' 117 GATCGGATCCTTAGAAGAAAAAGTCTGGTGAAGTTTC.

The sequence encoding US28-3XFLAG was cloned from TB40/EmCherry-US28-3XFLAG into the
 retroviral plasmid pBABE eGFP (a gift from Debu Chakravarti (Addgene plasmid #36999)) as

6

120	described previously ²⁵ . The Q5 site directed mutagenesis kit (New England Biotech) was used to
121	generate the US28-R129A mutant of this vector, which was verified by Sanger Sequencing.
122	Expression of US28 in these vectors is driven by the long terminal repeat and partial gag.
123	Generation of VSV-G pseudotyped lentiviral particles was conducted generally in line with the Broad
124	Institute Protocols. Five hundred thousand 293T cells were transfected with 1250 ng of lentiviral
125	expression vector, 625 ng of lentiviral packaging vector psPAX and 625 ng of envelope vector
126	pMD.2G (both gifts from S. Karniely, Weizmann Institute, Israel) using transfection reagent FuGene6
127	(Promega) according to the manufacturer's instructions. For generation of VSV-G pseudotyped
128	retrovirus particles, 1250 ng of the murine leukemia virus retroviral packaging vector KB4 46 (a gift
129	from H. Groom, University of Cambridge) was transfected along with 625 ng pMD.2G and 1250 ng
130	retroviral expression vector.
131	Lentiviral and retroviral transduction
132	Supernatants from transfected 293T cells were harvested at 36 and 60 hours post transfection,
133	filtered through a 0.45 μm syringe filter, and used to transduce THP-1 cells in the presence of 2

134 μ g/mL polybrene. When necessary, lentiviral titres were determined by in-house p24 enzyme-linked

immunosorbent assay (ELISA). For transduction with puromycin-resistance vectors, puromycin (2

136 μg/mL, Sigma) was added to media and refreshed every 2-5 days until all control non-transduced

137 THP-1 cells were dead. Similarly, where blasticidin-resistance vectors were used, blasticidin (1

μg/mL, Invivogen) was added to media. Emerald positive cells were sorted using a BD FACSAriaIII
 instrument.

140 Human cytomegaloviruses

Infection of monocytes and THP-1 cells were carried out at a multiplicity of infection (MOI) of 3 as
determined by titration on RPE-1 cells. TB40/E BAC4 strains were propagated in RPE-1 cells by
seeding 50% confluent T175 flasks with virus at an MOI of 0.1. Spread of virus was monitored for 2-6
weeks following inoculation by fluorescence microscopy, and infected monolayers were subcultured

145 twice during this period. When cells were 95-100% infected, supernatants were harvested on three 146 occasions spaced over 7 days and stored at -80°C. In the final harvest, the monolayer was scraped 147 and also stored at -80°C. After thawing the virus-containing media, cell debris was pelleted by centrifugation at 1500 x q for 20 minutes at RT. Then, the clarified supernatant was concentrated by 148 149 high speed centrifugation at $14500 \times g$ for 2 hours at 18° C. Virus-containing pellets were then 150 resuspended in X-vivo-15 media in aliquots at -80°C. 151 TB40/EmCherry-US28-3XFLAG and TB40/EmCherry-US28∆ have been described previously ¹⁹. TB40/Eqfp⁴⁷ and TB40/E BAC4 SV40 mCherry IE2-2A-GFP⁴⁸ were kind gifts from E.A. Murphy, SUNY 152 Upstate Medical University. TB40/E BAC4 IE2-eYFP has been described previously ^{49,50}. TB40/E BAC4 153 GATA2mCherry has been described previously⁵¹. TB40/E with deleted NF-KB sites in the MIEP at 154 positions -94, -157, -262 and -413, and the revertant virus, were a kind gift from Jeffery Meier and 155

156 Ming Li (University of Iowa, United States), and have been described previously⁵².

157 UV-inactivation of virus was performed by placing a 100 μL aliquot of virus in one well of a 24-well

plate and placing this within 10cm of a UV germicidal (254 nm) lamp for 15 minutes, which routinely

159 results in no detectable IE gene expression upon infection of Hff1 cells.

160 Immunofluorescence staining and image analysis

Cells were fixed with 2% paraformaldehyde for 15 minutes and permeablised with 0.1% Triton-X100 161 162 for 10 minutes at RT. Blocking and antibody incubations were performed in phosphate buffered 163 saline (PBS) with 1% bovine serum albumin and 5% normal goat serum. Antibodies used: anti-IFI16 (Santa Cruz sc-8023, 1:100), anti-FLAG (Sigma F1804, 1:1000), anti-MNDA (Cell Signaling Technology 164 3329, 1:100), anti-IE (Argene 11-003, 1:1000 or directly conjugated to FITC, 1:100), anti-GFP (directly 165 conjugated to FITC, Abcam ab6662, 1:200), anti-mCherry (Abcam ab167453, 1:500), anti-HLA DR 166 167 (conjugated to Brilliant Violet 421, Biolegend Clone L423 or Abcam ab92511 1:100). Cells were 168 imaged with a widefield Nikon TE200 microscope and images were processed using ImageJ. For 169 contingency analyses of IFI16 expression during experimental latency, cells were assigned as 'IFI16

positive/negative', and 'infected/uninfected' and then counted. These results were then analysed using Fischer's Exact statistical test for significance. For analysis of signal intensity, nuclear stained images were used to create a mask from which intensity values of the corresponding IFI16/MNDA stained image were derived using the Analyze Particles feature of ImageJ. Cells were assigned as infected or uninfected based on signal from the GFP/mCherry stain. The average signal intensity of uninfected cells was used to normalise the signal intensity in order to compare different fields of view.

177 Inhibitors

The c-fos inhibitor T5524 was purchased from Cayman Chemical, solubilised in DMSO and used at 10
μM. The Janus kinase inhibitor Ruxolitinib was purchased from Cell Guidance Systems, solubilised in
DMSO and used at 5 μM. The IKKα inhibitor/NF-κB pathway inhibitor BAY11-7082 was purchased
from Santa Cruz, solubilised in DMSO, and used at a concentration of 5 μM.

182 Western blotting

183 Except for US28 blots, cells were lysed directly in Laemmli Buffer and separated by SDS-PAGE. 184 Following transfer to nitrocellulose, the membrane was blocked in 5% milk in tris buffered saline 185 (TBS) with 0.1% Tween-20. Antibodies used: anti-IFI16 (Santa Cruz sc-8023, 1:500), anti-MNDA (Cell 186 Signaling Technology 3329, 1:250), anti-STAT1 (Cell Signaling Technology 14994, 1:1000), anti-187 phosphoSTAT1 Tyr701 (Cell Signaling Technology 9167, 1:1000, anti-beta actin (Abcam ab6276, 188 1:5000). For US28 blots, cells were pelleted, washed once in ice cold PBS, then lysed in native lysis buffer (25 mM Tris HCl pH 7.4, 150 mM NaCl, 1 mM EDTA, 1% NP40, 5% glycerol, plus protease 189 190 inhibitors) for 30 minutes, vortexing every 10 minutes. After the addition of non-reducing Laemmli 191 buffer, samples were heated at 42°C for 10 minutes and then separated by SDS-PAGE. Polyvinylidene difluoride membranes were used for transfer, and blocked membranes were incubated with the 192 rabbit anti-US28 antibody⁵³ (a gift from M. Smit, Vrije University) at 1:1000 dilution. To quantify 193 194 western blots, the Analyze Gels feature of Image J was used to plot the band intensities. Signal for

9

- 195 genes of interest were then normalised to US28-R129A cells and then normalised to either the
- 196 relative amounts of actin or STAT1, as described in the figure legends.

197 Flow cytometry

- 198 Transduced THP-1 cells and MIEP-reporter THP-1 cells were analysed on a BD Accuri Instrument. Live
- 199 cells were gated using forward and side scatter. Paraformaldehyde-fixed cells were stained using
- anti-HLA-DR APC conjugate (Biolegend, Clone L243, 1:50). Latently infected CD14⁺ monocytes were
- 201 fixed with 1% paraformaldehyde and stained using anti-HLA-DR Pacific blue conjugate (Biolegend,
- 202 Clone L243, 1:50) and anti-HLA-A,B,C, PE-Cy7 conjugate (Biolegend, Clone W6/32, 1:50), before
- 203 analysis on the Nxt Attune Instrument (Thermo Fisher).

204 RNA extraction, reverse transcription and quantitative PCR

- 205 RNA was extracted and purified using Direct-Zol RNA MiniPrep kit (Zymo Research) according to the
- 206 manufacturer's instructions. A total of 5 ng of purified RNA was used in RT-qPCR reactions,
- 207 performed using QuantiTect SYBR[®] Green RT-PCR Kit reagents (Qiagen) on a StepOne Real-Time PCR
- 208 instrument (Applied Biosystems). TATA-box binding protein (TBP) was used as a reference gene and
- 209 fold changes were analysed by the 2-ΔΔCt method. Reverse transcription was performed using the
- 210 Qiagen QuantiTect Reverse Transcription kit, and then cDNA was used in qPCR analysis using New
- 211 England Biotech LUNA SYBR Green qPCR reagents using TBP or GAPDH as a reference gene. Primers

212 used: US28 FW: AATCGTTGCGGTGTCTCAGT; US28 RV: TGGTACGGCAGCCAAAAGAT; MNDA FW:

- 213 GGAAGAAGCATCCATTAAGG; MNDA RV: GTTTGTCTAGACAGGCAAC; IFI16 FW: CTGCACCCTCCACAAG;
- 214 IFI16 RV: CCATGGCTGTGGACATG; TBP FW: CGGCTGTTTAACTTCGCTTC; TBP RV:
- 215 CACACGCCAAGAAACAGTGA; HLA-DRA FW: TGTAAGGCACATGGAGGTGA; HLA-DRA RV:
- 216 ATAGGGCTGGAAAATGCTGA; IE72 FW: GTCCTGACAGAACTCGTCAAA; IE72 RV:
- 217 TAAAGGCGCCAGTGAATTTTTCTTC; GAPDH FW TGCACCAACTGCTTAGC, GAPDH RV:

218 GGCATGGACTGTGGTCATGAG.

10

219 Sequence alignment

- US28 sequences were exported from GenBank (TB40/E BAC4 EF999921.1, AD169 FJ527563.1, Merlin
- 221 NC_006273.2, VHL/E MK425187.1) or from sequenced plasmids, and aligned using Clustal Omega⁵⁴
- https://www.ebi.ac.uk/Tools/msa/clustalo/ and the output format MView⁵⁵.
- 223 Cell lysis, digestion and cleanup for proteomic analysis
- 224 Cells were harvested by centrifugation and washed 3x in cold phosphate-buffered saline (PBS)
- 225 before finally pelleting into a low adhesion microfuge tube. Cell pellets were lysed in 2%/50 mM
- 226 Triethylamminium bicarbonate (TEAB) pH 8.5. Samples were quantified by BCA assay and 50µg of
- 227 each sample was taken and adjusted to the same volume with lysis buffer. Reduction and alkylation
- 228 was achieved by addition of 10 mM Tris (2-carboxyethyl)phosphine(TCEP) and 20 mM
- iodoacetamide for 20mins at room temperature in the dark followed by quenching with 10mM DTT.
- 230 Samples were further purified and digested using a modified Filtered Aided Sample Prep (FASP)
- 231 protocol. Briefly, samples were brought to 500 μL volume with 8 M urea/TEAB and loaded onto a
- 30kDa molecular weight cut-off (MWCO) ultrafiltration device. Samples were then washed 3 times
- with 8 M urea/TEAB followed by 3 times with 0.5% deoxycholate (SDC) /50mM TEAB. Samples were
- finally resuspended in ~50 μL of SDC /TEAB containing 1 μg trypsin and incubated overnight at 37°C.
- 235 After digestion samples were recovered from the filter device by centrifugation, acidified to
- precipitate SDC and cleaned up by two-phase partitioning into 2x volumes of ethyl acetate (repeated
- twice) before drying in a vacuum centrifuge.

238 TMT Labelling

Samples were resuspended in 20 µL 100 mM TEAB and to each tube 0.2 µg of a unique tandem mass
tag (TMT) label for each sample was added in 8.5 µL acetonitrile (ACN) and incubated for 1 h at room
temperature. TMT reactions were quenched by addition of 3 µL of 200 mM ammonium formate,
pooled and dried in a vacuum centrifuge. The sample was then resuspended in 800µL 0.1%
trifluoroacetic acid (TFA) and acidified to ~pH 2 with formic acid (FA) before performing a C18-solid

11

phase extraction (C18-SPE) using a Sep-Pak cartridge (Waters) attached to a vacuum manifold. C18
eluate was dried in a vacuum centrifuge and resuspended in 40 µL 200 mM ammonium formate, pH
10.

247 High pH Revered Phase Fractionation

248 Sample was injected onto an Ultimate 3000 RSLC UHPLC system (Thermo Fisher Scientific) equipped 249 with a 2.1 i.d. x25cm, 1.7µm particle Kinetix Evo C18 column (Phenomenex). Mobile phase consisted 250 of A: 3% ACN, B: ACN and C: 200mM ammonium formate pH 10. Isocratic conditions were 90% 251 A/10%C and C was maintained at 10% throughout the gradient elution. Separations were carried out 252 at 45°C. After loading at 200 μ L/min for 5 min and ramping the flow rate to 400 μ L/min over 5 min, 253 the gradient elution proceeded as follows: 0-19% B over 10 minutes (curve 3), 19-34% B over 254 14.25mins (curve 5), 34-50% B over 8.75mins (curve 5), followed by a 10 min wash at 90% B. UV 255 absorbance was monitored at 280 nm and 15 s fractions were collected into 96 well microplates 256 using the integrated fraction collector. Peptide containing fractions were then orthogonally recombined into 24 fractions and dried in a vacuum centrifuge and resuspended in 10 µL 5% DMSO 257 258 0.5% TFA for analysis.

259 LC-MS analysis

260 All samples were injected onto an Ultimate 3000 RSLC nano UHPLC equipped with a 300 μ m i.d. x 5 261 mm Acclaim PepMap µ-Precolumn (Thermo Fisher Scientific) and a 75 µm i.d. x50c m 2.1 µm particle 262 Acclaim PepMap RSLC analytical column. Loading solvent was 0.1% TFA, analytical solvent A: 0.1% FA and B: ACN+0.1% FA. All separations are carried out at 55°C. Samples were loaded at 10 μ L/min for 5 263 264 min in loading solvent before beginning the analytical gradient. For high pH reverse phase (RP) 265 fractions, a gradient of 3-5.6% B over 4 min, 5.6 – 32% B over 162 min, followed by a 5 min wash at 266 80% B and a 5 min wash at 90% B and equilibration at 3% B for 5 min. During the gradient the 267 Orbitrap Fusion mass spectrometer (Thermo Fisher Scientific) was set to acquire spectra according 268 to the settings given in supplementary file S6 "MS Settings".

12

269 Data Processing

- 270 All raw files were searched by Mascot within Proteome Discoverer 2.1 (Thermo Fisher Scientific)
- against the Swissprot human database and a database of common contaminants.
- 272 The search parameters were as follows. Enzyme: Trypsin. MS1 tol: 10 ppm. MS2 tol: 0.6 Da. Fixed
- 273 modifications: Carbamidomethyl cysteine, TMT peptide N-termini and lysine. Variable modification
- 274 oxidised methionine. MS3 reporter ion tol: 20 ppm, most confident centroid. Mascot Percolator was
- used to calculate peptide-spectrum match false discovery rate (PSM FDR).
- 276 Search results were further processed and filtered as follows: Peptides below a percolator FDR of
- 277 0.01% and proteins below the 0.01% protein FDR (calculated from a built-in decoy database search)
- were rejected. Protein groups were then generated using the strict parsimony principle. Peptides
- both unique and razor with a co-isolation threshold of 50 and an average signal-to-noise (s/n)
- threshold of 10 were used for quantification and a normalisation of these values to the total peptide
- amount in each channel was applied. Instances where a protein was identified but not quantified in
- all channels were rejected from further analysis. "Scaled" abundances of proteins provided by
- 283 Proteome Discoverer were used to derive ratios of abundance. Q values of significance between
- 284 groups were calculated by Benjamini-Hochberg correction of p values generated using the
- 285 moderated T-test LIMMA within the R environment.

286 Results

287 Proteomic analysis reveals US28-induced changes in host proteins in myeloid cells

US28-mediated signalling is critical to latency: US28-deleted viruses, or the loss of G-protein
 coupling by the US28 mutant R129A, result in lytic infection of undifferentiated myeloid cells
 ^{19,21,23,25}. Similarly, infection of US28-WT-expressing THP-1 cells with US28-deletion viruses leads to
 complementation and the establishment of latency; both the US28-R129A and empty vector cell
 lines fail to establish latent infection under these conditions ²¹. Therefore, to understand how US28-

13

293 WT alters the host cell environment to support latency, we analysed the total proteomes of

294 myelomonocytic THP-1 cell lines which express either US28-WT (sequence derived from the VHL/E

strain of HCMV), signalling mutant US28-R129A, or the empty vector (EV) control (Figures 1A, 1B, 1C,

296 File S5).

297 By using these three cell lines, we were able to identify host cell proteins modulated by US28

298 expression in a G-protein signalling dependent and independent manner. Changes in host protein

abundance common between US28-WT and US28-R129A when compared to empty vector (Figure

300 1B and 1C) represent signalling independent changes, and these include CD44 and CD82 proteins,

301 which are each downregulated by both sets of US28-expressing cells.

302 While we do not rule out that signalling independent changes in myeloid cells driven by US28 may

303 be important for HCMV latency, G-protein dependent signalling is absolutely required for latency,

and therefore we were particularly interested in the direct comparison of host protein abundances

in THP-1 cells expressing US28-WT and US28-R129A (Figure 1A). This comparison reveals 42 host

306 proteins whose expression is two-fold or more increased or decreased by US28-WT, and our

307 analyses focussed on these signalling-dependent changes.

308 One remarkable feature of many of the most downregulated proteins in Figure 1A is that they are

309 interferon-inducible (Figure 1D, Figure S1A). According to the Interferome database (v2.01,

310 <u>www.interferome.org</u>) ⁵⁶, two-thirds (27/40) of the most downregulated proteins (fold change 1.86-

fold or higher) we identified are Type I or Type II interferon-inducible (Figure 1D). In contrast, of the

40 proteins which showed no changes (fold change = 0) in abundance between US28-WT and US28-

313 R129A, 12/40 (30%) were included in the Interferome database (File S5). Importantly,

314 overexpression of the multipass membrane US28 proteins did not lead to induction of the unfolded

protein response or other endoplasmic stress related genes (Figure S1B), suggesting that the

316 changes identified in the screen are not general effects of protein overexpression.

Since STAT1 phosphorylation is common to both the Type I and Type II interferon signalling
pathways ⁵⁷, we examined total STAT1 and phosphorylated STAT1 in US28-WT cells. We found that
US28-WT cells had lower overall levels of STAT1 and phosphorylated STAT1 compared to cells
transduced with US28-R129A or empty vector control (Figure 1E and 1F). Furthermore, when
correcting for total levels of STAT1, we found that US28-WT cells had lower relative levels of
phosphorylated STAT1 compared with US28-R129A (Figure 1E and 1G). This could help explain why
US28-WT downregulated many interferon-inducible genes in our proteomic screen.

324 IFI16, MNDA, and HLA-DR are all downregulated by US28

325 Several interferon-inducible proteins showing decreased expression were of interest as potentially 326 important targets for US28 during HCMV latency. These included MNDA (9 unique peptides; 5.0-fold 327 downregulated compared to US28-R129A), IFI16 (4 unique peptides; 2.4-fold downregulated 328 compared to US28-R129A), and components of the MHC Class II HLA-DR complex (3-6 unique 329 peptides; between 1.9 and 2.2-fold downregulated compared to US28-R129A). We began by 330 confirming US28-WT-mediated downregulation of these proteins in independently-transduced 331 US28-expressing THP-1 cells. After generating these fresh US28-expressing cell lines, we checked 332 expression levels of US28-WT or US28-R129A by RT-qPCR (Figure S2A) and Western blot (Figure S2B 333 and S2C). RT-gPCR confirmed that IFI16, MNDA, and HLA-DRA transcripts are all downregulated in 334 US28-WT-expressing cells compared to those expressing the signalling mutant R129A (Figure 2A). 335 Subsequently, we confirmed this US28-WT mediated downregulation of IFI16 and MNDA at the 336 protein level by western blot (Figure 2B, 2C, S2D, S2E, S2F) or by flow cytometry for cell surface HLA-337 DR (Figure 2F and 2G), although both US28-WT and US28-R129A cells responded similarly to IFN-y 338 stimulation, by upregulating HLA-DR similar levels (Figure 2F). This suggests that, whilst US28 is able 339 to downregulate constitutive HLA-DR expression, it is unable to prevent its induction by IFN-y. 340 Furthermore, this downregulation was not due to strain-specific effects of US28 as US28-WT also 341 downregulated HLA-DR when the US28 sequence from the TB40/E strain of HCMV was used to 342 transduce THP-1 cells (Figure S3A-D).

15

Since US28 attenuates c-fos signalling²⁵ and STAT1 phosphorylation (Figure 1E, 1F, 1G), and both 343 HLA-DR and IFI16 are fos and STAT1-responsive genes^{58–62}, we hypothesised that one or both of 344 these mechanisms is responsible for US28-mediated downregulation of IFI16 and HLA-DR. To test 345 this, we treated empty vector, US28-WT, and US28-R129A expressing THP-1 cells with the Janus 346 347 Kinase inhibitor Ruxolitinib, or the c-fos inhibitor T-5224. Ruxolitinib partially downregulated IFI16 348 expression in all three cell types (Figure 2D), but downregulated HLA-DR only in the US28-R129A cell 349 line (Figure 2H), despite a complete block in STAT1 phosphorylation (Figure S2G). The c-fos inhibitor 350 reduced IFI16 and HLA-DR expression in comparison with DMSO controls in empty vector and R129A 351 cell lines, and in the case of empty vector, this drop in expression was almost down to the level in 352 untreated US28-WT cells (Figure 2E and 2H). Taken together, we think it likely that both c-fos and 353 STAT1 attenuation are important for US28-mediated downregulation of IFI16 and HLA-DR. 354 Interestingly, the c-fos inhibitor actually increased IFI16 expression in US28-WT expressing cells. We 355 think this could be due to a basal level of c-fos being required for the expression of a host gene, as 356 yet unidentified, that is need by US28-WT to attenuate multiple signalling pathways; one candidate gene for this is the AP-1 inducible phosphatase DUSP1⁶³ which will require further investigation. 357

358 Latent infection of monocytes is associated with the downregulation of IFI16, MNDA,

and HLA-DR

360 Having confirmed key observations from our proteomic data in transduced THP-1 cells 361 overexpressing US28 in isolation, we then sought to determine whether IFI16, MNDA, and HLA-DR 362 are also downregulated in an experimental model of latency in *ex vivo* primary CD14⁺ monocytes 363 where latency-associated expression of US28 is well established. To do this, we infected CD14⁺ 364 monocytes with TB40/E-BAC4 strains of HCMV engineered to express either GFP or mCherry as 365 markers for latent or lytic infection. Firstly, we analysed CD14⁺ monocytes infected with TB40/E SV40 366 mCherry/IE2-2A-GFP. This virus drives constitutive mCherry expression in all infected cells via the SV40 promoter, but GFP expression is restricted to lytically infected cells as a result of IE2 367

368	expression, which is linked to GFP by the self-cleaving peptide 2A. Therefore, we were able to
369	distinguish IE2-positive (lytic) from IE2-negative cells (one hallmark of latency) amongst infected,
370	mCherry positive cells. At four days post infection (d.p.i.), we fixed and immunostained the
371	monocytes for our cellular proteins of interest in mCherry positive, IE2-2A-GFP negative cells (Figure
372	3A). As a control, we also differentiated monocytes with phorbol 12-myristate 12-acetate (PMA),
373	which drives IE2-2A-GFP expression through differentiation-dependent reactivation. We found that
374	IFI16, MNDA, and HLA-DR were all downregulated in latently infected, mCherry positive but IE2-
375	negative, CD14 $^{\scriptscriptstyle +}$ monocytes. Importantly, this comparison held true when comparing infected
376	monocytes with uninfected monocytes that had not had contact with viral particles (Figure 3A, S4A).
377	We then sought to look at expression of these proteins at earlier time points. US28 is a virion-
378	associated protein ¹⁹ , and incoming US28 is reported to have rapid effects on host cells ²⁵ . We
379	speculated that the downregulation of IFI16, MNDA, and HLA-DR might occur early during the
380	establishment of latency. For these experiments, we used TB40/Egfp which marks infected cells with
381	GFP expression via the SV40 promoter and confirmed the establishment of latency in this system by
382	coculture of monocytes with fibroblasts either with or without PMA-induced reactivation (Figure
383	3B). In our latency system, we found a stark and specific loss of IFI16 in infected monocytes from 24
384	hours post infection (h.p.i.) (Figure 3C), a phenotype maintained at 48 and 72 h.p.i (Figure 3C) as
385	measured by immunofluorescence. We quantified these observations in several fields of view for
386	each of these three time points, and performed contingency analyses (Fisher's Exact), which
387	confirmed specific loss of IFI16 in latently infected cells (Figure 3D). Loss of IFI16 was observed in <i>ex</i>
388	vivo infected monocytes at these time points in a total of four separate donors with TB40/Egfp virus.
389	We found a partial downregulation of MNDA by 72 h.p.i (Figure 3E and 3F), with a very small
390	downregulation at 48 h.p.i and no effect at 24 h.p.i, suggesting that modulation of MNDA is delayed
391	compared with fellow PYHIN family member, IFI16. We also observed that HLA-DR, but not
392	corresponding MHC Class I HLA-A,B,C, were downregulated at 72 h.p.i specifically in GFP positive,

17

393 latently infected monocytes (Figure 3G and 3H). Therefore, IFI16, MNDA, and HLA-DR are indeed

- downregulated at early times during the establishment of latency, with IFI16 showing
- downregulation within 24 hours of infection.

396 The downregulation of IFI16 is dependent on viral US28

- Having confirmed that IFI16 is downregulated very early during latent infection of monocytes, we
- then sought to establish whether this effect is dependent on US28. We predicted this would be the
- 399 case because of the results of our US28 proteomic screen and the established functionality of
- 400 incoming virion-associated US28²⁵. We infected monocytes with either the US28
- 401 WTTB40/EmCherry-US28-3XFLAG HCMV (US38-3XF), or the corresponding US28 deletion virus
- 402 TB40/EmCherry-US28Δ (ΔUS28). These viruses establish latent and lytic infections, respectively, in
- 403 CD34⁺ progenitor cells, Kasumi-3 cells, and THP-1 cells ^{19,25}, and we confirmed these phenotypes are
- 404 also maintained in primary CD14⁺ monocytes by supernatant transfer to permissive fibroblasts
- 405 (Figure 4A). We were also able to detect US28 protein during the establishment of latency in
- 406 monocytes by immunostaining for the FLAG epitope tag on the C terminus of US28 (Figure 4B).
- 407 To determine if US28 specifically downregulates IFI16 in the context of infection, we compared the
- 408 expression of this cellular protein in monocytes infected with US28-3XF or ΔUS28. Consistent with
- 409 Figure 3B, we found that monocytes infected with the US28-3xF virus showed downregulation of
- 410 IFI16 at 24 and 48 h.p.i., while monocytes infected with ΔUS28 displayed robust IFI16 expression at
- 411 24 h.p.i. (Figure 4C and 4D) and only partial downregulation at 48 h.p.i (Figure 4C and 4E). These
- 412 data demonstrate that the early downregulation of IFI16 in CD14⁺ monocytes is dependent on US28.

413 IFI16 is downregulated by US28 only in undifferentiated myeloid cells

- 414 We previously showed that US28 modulates cellular signalling pathways in undifferentiated, but not
- 415 differentiated THP-1 cells ²¹. We were therefore curious as to whether the effects on IFI16
- 416 expression were dependent on cellular differentiation status. This is significant because
- 417 differentiated THP-1 cells and mature dendritic cells are permissive for HCMV lytic infection. To

18

418	analyse whether these effects are differentiation-dependent, we transduced THP-1 cells with a
419	lentiviral vector that co-expresses US28 and the fluorescent protein Emerald (US28-UbEm), or co-
420	expresses eGFP and Emerald (empty UbEm), as a control. For each population, we sorted the
421	Emerald-positive THP-1 cells by FACS (Figure S4B) and validated US28 expression by RT-qPCR (Figure
422	S4C). We treated half of these cells with PMA in order to induce cellular differentiation. We found
423	that undifferentiated US28-expressing THP-1 cells downregulated IFI16, but PMA-differentiated cells
424	did not downregulate IFI16 (Figure 5A), suggesting only latency-associated expression of US28
425	attenuates IFI16 expression.
426	We also analysed the effect of cellular differentiation on IFI16 expression following infection in
427	mature dendritic cells derived by treating <i>ex vivo</i> CD14 ⁺ monocytes with GM-CSF/IL-4/LPS. Again, we
428	found that undifferentiated infected CD14 $^{+}$ monocytes downregulate IFI16 in a US28-dependent
429	manner at 48 h.p.i, while infected mature dendritic cells do not downregulate IFI16 with WT or
430	ΔUS28 HCMV (Figure 5B). Taken together, our results indicate that US28 rapidly downregulates IFI16
431	during latent infection of monocytes, but not during lytic infection of mature dendritic cells.
432	Low levels of IFI16 are maintained during long term latency in monocytes and CD34^+
433	progenitor cells
434	We next assessed whether downregulation of IFI16 and MNDA occurs during long term maintenance
435	of latency; long term downregulation of HLA-DR is already known to be important for latent carriage
436	of HCMV ⁴¹ . We infected monocytes with HCMV that drives mCherry from GATA2 promoter, and

437 maintains this marker for far longer during latency than SV40 promoter-driven tags⁵¹. At 10 and 14

438 d.p.i., IFI16 remained absent and MNDA remained partially downregulated in infected cells (Figure

- 439 5C). We then repeated this analysis in primary CD34⁺ hematopoietic progenitor cells (HPCs), a site of
- 440 long-term *in vivo* latent carriage, as well as the Kasumi-3 cell line, an experimental model for HCMV
- 441 latency⁶⁴. Consistent with our observations in monocytes, and RNAseq experiments in cord blood
- 442 derived CD34⁺ cells⁶⁵, IFI16 levels were low or absent in almost all infected cells imaged at 4 and 10

19

days post infection (Figure 5D and 5E). Thus, it seems likely that downregulation of IFI16 is a 443

444 conserved process in cellular sites of HCMV latency.

467

IFI16 activates IE gene expression via NF-κB in myeloid cells 445

446 Having established that US28 downregulates IFI16 early during the establishment of latency, we 447 wanted to address why this may be beneficial to the virus for latent infection. One function of IFI16 is the sensing of viral DNA and subsequent induction of Type I interferon or interleukin-1-beta^{27,65.} 448 449 While we do not currently rule out a potential role for IFI16-mediated sensing of HCMV during 450 latency, we were more struck by the previous work identifying IFI16 as a modulator of host and viral transcription ^{32–34,36,37,66–71}. In particular, IFI16 is capable of activating the MIEP and driving IE gene 451 expression within the first 6 hours of lytic infection of fibroblasts ^{32,34}, though at later times IFI16 452 blocks early and late gene expression ^{34,37}. Given that a hallmark of HCMV latency is the suppression 453 of IE gene expression ⁷², we hypothesised that high levels of IFI16 might drive MIEP activity and IE 454 455 gene expression in undifferentiated myeloid lineage cells. To address this, we transduced and selected THP-1 cells with control empty vector (EV) or IFI16-overexpression lentiviruses to generate 456 457 control and IFI16-overexpressing cell lines. We validated IFI16 overexpression by western blot 458 (Figure 6A) and then infected these cell lines with recombinant HCMV carrying an IE2-YFP cassette to allow us to identify cells that express IE2⁵⁰. Undifferentiated THP-1 cells are an established model for 459 a number of aspects of HCMV latency⁷³, including the significant repression of IE2 expression ⁷³. 460 461 When we infected control and IFI16-overexpressing THP-1 cells with HCMV in five paired experiments, we found a consistent increase in the number of IE positive cells in IFI16-462 463 overexpressing THP-1 cells (Figure 6B and 6C), suggesting IFI16 overexpression drives IE protein 464 production in cells that would otherwise repress this viral protein. 465 Prior work has identified that IFI16 could activate IE gene expression in a UL83-dependent manner during lytic infection ^{32,37}, but since this tegument protein does not enter the nucleus in the CD34⁺ 466 progenitor cell model of HCMV latency ⁷⁴, we hypothesised that IFI16 could activate the MIEP

20

468 without additional virion components. To test this hypothesis, we utilized a THP-1 MIEP reporter system; THP-1 cells in which an integrated 1151 bp region of the MIEP drives the expression of eGFP 469 470 ⁴². In these undifferentiated THP-1 cells, the MIEP is epigenetically repressed unless stimulated (for example by differentiation)⁴². We treated these MIEP-eGFP THP-1 cells with control lentiviruses or 471 472 lentiviruses which drive the overexpression of IFI16, ensuring equivalent lentivirus infection of 473 reporter cells by p24 ELISA. These cultures were maintained for two weeks, after which we validated 474 IFI16 expression by immunofluorescence (Figure 6D) and then analysed eGFP expression by flow 475 cytometry (Figure 6E). We found that the IFI16-overexpressing cells had increased eGFP expression 476 compared with controls (Figure 6D and 6E), suggesting that IFI16 overexpressed in isolation and in the absence of additional HCMV components, drives MIEP activity. Furthermore, culturing THP-1 477 MIEP-eGFP reporter cells with supernatants from the empty vector or IFI16-overexpressing cell lines 478 479 in Figure 6A resulted in no significant MIEP activity, suggesting that the effect is mediated 480 intracellularly, and not by a secreted factor (Figure 6E). IFI16 activates NF-κB signalling in a number of contexts^{66,75}, and our previous work indicates that 481 US28-mediated attenuation of NF-KB signalling is important for the establishment of latency²¹. 482 483 Therefore, we hypothesised that IFI16 activates the MIEP via NF-KB. We found increased nuclear NF-484 κB localisation in IFI16 overexpressing cells (Figure 6F), and by using the NF-κB pathway inhibitor, 485 BAY11-7082, we were able to ameliorate the effect of IFI16 overexpression on IE (Figure 6G and 6I), suggesting that NF-κB plays an important role in this pathway. Finally, we infected IFI16-486 overeexpressing cells with HCMV that lacks NF-κB sites within the MIEP⁵² to check whether IFI16 487 488 exerts its effects via direct binding of NF-kB to the MIEP. In this case, IFI16 overexpression failed to 489 induce IE gene expression, unlike IFI16 cells infected with the revertant strain (Figure 6H and 6I). 490 Taken together, our data are consistent with the view that IFI16 activates IE gene expression in early 491 myeloid lineage cells by allowing NF-KB to bind at the MIEP.

492 Discussion

The viral GPCR US28 is expressed during both lytic and latent infection of HCMV. While US28 is
dispensable for lytic replication *in vitro* ^{76,77}, it is essential for the establishment and maintenance of
HCMV latency in early myeloid lineage cells ^{19,21,23,25}. This is attributable, in part, to the ability of
US28 to suppress the major immediate early promoter; a US28 function specific for undifferentiated
myeloid cells ^{18,21,23,25}.

498 We hypothesised that this ability of US28 to so profoundly regulate viral IE gene expression in 499 undifferentiated myeloid cells was likely via US28-mediated modulation of host protein abundance 500 and, therefore, we wanted to determine whether such US28-driven changes could be important for the establishment or maintenance of HCMV latency. While previous work has used targeted arrays 501 to assess US28-mediated effects on myeloid cells ^{21,23,25}; here we have used an unbiased proteomic 502 503 screen to understand how US28 reprograms host cells in order to support latent infection. Our screen compared host protein abundance in control THP-1 cells or THP-1 cells which express either 504 505 WT-US28 or the US28 signalling mutant, US28-R129A. As such, we could assess the signalling-506 dependent and signalling-independent effects of US28. We then chose to focus on signalling-507 dependent changes because G protein coupling via the residue R129A is essential for experimental 508 latency ^{21,25}. However, we predict that some of the signalling-independent changes driven by US28 509 could also be important for HCMV latency, since these changes included alterations in several cell-510 surface molecules such as co-stimulatory molecule CD82, adhesion molecule CD44, and in receptor 511 tyrosine kinase FLT3. The latter two cellular factors are implicated in myeloid cell differentiation, which is intimately linked with HCMV latency and reactivation ^{18,72,78–80}. As such, modulating these 512 513 cell-surface molecules could help to control interactions with immune effectors and cellular 514 differentiation-linked reactivation.

515 By looking at changes in host protein abundance between US28-WT and US28-R129A expressing 516 THP-1 cells, we found a number of significant changes in the host proteome which likely result

22

517 specifically from US28 signalling. Interestingly, we found US28-WT downregulated a large number of interferon-inducible proteins, including canonical interferon-stimulated genes (ISGs) like OAS2 and 518 519 IFITM3, as well as MNDA, IFI16, and several HLA-DR components. We found that levels of both total 520 STAT1 and phosphorylated STAT1 were reduced in US28-WT expressing cells, a mechanism that may 521 act in synergy with the US28-mediated attenuation of c-fos to downregulate interferon-inducible genes^{25,61}. Modulation of interferon signalling has not previously been reported for US28, but in the 522 523 context of the latently infected monocyte, a general block in downstream interferon signalling may be important for maintaining the polarisation of the monocyte ^{81,82}, or perhaps to avoid the anti-viral 524 activities of ISGs. We believe these questions merit further interrogation. 525 526 We chose to focus on the two PYHIN proteins and the set of HLA-DR components which are downregulated by US28. We confirmed downregulation of IFI16, MNDA, and HLA-DR in THP-1 cells 527 528 which overexpress US28 and recapitulated these effects in experimental latency in primary CD14⁺ 529 monocytes. HLA-DR was previously reported to be downregulated during experimental latency in 530 granulocyte-macrophage progenitor cells, which prevents CD4⁺T cell recognition and activation ^{41,83,84}. Whilst this down-regulation of MHC Class II involved the expression of the latency-associated 531 gene UL111A⁴¹, our data argue that viral US28 could also contribute to this phenotype. Little is 532 533 known about the function of MNDA, a myeloid specific PYHIN protein implicated in neutrophil cell death ⁸⁵ and monocyte transcriptional networks ⁸⁶. Ongoing work in our laboratory aims to identify 534 whether US28-mediated downregulation of MNDA during latent infection could benefit latent 535 carriage and/or reactivation. 536

Our results clearly characterised a rapid downregulation of IFI16 during the establishment of latency
in monocytes, which occurred within the first 24h of infection and was also maintained during long
term latency in monocytes and CD34⁺ HPCs. The early downregulation of IFI16 was clearly US28dependent as ΔUS28 virus failed to display immediate IFI16 down-regulation. However, we did
observe a partial downregulation of IFI16 in ΔUS28-infected monocytes at later time points of

23

infection. We think it likely that this involves an unidentified lytic-phase viral gene product, which
 may be required for overcoming the known IFI16-mediated restriction of HCMV lytic infection
 ^{30,34,36,37} and occurs as a result of ΔUS28 virus initiating a lytic infection in undifferentiated
 monocytes.

546 Our observation that the US28-dependent downregulation of IFI16 occurred rapidly (within 24h of 547 infection) may, in part, be attributable to incoming US28 which is functional ²⁵. IFI16 protein has a 548 short half-life of approximately 150 minutes in fibroblasts ⁸⁷ and therefore, incoming US28 protein 549 may rapidly target IFI16 transcription in latently infected monocytes, as it does in US28-expressing 550 THP-1 cells, resulting in loss of IFI16 within 24 hours of infection; this is then maintained by 551 subsequent latency-associated *de novo* US28 expression.

552 We found that preventing IFI16 expression has a clear benefit to the establishment of HCMV latency. 553 This contrasts with previous analyses of latency in other viral systems, where IFI16 expression is necessary to repress lytic viral transcription ^{39,40}. In our study, IFI16 overexpression activated MIEP 554 activity in the absence of additional viral proteins, and, furthermore, IFI16 overexpression increased 555 IE positive nuclei in latently infected THP-1 cells. IFI16 activates the MIEP during lytic infection ^{32,34}, 556 557 though in these cases an additional viral gene product, UL83, is thought to be required. Our results 558 suggest that UL83 is not required for IFI16-mediated activation of the MIEP in undifferentiated 559 myeloid cells, and suggest that IFI16 activates NF-KB to achieve this, as use of either an NF-KB 560 pathway inhibitor or deletion of NF-KB binding sites from the MIEP prevented IFI16-mediated IE 561 expression. We believe this provides one mechanism by which US28 blocks NF-KB activity early 562 during latency, a phenomenon we previously showed to be important for the establishment of latency in myeloid cells²¹. 563

Taken together, our results suggest that one of the early events in the establishment of latency in
 CD14⁺ monocytes is the US28-mediated targeting of interferon responsive genes, including the
 downregulation of IFI16, which serves to support the repression of the MIEP.

24

567 Acknowledgements

- 568 We would like to thank Linda Teague, Paula Rayner, Isobel Jarvis, and Roy Whiston for technical
- assistance and Dr. Ian Groves for critical reading of the manuscript. This work was funded by the
- 570 British Medical Research Council, Grant (Grant G0701279), the Wellcome Trust (Grant
- 571 109075/Z/15/A) and the Cambridge NIHR BRC Cell Phenotyping Hub. The funders had no role in
- 572 study design, data collection and interpretation, or the decision to submit the work for publication.

573 Declaration of Interests

574 The authors declare no competing interests.

575 Ethics statement

- 576 All human samples were obtained under ethical approval and after approval of protocols from the
- 577 Cambridgeshire 2 Research Ethics Committee (REC reference 97/092) and all protocols were
- 578 conducted in accordance with the Declaration of Helsinki. Informed written consent was obtained
- 579 from all of the volunteers included in this study prior to providing blood samples and all experiments
- 580 were carried out in accordance with the approved guidelines.

581 Author contributions

- 582 E.G.E., B.K., J.W., E.Y.L, G.X.S., P.J.L., and J.S. designed experiments. E.G.E., B.K., J.W., E.Y.L., E.P.,
- 583 G.X.S., and J.S. performed experiments and data analysis. E.G.E. and J.W. wrote the manuscript.
- 584 C.M.O., M.W., P.J.L, J.S. supervised the research. E.G.E., B.K., J.W., E.Y.L, M.W., C.M.O., P.J.L., and J.S.
- 585 edited the manuscript.

586

587 References

588 1. Hahn G, Jores R, Mocarski ES. Cytomegalovirus remains latent in a common precursor of

- dendritic and myeloid cells. *Proc Natl Acad Sci U S A*. 1998;95(7):3937-3942.
- 590 http://www.ncbi.nlm.nih.gov/pubmed/9520471. Accessed December 3, 2018.
- 591 2. Kondo K, Xu J, Mocarski ES. Human cytomegalovirus latent gene expression in granulocyte-
- 592 macrophage progenitors in culture and in seropositive individuals. *Proc Natl Acad Sci U S A*.
- 593 1996;93(20):11137-11142. http://www.ncbi.nlm.nih.gov/pubmed/8855322. Accessed August
- 594 11, 2017.
- 595 3. Taylor-Wiedeman J, Sissons JGP, Borysiewicz LK, Sinclair JH. Monocytes are a major site of
- 596 persistence of human cytomegalovirus in peripheral blood mononuclear cells. *J Gen Virol*.
- 597 1991;72(9):2059-2064. doi:10.1099/0022-1317-72-9-2059
- 598 4. Mendelson M, Monard S, Sissons P, Sinclair J. Detection of endogenous human
- 599 cytomegalovirus in CD34+ bone marrow progenitors. *J Gen Virol*. 1996;77(12):3099-3102.
 600 doi:10.1099/0022-1317-77-12-3099
- 5. Reeves MB, Sinclair JH. Circulating Dendritic Cells Isolated from Healthy Seropositive Donors
- 602 Are Sites of Human Cytomegalovirus Reactivation In Vivo. J Virol. 2013;87(19):10660-10667.
- 603 doi:10.1128/JVI.01539-13
- 604 6. Poole E, Juss JK, Krishna B, Herre J, Chilvers ER, Sinclair J. Alveolar Macrophages Isolated
- 605 Directly From Human Cytomegalovirus (HCMV)-Seropositive Individuals Are Sites of HCMV
- 606 Reactivation In Vivo. *J Infect Dis*. 2015;211(12):1936-1942. doi:10.1093/infdis/jiu837
- 607 7. Reeves MB, Breidenstein A, Compton T. Human cytomegalovirus activation of ERK and
- 608 myeloid cell leukemia-1 protein correlates with survival of latently infected cells. *Proc Natl*
- 609 Acad Sci U S A. 2012;109(2):588-593. doi:10.1073/pnas.1114966108
- 8. Reeves MB, Lehner PJ, Sissons JGP, Sinclair JH. An in vitro model for the regulation of human
- 611 cytomegalovirus latency and reactivation in dendritic cells by chromatin remodelling. J Gen
- 612 *Virol*. 2005;86(11):2949-2954. doi:10.1099/vir.0.81161-0

- 613 9. Jackson SE, Mason GM, Wills MR. Human cytomegalovirus immunity and immune evasion.
- 614 *Virus Res.* 2011;157(2):151-160. doi:10.1016/j.virusres.2010.10.031
- 615 10. Chan ST, Logan AC. The clinical impact of cytomegalovirus infection following allogeneic
- 616 hematopoietic cell transplantation: Why the quest for meaningful prophylaxis still matters.
- 617 Blood Rev. 2017;31(3):173-183. doi:10.1016/j.blre.2017.01.002
- 618 11. Carbone J. The Immunology of Posttransplant CMV Infection. *Transplantation*. 2016;100:S11-
- 619 S18. doi:10.1097/TP.00000000000001095
- 620 12. Reeves MB, MacAry PA, Lehner PJ, Sissons JGP, Sinclair JH. Latency, chromatin remodeling,
- 621 and reactivation of human cytomegalovirus in the dendritic cells of healthy carriers. *Proc Natl*
- 622 *Acad Sci*. 2005;102(11):4140-4145. doi:10.1073/pnas.0408994102
- 623 13. Goodrum FD, Jordan CT, High K, Shenk T. Human cytomegalovirus gene expression during
- 624 infection of primary hematopoietic progenitor cells: a model for latency. Proc Natl Acad Sci U

625 S A. 2002;99(25):16255-16260. doi:10.1073/pnas.252630899

- 626 14. Söderberg-Nauclér C, Fish KN, Nelson JA. Reactivation of Latent Human Cytomegalovirus by
- 627 Allogeneic Stimulation of Blood Cells from Healthy Donors. *Cell*. 1997;91(1):119-126.
- 628 doi:10.1016/S0092-8674(01)80014-3
- 15. Stinski MF, Isomura H. Role of the cytomegalovirus major immediate early enhancer in acute

630 infection and reactivation from latency. *Med Microbiol Immunol*. 2008;197(2):223-231.

631 doi:10.1007/s00430-007-0069-7

- 632 16. Nelson JA, Groudine M. Transcriptional regulation of the human cytomegalovirus major
- 633 immediate-early gene is associated with induction of DNase I-hypersensitive sites. *Mol Cell*
- 634 Biol. 1986;6(2):452-461. http://www.ncbi.nlm.nih.gov/pubmed/3023848. Accessed January
- 635 28*,* 2019.
- 636 17. Boshart M, Weber F, Jahn G, Dorsch-Häsler K, Fleckenstein B, Schaffner W. A very strong

- 637 enhancer is located upstream of an immediate early gene of human cytomegalovirus. *Cell*.
- 638 1985;41(2):521-530. http://www.ncbi.nlm.nih.gov/pubmed/2985280. Accessed January 28,
- 639 2019.
- 18. Elder E, Sinclair J. HCMV latency: what regulates the regulators? *Med Microbiol Immunol*.
- 641 February 2019. doi:10.1007/s00430-019-00581-1
- 642 19. Humby MS, O'Connor CM. Human Cytomegalovirus US28 Is Important for Latent Infection of
- 643 Hematopoietic Progenitor Cells. Longnecker RM, ed. *J Virol*. 2016;90(6):2959-2970.
- 644 doi:10.1128/JVI.02507-15
- 645 20. Beisser PS, Laurent L, Virelizier J-LL, Michelson S. Human cytomegalovirus chemokine
- 646 receptor gene US28 is transcribed in latently infected THP-1 monocytes. *J Virol*.
- 647 2001;75(13):5949-5957. doi:10.1128/JVI.75.13.5949-5957.2001
- 648 21. Krishna BA, Poole EL, Jackson SE, Smit MJ, Wills MR, Sinclair JH. Latency-associated
- 649 expression of human cytomegalovirus US28 attenuates cell signaling pathways to maintain
- 650 latent infection. Imperiale MJ, ed. *MBio*. 2017;8(6):e01754-17. doi:10.1128/mBio.01754-17
- 651 22. Hargett D, Shenk TE. Experimental human cytomegalovirus latency in CD14+ monocytes. *Proc*
- 652 Natl Acad Sci U S A. 2010;107(46):20039-20044. doi:10.1073/pnas.1014509107
- 23. Zhu D, Pan C, Sheng J, et al. Human cytomegalovirus reprogrammes haematopoietic
- 654 progenitor cells into immunosuppressive monocytes to achieve latency. *Nat Microbiol*.
- 655 2018;3(4):503-513. doi:10.1038/s41564-018-0131-9
- Krishna BA, Spiess K, Poole EL, et al. Targeting the latent cytomegalovirus reservoir with an
 antiviral fusion toxin protein. *Nat Commun*. 2017;8:14321. doi:10.1038/ncomms14321
- 658 25. Krishna BA, Humby MS, Miller WE, O'Connor CM. Human cytomegalovirus G protein-coupled
- 659 receptor US28 promotes latency by attenuating c-fos. *Proc Natl Acad Sci*. January
- 660 2019:201816933. doi:10.1073/pnas.1816933116

- 661 26. Krishna BA, Miller WE, O'Connor CM. US28: HCMV's Swiss Army Knife. Viruses. 2018;10(8).
- 662 doi:10.3390/v10080445
- 663 27. Orzalli MH, DeLuca NA, Knipe DM. Nuclear IFI16 induction of IRF-3 signaling during
- 664 herpesviral infection and degradation of IFI16 by the viral ICPO protein. *Proc Natl Acad Sci U S*
- 665 A. 2012;109(44):E3008-17. doi:10.1073/pnas.1211302109
- 666 28. Dutta D, Dutta S, Veettil MV, et al. BRCA1 Regulates IFI16 Mediated Nuclear Innate Sensing of
- 667 Herpes Viral DNA and Subsequent Induction of the Innate Inflammasome and Interferon-β
- 668 Responses. Feng P, ed. *PLOS Pathog*. 2015;11(6):e1005030.
- 669 doi:10.1371/journal.ppat.1005030
- 670 29. Kerur N, Veettil MV, Sharma-Walia N, et al. IFI16 Acts as a Nuclear Pathogen Sensor to Induce
- the Inflammasome in Response to Kaposi Sarcoma-Associated Herpesvirus Infection. *Cell Host Microbe*. 2011;9(5):363-375. doi:10.1016/j.chom.2011.04.008
- 673 30. Li T, Chen J, Cristea IM. Human cytomegalovirus tegument protein pUL83 inhibits IFI16-
- 674 mediated DNA sensing for immune evasion. *Cell Host Microbe*. 2013;14(5):591-599.
- 675 doi:10.1016/j.chom.2013.10.007
- 676 31. Alandijany T, Roberts APE, Conn KL, et al. Distinct temporal roles for the promyelocytic
- 677 leukaemia (PML) protein in the sequential regulation of intracellular host immunity to HSV-1
- 678 infection. *PLoS Pathog*. 2018;14(1):e1006769. doi:10.1371/journal.ppat.1006769
- 679 32. Cristea IM, Moorman NJ, Terhune SS, et al. Human Cytomegalovirus pUL83 Stimulates
- 680 Activity of the Viral Immediate-Early Promoter through Its Interaction with the Cellular IFI16
- 681 Protein. J Virol. 2010;84(15):7803-7814. doi:10.1128/JVI.00139-10
- 682 33. Diner BA, Lum KK, Toettcher JE, Cristea IM. Viral DNA sensors IFI16 and cyclic GMP-AMP
- 683 synthase possess distinct functions in regulating viral gene expression, immune defenses, and
- 684 apoptotic responses during herpesvirus infection. *MBio*. 2016;7(6):e01553-16.

685 doi:10.1128/mBio.01553-16

- 686 34. Gariano GR, Dell'Oste V, Bronzini M, et al. The Intracellular DNA Sensor IFI16 Gene Acts as
- 687 Restriction Factor for Human Cytomegalovirus Replication. Damania B, ed. *PLoS Pathog*.
- 688 2012;8(1):e1002498. doi:10.1371/journal.ppat.1002498
- 689 35. Diner BA, Lum KK, Javitt A, Cristea IM. Interactions of the Antiviral Factor Interferon Gamma-
- 690 Inducible Protein 16 (IFI16) Mediate Immune Signaling and Herpes Simplex Virus-1
- 691 Immunosuppression. *Mol Cell Proteomics*. 2015;14(9):2341-2356.
- 692 doi:10.1074/mcp.M114.047068
- 693 36. Johnstone RW, Kerry JA, Trapani JA. The human interferon-inducible protein, IFI 16, is a
- 694 repressor of transcription. *J Biol Chem*. 1998;273(27):17172-17177.
- 695 doi:10.1074/jbc.273.27.17172
- 696 37. Biolatti M, Dell'Oste V, Pautasso S, et al. Regulatory Interaction between the Cellular
- 697 Restriction Factor IFI16 and Viral pp65 (pUL83) Modulates Viral Gene Expression and IFI16
- 698 Protein Stability. Frueh K, ed. J Virol. 2016;90(18):8238-8250. doi:10.1128/JVI.00923-16
- 699 38. Roy A, Dutta D, Iqbal J, et al. Nuclear Innate Immune DNA Sensor IFI16 Is Degraded during
- 700 Lytic Reactivation of Kaposi's Sarcoma-Associated Herpesvirus (KSHV): Role of IFI16 in
- 701 Maintenance of KSHV Latency. J Virol. 2016;90(19):8822-8841. doi:10.1128/JVI.01003-16
- 702 39. Pisano G, Roy A, Ahmed Ansari M, Kumar B, Chikoti L, Chandran B. Interferon-γ-inducible
- 703 protein 16 (IFI16) is required for the maintenance of Epstein-Barr virus latency. *Virol J*.
- 704 2017;14(1):221. doi:10.1186/s12985-017-0891-5
- 40. Hotter D, Bosso M, Jønsson KL, et al. IFI16 Targets the Transcription Factor Sp1 to Suppress
- HIV-1 Transcription and Latency Reactivation. *Cell Host Microbe*. 2019;25(6):858-872.e13.
 doi:10.1016/J.CHOM.2019.05.002
- 708 41. Cheung AKL, Gottlieb DJ, Plachter B, et al. The role of the human cytomegalovirus UL111A

709		gene in down-regulating CD4+ T-cell recognition of latently infected cells: implications for
710		virus elimination during latency. <i>Blood</i> . 2009;114(19):4128-4137. doi:10.1182/blood-2008-
711		12-197111
712	42.	Van Damme E, Sauviller S, Lau B, et al. Glucocorticosteroids trigger reactivation of human
713		cytomegalovirus from latently infected myeloid cells and increase the risk for HCMV infection
714		in D+R+ liver transplant patients. <i>J Gen Virol</i> . 2015;96(Pt_1):131-143.
715		doi:10.1099/vir.0.069872-0
716	43.	Poole E, Reeves M, Sinclair JH. The Use of Primary Human Cells (Fibroblasts, Monocytes, and
717		Others) to Assess Human Cytomegalovirus Function. In: Methods in Molecular Biology
718		(Clifton, N.J.). Vol 1119. Humana Press, Totowa, NJ; 2014:81-98. doi:10.1007/978-1-62703-
719		788-4_6
720	44.	Timms RT, Duncan LM, Tchasovnikarova IA, et al. Haploid genetic screens identify an essentia
721		role for PLP2 in the downregulation of novel plasma membrane targets by viral E3 ubiquitin
722		ligases. PLoS Pathog. 2013;9(11):e1003772. doi:10.1371/journal.ppat.1003772
723	45.	van den Boomen DJH, Timms RT, Grice GL, et al. TMEM129 is a Derlin-1 associated ERAD E3
724		ligase essential for virus-induced degradation of MHC-I. Proc Natl Acad Sci U S A.
725		2014;111(31):11425-11430. doi:10.1073/pnas.1409099111
726	46.	Groom HCT, Boucherit VC, Makinson K, et al. Absence of xenotropic murine leukaemia virus-
727		related virus in UK patients with chronic fatigue syndrome. <i>Retrovirology</i> . 2010;7(1):10.
728		doi:10.1186/1742-4690-7-10
729	47.	Spector DJ, Yetming K. UL84-independent replication of human cytomegalovirus strain
730		TB40/E. Virology. 2010;407(2):171-177. doi:10.1016/J.VIROL.2010.08.029
731	48.	Poole E, Huang C, Forbester J, et al. An iPSC derived myeloid lineage model of herpes virus
732		latency and reactivation. Front Microbiol. 2019;10:2233. doi:10.3389/FMICB.2019.02233

733	49.	Weekes MP, Tan SYL, Poole E, et al. Latency-associated degradation of the MRP1 drug
734		transporter during latent human cytomegalovirus infection. Science. 2013;340(6129):199-
735		202. doi:10.1126/science.1235047
736	50.	Straschewski S, Warmer M, Frascaroli G, Hohenberg H, Mertens T, Winkler M. Human
737		cytomegaloviruses expressing yellow fluorescent fusion proteinscharacterization and use in
738		antiviral screening. PLoS One. 2010;5(2):e9174. doi:10.1371/journal.pone.0009174
739	51.	Elder E, Krishna B, Williamson J, et al. Monocytes latently infected with human
740		cytomegalovirus evade neutrophil killing. iScience. January 2019.
741		doi:10.1016/J.ISCI.2019.01.007
742	52.	Kew V, Yuan J, Meier J, Reeves M. Mitogen and Stress Activated Kinases Act Co-operatively
743		with CREB during the Induction of Human Cytomegalovirus Immediate-Early Gene Expression
744		from Latency. Murphy EA, ed. PLoS Pathog. 2014;10(6):e1004195.
745		doi:10.1371/journal.ppat.1004195
746	53.	Slinger E, Maussang D, Schreiber A, et al. HCMV-Encoded Chemokine Receptor US28
747		Mediates Proliferative Signaling Through the IL-6-STAT3 Axis. Sci Signal. 2010;3(133):ra58-
748		ra58. doi:10.1126/scisignal.2001180
749	54.	Madeira F, Park YM, Lee J, et al. The EMBL-EBI search and sequence analysis tools APIs in
750		2019. Nucleic Acids Res. 2019;47(W1):W636-W641. doi:10.1093/nar/gkz268
751	55.	Brown NP, Leroy C, Sander C. MView: a web-compatible database search or multiple
752		alignment viewer. Bioinformatics. 1998;14(4):380-381. doi:10.1093/bioinformatics/14.4.380
753	56.	Rusinova I, Forster S, Yu S, et al. Interferome v2.0: an updated database of annotated
754		interferon-regulated genes. Nucleic Acids Res. 2013;41(Database issue):D1040-6.
755		doi:10.1093/nar/gks1215
756	57.	Rauch I, Müller M, Decker T. The regulation of inflammation by interferons and their STATs.

757 JAK-STAT. 2013;2(1):e23820. doi:10.4161/jkst.23820

- 758 58. Ono SJ, Liou HC, Davidon R, Strominger JL, Glimcher LH. Human X-box-binding protein 1 is
- 759 required for the transcription of a subset of human class II major histocompatibility genes
- and forms a heterodimer with c-fos. *Proc Natl Acad Sci U S A*. 1991;88(10):4309-4312.
- 761 doi:10.1073/pnas.88.10.4309
- 762 59. Ono SJ, Bazil V, Levi BZ, Ozato K, Strominger JL. Transcription of a subset of human class II
- 763 major histocompatibility complex genes is regulated by a nucleoprotein complex that
- contains c-fos or an antigenically related protein. *Proc Natl Acad Sci*. 1991;88(10):4304-4308.
- 765 doi:10.1073/pnas.88.10.4304
- 60. Lee YJ, Benveniste EN. Stat1 alpha expression is involved in IFN-gamma induction of the class

767 II transactivator and class II MHC genes. *J Immunol*. 1996;157(4):1559-1568.

768 http://www.ncbi.nlm.nih.gov/pubmed/8759739. Accessed September 25, 2019.

- 769 61. Clarke CJP, Apostolidis V, Hii LLP, Gough DJ, Trapani JA, Johnstone RW. Critical role of the
- 770 transcription factor AP-1 for the constitutive and interferon-induced expression of IFI 16. J

771 *Cell Biochem*. 2003;89(1):80-93. doi:10.1002/jcb.10475

772 62. Dell'Oste V, Gatti D, Giorgio AG, Gariglio M, Landolfo S, De Andrea M. The interferon-

inducible DNA-sensor protein IFI16: a key player in the antiviral response. *New Microbiol*.

774 2015;38(1):5-20. http://www.ncbi.nlm.nih.gov/pubmed/25742143. Accessed September 25,

- 775 2019.
- 776 63. Kristiansen M, Hughes R, Patel P, Jacques TS, Clark AR, Ham J. Mkp1 Is a c-Jun Target Gene
- 777 That Antagonizes JNK-Dependent Apoptosis in Sympathetic Neurons. J Neurosci.
- 778 2010;30(32):10820-10832. doi:10.1523/JNEUROSCI.2824-10.2010
- 779 64. O'Connor CM, Murphy EA. A Myeloid Progenitor Cell Line Capable of Supporting Human
- 780 Cytomegalovirus Latency and Reactivation, Resulting in Infectious Progeny. J Virol.

- 781 2012;86(18):9854-9865. doi:10.1128/JVI.01278-12
- 782 65. Ansari MA, Singh VV, Dutta S, et al. Constitutive Interferon-Inducible Protein 16-
- 783 Inflammasome Activation during Epstein-Barr Virus Latency I, II, and III in B and Epithelial
- 784 Cells. J Virol. 2013;87(15):8606-8623. doi:10.1128/JVI.00805-13
- 785 66. Caposio P, Gugliesi F, Zannetti C, et al. A novel role of the interferon-inducible protein IFI16 as
- inducer of proinflammatory molecules in endothelial cells. J Biol Chem. 2007;282(46):33515-
- 787 33529. doi:10.1074/jbc.M701846200
- 788 67. Orzalli MH, Conwell SE, Berrios C, DeCaprio JA, Knipe DM. Nuclear interferon-inducible
- 789 protein 16 promotes silencing of herpesviral and transfected DNA. *Proc Natl Acad Sci*.
- 790 2013;110(47):E4492-E4501. doi:10.1073/pnas.1316194110
- 791 68. Thompson MR, Sharma S, Atianand M, et al. Interferon γ-inducible protein (IFI) 16
- 792 transcriptionally regulates type I interferons and other interferon-stimulated genes and
- controls the interferon response to both DNA and RNA viruses. *J Biol Chem*.
- 794 2014;289(34):23568-23581. doi:10.1074/jbc.M114.554147
- 795 69. Johnson KE, Bottero V, Flaherty S, Dutta S, Singh VV, Chandran B. IFI16 Restricts HSV-1
- 796 Replication by Accumulating on the HSV-1 Genome, Repressing HSV-1 Gene Expression, and
- 797 Directly or Indirectly Modulating Histone Modifications. Everett RD, ed. *PLoS Pathog*.
- 798 2014;10(11):e1004503. doi:10.1371/journal.ppat.1004503
- 70. Song LL, Ponomareva L, Shen H, Duan X, Alimirah F, Choubey D. Interferon-Inducible IFI16, a
- 800 Negative Regulator of Cell Growth, Down-Regulates Expression of Human Telomerase
- 801 Reverse Transcriptase (hTERT) Gene. Blagosklonny M V., ed. *PLoS One*. 2010;5(1):e8569.
- 802 doi:10.1371/journal.pone.0008569
- Sponza S, De Andrea M, Mondini M, Gugliesi F, Gariglio M, Landolfo S. Role of the interferoninducible IFI16 gene in the induction of ICAM-1 by TNF-alpha. *Cell Immunol*. 2009;257(1-

- 805 2):55-60. doi:10.1016/j.cellimm.2009.02.007
- 806 72. Sinclair J, Reeves M. The intimate relationship between human cytomegalovirus and the
- dendritic cell lineage. Front Microbiol. 2014;5(AUG):389. doi:10.3389/fmicb.2014.00389
- 808 73. Lau B, Poole E, Van Damme E, et al. Human cytomegalovirus miR-UL112-1 promotes the
- 809 down-regulation of viral immediate early-gene expression during latency to prevent T-cell
- 810 recognition of latently infected cells. J Gen Virol. 2016;97(9):2387-2398.
- 811 doi:10.1099/jgv.0.000546
- 812 74. Lee J-H, Kalejta RF. Human Cytomegalovirus Enters the Primary CD34+ Hematopoietic
- 813 Progenitor Cells where it Establishes Latency by Macropinocytosis. *J Virol*. May 2019.
- 814 doi:10.1128/JVI.00452-19
- 815 75. Dunphy G, Flannery SM, Almine JF, et al. Non-canonical Activation of the DNA Sensing
- Adaptor STING by ATM and IFI16 Mediates NF-κB Signaling after Nuclear DNA Damage. *Mol*
- 817 *Cell*. 2018;71(5):745-760.e5. doi:10.1016/j.molcel.2018.07.034
- 818 76. Noriega VM, Gardner TJ, Redmann V, Bongers G, Lira SA, Tortorella D. Human
- 819 cytomegalovirus US28 facilitates cell-to-cell viral dissemination. Viruses. 2014;6(3):1202-
- 820 1218. doi:10.3390/v6031202
- 821 77. Vieira J, Schall TJ, Corey L, Geballe AP. Functional analysis of the human cytomegalovirus
- 822 US28 gene by insertion mutagenesis with the green fluorescent protein gene. *J Virol*.
- 823 1998;72(10):8158-8165. http://www.ncbi.nlm.nih.gov/pubmed/9733857. Accessed August
- 824 16, 2017.
- 78. Zhang G, Zhang H, Liu Y, et al. CD44 clustering is involved in monocyte differentiation. *Acta Biochim Biophys Sin (Shanghai)*. 2014;46(7):540-547. doi:10.1093/abbs/gmu042
- 827 79. Schmid MA, Kingston D, Boddupalli S, Manz MG. Instructive cytokine signals in dendritic cell
- 828 lineage commitment. *Immunol Rev.* 2010;234(1):32-44. doi:10.1111/j.0105-

829 2896.2009.00877.x

- 830 80. Watowich SS, Liu Y-J. Mechanisms regulating dendritic cell specification and development.
- 831 Immunol Rev. 2010;238(1):76-92. doi:10.1111/j.1600-065X.2010.00949.x
- 832 81. Garrett S, Dietzmann-Maurer K, Song L, Sullivan KE. Polarization of primary human
- 833 monocytes by IFN-gamma induces chromatin changes and recruits RNA Pol II to the TNF-
- alpha promoter. *J Immunol*. 2008;180(8):5257-5266.
- http://www.ncbi.nlm.nih.gov/pubmed/18390706. Accessed April 16, 2019.
- 836 82. Chistiakov DA, Myasoedova VA, Revin V V., Orekhov AN, Bobryshev Y V. The impact of
- interferon-regulatory factors to macrophage differentiation and polarization into M1 and M2.
- 838 *Immunobiology*. 2018;223(1):101-111. doi:10.1016/j.imbio.2017.10.005
- 839 83. Slobedman B, Mocarski ES, Arvin AM, Mellins ED, Abendroth A. *Latent Cytomegalovirus*
- 840 Down-Regulates Major Histocompatibility Complex Class II Expression on Myeloid Progenitors.
- 841 Vol 100.; 2002:2867-2873. doi:10.1182/blood.V100.8.2867
- 842 84. Jenkins C, Garcia W, Godwin MJ, et al. Immunomodulatory Properties of a Viral Homolog of
- 843 Human Interleukin-10 Expressed by Human Cytomegalovirus during the Latent Phase of

844 Infection. J Virol. 2008;82(7):3736-3750. doi:10.1128/JVI.02173-07

- 845 85. Fotouhi-Ardakani N, Kebir D El, Pierre-Charles N, et al. Role for Myeloid Nuclear
- 846 Differentiation Antigen in the Regulation of Neutrophil Apoptosis during Sepsis. Am J Respir
- 847 *Crit Care Med*. 2010;182(3):341-350. doi:10.1164/rccm.201001-00750C
- 848 86. Suzuki T, Nakano-Ikegaya M, Yabukami-Okuda H, et al. Reconstruction of monocyte
- transcriptional regulatory network accompanies monocytic functions in human fibroblasts.
- 850 Bonecchi R, ed. *PLoS One*. 2012;7(3):e33474. doi:10.1371/journal.pone.0033474
- 851 87. Orzalli MH, Broekema NM, Knipe DM. Relative Contributions of Herpes Simplex Virus 1 ICP0
- and vhs to Loss of Cellular IFI16 Vary in Different Human Cell Types. J Virol. 2016;90(18):8351-

853 8359. doi:10.1128/JVI.00939-16

854

37

856 Figure legends

857 Figure 1. US28 induces changes in the host proteome of THP-1 cells

858 A, B, C) THP-1 cells expressing empty vector, US28-WT, and US28-R129A were subject to total cell proteomic analysis using a TMT labelling approach as described in Materials and Methods. Each dot 859 represents one human protein and is shown in grey if its abundance changes by a factor of less than 860 861 2, in red if between 2- and 4-fold, and in purple if greater than 4-fold. The exception is components 862 of the HLA-DR complex which are represented by pink triangles. The dotted line represents a significance threshold of q = 0.001; q < 0.001 is considered significant. Q values of significance 863 864 between groups were calculated by Benjamini-Hochberg correction of p values generated using the 865 moderated T-test LIMMA within the R environment. Comparison of A) US28-WT and US28-R129A, 866 B) US28-WT and empty vector, and C) US28-R129A and empty vector. In each case, the relative 867 abundance of human proteins MNDA and IFI16 is marked with an arrow. D) Analysis of the top 40-868 downregulated proteins identified in (A). After filtering for changes with a q <0.001, the gene names 869 were entered in the Interferome database. Proteins that are induced by Type I and/or Type II interferon are depicted in the Venn diagram. Proteins we identified that are not interferon-inducible 870 are listed below ('Neither'). E) Lysates from THP-1 cells expressing Empty Vector (EV), US28-WT and 871 872 US28-R129A were assessed by western blot for phospho-STAT1 (Tyr701), total STAT1, and beta-actin 873 (loading control). F and G) Quantification of STAT1 and phospho-STAT1 band intensity from two western blots from two independent samples of transduced THP-1 cells. F) Quantification of the 874 875 indicated protein levels normalised to actin. G) Quantification of phospho-STAT1 levels relative to 876 total STAT1.

877 Figure 2. US28-expressing cell lines downregulate IFI16, MNDA, and HLA-DR

A) Relative RNA expression of IFI16, MNDA, and HLA-DR in US28-expressing THP-1 cells. Levels of

879 IFI16, MNDA, HLA-DRA were normalised to TBP and then to US28-R129A using the $\Delta\Delta C_t$ method. B

and C) Lysates from US28-WT and US28-R129A expressing cells were analysed by western blot for

881 IFI16 (B) and MNDA (C) expression; actin is shown as a loading control. D) Empty vector (EV), US28-882 WT, and US28-R129A expressing cells were treated with 5 μ M ruxolitinib (+), or an equivalent 883 concentration of DMSO (-), for 48 hours, before analysis for IFI16 expression by western blot, using actin as a loading control. E) As D), except cells were treated with T-5224 at 10 μ M, DMSO, or left 884 885 untreated. F) US28-expressing cells were maintained in culture media only (left panel) or treated 886 with 1 ng/mL of IFN-γ (right panel) for 24 hours before staining for cell-surface HLA-DR by flow-887 cytometry. Staining was performed in triplicate for untreated cells and the mean of these is 888 experiments is presented in G) as mean fluorescence intensity with standard deviation. Statistical analysis was performed by one-way ANOVA; *** P<0.001. H) As D and E), except cells were analysed 889 890 for cell-surface HLA-DR by flow-cytometry and results are presented as mean fluorescence intensity 891 with 95% confidence intervals. Statistical analysis was performed by two-way ANOVA using Boniferri's multiple comparison test; ns P>0.01, **** P<0.0001. 892 Figure 3. IFI16, MNDA, and HLA-DR are downregulated in latently infected CD14⁺ 893

894 monocytes.

895 Primary CD14⁺ monocytes were isolated from peripheral blood or apheresis cones as described in 896 Materials and Methods. These cells were then infected using BAC-derived strains of TB40/E. A) 897 CD14⁺ monocytes latently infected with TB40/E SV40-mCherry IE2-2A-GFP stained by 898 immunofluorescence for IFI16, MNDA, or HLA-DR as indicated at four d.p.i and imaged by widefield 899 fluorescence microscopy. Top left image: Uninfected monocytes. Second from the left: Monocytes 900 were treated +PMA (to permit lytic infection). mCherry (red) serves as a marker for infection and 901 GFP (green) denotes expression from the IE2-2A-GFP cassette. Remaining panels: Monocytes were 902 cultured in the absence of PMA. The absence of green fluorescence results from suppressed 903 expression of the IE2-2A-GFP cassette and scored as IE negative. The magnification is indicated (40X 904 or 20X). White arrows indicate corresponding cells in the upper and lower panels. B) Validation of 905 experimental latency using TB40/Eqfp virus. CD14⁺ Monocytes were infected and allowed to

39

906 establish latency for 4 days (left panel, 10X magnification). Citrate wash buffer was used to remove 907 externally bound virions. These latently infected cells were cultured -/+PMA for 3 days, and at 7 908 d.p.i, Hff-1 cells were added to the culture to demonstrate production of infectious virions. Transfer 909 of virus to Hff-1 was monitored by fluorescence microscopy up to 13 d.p.i., and infected Hff-1 foci 910 were counted and summed across the experiment (three wells of CD14⁺ monocytes per condition, 911 graphed). C) CD14⁺ monocytes infected with TB40/Egfp stained by immunofluorescence for IFI16 at 912 24, 48, and 72 h.p.i. and imaged as before using 60X magnification. D) Quantification of IFI16 913 positive and negative monocytes in the uninfected and infected populations from two donors per 914 time point. Raw numbers of cells are indicated in white text. Fisher's exact test indicates a statistically significant difference between uninfected and infected populations for each time point 915 916 (P<0.0001). E) CD14⁺ monocytes infected with TB40/Egfp were stained by immunofluorescence for 917 MNDA at the indicated times and imaged as before using 60X magnification. F) Quantification of the 918 signal intensity from infected monocytes at the indicated time points (n=9,7,10, respectively). MNDA 919 signal intensity in each nucleus was normalised to the average of uninfected monocytes from each 920 field of view. A t-test with Welch's correction was used to determine statistical significance. ns, not significant, *P<0.05, **P<0.01 G) CD14⁺ monocytes infected with TB40/Eqfp (+/- UV inactivation) 921 922 were analysed for HLA-ABC and HLA-DR expression at three d.p.i. by flow cytometry. The gating 923 strategy for identifying infected cells (GFP⁺) is shown. H) Histogram showing HLA-ABC and HLA-DR 924 staining in HCMV-uninfected GFP-negative (grey) monocytes, and latently infected GFP positive 925 (green) monocytes.

926 Figure 4: IFI16 is rapidly downregulated in a US28-dependent manner during latent927 infection

CD14⁺ monocytes were infected with either US28 WT TB40/EmCherry-US28-3XFLAG HCMV or the
 ΔUS28. A) Validation of the latent and lytic phenotypes associated with US28-3xF and ΔUS28
 monocyte infections, respectively. At 7 d.p.i., supernatant from infected CD14⁺ cells (upper panel)

931	were transferred to Hff1 cells (middle brightfield and lower mCherry panels) and formation of
932	plaques was monitored and imaged at 20X magnification. B) Detection of US28-3XFLAG during the
933	establishment of latency in CD14 $^{+}$ monocytes. At 2 d.p.i. US28-3xF or Δ US28-infected CD14 $^{+}$
934	monocytes were fixed and stained by immunofluorescence for US28-3XFLAG using an anti-FLAG
935	antibody and imaged at 40X magnification. C) US28-3xF and Δ US28-infected monocytes were
936	stained by immunofluorescence for IFI16 at the indicated times and imaged using 40X magnification.
937	White arrows indicate corresponding cells. D and E) IFI16 signal intensity in each nucleus was
938	normalised to the average of the uninfected cells in a field of view. The results of three fields of view
939	were then averaged to derive the resulting average signal intensities for each subpopulation of
940	monocytes at the indicated time points infected with US28-3xF or Δ US28 HCMV. Statistical
941	significance was determined using one-way ANOVA. *** indicates P<0.001, ** indicates P<0.01, and
942	* indicates P<0.05.
943	Figure 5: IFI16 downregulation is maintained during long term latency of
944	undifferentiated monocytes and CD34 ⁺ progenitor cells.
945	A) US28 expressing and empty vector THP-1 cells were either left untreated or treated with PMA for
946	48 hours before cell lysates were harvested. These lysates were then subject to western blotting for
947	IFI16 and actin as a loading control, with molecular weight markers annotated. B) At 48 h.p.i, either
948	undifferentiated CD14 $^{\scriptscriptstyle +}$ monocytes, or monocytes pre-differentiated for 7 days with GM-CSF/IL-
949	4/LPS, were fixed and stained for IFI16 and imaged as before at 40X magnification. White arrows
950	indicate corresponding infected cells. C) CD14+ monocytes were infected with HCMV
951	GATA2mCherry, or left uninfected. At the indicated times, cells were fixed and stained for IFI16 or
952	MNDA and imaged as before. D) Primary CD34 * hematopoietic progenitor cells from two donors, or
953	Kasumi-3 cells, were infected infected with HCMV GATA2mCherry, or left uninfected. At the
954	indicated times, cells were fixed and stained for IFI16 and imaged as before. E) Quantification of at
955	least 3 fields of view from D), presented as the proportion of cells with low IFI16 expression in the

41

956 infected, mCherry positive and uninfected, mCherry negative populations. UI – uninfected.

957 Statistical analysis was performed by Fischer's Exact test on the total number of cells in each

958 category. **** indicates P<0.0001, *** indicates P<0.001, ** indicates P<0.01.

959 Figure 6: Overexpression of IFI16 in monocytic cells leads to MIEP activation and IE

960 gene expression via NF-κB

961 A) THP-1 cells were transduced with empty vector (EV) or IFI16-overexpression lentiviruses and 962 following blasticidin selection, IFI16 overexpression was confirmed by western blot. Actin is shown 963 as a loading control. B and C) Empty vector or IFI16-overexpressing cells (from A) were infected with 964 TB40/E IE2-eYFP virus, and IE2-eYFP positive nuclei were imaged and counted by fluorescence 965 microscopy. C) Cumulative results from five paired experiments, which were analysed by paired two-966 tailed Student's t-test; * P<0.05. D) EV and IFI16 lentivirus concentration was determined by p24 967 ELISA (data not shown) and 15 ng p24 equivalents of each lentivirus was used to transduce MIEP-968 eGFP THP-1 cells. Cells were maintained for two weeks in culture, and IFI16 overexpression was 969 validated by immunofluorescence. E) Left hand comparison: cells described in (D) were assessed for 970 eGFP fluorescence by flow cytometry. Right hand comparison: non-transduced MIEP-eGFP 971 expressing cells were incubated with supernatants from cells described in (A) for two days. eGFP 972 expression was quantified by flow cytometry. A statistical comparison of the median fluorescence 973 intensity was performed using two-tailed Mann-Whitney test; ns, not significant and * P<0.05. F) 974 Empty vector or IFI16-overexpressing cells were fixed and stained for NF-κB, with Hoechst as a 975 nuclear stain, at 40X magnification to assess levels of nuclear NF-kB. G) Empty vector or IFI16-976 overexpressing cells were infected with TB40/E IE2-eYFP virus in the presence of the IKKα inhibitor 977 BAY11-7082 which inhibits the NF-κB pathway, or DMSO as a control. IE2-eYFP positive nuclei were 978 imaged and counted by fluorescence microscopy at 48 hours post infection. H) Empty vector or 979 IFI16-overexpressing cells were infected with a revertant WT-like TB40/E at MOI 3, TB40/E with NF-980 κB binding sites deleted from the MIEP (ΔNF-κB) at MOI 3 or MOI 15. At 48 h.p.i., cells were fixed

981	and stained for IE and the number of IE positive nuclei were counted. Graph shows the results of
982	three experiments and statistical analysis by 2-way ANOVA using Sidak's multiple comparison test.
983	** P<0.01, ns, P > 0.05. I) Empty vector or IFI16-overexpressing cells were infected as per F) and H) at
984	MOI 3, but cells were instead analysed for IE72 expression by RT-qPCR. PCR products were then run
985	on a 2% (upper panel, IE72) or 1.2% (lower panel, GAPDH) agarose gel. UI refers to uninfected cells,
986	DMSO is the solvent control, BAY refers to BAY11-7082, Rev refers to the revertant TB40/E and Δ NF-
987	κB to the NF- κB binding site mutant virus. The positive control (+ve ctrl) was HCMV-infected PMA-
988	differentiated monocytes. Molecular weight markers (M) are annotated.
989	
990	Figure S1. US28 expression induces IFN-inducible genes, but not ER stress-related
991	genes
992	A) Changes in interferon inducible genes identified in Figure 1D, and other canonical ISGs, in US28-
993	WT with respect to US28-R129A. Green bars indicate changes with a q value of <0.001. D) Heat map

of the changes in canonical ER stress-related genes induced by US28-WT or US28-R129A expression

- as per the proteomic screens in Figure 1A, B, C. HUGO gene symbols are listed followed by a
- common gene name, if applicable. An outgroup of genes that are regulated by US28 (IFI16, MNDA,
- 997 FLT3) is included for comparison.

998 Figure S2. US28-expressing cell lines downregulate IFI16, MNDA, and HLA-DR

- A) Empty vector, US28-WT and US28-R129A-expressing THP-1 cells were regenerated in
- 1000 independent transductions using the same expression vectors as used for the proteomic screen
- 1001 (Figure 1). US28 expression was validated by RT-qPCR, with US28 RNA normalised to TATA-box
- binding protein (TBP) and presented as $2^{-\Delta Ct}$. B) Cells from A were lysed and subject to western blot
- 1003 for US28, and actin as a loading control. C) Quantification of three western blots for US28
- 1004 expression. C and D) Lysates prepared from cells in (A) were analysed by western blot for IFI16 (C)

and MNDA (D) expression; actin is shown as a loading control. Note that panel E is from the same

1006 membrane as Figure 1C. F) Quantification of 5 and 4 independent western blots for IFI16 and MNDA,

1007 respectively. G) Cells from A) were treated with ruxolitinib as per figure 2D, or left untreated. Lysates

1008 from these cells were analysed by western blot for phosphorylated STAT1, total STAT1, or actin as a

1009 loading control.

1010 Figure S3. Strain-dependent differences in US28 do not affect downregulation of

- 1011 interferon-inducible genes
- 1012 A) Sequences encoding US28 from the indicated HCMV strains or plasmids were aligned using Clustal
- 1013 Omega. B) Retroviral plasmids encoding US28-WT (from TB40/E) or R129A, each with a C-terminal
- 1014 3XFLAG tag, and an eGFP marker, were used to transduce THP-1 cells. They were then subject to
- 1015 immunofluorescence staining for the 3XFLAG tag. C and D) Cells from B were stained for cell-surface
- 1016 HLA-DR by flow cytometry. D) Mean fluorescence intensity of the US28-WT and US28-R129A cell
- 1017 lines. Statistical analysis by Student's t test; ** P<0.01.

1018 Figure S4 Downregulation of IFI16, MNDA, and HLA-DR is not simply a bystander

1019 effect of contact with viral particles.

1020 A) CD14⁺ monocytes were left uninfected, or infected with HCMV for 24 hours before fixing and

- staining for the indicated proteins, and imaging as before. B) The sequence encoding US28 from
- 1022 VHL/E was cloned into the lentiviral plasmid pUbEm (US28-UbEm), and this or empty UbEm plasmid
- 1023 was used to transduce THP-1 cells, which were subsequently cell-sorted for Emerald expression. C)
- 1024 US28 expression was validated in the cells from (B) by RT-qPCR. US28 RNA was normalised to cellular
- 1025 TBP and presented as $2^{-\Delta Ct}$.
- 1026 File S5: US28 proteome in THP-1 cells

Tab 1: Data: THP-1 cells expressing empty vector, US28-WT, and US28-R129A were subject to total
cell proteomic analysis using a TMT labelling approach as described in Materials and Methods. This

1029	file lists all genes identified in this proteomic screen, including their Uniprot Accession number,
1030	HUGO gene symbol, fold changes in abundance between cell lines, and q values of statistical
1031	significance. Tab 2: Interferome Top 40 Downreg. Gene names, fold changes, and q values of the top
1032	40-most downregulated genes (US28-WT vs US28-R129A) are presented, along with whether they
1033	are included in the Interferome database as being Type I or Type II interferon-inducible (marked with
1034	'y'). Tab 3: Interferome Zero-Change 40: Gene names, fold changes, and q values of the genes with a
1035	fold change value of zero (US28-WT vs US28-R129A) are presented, along with whether they are
1036	included in the Interferome database as being Type I or Type II interferon-inducible (marked with
1037	'y').
1038	
1039	File S6: Schematic showing mass spectrometry settings for experiments presented in

- **1040** Figure 1 and File S5
- 1041



C

US28 R129A vs Empty Vector 10-< 2-fold change 2-fold to 4-fold change > 4-fold change Statistical confidence (-log₁₀q value) 8 **HLA-DR** components D 6 Type II Туре CTH GLUL HLA-DRA ALDH2 ALOX5AP MNDA HLA-DRB1-1 CAND2 HLA-DRB1-15 DDX3Y 4 IFITM3 HLA-DRB5 SYNPO2 IFI16 MYO1G ME1 **IFI16** KYNU MARCKS PADI2 PLD4 MNDA OAS2 CKB RPS4Y1 S100P 2 SULT1A1 RNASE2



Fold change abundance (log₂US28-WT/Empty Vector)





















U



















72 h.p.i.













Hff1

Primary CD14⁺ monocytes



24 h.p.i.

48 h.p.i.

bioRxiv preprint doi: https://doi.org/10.1101/791673; this version posted October 4, 2019. The copyright holder for this preprint (w ch was not certified by peer review) is the author/funder. All rights reserved. No reuse allowed without permission.



TB40/EmCherry US28Δ





TB40/EmCherry US28-3XFLAG

TB40/EmCherry US28Δ



Hoechst



mCherry





Merge



TB40/E GATA2 mCherry







Loss of IFI16 expression during latency







Empty vector IFI16





D

4 - St



MIEP-driven GFP expression







Ε



