

Predicting cellular position in the *Drosophila* embryo from Single-Cell Transcriptomics data

Jovan Tanevski*^{1,2}, Thin Nguyen*³, Buu Truong*⁴, Nikos Karaiskos⁵, Mehmet Eren Ahsen^{6,7}, Xinyu Zhang⁸, Chang Shu⁸, Ke Xu⁸, Xiaoyu Liang⁸, Ying Hu⁹, Hoang V.V. Pham⁴, Li Xiaomei⁴, Thuc D. Le⁴, Adi L. Tarca¹⁰, Gaurav Bhatti¹⁰, Roberto Romero¹¹, Nestoras Karathanasis¹², Phillipe Loher¹², Yang Chen¹³, Zhengqing Ouyang¹⁴, Disheng Mao¹⁵, Yuping Zhang¹⁵, Maryam Zand¹⁶, Jianhua Ruan¹⁶, Christoph Hafemeister¹⁷, Peng Qiu^{18,19}, Duc Tran²⁰, Tin Nguyen²⁰, Attila Gabor¹, Thomas Yu²¹, Enrico Glaab²², Roland Krause²³, Peter Banda²³, DREAM SCTC Consortium[†], Gustavo Stolovitzky²⁴, Nikolaus Rajewsky^{‡5}, Julio Saez-Rodriguez^{‡1,25}, and Pablo Meyer^{‡24}

¹Institute for Computational Biomedicine, Faculty of Medicine, Heidelberg University Hospital and Heidelberg University, Heidelberg, Germany

²Department of Knowledge Technologies, Jožef Stefan Institute, Ljubljana, Slovenia

³Deakin University, Geelong, Australia

⁴University of South Australia, Mawson Lakes, Australia

⁵Berlin Institute for Medical Systems Biology, Max Delbrück Center for Molecular Medicine in the Helmholtz Association, Berlin, Germany

⁶Icahn School of Medicine at Mount Sinai, New York City, NY, USA

⁷University of Illinois, Urbana-Champaign, IL, USA

⁸Department of Psychiatry, Yale School of Medicine, New Haven, CT, USA

⁹Center for Biomedical Informatics & Information Technology, National Cancer Institute, MD, USA

¹⁰Wayne State University, Detroit, MI, USA

¹¹Perinatology Research Branch, NICHD/NIH/DHHS, Bethesda, MD, and Detroit, MI, USA

¹²Computational Medicine Center, Thomas Jefferson University, Philadelphia, PA, USA

¹³The Jackson Laboratory for Genomic Medicine, Farmington, CT, USA

¹⁴University of Massachusetts, Amherst, MA, USA

¹⁵University of Connecticut, CT, USA

¹⁶University of Texas at San Antonio, TX, USA

¹⁷New York Genome Center, New York City, NY, USA

¹⁸Georgia Institute of Technology, Atlanta, GA, USA

¹⁹Emory University, Atlanta, GA, USA

²⁰University of Nevada, Reno, NV, USA

²¹Sage Bionetworks, Seattle, WA USA

²²Biomedical Data Science Group, Luxembourg Centre for Systems Biomedicine, University of Luxembourg, Esch-sur Alzette, Luxembourg

²³Bioinformatics Core Group, Luxembourg Centre for Systems Biomedicine, University of Luxembourg, Esch-sur Alzette, Luxembourg

²⁴IBM T.J. Watson Research Center, Yorktown Heights, NY, USA

²⁵Joint Research Centre for Computational Biomedicine, Faculty of Medicine, RWTH Aachen University, Aachen, Germany

*These authors contributed equally

†DREAM SCTC Consortium authors and affiliations are listed in the supplementary material

‡To whom correspondence should be addressed; pmeyerr@us.ibm.com

Abstract

Single-cell RNA-seq (scRNAseq) technologies are rapidly evolving and a growing number of datasets are now available. While very informative, in standard scRNAseq experiments the spatial organization of the cells in the organism or tissue of origin is lost. Conversely, spatial RNA-seq technologies designed to keep the localization of the cells have limited throughput and gene coverage. Mapping scRNAseq to data of genes with spatial information can thus increase coverage while providing spatial location. However, methods to perform such a mapping are still in their infancy and have not been benchmarked in an unbiased manner. To bridge the gap, we organized the DREAM Single-Cell Transcriptomics challenge to evaluate methods for reconstructing the spatial arrangement of single cells from single-cell RNA sequencing data. The challenge focused on the spatial reconstruction of cells from the *Drosophila* embryo from single-cell transcriptomic and, leveraging as gold standard, *in situ* hybridization data of a set of selected driver genes from the Berkeley *Drosophila* Transcription Network Project reference atlas. The 34 participating teams used an array of different algorithms for gene selection and location prediction. We devised a novel scoring and cross-validation scheme to evaluate the robustness of the best performing algorithms. Participants were able to correctly and robustly localize rare subpopulations of cells, accurately mapping both spatially co-localized and scattered groups of cells. The selection of predictor genes was essential for accurately locating the cells in the embryo. Among the most frequently selected set of genes we measured a relatively high expression entropy, high spatial clustering and the presence of prominent developmental genes such as gap and pair-ruled genes and tissue defining markers.

1 Introduction

1 The recent technological advances in single-cell sequencing technologies have revolutionized
2 the biological sciences. In particular single-cell RNA sequencing (scRNAseq) methods allow
3 transcriptome profiling in a highly parallel manner, resulting in the quantification of thousands of
4 genes across thousands of cells of the same tissue. However, with a few exceptions [1, 2, 3, 4, 5]
5 current high-throughput scRNAseq methods share the drawback of losing the information relative
6 to the spatial arrangement of the cells in the tissue during the cell dissociation step.

7 One way of regaining spatial information computationally is to appropriately combine the single-
8 cell RNA dataset at hand with a reference database, or atlas, containing spatial expression patterns
9 for several genes across the tissue. This approach was pursued in a few studies [6, 7, 8, 9, 10].
10 Achim *et al* identified the location of 139 cells using 72 reference genes with spatial information
11 from whole mount *in situ* hybridization (WMISH) of a marine annelid and Satija *et al* developed
12 the *Seurat* algorithm to predict position of 851 zebrafish cells based on their scRNAseq data and
13 spatial information from *in situ*-hybridizations of 47 genes in ZFIN collection [11]. In both cases,
14 cell positional predictions stabilized after the inclusion of 30 reference genes. Karaiskos *et al*
15 reconstructed the early *Drosophila* embryo at single-cell resolution and while the authors were
16 successful in their reconstruction, their approach did not lead to a predictive algorithm and mainly
17 centered around maximizing the correlation between scRNAseq data and the expression patterns
18 from *in situ*-hybridizations of 84 mapped genes in The Berkeley *Drosophila* Transcription Network
19 Project (BDTNP). In this project, *in situ* hybridization data was collected resulting in a quantitative
20 high-resolution gene expression reference atlas [12]. Indeed, Karaiskos *et al* showed that the
21 combinatorial expression of these 84 BDTNP markers suffice to uniquely classify almost every cell
22 to a position within the embryo.

23 In the absence of a reference database, it is also possible to regain spatial information compu-
24 tationally solely from the transcriptomics data by leveraging general knowledge about statistical
25 properties of spatially mapped genes against the statistical properties of the single-cell RNA dataset
26 [13, 14]. Bageritz *et al.* were able to reconstruct the expression map of a *Drosophila* wing disc
27 using scRNAseq data by correlation analysis. They exploited the coexpression of non-mapping

28 genes to a few mapping genes with known expression patterns, to predict the spatial expression
29 patterns of 824 genes [13]. Nitzan *et al.* exploited the knowledge of the distribution of distances
30 between mapping genes in physical space to predict the possible locations of cells based on the
31 distribution of distances between genes in the expression space. Following this approach, they
32 were able to successfully reconstruct the locations of cells of the *Drosophila* and zebrafish embryos
33 from scRNAseq data [14]. Although these approaches have indicated important steps to reconstruct
34 the position of a cell in a tissue from their RNAseq expression, a global assessment is needed
35 to evaluate the methods used and the number and nature of the genes with spatial expression
36 information required for correctly assigning a location to each cell.

37 With this purpose in mind, and to catalyze the development of new methods to predict the
38 location of cells from scRNAseq data we organized the DREAM Single cell transcriptomics
39 challenge which ran from September through November 2018. DREAM challenges are a platform
40 for crowdsourcing collaborative competitions[15] where a rigorous evaluation of each submitted
41 solution allows for the comparison of their performance. The quality and reproducibility of each
42 provided solution is also ensured. The combination of the individual solutions, i.e., the different
43 approaches and insights to a common problem, leads to an overall wisdom-of-the-crowds (WOC)
44 solution, with generally superior performance to any individual solution, from where collective
45 insights can be garnered. We set up the challenge with 3 goals in mind. First, we used the data
46 from Karaiskos *et al* to foster the design of a variety of algorithms and objectively tested how
47 well they could predict the localization of the cells. Second, we evaluated how the predictive
48 performance of the algorithms was impacted by the number of reference genes from BDTNP
49 with *in situ* hybridization information included in the predictions. Third, we investigated how the
50 biological information carried in the selected genes was implemented in the algorithms to determine
51 embryonic patterning.

52 The challenge, a first of its kind for single cell data, consisted of predicting the position of
53 1297 cells among 3039 *Drosophila melanogaster* embryonic locations for one half of a stage 6
54 pre-gastrulation embryo from their scRNAseq data (Figure 1A) [10]. At this stage cells in the
55 embryo are positioned in a single two dimensional sheet following a bilateral symmetry, so that
56 only positions in one half of the embryo were considered - accounting for the 3039 locations.
57 Participants used the scRNAseq data for each of the 1297 cells obtained from the dissociation
58 of 100-200 stage 6 embryos and the spatial expression patterns from *in situ*-hybridizations of 84
59 genes in the BDTNP database [12]. Gene determinants of different tissues such as neurectoderm,
60 dorsal ectoderm, mesoderm, yolk and pole cells were provided as a hint. To aid the development
61 of prediction algorithms, we provided (when available) the regulatory relationship -positive or
62 negative- between the 84 genes in the *in situ*-hybridizations and the rest of the genes. We asked
63 participants to provide an ordered list of 10 most probable locations in the embryo predicted for
64 each of the 1297 cells using the expression patterns from (i) 60 genes out of the 84 in subchallenge
65 1, (ii) 40 genes out of the 84 in subchallenge 2, and (iii) 20 genes out of the 84 in subchallenge
66 3. The predictions were compared to the ground truth location determined by calculating the
67 maximum correlation using all 84 *in situs* [10]. We received submissions from 34 teams, and
68 the overall analysis of the results showed that the selection of genes is essential for accurately
69 locating the cells in the embryo. The most selected genes had a relatively high expression entropy,
70 showed high spatial clustering and featured developmental genes such as gap and pair-ruled genes
71 in addition to tissue defining markers.




Figure 1: Overview of the challenge and results. **A.** In the DREAM Single-Cell Transcriptomics Challenge participants were asked to map the location of 1297 cells to 3039 location bins of an embryo of *Drosophila melanogaster*, by combining the scRNAseq measurements of 8924 genes for each cell and the spatial expression patterns from *in situ* hybridization of 60, 40 or 20 genes, for subchallenge 1, 2 and 3 respectively, for each embryonic location bin, selected from a total of 84 driver-genes. **B.** Ranking of the top 10 best performing teams and a wisdom of the crowds (WOC, in *italic*) solution, based on results from a post challenge cross-validated selection and prediction performance measured with three complementary scoring metrics. The boxplots show the distribution of ranks for each team on the 10 test folds. The rank for each fold is calculated as the average of the ranking on each scoring metric.

72 2 Results

73 2.1 Challenge setup

74 A distinctive feature of the single cell transcriptomics challenge was the public availability of
 75 the entire dataset and the ground truth locations produced by DistMap, a method using the *in*
 76 *situ*-hybridizations available at BDTNP [12], published together with the data [10]. We took three
 77 actions to mitigate the issue of not having a blinded ground truth. First, for the purpose of predictor
 78 gene selection, we allowed the use of scRNA-Seq data and biological information from other
 79 databases, but prohibited the use of *in situ* data. Second, to assess the quality of predictions, we
 80 devised three scores (detailed in the Methods section) that were not disclosed to the participants
 81 during the challenge. The scores measured not only the accuracy of the predicted location, but also
 82 how well the expression in the cell at the predicted location correlates with the expression from the
 83 reference atlas, the variance of the predicted locations for each cell, and how well the gene-wise
 84 spatial patterns were reconstructed. Finally, we devised a post-challenge cross-validation scheme to
 85 evaluate the soundness and robustness of the methods.

86 The challenge was organized in two rounds, a leaderboard round, and a final round. During the
 87 leaderboard round the participants were able to obtain scores for five submitted solutions before
 88 submitting a single solution in the final round. We received submissions from 40 teams in the
 89 leaderboard round and 34 submissions in the final round. Out of the 34 teams that made submissions
 90 in the final round, 29 followed up with public write-ups of their approaches and source code. For

Table 1: Best mean score for metrics s_1 , s_2 and s_3 achieved by the teams (Thin Nguyen, WhatATeam and OmicsEngineering) and the WOC solution. The standard deviation of scores across folds are in parenthesis. For more details on the scoring metrics see the Methods section.

	s_1		s_2		s_3	
	Teams	WOC	Teams	WOC	Teams	WOC
Subchallenge 1	0.76 (±0.04)	0.73(±0.04)	2.52 (±0.28)	2.16(±0.20)	0.59(±0.01)	0.62 (±0.01)
Subchallenge 2	0.69(±0.03)	0.70 (±0.05)	1.16(±0.12)	1.84 (±0.26)	0.67 (±0.02)	0.65(±0.01)
Subchallenge 3	0.65(±0.05)	0.68 (±0.03)	0.88(±0.13)	1.42 (±0.16)	0.79 (±0.02)	0.71(±0.01)

91 subchallenges 1 and 3 we were able to determine a clear best performer, but for subchallenge 2,
92 there were two top ranked teams with statistically indistinguishable difference in performance (see
93 Supplementary Figures S1,S2 and S3).

94 As stated, given that the ground truth for this challenge was publicly available and to avoid
95 over-fitting, we decided to invite the top 10 performing teams to contribute to a post-challenge
96 collaborative analysis phase to assess the soundness and stability of their gene selection and
97 cell location prediction. Consequently, teams were tasked to provide predictions for a 10-fold
98 cross-validation (CV) scenario, under the same conditions as for the challenge phase. The folds
99 were extracted from the same RNA-seq dataset as in the challenge and every team used the same
100 assignment of cells to folds. We evaluated the performance of the teams using the same scoring
101 approach as in the challenge. To ensure the validity of the findings we decided to perform all further
102 analysis and interpretation only from the results of the post-challenge phase.

103 2.2 Overview of results

104 Interestingly, for subchallenge 1 and 2, when participants had to use 60 or 40 genes for their
105 predictions, the ranking of the best performing teams in the CV scenario did not change significantly
106 compared to the challenge (Figure 1B cf. Figures S1 and S2). This was not the case in subchallenge
107 3 as no particular team from the top 10 outperformed in a statistically significant way the others
108 when using 20 genes for their predictions. The results from the cross-validation showed that the
109 approaches generalize well, i.e. the gene selection is performed consistently across the folds and
110 the variance of the achieved scores across the folds is small for all teams (Figure S4). For each
111 subchallenge we combined the gene selection and location predictions from the top 10 participants
112 into a WOC solution (see details below) that performed better compared to the individual solutions
113 (Figure 1B). The scores obtained by the best performing teams and the WOC solution are shown in
114 Table 1.

115 A summary of the methods used by participants for gene selection and location prediction can
116 be seen in Table S2. The most frequently used method by participants for location prediction was a
117 similarity based prediction, such as the maximum Matthews correlation coefficient between the
118 binarized transcriptomics and the *in situ* that was proposed by Karaiskos et al. [10]. Another well
119 performing approach was combining the predictions of a machine learning model and the Matthews
120 correlation coefficients. The models were trained to predict either the coordinates of each cell or
121 the binarized values of the selected *in situ* given transcriptomics data as input. The predictions
122 were then made by selecting the location bins that corresponded to the nearest neighbors of the
123 predicted values.

124 The most frequently used method by participants for gene selection was unsupervised or
125 supervised feature importance estimation and ranking. For example, in a supervised feature
126 importance estimation approach a machine learning model is trained to predict the coordinates of
127 each cell, given the transcriptomics data at input, that is, the genes with available *in situ* hybridization
128 measurements or all genes. Different machine learning models were trained such as Random Forest




Figure 2: **Wisdom of crowds location prediction.** The location predictions for each cell by the top performing teams in the post-challenge cross-validation phase were aggregated in a wisdom of the crowds solution based on a k-means clustering approach.

129 (BCBU, OmicsEngineering) or a neural network (DeepCMC, NAD). There were examples of
130 unsupervised feature importance estimation and ranking by expression based clustering (NAD,
131 Christoph Hafemeister, MLB), or a greedy feature selection based on predictability of expression
132 from other genes (WhatATeam). Background knowledge about location specific marker genes, or
133 the expected number of location clusters, was used by a small number of teams (WhatATeam and
134 NAD) to inform the gene selection. Given the diversity of approaches to gene selection, we focused
135 our analysis on better understanding the properties of frequently selected genes and providing
136 recommendations for future experimental designs.

137 **2.3 Analysis of the location prediction**

138 A recurrent observation across DREAM challenges is that an ensemble of individual predictions
139 performs usually better and is more robust than any individual method [16, 17]. This phenomenon,
140 common also in other contexts, is denoted as the wisdom-of-the-crowds (WOC) [15]. In a typical
141 challenge, individual methods output a single probability reflecting the likelihood of occurrence of
142 an event. The WOC prediction is then constructed in an unsupervised manner by averaging the
143 predictions of individual methods.

144 Given that in the single cell RNAseq prediction challenge participants had to submit 10 positions
145 per cell, we developed a novel method that is based on k-means clustering to generate the WOC
146 predictions. A diagram of the k-means approach is given in Figure 2 where for each single cell
147 we first used k-means clustering to cluster the locations predicted by the individual teams [18]
148 where the euclidean distance between the locations was used as the distance metric. In order to
149 find the optimal k , we used the elbow method, i.e. we chose a k that saturates the sum of squares
150 between clusters [19]. Note that each cluster consists of a group of locations and each location
151 is predicted by one or more teams. Hence, for each cluster we calculated the average frequency
152 that its constituent locations are predicted by individual teams. We then picked the cluster with
153 the highest average frequency as our final cluster and ranked each location in this cluster based on
154 how frequently it was predicted by individual methods. For each cell, the final prediction of the
155 proposed WOC method consisted of the top 10 locations based on the above ranking. The k-means
156 approach is based on the intuition that a single cell belongs to one location and its expression is
157 mostly similar to that of cells in locations surrounding it.

158 The WOC location prediction approach does not take the genes used by the teams to make the
159 predictions into account. However, after the WOC predictions are generated, in order to score them,
160 we needed a list of genes for every subchallenge. To this end we used a WOC approach to gene
161 selection (see the following section for more details) and used the most frequently selected genes
162 per challenge. As reported above, the WOC solution performed better compared to the individual
163 solutions (Figure 1B).

164 **2.4 Analysis of selected genes**

165 The selection of a subset of *in situs* used for cell location prediction was the hallmark that differen-
166 tiated the subchallenges. It is unfeasible to evaluate all subsets of 20, 40 or 60 genes from the 84
167 due to the immense number of possible combinations of genes. Different approaches and heuristics
168 can be used to select a subset of genes and the most frequent among the top 10 ranked teams were
169 based on model based feature ranking algorithms, using normalized transcriptomics data (for more
170 details see Table S2). However, if a subset of genes is selected as a candidate for solving the general
171 task of location prediction, it should be consistently identified when similar sets of single cells are
172 used as inputs. Therefore, we analyzed the consistency of gene selection for each team across folds
173 by 10-fold cross-validation. More importantly, we were interested in subsets of genes that were
174 consistently selected by multiple teams as this could underlie biological relevance.

175 The approaches for selecting genes taken by the top 10 teams resulted in consistent selection
176 across folds, significantly better than random, for all subchallenges. Indeed, all of the pairwise
177 Jaccard similarities of sets of selected genes for all teams were significantly higher than the expected
178 Jaccard similarity of a random pair of subset of genes (see Supplementary Figure S4). Importantly,
179 we measured an observable increase in variance and decrease of mean similarity as the number of
180 selected genes decreased.

181 For each subchallenge we counted the number of times that the genes were selected by all teams
182 in all folds. The genes, ordered by the frequency of selection in all subchallenges are depicted in
183 Figure 3A. Forty percent of the top 20, 67% of the top 40 and 81% of the top 60 most frequently
184 selected genes are the same for all three subchallenges (Figure 3B). The ranks assigned to all genes
185 in the three subchallenges are highly correlated. Namely, the rank correlations range from 0.69
186 between subchallenges 1 and 3, to 0.83 between subchallenges 1 and 2, and subchallenges 2 and 3.
187 Figure 3C shows a plot of the Jaccard similarity of the sets of top-k most frequently selected genes
188 for pairs of subchallenges. We observe that a high proportion of genes are consistently selected
189 across subchallenges. The lists of most frequently selected 60, 40 and 20 genes in subchallenges 1,
190 2 and 3 respectively are available in the supplementary material (Table S3).

191 We conclude that the gene selection is not only consistent by team across folds, but also across
192 teams and subchallenges. This finding outlines a direction for further analysis, namely the validation
193 of the predictive performance and analysis of the common properties of the most frequently selected
194 genes.

195 **2.4.1 Validation of frequently selected genes**

196 We defined a simple procedure to obtain a WOC gene selection for each of the subchallenges. It
197 consisted on selecting the most frequently selected genes for each subchallenge (different colored
198 bars in Figure 3A). For example, for subchallenge 1 we chose the 60 most frequently selected genes
199 looking only at the heights of blue portion of the bar. Interestingly, the 20 most frequently selected
200 genes in subchallenge 3 are included in the list of 40 most selected genes in subchallenge 2 (except
201 for *Doc2*), conversely included in the list of 60 most selected genes in subchallenge 1.




Figure 3: Analysis of gene selection. The results in all figures were generated from the genes that were selected by the top performing teams in the post-challenge cross-validation scenario. **A.** Frequency of selected genes in subchallenge 1 (blue), subchallenge 2 (green) and subchallenge 3 (red). The genes are ordered according to their cumulative frequency. **B.** Venn diagrams of the most frequently selected genes in the subchallenges with cutoff at 20, 40 and 60 most frequently selected genes, corresponding to the number of genes required for each subchallenge. **C. Left,** the similarity of most frequently selected genes for pairs of subchallenges. The Jaccard similarity measures the ratio of the size of the intersection and the union of two sets $J(A,B) = \frac{|A \cap B|}{|A \cup B|}$. **Right,** table of correlations between gene rankings (by frequency) for pairs of subchallenges. **D.** Validation of the performances of the wisdom of the crowds (WOC) selection of genes, i.e the most frequently selected 60, 40 and 20 genes in the respective subchallenges. The violin plots represent null distribution of scores obtained by 100 randomly selected sets of 60, 40 and 20 genes using DistMap. The red dots represents the performance obtained by using DistMap with the WOC selection of genes.

202 To validate the predictive performance of the WOC gene selection, we predicted the cell
 203 locations using DistMap and scored the predictions using the same scoring metrics as for the
 204 challenge, estimating the significance of the scores through generated null distributions of scores
 205 for each subchallenge. The null distribution of the scores was generated by scoring the DistMap
 206 location prediction using 100 different sets of randomly selected genes. For each subchallenge and
 207 each score we estimated the empirical distribution function and then calculated the percentile of the
 208 values of the scores obtained with the WOC gene selection.

209 The null distributions and the values of the scores obtained with the WOC gene selection
 210 are shown in Figure 3D. All values of the scores for subchallenge 1 fall in the 99th percentile.
 211 For subchallenge 2 s_1 and s_3 fall into the 92nd percentile and s_2 in the 100th percentile. For
 212 subchallenge 3 all scores fall in the 100th percentile. Overall the performance of DistMap with the
 213 WOC selected genes performs significantly better than a random selection of genes. The actual
 214 values of the scores are on par with those achieved by the top 10 teams in the challenge.

215 2.4.2 Properties of frequently selected genes

216 We conjectured that the most frequently selected genes should carry enough information content
 217 collectively to uniquely encode a cell's location. Furthermore, genes should also contain location
 218 specific information, i.e. their expression should cluster well in space. To quantify these features,
 219 we calculated the entropy and the join count statistic for spatial autocorrelation of the *in situ* (see
 220 Figure 4A and Methods for description). We observed that most of the *in situ* genes have relatively
 221 high entropy as observed by the high density in the upper part of the plots and show high spatial
 222 clustering, i.e. show values of the join count test statistic lower than zero.

223 To test our conjectures of high entropy and spatial correlation we tested the significance of the
 224 shift of the values between the WOC selected genes and the non-selected genes from all *in situ* for
 225 each subchallenge. Since the Shapiro-Wilk test of normality rejected the null-hypothesis for both
 226 entropy and join count metrics ($p < 2.3 \cdot 10^{-6}$ and $p < 1.8 \cdot 10^{-15}$) that their values are distributed
 227 normally for the *in situ* genes, we opted for a nonparametric, one sided Mann-Whitney U test. We
 228 observed significant value shift for the autocorrelation statistic for all subchallenges 1 to 3 (see
 229 bottom of Figure 4A right red part of violin plots and table). Although we see a decrease of the




Figure 4: **Properties of selected genes.** **A.** Double violin plots of the distribution of entropy and spatial autocorrelation statistic of *Left, green* all *in situ* calculated on all embryonic location bins and *Right, red* the most frequently selected 60, 40 and 20 genes in the respective subchallenges. [bottom table] p-values of a one sided Mann-Whitney U test of location shift comparing the selected (red part of the violin plot) genes vs the non-selected genes. **B.** *Top left*, visualization of the transcriptomics data containing only the most frequently selected 60 genes from subchallenge 1 by the top performing teams (embedding to 2D by t-SNE). Each point (cell) is filled with the color of the cluster that it belongs to (density-based clustering with DBSCAN). *Top right*, spatial mapping of the cells in the *Drosophila* embryo as assigned by DistMap using only the 60 most frequently selected genes from subchallenge 1. The color of each point corresponds to the color of the cluster from the t-SNE visualization. *Bottom*, highlighted (red) location mapping of cells in the *Drosophila* embryo for each cluster separately.

Table 2: Correlations of transcriptomics to *in situ* properties of the genes where both measurements are available. σ^2 - variance of a gene across cells, c_v - coefficient of variation, 0 - number of cells with zero expression, H_b - entropy of binarized expression, H - entropy, Z - join count test statistic

		<i>in situ</i>	
		H	Z
ρ			
scRNAseq	σ^2	0.50	0.18
	c_v	-0.69	0.26
	0	-0.64	0.29
	H_b	0.72	-0.30

230 statistical significance of the mean value shift for the distribution of values of the entropy of the
 231 selected subsets of genes, the shift is significant for all subchallenges and at the same time, we
 232 observe that tail of the distribution shortens.

233 To test whether the information relative to different cell types is retained with the selected
 234 subset of 60, 40 or 20 WOC selected genes, we embedded the cells into 2D space using t-distributed
 235 stochastic embedding (t-SNE) [20] aiming for high accuracy ($\theta = 0.01$), Figure 4B and Figure S5.
 236 We then clustered the t-SNE embedded data using density-based spatial clustering of applications
 237 with noise (DBSCAN) [21]. DBSCAN determines the number of clusters in the data automatically
 238 based on the density of points in space. The minimum number of cells in a local neighborhood was
 239 set to 10 and the parameter $\epsilon = 3.5$ was selected by determining the elbow point in a plot of sorted
 240 distances of each cell to its 10th nearest neighbor. We found that the 9 prominent cell clusters
 241 identified in the study by Karaikos *et al.* [10] are preserved in our t-SNE embedding and clustering
 242 experiments when considering the most frequently selected 60 or 40 genes from subchallenges 1
 243 and 2. The number of clusters of cells with specific localization is reduced when considering the
 244 most frequently selected 20 genes from subchallenge 3.

245 We next associated the properties of the *in situs* that were found to be indicative of good perfor-
 246 mance in the task of location prediction with statistical properties of the genes in the transcriptomics
 247 data. Our goal was to discover statistical properties of the transcriptomics data that might inform
 248 future experimental designs when selecting target genes for *in situ* hybridizations. We calculated
 249 statistical features across cells for the subset of genes from the transcriptomics data for which we
 250 also have *in situ* measurements. These include the variance of gene expression σ^2 across cells,
 251 the coefficient of variation $c_v = \frac{\sigma}{\mu}$, the number of cells with expression zero 0 and the entropy of
 252 binarized expression H_b . We then calculated the correlation across genes for each of these metrics
 253 and the measured spatial properties of interest of the *in situs*, i.e entropy H and the value of the
 254 joint count statistic Z (see Table 2). Although the selection of highly variable genes was one of
 255 the approaches used by some of the top 10 teams, the variance for each gene in the scRNAseq
 256 expression, although highly correlated to the entropy of the corresponding *in situ* measurements of
 257 that gene, it is less correlated than other properties. Also, we observed that the positive correlation
 258 of the entropy to the variance of each gene, becomes a negative correlation against their coefficient
 259 of variation. This negative correlation can have two sources, the genes with high entropy may have
 260 low standard deviation or high mean expression. Since we observe positive correlation of entropy
 261 to the variance of expression, we can conclude that the negative correlation is a result of highly
 262 expressed genes. Since a known drawback of scRNAseq is a high number of dropout events for
 263 lowly expressed genes [22], this observation is further supported by the negative correlation of the
 264 entropy and the number of cells with zero expression. We observed the highest correlation of *in*
 265 *situ* entropy to the entropy of the binarized expression. Regarding the spatial autocorrelation, all
 266 statistical features of the transcriptomics were only slightly positively correlated to the join count
 267 statistic except for the entropy of binarized expression which had slightly negative correlation.

268 3 Discussion

269 In this paper we report the results of a crowdsourcing effort organized as a DREAM challenge,
 270 around the issue of predicting the spatial arrangement of cells in a tissue from scRNAseq data.
 271 Analysis of the top performing methods and their performance provided us a number of unbiased
 272 insights. First, it unveiled a connection in the cell-to-cell variability in *Drosophila* embryo gene
 273 expression and the selection of the best genes for predicting the localization of a cell in the embryo
 274 from their scRNAseq expression. The most selected genes had a relatively high entropy, hence high
 275 variance and expression while also showing high spatial clustering. The smaller the number of
 276 selected genes, i.e going from subchallenge 1 to 3, the more these features became apparent. The
 277 observed advantage of genes with high overall expression in cells might lead to less dropout counts
 278 in the scRNAseq data, a known disadvantage of the technology, leading to more accuracy in the cell
 279 placement. We also found that the 9 prominent spatially distinct cell clusters previously identified
 280 [10] are preserved when considering the most frequently selected 60 or 40 genes, but the number
 281 of clusters is reduced when considering only the most frequently selected 20 genes. This finding
 282 is in line with the conclusions of Howe et al. [11] where in a related task of location prediction
 283 the performance stabilized after the inclusion of 30 genes in a related experiment. The WOC gene
 284 selection and the k-means clustered WOC model for cell localization performed comparably or
 285 better than the participant's models, showing once more the advantage of the wisdom-of-the-crowds.

286 Given that it has been shown that positional information of the anterior-posterior (A-P) axis is
 287 encoded as early in the embryonic development as when the expression of the gap genes occurs
 288 [23, 24], we thought that it should be possible to implement in algorithms for this challenge the
 289 information contained in the regulatory networks of *Drosophila* development [25]. Although only a
 290 small number of participants, among them the best performers, directly used biological information
 291 related to the regulation of the genes or their connectivity, the most frequently selected genes in
 292 all 3 subchallenges have interesting biological properties. Indeed, gap genes such as *giant* (*gt*),
 293 *kruppel* (*kr*), *knirps* (*kni*) were selected in all 3 subchallenges (see Figure 5 and Table S3 that also
 294 includes *kni*-like *knrl*) although *tailless* (*tll*) and *hunchback* (*hb*) were not. Along the A-P axis,
 295 maternally provided *bicoid* (*bcd*) and *caudal* (*cad*) first establish the expression patterns of gap and




Figure 5: Gene regulatory network of early *Drosophila* development. Not all regulations are represented, nor pair ruled genes *odd* & *prd*. Frequently selected genes are represented in bold.

296 terminal class factors, such as *hb*, *gt*, *kr* and *kni*. These A-P early regulators then collectively direct
297 transcription of A-P pair-rule factors, such as *even-skipped* (*eve*), *fushi-tarazu* (*ftz*), *hairy* (*h*), *odd*
298 *skipped*, (*odd*), *paired* (*prd*) and *runt* (*run*) which in turn cross-regulate each other. Not being part
299 of the *in situs*, neither *bcd*, nor *cad* were selected but *ama* sitting near *bcd* in the genome might
300 have been selected for its similar expression properties. Furthermore, we also find that pair-rule
301 genes were most prominently selected in subchallenges 1 (*eve*, *odd*, *prd*, the Paired-like *bcd* and
302 *bcd*) and 2 (*h*, *ftz* and *run*). A similar cascade of maternal and zygotic factors controls patterning
303 along the dorsal-ventral (D-V) axis were *dorsal* (*d*), *snail* (*sna*) and *twist* (*twi*) specify mesoderm
304 and the pair rule factors *eve* and *ftz* specify location along the trunk of the A-P axis. Again, *sna* and
305 *twi* were selected in all subchallenges and *d* in subchallenges 1 and 2. These selected transcription
306 factors specify distinct developmental fates and can act via different cis-regulatory modules but
307 their quantitative differences in relative levels of binding to shared targets correlates with their
308 known biological and transcriptional regulatory specificities [26]. The rest of the selected genes
309 were the homeobox genes (*nub*, *antp*) and differentiators of tissue such as mesoderm (*ama*, *mes2*,
310 *zfh1*), ectoderm (*doc2* and *doc3*), neural tissue (*noc*, *oc*, *rho*) and EGFR pathway (*rho*, *edl*). The
311 complete lists of most frequently selected genes are available in Table S3.

312 Since the ground truth of single cell locations was publicly available, the organization of this
313 DREAM challenge brought risks that, given the importance of the scientific question asked, we
314 thought worth taking. However, without the post-challenge phase it would have been impossible to
315 distinguish the robust and sound methods from methods that were overfitting the results. Overall,
316 the single cell transcriptomics challenges unveils not only the best gene-selection methods and
317 prediction approaches to localize a cell in the *Drosophila* embryo, but also explains the biological
318 and statistical properties of the genes selected for the predictions. Further identification of additional
319 properties such as spatially autocorrelated genes might require the use of alternative scRNAseq
320 focused approaches [13, 14]. However, we think that the approach defined here could be used or
321 adapted when performing similar cell-placing tasks in other organisms, including human tissues.
322 Given the importance of spatial arrangements for disease development and treatment, we foresee
323 an application of these methods to medical questions as well.

324 References

- 325 [1] Anna K Casasent, Aislyn Schalck, Ruli Gao, Emi Sei, Annalyssa Long, William Pangburn,
326 Tod Casasent, Funda Meric-Bernstam, Mary E Edgerton, and Nicholas E Navin. Multiclonal
327 invasion in breast tumors identified by topographic single cell sequencing. *Cell*, 172(1-2):205–
328 217, 2018.
- 329 [2] Patrik L Ståhl, Fredrik Salmén, Sanja Vickovic, Anna Lundmark, José Fernández Navarro,
330 Jens Magnusson, Stefania Giacomello, Michaela Asp, Jakub O Westholm, Mikael Huss, et al.
331 Visualization and analysis of gene expression in tissue sections by spatial transcriptomics.
332 *Science*, 353(6294):78–82, 2016.
- 333 [3] Ditte Lovatt, Brittani K Ruble, Jaehee Lee, Hannah Dueck, Tae Kyung Kim, Stephen Fisher,
334 Chantal Francis, Jennifer M Spaethling, John A Wolf, M Sean Grady, et al. Transcriptome in
335 vivo analysis (tiva) of spatially defined single cells in live tissue. *Nature methods*, 11(2):190,
336 2014.
- 337 [4] Samuel G Rodriques, Robert R Stickels, Aleksandrina Goeva, Carly A Martin, Evan Murray,
338 Charles R Vanderburg, Joshua Welch, Linlin M Chen, Fei Chen, and Evan Z Macosko. Slide-
339 seq: A scalable technology for measuring genome-wide expression at high spatial resolution.
340 *Science*, 363(6434):1463–1467, 2019.

- 341 [5] Yang Liu, Mingyu Yang, Yanxiang Deng, Graham Su, Cindy C Guo, Di Zhang, Dongjoo Kim,
342 Zhiliang Bai, Yang Xiao, and Rong Fan. High-spatial-resolution multi-omics atlas sequencing
343 of mouse embryos via deterministic barcoding in tissue. *bioRxiv*, page 788992, 2019.
- 344 [6] Kaia Achim, Jean-Baptiste Pettit, Luis R Saraiva, Daria Gavriouchkina, Tomas Larsson,
345 Detlev Arendt, and John C Marioni. High-throughput spatial mapping of single-cell rna-seq
346 data to tissue of origin. *Nature biotechnology*, 33(5):503, 2015.
- 347 [7] Rahul Satija, Jeffrey A Farrell, David Gennert, Alexander F Schier, and Aviv Regev. Spatial
348 reconstruction of single-cell gene expression data. *Nature biotechnology*, 33(5):495, 2015.
- 349 [8] Keren Bahar Halpern, Rom Shenhav, Orit Matcovitch-Natan, Beáta Tóth, Doron Lemze,
350 Matan Golan, Efi E Massasa, Shaked Baydatch, Shanie Landen, Andreas E Moor, et al.
351 Single-cell spatial reconstruction reveals global division of labour in the mammalian liver.
352 *Nature*, 542(7641):352, 2017.
- 353 [9] Tim Stuart, Andrew Butler, Paul Hoffman, Christoph Hafemeister, Efthymia Papalexi,
354 William M Mauck III, Yuhan Hao, Marlon Stoeckius, Peter Smibert, and Rahul Satija.
355 Comprehensive integration of single-cell data. *Cell*, 2019.
- 356 [10] Nikos Karaiskos, Philipp Wahle, Jonathan Alles, Anastasiya Boltengagen, Salah Ayoub,
357 Claudia Kipar, Christine Kocks, Nikolaus Rajewsky, and Robert P Zinzen. The drosophila
358 embryo at single-cell transcriptome resolution. *Science*, 358(6360):194–199, 2017.
- 359 [11] Douglas G Howe, Yvonne M Bradford, Tom Conlin, Anne E Eagle, David Fashena, Ken
360 Frazer, Jonathan Knight, Prita Mani, Ryan Martin, Sierra A Taylor Moxon, et al. Zfin, the
361 zebrafish model organism database: increased support for mutants and transgenics. *Nucleic
362 acids research*, 41(D1):D854–D860, 2012.
- 363 [12] Charless C Fowlkes, Cris L Luengo Hendriks, Soile VE Keränen, Gunther H Weber, Oliver
364 Rübél, Min-Yu Huang, Sohail Chatoor, Angela H DePace, Lisa Simirenko, Clara Henriquez,
365 et al. A quantitative spatiotemporal atlas of gene expression in the drosophila blastoderm.
366 *Cell*, 133(2):364–374, 2008.
- 367 [13] Josephine Bageritz, Philipp Willnow, Erica Valentini, Svenja Leible, Michael Boutros, and
368 Aurelio A Teleman. Gene expression atlas of a developing tissue by single cell expression
369 correlation analysis. *Nature Methods*, 16(8):750, 2019.
- 370 [14] Mor Nitzan, Nikos Karaiskos, Nir Friedman, and Nikolaus Rajewsky. Charting a tissue from
371 single-cell transcriptomes. *bioRxiv*, page 456350, 2018.
- 372 [15] Julio Saez-Rodriguez, James C Costello, Stephen H Friend, Michael R Kellen, Lara Man-
373 gravite, Pablo Meyer, Thea Norman, and Gustavo Stolovitzky. Crowdsourcing biomedical
374 research: leveraging communities as innovation engines. *Nature Reviews Genetics*, 17(8):470–
375 486, 2016.
- 376 [16] Daniel Marbach, James C Costello, Robert Küffner, Nicole M Vega, Robert J Prill, Diogo M
377 Camacho, Kyle R Allison, Andrej Aderhold, Richard Bonneau, Yukun Chen, et al. Wisdom
378 of crowds for robust gene network inference. *Nature methods*, 9(8):796, 2012.
- 379 [17] Michael P Menden, Dennis Wang, Mike J Mason, Bence Szalai, Krishna C Bulusu, Yuanfang
380 Guan, Thomas Yu, Jaewoo Kang, Minji Jeon, Russ Wolfinger, et al. Community assessment
381 to advance computational prediction of cancer drug combinations in a pharmacogenomic
382 screen. *Nature communications*, 10(1):2674, 2019.

- 383 [18] John A Hartigan and Manchek A Wong. Algorithm as 136: A k-means clustering algorithm.
384 *Journal of the Royal Statistical Society. Series C (Applied Statistics)*, 28(1):100–108, 1979.
- 385 [19] Trupti M Kodinariya and Prashant R Makwana. Review on determining number of cluster in
386 k-means clustering. *International Journal*, 1(6):90–95, 2013.
- 387 [20] Laurens van der Maaten and Geoffrey Hinton. Visualizing data using t-SNE. *Journal of*
388 *Machine Learning Research*, 9:2579–2605, 2008.
- 389 [21] Martin Ester, Hans-Peter Kriegel, Jörg Sander, and Xiaowei Xu. A density-based algorithm
390 for discovering clusters in large spatial databases with noise. In *Proceedings of the Second*
391 *International Conference on Knowledge Discovery and Data Mining*, KDD’96, pages 226–
392 231, 1996.
- 393 [22] Peter V Kharchenko, Lev Silberstein, and David T Scadden. Bayesian approach to single-cell
394 differential expression analysis. *Nature methods*, 11(7):740, 2014.
- 395 [23] Julien O Dubuis, Gašper Tkačik, Eric F Wieschaus, Thomas Gregor, and William Bialek. Posi-
396 tional information, in bits. *Proceedings of the National Academy of Sciences*, 110(41):16301–
397 16308, 2013.
- 398 [24] Mariela D Petkova, Gašper Tkačik, William Bialek, Eric F Wieschaus, and Thomas Gregor.
399 Optimal decoding of cellular identities in a genetic network. *Cell*, 176(4):844–855, 2019.
- 400 [25] David Umulis, Michael B O’Connor, and Hans G Othmer. Robustness of embryonic spatial
401 patterning in drosophila melanogaster. *Current topics in developmental biology*, 81:65–111,
402 2008.
- 403 [26] Stewart MacArthur, Xiao-Yong Li, Jingyi Li, James B Brown, Hou Cheng Chu, Lucy Zeng,
404 Brandi P Grondona, Aaron Hechmer, Lisa Simirenko, Soile VE Keränen, et al. Developmental
405 roles of 21 drosophila transcription factors are determined by quantitative differences in
406 binding to an overlapping set of thousands of genomic regions. *Genome biology*, 10(7):R80,
407 2009.
- 408 [27] Andrew D. Cliff and John K. Ord. *Spatial autocorrelation*. Pion London, 1973.
- 409 [28] Robert R. Solak and Neal L. Oden. Spatial autocorrelation in biology: 1. methodology.
410 *Biological Journal of the Linnean Society*, 10(2):199–228, 1978.

411 **4 Methods**

412 **4.1 Scoring**

413 We scored the submissions for the three subchallenges using three metrics s_1 , s_2 and s_3 . s_1 measured
414 how well the expression of the cell at the predicted location correlates to the expression from the
415 reference atlas and included the variance of the predicted locations for each cell. While s_2 measured
416 the accuracy of the predicted location and s_3 measured how well the gene-wise spatial patterns
417 were reconstructed.

418 Let c represent the index of a cell, given in the transcriptomics data in the challenge where
419 $1 \leq c \leq 1297$. Each cell c is located in a bin $\epsilon_c \in \{1..3039\}$ at a position with coordinates
420 $r(\epsilon_c) = (x_c, y_c, z_c)$. Each cell is associated with a binarized expression profile $t_c = (t_{c1}, t_{c2}, \dots, t_{cE})$,
421 where $1 \leq E \leq 8924$, and a corresponding binarized *in situ* profile $f_c = (f_{c1}, f_{c2}, \dots, f_{cK})$, where
422 the maximum possible value of K for which we have *in situ* information is $K = 84$. For different

423 subchallenges we consider $K \in \{20, 40, 60\}$. Using K selected genes the participants were asked to
 424 provide an ordered list of 10 most probable locations for each cell. We represent with the mapping
 425 function $A(c, i, K)$ the value of the predicted i -th most probable location for cell c using K *in situ*.

426 For the first scoring metric s_1 we calculated the weighted average of the Mathews correlation
 427 coefficient (MCC) between the *in situ* profile of the ground truth cell location f_{ϵ_c} and the *in situ*
 428 profile of the most probable predicted location $f_{A(c,1,K)}$ for that cell

$$s_1 = \sum_{c=1}^N \frac{p_K(c, A)}{\sum_{i=1}^N p_K(i, A)} MCC(f_{A(c,1,K)}, f_{\epsilon_c}),$$

429 where N is the total number of cells with predicted locations.

430 The Mathews correlation coefficient, or ϕ coefficient, is calculated from the contingency table
 431 obtained by correlating two binary vectors. The MCC is weighted by the inverse of the distance of
 432 the predicted most probable locations to the ground truth location $p_K(c)$. The weights are calculated
 433 as $p_K(c, A) = \frac{\widetilde{d_{84}(c, A)}}{\widetilde{d_K(c, A)}}$, where $d_K(c, A) = \frac{1}{10} \sum_{i=1}^{10} \|r(A(c, i, K)) - r(\epsilon_c)\|_2$, $\widetilde{d_{84}(c, A)}$ is the value of
 434 $d_K(c, A)$ using the ground truth most probable locations assigned with $K = 84$ using DistMap, and
 435 $\|\cdot\|_2$ is the Euclidean norm.

The second metric s_2 is simply the average inverse distance of the predicted most probable
 locations to the ground truth location

$$s_2 = \frac{1}{N} \sum_{c=1}^N p_K(c, A).$$

436 Finally, the third metric s_3 measures the accuracy of reconstructed gene-wise spatial patterns

$$s_3 = \sum_{s=1}^K \frac{MCC(t_{cs}, f_{\epsilon_c s})_{\forall c}}{\sum_{i=1}^K MCC(t_{ci}, f_{\epsilon_i r})_{\forall c}} MCC(t_{cs}, f_{A(c,1,K)s})_{\forall c},$$

437 where $\forall c$ denotes that the MCC is calculated cell wise for each gene.

438 For 287 out of the 1297 cells, the ground truth location predictions were ambiguous, i.e., the
 439 MCC scores were identical for multiple locations. These cells were removed both from the ground
 440 truth and the submissions before calculating the scores.

441 The teams were ranked according to each score independently. The final assigned rank r_t
 442 for team t was calculated as the average rank across scores. Teams were ranked based on the
 443 performance as measured by the three scores on 1000 bootstrap replicates of the submitted solutions.
 444 The three scores were calculated for each bootstrap. The teams were then ranked according to
 445 each score. These ranks were then averaged to obtain a final rank for each team on that bootstrap.
 446 The winner for each subchallenge was the team that achieved the lowest ranks. We calculated the
 447 Bayes factor of the bootstrap ranks for the top performing teams. Bayesian factor of 3 or more was
 448 considered as a significantly better performance. The Bayes factor of the 1000 bootstrapped ranks
 449 of teams T_1 and T_2 was calculated as

$$BF(T_1, T_2) = \frac{\sum_{i=1}^{1000} \mathbf{1}(r(T_1)_i < r(T_2)_i)}{\sum_{i=1}^{1000} \mathbf{1}(r(T_1)_i > r(T_2)_i)},$$

450 where $r(T_1)_i$ is the rank of team T_1 on the i -th bootstrap, $r(T_2)_i$ is the rank of team T_2 on the i -th
 451 bootstrap, and $\mathbf{1}$ is the indicator function.

452 4.2 Entropy and spatial autocorrelation

The entropy of a binarized *in situ* measurements of gene G was calculated as

$$H(G) = -p \log_2 p - (1 - p) \log_2 (1 - p),$$

453 where p is the probability of gene G to have value 1. In other words, p is the fraction of cells where
 454 G is expressed.

455 The join count statistic is a measure of a spatial autocorrelation of a binary variable. We will
 456 refer to the binary expression 1 and 0 as black (B) and white (W). Let n_B be the number of bins
 457 where G is expressed ($G = B$), and $n_W = n - n_B$ the number of bins where G is not expressed
 458 ($G = W$). Two neighboring spatial bins can form join of type $J \in \{WW, BB, BW\}$.

459 We are interested in the distribution of BW joins. If a gene has a lower number of BW joins
 460 than the expected number of BW, then the gene is positively spatially autocorrelated, i.e., the gene is
 461 highly clustered. Contrarily, higher number of BW joins points towards negative spatial correlation,
 462 i.e. dispersion.

Following Cliff and Ord [27] and Sokal and Oden [28], the expected count of BW joins is

$$\mathbb{E}[BW] = \frac{1}{2} \sum_i \sum_j \frac{w_{ij} n_B^2}{n^2},$$

where the spatial connectivity matrix w is defined as

$$w_{ij} = \begin{cases} 1 & \text{if } i \neq j \text{ and } j \text{ is in the list of 10 nearest neighbors of } i \\ 0 & \text{otherwise} \end{cases}$$

The variance of BW joins is

$$\sigma_{BW}^2 = \mathbb{E}[BW^2] - \mathbb{E}[BW]^2.$$

where the term $\mathbb{E}[BW^2]$ is calculated as

$$\mathbb{E}[BW^2] = \frac{1}{4} \left(\frac{2x_2 n_B n_W}{n^2} + \frac{(x_3 - 2x_2) n_B n_W (n_B + n_W - 2)}{n^3} + \frac{4(x_1^2 + x_2 - x_3) n_B^2 n_W^2}{n^4} \right),$$

463 where $x_1 = \sum_i \sum_j w_{ij}$, $x_2 = \frac{1}{2} \sum_i \sum_j (w_{ij} - w_{ji})^2$, $x_3 = \sum_i (\sum_j w_{ij} + \sum_j w_{ji})^2$.

464 Note that the connectivity matrix w can also be asymmetric, since it is defined by the nearest
 465 neighbor function.

Finally, the observed BW counts are

$$BW = \frac{1}{2} \sum_i \sum_j w_{ij} (G_i - G_j)^2.$$

The join counts test statistic is then defined as

$$Z(BW) = \frac{BW - \mathbb{E}[BW]}{\sqrt{\sigma_{BW}^2}},$$

466 which is assumed to be asymptotically normally distributed under the null hypothesis of no spatial
 467 autocorrelation. Negative values of the Z statistic represent positive spatial autocorrelation, or
 468 clustering, of gene G . Positive values of the Z statistic represent negative spatial autocorrelation, or
 469 dispersion, of gene G .

470 4.3 Implementation details

471 The challenge scoring was implemented and run in R version 3.5, the post analysis was performed
 472 with R version 3.6 and the core tidyverse packages. We used the publicly available implemen-
 473 tation of DistMap (<https://github.com/rajewsky-lab/distmap>). MCC calculated
 474 with R package mcor (0.4.4). t-SNE embedding and visualization produced with R package
 475 Rtsne (0.15). DBSCAN clustering with R package dbscan (1.1-4).

476 **4.4 Code availability**

477 <https://github.com/dream-sctc/Scoring>

478 **4.5 Data description**

479 **Reference Database** The reference database comes from the Berkeley *Drosophila* Transcription
480 Network Project. The *in situ* expression of 84 genes (columns) is quantified across the 3039
481 *Drosophila* embryonic locations (rows) for raw data and for binarized data. The 84 genes were
482 binarized by manually choosing thresholds for each gene.

483 **Spatial coordinates** One half of *Drosophila* embryo has 3039 cells places as x, y and z (columns)
484 for a total of 3039 embryo locations (rows) and a total of 3039.3 coordinates.

485 **Single cell RNA sequencing** The single-cell RNA sequencing data is provided as a matrix with
486 8924 genes as rows and 1297 cells as columns. In the raw version of the matrix, the entries are the
487 raw unique gene counts (quantified by using unique molecular identifiers – UMI). The normalized
488 version is obtained by dividing each entry by the total number of UMIs for that cell, adding a
489 pseudocount and taking the logarithm of that. All entries are finally multiplied by a constant. For a
490 given gene and only considering the Drop-seq cells expressing it we computed a quantile value
491 above (below) which the gene would be designated ON (OFF). We sampled a series of quantile
492 values and each time the gene correlation matrix based on this binarized version of normalized data
493 versus the binarized BDTNP atlas was computed and compared by calculating the mean square
494 root error between the elements of the lower triangular matrices. Eventually, the quantile value
495 0.23 was selected, as it was found to minimize the distance between the two correlation matrices.

496 The short sequences for each of the 1297 cells in the raw and normalized data are the cell
497 barcodes.

498 **5 Acknowledgments**

499 This research was funded in part by PROACTIVE 2017 “From Single-Cell to Multi-Cells Informa-
500 tion Systems Analysis” (C92F17003530005 Department of Information Engineering, University of
501 Padova) for B.D.C.; National Institutes of Health grant number U54CA21729 for J.R.; ICMR JRF
502 (Indian Council of Medical Research - Junior Research Fellowship) for S.A. and X.W. was funded
503 by the National Natural Science Foundation of China (No.61702421 and No.61772426).

504 **6 Author contributions**

505 Conceptualization, N.K., N.R., J.S.R., G.S., and P.M.; Methodology, J.S., M.E.A., G.S., and P.M.;
506 Software, J.T., and M.E.A.; Formal Analysis, J.T., M.E.A., G.S. and P.M.; Writing - Original Draft,
507 J.T. and P.M.; Writing - Supervision, J.S.R., G.S., and P.M.

508 **7 Competing interests**


509 The authors declare no competing interests.

510 **8 Materials and Correspondence**

511 Requests for data, resources, and or reagents should be directed to Pablo Meyer (pmeyerr@us.ibm.com).

A

bioRxiv preprint doi: <https://doi.org/10.1101/796029>; this version posted October 10, 2019. The copyright holder has placed this preprint (which was not certified by peer review) in the Public Domain. It is no longer restricted by copyright. Anyone can legally share, reuse, remix, or adapt this material for any purpose without crediting the original authors.





in-situ hybridization of 84 driver genes
BDTNP

transcriptomes of 1297 single cells
Drop-seq, 8924 genes

Select
 ○ 60 genes
 ⊗ 40 genes
 ○ 20 genes


predict location




3039 location bins

B


Subchallenge 1



Subchallenge 2




Subchallenge 3



Single cell transcriptome


T

Team 1




+


Team M



Aggregated predictions











K-means clustering



- gold
- K_1
- K_2
- K_3
- K_4
- K_5
- K_6
- K_7
- K_8
- not_assigned

1. Pick the cluster with most frequently assigned locations by teams
2. Pick the 10 most frequently selected locations in that cluster



A**B**

p-value	Subchallenge 1	Subchallenge 2	Subchallenge 3
Entropy	0.0005	0.0212	0.1554
Autocorrelation	0.0044	0.0042	0.0216

maternal genes

cad | *bcd* | *nos*

gap genes

gt | *tll* → ***hb*** | *kr* | ***kni***

pair-rule genes

ftz ← ***eve*** | *run* | ***h***

segment polarity
genes

en → *hh* → *wg*

